ON PREHOMOGENEITY OF A RANK VARIETY

MASAYA OUCHI, MICHIO HAMADA, AND TATSUO KIMURA

(Communicated by Lev Borisov)

ABSTRACT. If a linear algebraic group G acts on $M(m, n)$, then it also acts on a rank variety $M^{(r)}(m, n) = \{X \in M(m, n) \mid \text{rank } X = r\}$. In this paper, we give the necessary and sufficient condition that this variety has a Zariski-dense G -orbit. We consider everything over the complex number field $\mathbb{C}$.

INTRODUCTION

Let G be a linear algebraic group. Let $\rho : G \rightarrow GL(m)$ and $\sigma : G \rightarrow GL(n)$ be its rational representations over $\mathbb{C}$. Then G acts on $M(m, n)$ by $\rho \otimes \sigma$, i.e., $X \mapsto \rho(g)X^t\sigma(g)$ ($X \in M(m, n)$, $g \in G$). By this action, G also acts on a rank variety $M^{(r)}(m, n) = \{X \in M(m, n) \mid \text{rank } X = r\}$. On the other hand, if a rational representation $\tau : H \rightarrow GL(V)$ of a linear algebraic group H on a finite-dimensional vector space V has a Zariski-dense H -orbit, we call a triplet (H, τ, V) a prehomogeneous vector space (abbrev. PV). A point of the Zariski-dense H -orbit is called a generic point, and the isotropy subgroup at a generic point is called a generic isotropy subgroup. For the basic facts of PVs, see [K]. In this paper, we shall prove the following theorem.

Theorem 0.1. *The following assertions are equivalent.*

- (1) $M^{(r)}(m, n)$ has a Zariski-dense G -orbit by the action $\rho \otimes \sigma$.
- (2) $(G \times GL(r), \rho \otimes \Lambda_1 + \sigma \otimes \Lambda_1^*, M(m, r) \oplus M(n, r))$ is a PV.

Here the action of (2) is given by $(X, Y) \mapsto (\rho(g)X^tA, \sigma(g)YA^{-1})$ for $(X, Y) \in M(m, r) \oplus M(n, r)$ and $(g, A) \in G \times GL(r)$. Note that we have $m \geq r$ and $n \geq r$ in (2).

1. PROOF OF THEOREM

The following lemma is the key for our proof.

Lemma 1.1 (M. Sato). *Assume that an algebraic group G acts on both of two irreducible algebraic varieties W and W' . Let $\varphi : W \rightarrow W'$ be a morphism satisfying*

- (I) $\varphi(gw) = g\varphi(w)$ ($g \in G, w \in W$),
- (II) $\varphi(W) = W'$.

Received by the editors May 26, 2011.

2010 *Mathematics Subject Classification.* Primary 11S90; Secondary 15A03.

Key words and phrases. Rank, Zariski-dense orbit, prehomogeneous vector space.

©2012 American Mathematical Society
 Reverts to public domain 28 years from publication

Then the following assertions are equivalent:

- (1) $W = \overline{G \cdot w}$ for some $w \in W$; that is, W is G -prehomogeneous.
- (2) (a) $W' = \overline{G \cdot w'}$ for some $w' \in W'$.
 (b) For the above point $w' \in W'$ in (a), there exists a point $w \in \varphi^{-1}(w')$ such that $\varphi^{-1}(w') = \overline{G_{w'} \cdot w}$, where $G_{w'} = \{g \in G : gw' = w'\}$ is the isotropy subgroup of G at w' .

Proof. For the proof, see Lemmas 7.2 and 7.6 in [K]. □

Now we shall prove Theorem 0.1. Put $W = \{(X, Y) \in M(m, r) \oplus M(n, r) \mid \text{rank } X = \text{rank } Y = r\}$. Then clearly $(G \times GL(r), \rho \otimes \Lambda_1 + \sigma \otimes \Lambda_1^*, M(m, r) \oplus M(n, r))$ is a PV if and only if W is $G \times GL(r)$ -prehomogeneous. Note that $\text{rank } X^t Y = r$ for any $(X, Y) \in W$. Hence we have a $(G \times GL(r))$ -equivariant map $\varphi : W \rightarrow M^{(r)}(m, n)$ by $(X, Y) \mapsto X^t Y$. Note that $GL(m) \times GL(n)$ also acts on both W and $M^{(r)}(m, n)$. Since φ is also $(GL(m) \times GL(n))$ -equivariant and $M^{(r)}(m, n)$ is a single $(GL(m) \times GL(n))$ -orbit, we see that φ is surjective. For $Z_0 = \varphi(X_0, Y_0) \in M^{(r)}(m, n)$, we shall show that the fiber $\varphi^{-1}(Z_0)$ is a principal $GL(r)$ -orbit. Assume that $(X, Y) \in \varphi^{-1}(Z_0)$. If $X = (x_1 | \cdots | x_r) \in M(m, r)$, let $\langle X \rangle$ be an r -dimensional subspace of $\mathbb{C}^m$ generated by column vectors $x_1, \dots, x_r$. Since we have $X^t Y = (\sum_{j=1}^r y_{1j} x_j | \cdots | \sum_{j=1}^r y_{nj} x_j)$ for $Y = (y_{ij})$ and $\text{rank } X = \text{rank } X^t Y (= r)$, we have $\langle X \rangle = \langle X^t Y \rangle = \langle Z_0 \rangle = \langle X_0^t Y_0 \rangle = \langle X_0 \rangle$, and hence there exists uniquely $A \in GL(r)$ satisfying $X = X_0 {}^t A$. Then we have $X_0 {}^t Y_0 = X^t Y = X_0 {}^t A^t Y$ and since $\text{rank } X_0 = r$, we obtain ${}^t Y_0 = {}^t A^t Y$, i.e., $Y = Y_0 A^{-1}$. This shows that the fiber $\varphi^{-1}(Z_0)$ is a principal $GL(r)$ -orbit. Since $GL(r)$ is contained in the isotropy subgroup of Z_0 , by Lemma 1.1, we obtain Theorem 0.1.

Note that (2) in Theorem 0.1 is equivalent to:

- (3) $(G \times GL(r), \rho \otimes \Lambda_1, M(m, r))$ is a PV with a generic isotropy subgroup H , and $(H, \sigma \otimes \Lambda_1^*, M(n, r))$ is a PV.

Example 1.2. Let $\rho : Spin(10) \rightarrow GL(16)$ be a half-spin representation of the spin group $Spin(10)$. Then $Spin(10) \times GL(14)$ acts on $M(16, 14)$ and $M^{(r)}(16, 14)$ has the Zariski-dense $(Spin(10) \times GL(14))$ -orbit if and only if $r = 1, 2, 3, 13, 14$. By (3), $(Spin(10) \times GL(14) \times GL(r), \text{a half-spin rep.} \otimes 1 \otimes \Lambda_1 + 1 \otimes \Lambda_1 \otimes \Lambda_1^*, M(16, r) \oplus M(14, r))$ is a PV if and only if $(Spin(10) \times GL(r), \text{a half-spin rep.} \otimes \Lambda_1, M(16, r))$ is a PV since the latter is a trivial PV (see [K]). So by [SK], we have our result. Note that the case $r = 13$ (resp. $r = 14$) is a castling transform of the case $r = 3$ (resp. $r = 2$).

Remark 1.3. Professors Shigefumi Mori and Yasuo Teranishi proved that (2) in Theorem 0.1 is equivalent to (4) : $U = \{(\langle X \rangle, \langle Y \rangle, X^t Y) \in \text{Grass}_r(V(m)) \oplus \text{Grass}_r(V(n)) \oplus M(m, n) \mid (X, Y) \in W\}$ is G -prehomogeneous, and later Professor Masanobu Taguchi proved that it is equivalent to (5) : $U' = \{(\langle X \rangle, X^t Y) \in \text{Grass}_r(V(m)) \oplus M(m, n) \mid (X, Y) \in W\}$ is G -prehomogeneous (see [KKT1, Remark 1.38]).

REFERENCES

- [K] T. Kimura, Introduction to prehomogeneous vector spaces, Translations of Mathematical Monographs, **215**, American Mathematical Society, Providence, RI, 2003. MR1944442 (2003k:11180)
- [KKTl] T. Kimura, S. Kasai, M. Taguchi and M. Inuzuka, Some P.V.-equivalence and a classification of 2-simple prehomogeneous vector spaces of type II, Trans. Amer. Math. Soc. **308** (1988), 433–494. MR951617 (89k:11101)
- [SK] M. Sato and T. Kimura, A classification of irreducible prehomogeneous vector spaces and their relative invariants, Nagoya Math. J. **65** (1977), 1–155. MR0430336 (55:3341)

INSTITUTE OF MATHEMATICS, UNIVERSITY OF TSUKUBA, IBARAKI, 305-8571, JAPAN
E-mail address: `msy2000@math.tsukuba.ac.jp`

INSTITUTE OF MATHEMATICS, UNIVERSITY OF TSUKUBA, IBARAKI, 305-8571, JAPAN
E-mail address: `mhamada@math.tsukuba.ac.jp`

INSTITUTE OF MATHEMATICS, UNIVERSITY OF TSUKUBA, IBARAKI, 305-8571, JAPAN
E-mail address: `kimurata@math.tsukuba.ac.jp`