HARDY TYPE INEQUALITIES RELATED TO CARNOT-CARATHÉODORY DISTANCE ON THE HEISENBERG GROUP

QIAO-HUA YANG

(Communicated by Mei-Chi Shaw)

ABSTRACT. Being motivated by a representation formula associated with the Korányi-Folland nonisotropic gauge proved by Cohn and Lu, we prove an analogous representation formula related to the Carnot-Carathéodory distance on the Heisenberg group. Using this formula, we obtain some Hardy inequalities associated with the Carnot-Carathéodory distance on such groups.

1. INTRODUCTION

It is well known that Hardy's inequality and its generalization in $\mathbb{R}^N$ play an important role in many areas of mathematics. In the case of the Heisenberg group $\mathbb{H}_n$, Garofalo and Lanconelli (cf. [9]) first proved the following Hardy inequality:

$$(1.1) \quad \int_{\mathbb{H}_n} |\nabla_H u|^2 \geq \frac{(Q-2)^2}{4} \int_{\mathbb{H}_n} \frac{u^2}{d^2} |\nabla_H d|^2, \quad u \in C_0^\infty(\mathbb{H}_n \setminus \{e\}),$$

where e is the origin of $\mathbb{H}_n$, $d = (|z|^4 + t^2)^{\frac{1}{4}}$ is the Korányi-Folland nonisotropic gauge induced by the fundamental solution and $Q = 2n + 2$ is the homogeneous dimension of $\mathbb{H}_n$ (see also [10]). P. Niu et al. ([12]) generalized inequality (1.1) to the nonlinear cases. That is, for $1 < p < Q$, we have

$$(1.2) \quad \int_{\mathbb{H}_n} |\nabla_H u|^p \geq \left(\frac{Q-p}{p} \right)^p \int_{\mathbb{H}_n} \frac{|u|^p}{d^p} |\nabla_H d|^p, \quad u \in C_0^\infty(\mathbb{H}_n \setminus \{e\}).$$

To obtain (1.2), they used a Picone type identity on $\mathbb{H}_n$. D'Ambrosio (cf. [3, 4, 5]) extended (1.2) to some degenerate elliptic differential operators such as the Heisenberg-Greiner operator, a Baouendi-Grushin type operator and the sub-Laplacian on Carnot groups. More recently, Danielli, Garofalo and Phuc (cf. [6, 7]) consider various types of Hardy-Sobolev inequalities on a Carnot-Carathéodory space. They use the fundamental solution of the corresponding p -Laplace operator and generalize (1.2) to Carnot groups of arbitrary step.

The aim of this paper is to prove an analogous Hardy-type inequality (1.2) on $\mathbb{H}_n$, where the Korányi-Folland nonisotropic gauge is replaced by the Carnot-Carathéodory distance d_{cc} on $\mathbb{H}_n$ (we refer to [1, 11] for more information about

Received by the editors September 10, 2009 and, in revised form, September 2, 2010 and June 14, 2011.

2010 *Mathematics Subject Classification.* Primary 22E25; Secondary 26D10.

Key words and phrases. Hardy inequalities, Heisenberg group, Carnot-Carathéodory distance.

This work was supported by the National Science Foundation of China under Grant No. 10671009.

this distance). To our knowledge, little is known about Hardy inequalities related to such a distance. Since d_{cc} is not differentiable in the center of $\mathbb{H}_n$, it seems that the methods used in [3, 5, 7, 12] do not work for such a distance. To do so, we prove a representation formula associated with d_{cc} , and the idea is due to Cohn and Lu ([2]). The main result is the following theorem.

Theorem 1.1. *Let $1 < p < Q$ and $u \in C_0^\infty(\mathbb{H}_n)$. Then*

$$(1.3) \quad \int_{\mathbb{H}_n} |\nabla_H u|^p \geq \left(\frac{Q-p}{p} \right)^p \int_{\mathbb{H}_n} \frac{|u|^p}{d_{cc}^p}.$$

The Hardy type inequalities with weights imply the following Hardy-Rellich type inequalities (see e.g. [14] for analogous inequalities on $\mathbb{R}^N$).

Theorem 1.2. *Let $\mathbb{Z} = \{(z, t) \in \mathbb{H}_n : z = 0\}$ be the center of $\mathbb{H}_n$. It follows, for all $u \in C_0^\infty(\mathbb{H}_n \setminus \mathbb{Z})$, that*

$$(1.4) \quad \int_{\mathbb{H}_n} |\Delta_H u|^2 \geq \left(\frac{Q(Q-4)}{4} \right)^2 \int_{\mathbb{H}_n} \frac{u^2}{d_{cc}^4}, \quad n \geq 2,$$

and

$$(1.5) \quad \int_{\mathbb{H}_n} |\Delta_H u|^2 \geq \frac{Q^2}{4} \int_{\mathbb{H}_n} \frac{|\nabla_H u|^2}{d_{cc}^2}, \quad n \geq 3.$$

Finally, we obtain some Hardy inequalities involving the distance from the boundary which generalize a result of D'Ambrosio (cf. [5]).

Theorem 1.3. *Let $p > 1$ and $u \in C_0^\infty(B_{cc}(e, \rho))$, where $B_{cc}(e, \rho)$ is the Carnot-Carathéodory metric ball centered at the origin. Then*

$$(1.6) \quad \int_{B_{cc}(e, \rho)} |\nabla_H u|^p \geq \left(\frac{p-1}{p} \right)^p \int_{B_{cc}(e, \rho)} \frac{|u|^p}{(\rho - d_{cc})^p}.$$

2. NOTATION AND PRELIMINARIES

Let $\mathbb{H}_n = (\mathbb{C}^n \times \mathbb{R}, \circ)$ be the $(2n+1)$ -dimensional Heisenberg group whose group structure is given by

$$(z, t) \circ (z', t') = (z + z', t + t' + 2\operatorname{Im}\langle z, z' \rangle),$$

where $z = (z_1, \dots, z_n)$, $z' = (z'_1, \dots, z'_n) \in \mathbb{C}^n$, $z_j = x_j + iy_j$ ($x_j, y_j \in \mathbb{R}$) and $\langle z, z' \rangle = \sum_{j=1}^n z_j \cdot \bar{z}'_j$. The vector fields

$$X_j = \frac{\partial}{\partial x_j} + 2y_j \frac{\partial}{\partial t}, \quad Y_j = \frac{\partial}{\partial y_j} - 2x_j \frac{\partial}{\partial t}$$

($j = 1, \dots, n$) are left invariant and generate the Lie algebra of $\mathbb{H}_n$. The commutators of these vector fields satisfy

$$[X_j, Y_j] = -4 \frac{\partial}{\partial t}, \quad j = 1, 2, \dots, n$$

with all other brackets equal to zero. We denote by $\mathbb{Z} = \{(z, t) \in \mathbb{H}_n : z = 0\}$ the center of $\mathbb{H}_n$. Then Kohn's sub-Laplacian on $\mathbb{H}_n$ is

$$\Delta_H = \sum_{j=1}^n (X_j^2 + Y_j^2) = \sum_{j=1}^n \left(\frac{\partial^2}{\partial x_j^2} + \frac{\partial^2}{\partial y_j^2} \right) + 4|z|^2 \frac{\partial^2}{\partial t^2} + 4 \sum_{k=1}^n (y_j \frac{\partial}{\partial x_j} - x_j \frac{\partial}{\partial y_j}) \frac{\partial}{\partial t},$$

and the horizontal gradient is the $(2n)$ -dimensional vector given by

$$\nabla_H = (X_1, \dots, X_n, Y_1, \dots, Y_n).$$

We call a curve $\gamma : [a, b] \rightarrow \mathbb{H}_n$ a horizontal curve connecting two points $\xi, \eta \in \mathbb{H}_n$ if $\gamma(a) = \xi$, $\gamma(b) = \eta$ and $\gamma'(s) \in \text{span}\{X_1, \dots, X_n, Y_1, \dots, Y_n\}$ for all s . Then the Carnot-Carathéodory distance between ξ, η is defined as

$$d_{cc}(\xi, \eta) = \inf_{\gamma} \int_a^b \|\gamma'(s)\| ds,$$

where the infimum is taken over all horizontal curves γ connecting ξ and η . It is known that any two points ξ, η on $\mathbb{H}_n$ can be joined by a horizontal curve of finite length and then d_{cc} is a metric on $\mathbb{H}_n$. An important feature of this distance function is that the distance and thus the associated metric balls are left-invariant. With this norm, we can define the metric ball centered at the origin and with radius ρ associated with this metric by

$$B_{cc}(e, \rho) = \{\eta : d_{cc}(e, \eta) < \rho\}$$

and the unit sphere $\Sigma = \partial B_{cc}(e, 1)$. For simplicity, we write $d_{cc}(\xi) = d_{cc}(e, \xi)$.

For each real number $\lambda > 0$, there is a dilation naturally associated with the group structure which is usually denoted as $\delta_\lambda(z, t) = (\lambda z, \lambda^2 t)$. The Jacobian determinant of δ_λ is λ^Q , where $Q = 2n + 2$ is the homogeneous dimension of H_n . For simplicity, we use the notation $\lambda(z, t) = (\lambda z, \lambda^2 t)$. The Carnot-Carathéodory distance d_{cc} satisfies

$$d_{cc}(\lambda(z, t)) = \lambda d_{cc}(z, t), \quad \lambda > 0.$$

Given any $\xi = (z, t) \in \mathbb{H}_n$, set $z^* = \frac{z}{d_{cc}(z, t)}$, $t^* = \frac{t}{d_{cc}(z, t)^2}$ and $\xi^* = (z^*, t^*)$. The polar coordinates on $\mathbb{H}_n$ associated with d_{cc} are the following (cf. [8], Proposition 1.15 or [11], via the coarea formula):

$$\int_{\mathbb{H}_n} f(z, t) dx dt = \int_0^\infty \int_\Sigma f(\lambda(z^*, t^*)) \lambda^{Q-1} d\sigma d\lambda$$

for all $f \in L^1(\mathbb{H}_n)$.

Set

$$\mu(\theta) = \frac{2\theta - \sin 2\theta}{2 \sin^2 \theta} : (-\pi, \pi) \rightarrow \mathbb{R}.$$

μ is a diffeomorphism of the interval $(-\pi, \pi)$ onto $\mathbb{R}$ (cf. [1]). We denote by μ^{-1} the inverse function of μ . The Carnot-Carathéodory distance d_{cc} satisfies (cf. [1])

$$(2.1) \quad d_{cc}(z, t) = \begin{cases} \pi|t|, & z = 0, \\ \frac{\theta}{\sin \theta} \|z\|, & z \neq 0, \end{cases}$$

where

$$(2.2) \quad \theta = \mu^{-1} \left(\frac{t}{\|z\|^2} \right) \quad \text{and} \quad \|z\|^2 = \sum_{j=1}^n \|z_j\|^2.$$

On the other hand, if we set

$$\Omega = \{(B_1, \dots, B_n, A_1, \dots, A_n, \phi, \rho) : \sum_{j=1}^n (A_j^2 + B_j^2) = 1, -2\pi \leq \phi\rho \leq 2\pi, \rho \geq 0\}$$

and define $\Phi : \Omega \rightarrow \mathbb{H}_n$ by $\Phi(B_1, \dots, B_n, A_1, \dots, A_n, \phi, \rho) = (z, t)$, where

$$(2.3) \quad \begin{cases} x_j(B_1, \dots, B_n, A_1, \dots, A_n, \phi, \rho) = \frac{A_j(1 - \cos \phi\rho) + B_j \sin \phi\rho}{\phi} \\ y_j(B_1, \dots, B_n, A_1, \dots, A_n, \phi, \rho) = \frac{-B_j(1 - \cos \phi\rho) + A_j \sin \phi\rho}{\phi} \\ t(B_1, \dots, B_n, A_1, \dots, A_n, \phi, \rho) = 2 \frac{\phi\rho - \sin \phi\rho}{\phi^2} \end{cases}$$

($j = 1, 2, \dots, n$), then the range of Φ is $\mathbb{H}_n$ and the center $\mathbb{Z}$ is just the set of points $\Phi(B_1, \dots, B_n, A_1, \dots, A_n, \phi, \rho)$ with $\phi\rho = \pm 2\pi$. Furthermore, if one fixes $\rho > 0$, equation (2.3) parameterizes $\partial B_{cc}(e, \rho)$ (cf. [11]).

From (2.3), we have

$$(2.4) \quad \|z\| = \sqrt{\frac{2(1 - \cos \phi\rho)}{\phi^2}} = \sqrt{\frac{4 \sin^2 \frac{\phi\rho}{2}}{\phi^2}} = \frac{2 \sin \frac{\phi\rho}{2}}{\phi}$$

and

$$\mu(\theta) = \frac{t}{\|z\|^2} = \frac{\phi\rho - \sin \phi\rho}{2 \sin^2 \frac{\phi\rho}{2}} = \mu\left(\frac{\phi\rho}{2}\right).$$

Therefore,

$$(2.5) \quad \theta = \frac{\phi\rho}{2} \quad \text{when} \quad -2\pi < \phi\rho < 2\pi$$

since μ is a diffeomorphism of the interval $(-\pi, \pi)$ onto $\mathbb{R}$.

3. MAIN RESULTS

To prove the main result, we first need the following representation formula, and the basic idea of the proof is the same as Cohn and Lu's (see [2]).

Lemma 3.1. *Let $R_2 > R_1 > 0$ and $f \in C^1(B_{cc}(e, R_2) \setminus B_{cc}(e, R_1))$. Then*

$$(3.1) \quad \int_{\Sigma} f(R_2 \xi^*) d\sigma - \int_{\Sigma} f(R_1 \xi^*) d\sigma = \int_{B_{cc}(e, R_2) \setminus B_{cc}(e, R_1)} \langle \nabla_H f, \nabla_H d_{cc} \rangle \cdot \frac{1}{d_{cc}^{Q-1}} d\xi.$$

Proof. Let ξ^* be a point on the sphere, that is, $\xi^* = (z^*, t^*)$, where $d_{cc}(z^*, t^*) = 1$. We consider for $0 < R_1 < R_2$ the following difference using the fundamental theorem of calculus:

$$(3.2) \quad \begin{aligned} & \int_{\Sigma} f(R_2 \xi^*) d\sigma - \int_{\Sigma} f(R_1 \xi^*) d\sigma = \int_{\Sigma} \int_{R_1}^{R_2} \frac{d}{d\rho} f(\rho \xi^*) d\rho d\sigma \\ & = \int_{\Sigma} \int_{R_1}^{R_2} \left(\sum_{j=1}^n \frac{\partial f(\xi)}{\partial x_j} \cdot \frac{\partial x_j}{\partial \rho} + \sum_{j=1}^n \frac{\partial f(\xi)}{\partial y_j} \cdot \frac{\partial y_j}{\partial \rho} + \frac{\partial f(\xi)}{\partial t} \cdot \frac{\partial t}{\partial \rho} \right) d\rho d\sigma, \end{aligned}$$

where $\xi = (x, t) = \rho\xi^*$. Using equation (2.3), we have

$$\begin{aligned}
 (3.3) \quad & \sum_{j=1}^n \frac{\partial f(\xi)}{\partial x_j} \cdot \frac{\partial x_j}{\partial \rho} + \sum_{j=1}^n \frac{\partial f(\xi)}{\partial y_j} \cdot \frac{\partial y_j}{\partial \rho} + \frac{\partial f(\xi)}{\partial t} \cdot \frac{\partial t}{\partial \rho} \\
 &= \sum_{j=1}^n \frac{\partial f(\xi)}{\partial x_j} \cdot (A_j \sin \phi \rho + B_j \cos \phi \rho) + \sum_{j=1}^n \frac{\partial f(\xi)}{\partial y_j} \cdot (-B_j \sin \phi \rho + A_j \cos \phi \rho) \\
 &+ \frac{\partial f(\xi)}{\partial t} \cdot \frac{2 - 2 \cos \phi \rho}{\phi} \\
 &= \sum_{j=1}^n [X_j f \cdot (A_j \sin \phi \rho + B_j \cos \phi \rho) + Y_j f \cdot (-B_j \sin \phi \rho + A_j \cos \phi \rho)] + \frac{\partial f(\xi)}{\partial t} \\
 &\cdot \left\{ \frac{2 - 2 \cos \phi \rho}{\phi} - \sum_{j=1}^n [2y_j (A_j \sin \phi \rho + B_j \cos \phi \rho) - 2x_j (-B_j \sin \phi \rho + A_j \cos \phi \rho)] \right\}.
 \end{aligned}$$

Again using (2.3), we have

$$\begin{aligned}
 (3.4) \quad & \sum_{j=1}^n [2y_j (A_j \sin \phi \rho + B_j \cos \phi \rho) - 2x_j (-B_j \sin \phi \rho + A_j \cos \phi \rho)] \\
 &= 2 \sum_{j=1}^n \frac{(-B_j(1 - \cos \phi \rho) + A_j \sin \phi \rho)(A_j \sin \phi \rho + B_j \cos \phi \rho)}{\phi} \\
 &- 2 \sum_{j=1}^n \frac{(A_j(1 - \cos \phi \rho) + B_j \sin \phi \rho)(-B_j \sin \phi \rho + A_j \cos \phi \rho)}{\phi} \\
 &= 2 \sum_{j=1}^n \frac{(A_j^2 + B_j^2)(1 - \cos \phi \rho)}{\phi} = 2 \frac{1 - \cos \phi \rho}{\phi}.
 \end{aligned}$$

Combining (3.2), (3.3) and (3.4), we obtain

$$\begin{aligned}
 & \int_{\Sigma} f(R_2 \xi^*) d\sigma - \int_{\Sigma} f(R_1 \xi^*) d\sigma \\
 &= \int_{\Sigma} \int_{R_1}^{R_2} [X_j f \cdot (A_j \sin \phi \rho + B_j \cos \phi \rho) + Y_j f \cdot (-B_j \sin \phi \rho + A_j \cos \phi \rho)] dp d\sigma.
 \end{aligned}$$

Rewriting the last expression into a solid integral using the polar coordinates over $\mathbb{H}_n$, we get

$$\begin{aligned}
 & \int_{\Sigma} f(R_2 \xi^*) d\sigma - \int_{\Sigma} f(R_1 \xi^*) d\sigma \\
 &= \int_{B_{cc}(e, R_2) \setminus B_{cc}(e, R_1)} \sum_{j=1}^n \frac{(A_j \sin \phi \rho + B_j \cos \phi \rho) X_j f + (-B_j \sin \phi \rho + A_j \cos \phi \rho) Y_j f}{d_{cc}(\xi)^{Q-1}} d\xi.
 \end{aligned}$$

To finish our proof, it is enough to show that

$$X_j d_{cc}(\xi) = A_j \sin \phi \rho + B_j \cos \phi \rho, \quad Y_j d_{cc}(\xi) = -B_j \sin \phi \rho + A_j \cos \phi \rho$$

($j = 1, \dots, n$) in $\mathbb{H}_n \setminus \mathbb{Z}$. This is done by the following Lemma 3.2. The proof of Lemma 3.1 is now completed. $\square$

Lemma 3.2. *It follows that, for $\xi = (z, t) \in \mathbb{H}_n \setminus \mathbb{Z}$,*

$$X_j d_{cc}(\xi) = A_j \sin \phi \rho + B_j \cos \phi \rho, \quad Y_j d_{cc}(\xi) = -B_j \sin \phi \rho + A_j \cos \phi \rho \\ (j = 1, \dots, n).$$

Proof. Recall that if $z \neq 0$, then

$$d_{cc}(\xi) = d_{cc}(z, t) = \frac{\theta}{\sin \theta} \|z\|,$$

where $\theta = \mu^{-1}(t/\|z\|^2)$. A simple calculation shows that, for $j = 1, \dots, n$,

$$\mu'(\theta) = \frac{2 \sin \theta - 2\theta \cos \theta}{\sin^3 \theta}, \quad \frac{\partial \theta}{\partial x_j} = -\frac{2tx_j}{\|z\|^4} \cdot \frac{1}{\mu'(\theta)} = -\frac{tx_j}{\|z\|^4} \cdot \frac{\sin^3 \theta}{\sin \theta - \theta \cos \theta}, \\ \frac{\partial \theta}{\partial y_j} = -\frac{ty_j}{\|z\|^4} \cdot \frac{\sin^3 \theta}{\sin \theta - \theta \cos \theta}, \quad \frac{\partial \theta}{\partial t} = \frac{1}{\|z\|^2} \cdot \frac{\sin^3 \theta}{2 \sin \theta - 2\theta \cos \theta}.$$

Therefore, for $\xi = (z, t) \in \mathbb{H}_n \setminus \mathbb{Z}$,

$$\begin{aligned} \frac{\partial d_{cc}(\xi)}{\partial x_j} &= \frac{\partial}{\partial x_j} \left(\frac{\theta}{\sin \theta} \|z\| \right) = \frac{x_j}{\|z\|} \cdot \frac{\theta}{\sin \theta} + \|z\| \cdot \frac{\sin \theta - \theta \cos \theta}{\sin^2 \theta} \cdot \frac{\partial \theta}{\partial x_j} \\ &= \frac{x_j}{\|z\|} \cdot \frac{\theta}{\sin \theta} - \frac{tx_j}{\|z\|^3} \cdot \sin \theta = \frac{x_j}{\|z\|} \cdot \frac{\theta}{\sin \theta} - \frac{x_j}{\|z\|} \cdot \mu(\theta) \sin \theta \\ &= \frac{x_j}{\|z\|} \cdot \cos \theta. \end{aligned}$$

Similarly,

$$\begin{aligned} \frac{\partial d_{cc}(\xi)}{\partial y_j} &= \frac{\partial}{\partial y_j} \left(\frac{\theta}{\sin \theta} \|z\| \right) = \frac{y_j}{\|z\|} \cdot \frac{\theta}{\sin \theta} + \|z\| \cdot \frac{\sin \theta - \theta \cos \theta}{\sin^2 \theta} \cdot \frac{\partial \theta}{\partial y_j} \\ &= \frac{y_j}{\|z\|} \cdot \cos \theta, \end{aligned}$$

$$\frac{\partial d_{cc}(\xi)}{\partial t} = \frac{\partial}{\partial t} \left(\frac{\theta}{\sin \theta} \|z\| \right) = \|z\| \cdot \frac{\sin \theta - \theta \cos \theta}{\sin^2 \theta} \cdot \frac{\partial \theta}{\partial t} = \frac{1}{\|z\|} \cdot \frac{\sin \theta}{2}.$$

Thus

$$X_j d_{cc}(\xi) = \left(\frac{\partial}{\partial x_j} + 2y_j \frac{\partial}{\partial t} \right) d_{cc}(\xi) = \frac{x_j}{\|z\|} \cdot \cos \theta + \frac{y_j}{\|z\|} \cdot \sin \theta.$$

Using equations (2.3) and (2.4), we get

$$\begin{aligned} X_j d_{cc}(\xi) &= \frac{\phi}{2 \sin \frac{\phi \rho}{2}} \left(\frac{A_j(1 - \cos \phi \rho) + B_j \sin \phi \rho}{\phi} \cos \theta \right. \\ &\quad \left. + \frac{-B_j(1 - \cos \phi \rho) + A_j \sin \phi \rho}{\phi} \sin \theta \right) \\ &= \left(A_j \sin \frac{\phi \rho}{2} + B_j \cos \frac{\phi \rho}{2} \right) \cos \theta + \left(-B_j \sin \frac{\phi \rho}{2} + A_j \cos \frac{\phi \rho}{2} \right) \sin \theta \\ &= A_j \sin \left(\frac{\phi \rho}{2} + \theta \right) + B_j \cos \left(\frac{\phi \rho}{2} + \theta \right). \end{aligned}$$

Therefore we obtain, by (2.5),

$$X_j d_{cc}(\xi) = A_j \sin \phi \rho + B_j \cos \phi \rho.$$

On the other hand, we have, again using equations (2.3), (2.4) and (2.5),

$$\begin{aligned}
 Y_j d_{cc}(\xi) &= \left(\frac{\partial}{\partial y_j} - 2x_j \frac{\partial}{\partial t} \right) d_{cc}(\xi) = \frac{y_j}{\|z\|} \cdot \cos \theta - \frac{x_j}{\|z\|} \cdot \sin \theta \\
 &= \frac{\phi}{2 \sin \frac{\phi \rho}{2}} \left(\frac{-B_j(1 - \cos \phi \rho) + A_j \sin \phi \rho}{\phi} \cos \theta \right. \\
 &\quad \left. - \frac{A_j(1 - \cos \phi \rho) + B_j \sin \phi \rho}{\phi} \sin \theta \right) \\
 &= \left(-B_j \sin \frac{\phi \rho}{2} + A_j \cos \frac{\phi \rho}{2} \right) \cos \theta - \left(A_j \sin \frac{\phi \rho}{2} + B_j \cos \frac{\phi \rho}{2} \right) \sin \theta \\
 &= A_j \cos \left(\frac{\phi \rho}{2} + \theta \right) - B_j \sin \left(\frac{\phi \rho}{2} + \theta \right) \\
 &= A_j \cos \phi \rho - B_j \sin \phi \rho.
 \end{aligned}$$

This completes the proof of Lemma 3.2. $\square$

Remark 3.3. Putting $f = u \in C_0^\infty(\mathbb{H}_n)$ in Lemma 3.1 and letting $R_2 \rightarrow \infty$ and $R_1 \rightarrow 0+$, we have

$$|\Sigma|u(e) = - \int_{\mathbb{H}_n} \langle \nabla_H u, \nabla_H d_{cc} \rangle \cdot \frac{1}{d_{cc}^{Q-1}} d\xi,$$

where $|\Sigma| = \int_\Sigma d\sigma$ (see [2] for an analogous representation formula associated with the Korányi-Folland nonisotropic gauge). We note the representation formula in Lemma 3.1 is also valid for $f \in C(\mathbb{H}_n) \cap C^1(\mathbb{H}_n \setminus \mathbb{Z})$. In fact, since $\xi \in \mathbb{Z}$ iff $\lambda \xi \in \mathbb{Z}$ for all $\lambda > 0$, one has, for all $f \in C(\mathbb{H}_n) \cap C^1(\mathbb{H}_n \setminus \mathbb{Z})$, as in the proof of Lemma 3.1,

$$\begin{aligned}
 &\int_{\Sigma \setminus (\Sigma \cap \mathbb{Z})} f(R_2 \xi^*) d\sigma - \int_{\Sigma \setminus (\Sigma \cap \mathbb{Z})} f(R_1 \xi^*) d\sigma \\
 &= \int_{(B_{cc}(e, R_2) \setminus \mathbb{Z}) \setminus (B_{cc}(e, R_1) \setminus \mathbb{Z})} \frac{\langle \nabla_H f, \nabla_H d_{cc} \rangle}{d_{cc}^{Q-1}} d\xi,
 \end{aligned}$$

where $R_2 > R_1 > 0$. Note that $\Sigma \cap \mathbb{Z} = \{(\mathbf{0}, 1/\pi), (\mathbf{0}, -1/\pi)\}$. So

$$\int_{\Sigma \setminus (\Sigma \cap \mathbb{Z})} f(R_k \xi^*) d\sigma = \int_\Sigma f(R_k \xi^*) d\sigma < \infty, \quad k = 1, 2,$$

since $f \in C(\mathbb{H}_n)$. These imply that

$$\begin{aligned}
 &\left| \int_{(B_{cc}(e, R_2) \setminus \mathbb{Z}) \setminus (B_{cc}(e, R_1) \setminus \mathbb{Z})} \frac{\langle \nabla_H f, \nabla_H d_{cc} \rangle}{d_{cc}^{Q-1}} d\xi \right| \\
 &= \left| \int_{\Sigma \setminus (\Sigma \cap \mathbb{Z})} f(R_2 \xi^*) d\sigma - \int_{\Sigma \setminus (\Sigma \cap \mathbb{Z})} f(R_1 \xi^*) d\sigma \right| \\
 &= \left| \int_\Sigma f(R_2 \xi^*) d\sigma - \int_\Sigma f(R_1 \xi^*) d\sigma \right| < \infty.
 \end{aligned}$$

Therefore,

$$\int_{(B_{cc}(e, R_2) \setminus \mathbb{Z}) \setminus (B_{cc}(e, R_1) \setminus \mathbb{Z})} \frac{\langle \nabla_H f, \nabla_H d_{cc} \rangle}{d_{cc}^{Q-1}} d\xi = \int_{B_{cc}(e, R_2) \setminus B_{cc}(e, R_1)} \frac{\langle \nabla_H f, \nabla_H d_{cc} \rangle}{d_{cc}^{Q-1}} d\xi$$

since $\mathbb{Z}$ is a measure-zero set. This means that the representation formula in Lemma 3.1 is valid for $f \in C(\mathbb{H}_n) \cap C^1(\mathbb{H}_n \setminus \mathbb{Z})$.

Proof of Theorem 1.1. Let $\epsilon > 0$. Then $0 \leq u_\epsilon := (|u|^2 + \epsilon^2)^{p/2} - \epsilon^p \in C_0^\infty(\mathbb{H}_n)$. In fact, u_ϵ has the same support as u . Putting $f = u_\epsilon d_{cc}^{Q-p}$ in Lemma 3.1 and letting $R_2 \rightarrow \infty$ and $R_1 \rightarrow 0+$, we get, by Remark 3.3, since $d_{cc} \in C(\mathbb{H}_n) \cap C^\infty(\mathbb{H}_n \setminus \mathbb{Z})$ and $d_{cc}(e) = 0$,

$$\int_{\mathbb{H}_n} \langle \nabla_H u_\epsilon, \nabla_H d_{cc} \rangle \cdot \frac{1}{d_{cc}^{p-1}} + (Q-p) \int_{\mathbb{H}_n} \frac{u_\epsilon}{d_{cc}^p} = 0.$$

Here we use the fact that $|\nabla_H d_{cc}(\xi)| = 1$ when $\xi \in \mathbb{H}_n \setminus \mathbb{Z}$ (cf. [11]). By Hölder's inequality:

$$\begin{aligned} (Q-p) \int_{\mathbb{H}_n} \frac{u_\epsilon}{d_{cc}^p} &= -p \int_{\mathbb{H}_n} (|u|^2 + \epsilon^2)^{(p-2)/2} u \langle \nabla_H u, \nabla_H d_{cc} \rangle \cdot \frac{1}{d_{cc}^{p-1}} \\ &\leq p \int_{\mathbb{H}_n} \frac{(|u|^2 + \epsilon^2)^{(p-2)/2} |u| \cdot |\nabla_H u|}{d_{cc}^{p-1}} \\ &\leq p \left(\int_{\mathbb{H}_n} \frac{(|u|^2 + \epsilon^2)^{p(p-2)/2(p-1)} |u|^{\frac{p}{p-1}}}{d_{cc}^p} \right)^{\frac{p-1}{p}} \left(\int_{\mathbb{H}_n} |\nabla_H u|^p \right)^{\frac{1}{p}}. \end{aligned}$$

By dominated convergence, letting $\epsilon \rightarrow 0+$, canceling and raising both sides to the power p , we get (1.3). The proof of Theorem 1.1 is complete. $\square$

Remark 3.4. Let $V^{1,p}(\mathbb{H}_n)$ denote the space

$$V^{1,p}(\mathbb{H}_n) = \{f \in C(\mathbb{H}_n) \cap C^1(\mathbb{H}_n \setminus \mathbb{Z}) : \int_{\mathbb{H}_n} \frac{|u|^p}{d_{cc}^p} < +\infty, \int_{\mathbb{H}_n} |\nabla_H f|^p < +\infty\}.$$

In this remark, we shall show that the following Hardy inequalities hold for all $f \in V^{1,p}(\mathbb{H}_n)$:

$$\int_{\mathbb{H}_n} |\nabla_H f|^p \geq \left(\frac{Q-p}{p} \right)^p \int_{\mathbb{H}_n} \frac{|f|^p}{d_{cc}^p}.$$

Furthermore, the constant $\left(\frac{Q-p}{p} \right)^p$ is sharp in the sense of

$$\left(\frac{Q-p}{p} \right)^p = \inf_{f \in V^{1,p}(\mathbb{H}_n) \setminus \{0\}} \frac{\int_{\mathbb{H}_n} |\nabla_H f|^p}{\int_{\mathbb{H}_n} \frac{|f|^p}{d_{cc}^p}}.$$

Step 1. We shall show that if $f \in V^{1,p}(\mathbb{H}_n)$, then

$$(3.5) \quad \lim_{R \rightarrow +\infty} R^{Q-p} \int_{\Sigma} |f(R\xi^*)|^p d\sigma = 0.$$

We note from Lemma 3.1 that

$$\begin{aligned}
 & \left| R_2^{Q-p} \int_{\Sigma} |f(R_2 \xi^*)|^p d\sigma - R_1^{Q-p} \int_{\Sigma} |f(R_1 \xi^*)|^p d\sigma \right| \\
 &= \left| \int_{B_{cc}(e, R_2) \setminus B_{cc}(e, R_1)} \langle \nabla_H(|f|^p d_{cc}^{Q-p}), \nabla_H d_{cc} \rangle \cdot \frac{1}{d_{cc}^{Q-1}} d\xi \right| \\
 &\leq p \int_{B_{cc}(e, R_2) \setminus B_{cc}(e, R_1)} \frac{|f|^{p-1} |\nabla_H f|}{d_{cc}^{p-1}} d\xi + (Q-p) \int_{B_{cc}(e, R_2) \setminus B_{cc}(e, R_1)} \frac{|f|^p}{d_{cc}^p} d\xi \\
 &\leq p \left(\int_{B_{cc}(e, R_2) \setminus B_{cc}(e, R_1)} \frac{|f|^p}{d_{cc}^p} \right)^{\frac{p-1}{p}} \left(\int_{B_{cc}(e, R_2) \setminus B_{cc}(e, R_1)} |\nabla_H f|^p \right)^{\frac{1}{p}} \\
 &\quad + (Q-p) \int_{B_{cc}(e, R_2) \setminus B_{cc}(e, R_1)} \frac{|f|^p}{d_{cc}^p} d\xi.
 \end{aligned}$$

Thus,

$$\left| R_2^{Q-p} \int_{\Sigma} |f(R_2 \xi^*)|^p d\sigma - R_1^{Q-p} \int_{\Sigma} |f(R_1 \xi^*)|^p d\sigma \right| \rightarrow 0$$

as $R_1, R_2 \rightarrow +\infty$. This shows that $R^{Q-p} \int_{\Sigma} |f(R \xi^*)|^p d\sigma$ converges to a finite constant $f_{\infty} \geq 0$ as $R \rightarrow +\infty$. One must have $f_{\infty} = 0$ for

$$\int_{\mathbb{H}_n} \frac{|f|^p}{d_{cc}^p} d\xi = \int_0^{\infty} \int_{\Sigma} |f(\lambda \xi^*)|^p \lambda^{Q-p-1} d\sigma d\lambda = \int_0^{\infty} \frac{\lambda^{Q-p} \int_{\Sigma} |f(\lambda \xi^*)|^p d\sigma}{\lambda} d\lambda < +\infty.$$

Therefore, (3.5) holds.

Step 2. Replace f by $d_{cc}^{Q-p}|f|^p$ in Lemma 3.1 and let $R_2 \rightarrow \infty$ and $R_1 \rightarrow 0+$. Using the result of Step 1 and following the proof of Theorem 1.1, one can obtain the Hardy inequalities as expected.

We now show that the constant $\left(\frac{Q-p}{p}\right)^p$ is sharp. The proof is similar to that of Lemma 2.3 in [13]. Let

$$f_{\varepsilon}(z, t) = \begin{cases} 1, & d_{cc} \leq 1; \\ d_{cc}^{-(Q-p)/p-\varepsilon}, & d_{cc} > 1. \end{cases}$$

We note that this family of functions can be approximated by the functions in $V^{1,p}(\mathbb{H}_n)$. In addition,

$$\int_{\mathbb{H}_n} \frac{|f_{\varepsilon}|^p}{d_{cc}^p} d\xi = \frac{|\Sigma|}{Q-p} + \frac{|\Sigma|}{\varepsilon p}$$

and

$$\int_{\mathbb{H}_n} |\nabla_H f_{\varepsilon}|^p d\xi = \left(-\frac{Q-p}{p} - \varepsilon\right)^p \cdot \frac{|\Sigma|}{\varepsilon p}.$$

Passing to the limit as $\varepsilon \rightarrow 0$, one can see that the constant $\left(\frac{Q-p}{p}\right)^p$ is sharp. We note that since the Carnot-Carathéodory distance is not differentiable in $\mathbb{Z}$, it is unknown whether f_{ε} , the family of functions above, can be approximated by the functions in $C_0^{\infty}(\mathbb{H}_n)$. So, we do not know whether the constant $\left(\frac{Q-p-\alpha}{p}\right)^p$ is sharp for all $f \in C_0^{\infty}(\mathbb{H}_n)$ in Theorem 1.1.

Remark 3.5. Let $\alpha < Q - p$. Put $f = u_\epsilon d_{cc}^{Q-p-\alpha}$ in Lemma 3.1 and let $R_2 \rightarrow \infty$ and $R_1 \rightarrow 0+$. We have

$$\begin{aligned} (Q - p - \alpha) \int_{\mathbb{H}_n} \frac{u_\epsilon}{d_{cc}^{p+\alpha}} &= -p \int_{\mathbb{H}_n} (|u|^2 + \epsilon^2)^{(p-2)/2} u \langle \nabla_H u, \nabla_H d_{cc} \rangle \cdot \frac{1}{d_{cc}^{p+\alpha-1}} \\ &\leq p \int_{\mathbb{H}_n} \frac{(|u|^2 + \epsilon^2)^{(p-2)/2} |u| \cdot |\nabla_H u|}{d_{cc}^{p+\alpha-1}} \\ &\leq p \left(\int_{\mathbb{H}_n} \frac{(|u|^2 + \epsilon^2)^{p(p-2)/2(p-1)} |u|^{\frac{p}{p-1}}}{d_{cc}^{p+\alpha}} \right)^{\frac{p-1}{p}} \left(\int_{\mathbb{H}_n} \frac{|\nabla_H u|^p}{d_{cc}^\alpha} \right)^{\frac{1}{p}}. \end{aligned}$$

As in the proof of Theorem 1.1, we obtain the following Hardy-type inequalities with weights:

$$(3.6) \quad \int_{\mathbb{H}_n} \frac{|\nabla_H u|^p}{d_{cc}^\alpha} \geq \left(\frac{Q - p - \alpha}{p} \right)^p \int_{\mathbb{H}_n} \frac{|u|^p}{d_{cc}^{p+\alpha}}, \quad \alpha < Q - p.$$

Proof of Theorem 1.2. Notice that

$$- \int_{\mathbb{H}_n} u \Delta_H u \cdot \frac{1}{d_{cc}^2} = -\frac{1}{2} \int_{\mathbb{H}_n} \Delta_H u^2 \cdot \frac{1}{d_{cc}^2} + \int_{\mathbb{H}_n} \frac{|\nabla_H u|^2}{d_{cc}^2}.$$

Through integration by parts, since $u \in C_0^\infty(\mathbb{H}_n \setminus \mathbb{Z})$ and d_{cc} is C^1 in $\mathbb{H}_n \setminus \mathbb{Z}$ (cf. [11]),

$$-\frac{1}{2} \int_{\mathbb{H}_n} \Delta_H u^2 \cdot \frac{1}{d_{cc}^2} = - \int_{\mathbb{H}_n} \langle \nabla_H u^2, \nabla_H d_{cc} \rangle \cdot \frac{1}{d_{cc}^3} d\xi.$$

Putting $f = u^2 d_{cc}^{Q-4}$ in Lemma 3.1 and letting $R_2 \rightarrow \infty$ and $R_1 \rightarrow 0+$, we have

$$- \int_{\mathbb{H}_n} \langle \nabla_H u^2, \nabla_H d_{cc} \rangle \cdot \frac{1}{d_{cc}^3} d\xi = (Q - 4) \int_{\mathbb{H}_n} \frac{u^2}{d_{cc}^4} d\xi.$$

Thus, for all $\varepsilon > 0$,

$$\begin{aligned} (3.7) \quad - \int_{\mathbb{H}_n} u \Delta_H u \cdot \frac{1}{d_{cc}^2} &= \int_{\mathbb{H}_n} \frac{|\nabla_H u|^2}{d_{cc}^2} + (Q - 4) \int_{\mathbb{H}_n} \frac{u^2}{d_{cc}^4} d\xi \\ &\leq \left(\int_{\mathbb{H}_n} \frac{u^2}{d_{cc}^4} \right)^{\frac{1}{2}} \left(\int_{\mathbb{H}_n} |\Delta_H u|^2 \right)^{\frac{1}{2}} \\ &\leq \frac{\varepsilon}{2} \int_{\mathbb{H}_n} \frac{u^2}{d_{cc}^4} + \frac{1}{2\varepsilon} \int_{\mathbb{H}_n} |\Delta_H u|^2. \end{aligned}$$

Choosing $\varepsilon = \frac{Q(Q-4)}{4}$ in (3.7), we have

$$(3.8) \quad \int_{\mathbb{H}_n} |\Delta_H u|^2 + \frac{Q(Q-4)^2(Q-8)}{16} \int_{\mathbb{H}_n} \frac{u^2}{d_{cc}^4} \geq \frac{Q(Q-4)}{2} \int_{\mathbb{H}_n} \frac{|\nabla_H u|^2}{d_{cc}^2}.$$

Since (see (3.6))

$$(3.9) \quad \int_{\mathbb{H}_n} \frac{|\nabla_H u|^2}{d_{cc}^2} \geq \frac{(Q-4)^2}{4} \int_{\mathbb{H}_n} \frac{u^2}{d_{cc}^4}, \quad n \geq 2,$$

we obtain, by (3.8),

$$\int_{\mathbb{H}_n} |\Delta_H u|^2 \geq \left(\frac{Q(Q-4)}{4} \right)^2 \int_{\mathbb{H}_n} \frac{u^2}{d_{cc}^4}.$$

When $Q \geq 8$, i.e. $n \geq 3$, again using (3.7) and (3.8), we obtain

$$\frac{Q(Q-4)}{2} \int_{\mathbb{H}_n} \frac{|\nabla_H u|^2}{d_{cc}^2} \leq \int_{\mathbb{H}_n} |\Delta_H u|^2 + \frac{Q(Q-8)}{4} \int_{\mathbb{H}_n} \frac{|\nabla_H u|^2}{d_{cc}^2}.$$

The desired result follows. $\square$

Proof of Theorem 1.3. As in the proof of Theorem 1.1, putting $f = \frac{u_\epsilon d_{cc}^{Q-1}}{(\rho - d_{cc})^{p-1}}$ in Lemma 3.1 and letting $R_2 = \rho$ and $R_1 \rightarrow 0+$, we get

$$\begin{aligned} & (Q-1) \int_{B_{cc}(e,\rho)} \frac{u_\epsilon}{d_{cc}(\rho - d_{cc})^{p-1}} + (p-1) \int_{B_{cc}(e,\rho)} \frac{u_\epsilon}{(\rho - d_{cc})^p} \\ &= -p \int_{B_{cc}(e,\rho)} (|u|^2 + \epsilon^2)^{(p-2)/2} u \langle \nabla_H u, \nabla_H d_{cc} \rangle \cdot \frac{1}{(\rho - d_{cc})^{p-1}} \\ &\leq p \int_{B_{cc}(e,\rho)} \frac{(|u|^2 + \epsilon^2)^{(p-2)/2} |u| \cdot |\nabla_H u|}{(\rho - d_{cc})^{p-1}} \\ &\leq p \left(\int_{B_{cc}(e,\rho)} \frac{(|u|^2 + \epsilon^2)^{p(p-2)/2(p-1)} |u|^{\frac{p}{p-1}}}{(\rho - d_{cc})^p} \right)^{\frac{p-1}{p}} \left(\int_{B_{cc}(e,\rho)} |\nabla_H u|^p \right)^{\frac{1}{p}}. \end{aligned}$$

Letting $\epsilon \rightarrow 0+$, we have, by dominated convergence,

$$\begin{aligned} & \left(\int_{B_{cc}(e,\rho)} \frac{|u|^p}{(\rho - d_{cc})^p} \right)^{\frac{p-1}{p}} \left(\int_{B_{cc}(e,\rho)} |\nabla_H u|^p \right)^{\frac{1}{p}} \\ &\geq \frac{p-1}{p} \int_{B_{cc}(e,\rho)} \frac{|u|^p}{(\rho - d_{cc})^p} \\ &\quad + \frac{Q-1}{p} \int_{B_{cc}(e,\rho)} \frac{|u|^p}{d_{cc}(\rho - d_{cc})^{p-1}} \\ &\geq \frac{p-1}{p} \int_{B_{cc}(e,\rho)} \frac{|u|^p}{(\rho - d_{cc})^p}. \end{aligned}$$

Canceling and raising both sides to the power p , we get (1.6). $\square$

ACKNOWLEDGEMENT

The authors thank the referee for a careful reading and very useful comments which improved the final version of this paper.

REFERENCES

- [1] Beals R., Gaveau B. and Greiner P. C., Hamilton-Jacobi theory and the heat kernel on Heisenberg groups, *J. Math. Pure Appl.*, **79** (2000), 633-689. MR1776501 (2001g:35047)
- [2] Cohn, W. and Lu, G. Best constants for Moser-Trudinger inequalities on the Heisenberg group, *Indiana Univ. Math. J.*, **50** (2001), 1567-1591. MR1889071 (2003i:46032)
- [3] D'Ambrosio L., Some Hardy inequalities on the Heisenberg group, *Differential Equations*, **40** (2004), 552-564. MR2153649 (2007d:26016)
- [4] D'Ambrosio L., Hardy inequalities related to Grushin type operators, *Proc. Amer. Math. Soc.*, **132** (2004), 725-734. MR2019949 (2005c:35050)
- [5] D'Ambrosio L., Hardy-type inequalities related to degenerate elliptic differential operators, *Ann. Sc. Norm. Super. Pisa Cl. Sci.*, (5) **IV** (2005), 451-486. MR2185865 (2006j:35027)

- [6] Danielli D., Garofalo N., and Phuc N. C., Inequalities of Hardy-Sobolev type in Carnot-Carathéodory spaces, Sobolev Spaces in Mathematics, vol. I (dedicated to the centenary of Sergey Sobolev), International Mathematical Series, Vol. 8, Springer, 2009. MR2508841 (2010i:46056)
- [7] Danielli D., Garofalo N., and Phuc N. C., Hardy-Sobolev type inequalities with sharp constants in Carnot-Carathéodory spaces, Potential Anal., **34** (2011), 223-242. MR2782971
- [8] Folland G. B. and Stein E. M., Hardy spaces on homogeneous groups, Princeton University Press, Princeton, NJ, 1982. MR657581 (84h:43027)
- [9] Garofalo N. and Lanconelli E., Frequency functions on the Heisenberg group, the uncertainty principle and unique continuation, Ann. Inst. Fourier (Grenoble), **40** (1990), 313-356. MR1070830 (91i:22014)
- [10] Goldstein J. A. and Zhang Q. S., On a degenerate heat equation with a singular potential, J. Funct. Anal., **186** (2001), 342-359. MR1864826 (2002k:35179)
- [11] Monti R., Some properties of Carnot-Carathéodory balls in the Heisenberg group, Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl., **11**(3) (2000), 155-167. MR1841689 (2002c:53048)
- [12] Niu P., Zhang H. and Wang Y., Hardy type and Rellich type inequalities on the Heisenberg group, Proc. Amer. Math. Soc., **129** (2001), 3623-3630. MR1860496 (2002i:26009)
- [13] Secchi S., Smets D., and Willem M., Remarks on a Hardy-Sobolev inequality, C. R. Acad. Sci. Paris, Ser. I, **336** (2003), 811-815. MR1990020 (2004d:35050)
- [14] Tertikas A. and Zographopoulos N. B., Best constants in the Hardy-Rellich inequalities and related improvements, Adv. Math., **209** (2007), 407-459. MR2296305 (2007m:26014)

SCHOOL OF MATHEMATICS AND STATISTICS, WUHAN UNIVERSITY, WUHAN, 430072, PEOPLE'S REPUBLIC OF CHINA

E-mail address: qiaohyang2465@yahoo.com.cn