ON RADIAL AND POLAR BLASCHKE-MINKOWSKI HOMOMORPHISMS

CHANG-JIAN ZHAO

(Communicated by Alexander N. Dranishnikov)

ABSTRACT. Brunn-Minkowski type inequalities for radial Blaschke-Minkowski homomorphisms of star bodies and polar Blaschke-Minkowski homomorphisms of convex bodies are established.

The setting for this paper is n -dimensional Euclidean space $\mathbb{R}^n$ ($n > 2$). Let $\mathcal{C}^n$ denote the set of nonempty convex figures (compact, convex subsets) and let $\mathcal{K}^n$ denote the subset of $\mathcal{C}^n$ consisting of all convex bodies (compact, convex subsets with nonempty interiors) in $\mathbb{R}^n$. We reserve the letter u for unit vectors, and the letter B is reserved for the unit ball centered at the origin. The surface of B is S^{n-1} . The volume of the unit n -ball is denoted by ω_n . For $u \in S^{n-1}$, let E_u denote the hyperplane, through the origin, that is orthogonal to u . We will use K^u to denote the image of K under an orthogonal projection onto the hyperplane E_u .

We use $V(K)$ for the n -dimensional volume of a convex body K . Let $h(K, \cdot) : S^{n-1} \rightarrow \mathbb{R}$ denote the support function of $K \in \mathcal{K}^n$; i.e. for $u \in S^{n-1}$,

$$h(K, u) = \text{Max}\{u \cdot x : x \in K\},$$

where $u \cdot x$ denotes the usual inner product of u and x in $\mathbb{R}^n$.

Let δ denote the Hausdorff metric on $\mathcal{K}^n$, i.e., for $K, L \in \mathcal{K}^n$, $\delta(K, L) = |h_K - h_L|_\infty$, where $|\cdot|_\infty$ denotes the sup-norm on the space of continuous functions $C(S^{n-1})$.

Associated with a compact subset K of $\mathbb{R}^n$, which is star-shaped with respect to the origin, is its radial function $\rho(K, \cdot) : S^{n-1} \rightarrow \mathbb{R}$, defined for $u \in S^{n-1}$, by

$$\rho(K, u) = \text{Max}\{\lambda \geq 0 : \lambda u \in K\}.$$

If $\rho(K, \cdot)$ is positive and continuous, K will be called a star body. Let $\mathcal{S}^n$ denote the set of star bodies in $\mathbb{R}^n$. Let $\tilde{\delta}$ denote the radial Hausdorff metric, i.e., if $K, L \in \mathcal{S}^n$, then $\tilde{\delta}(K, L) = |\rho_K - \rho_L|_\infty$.

If K and L are convex bodies in $\mathbb{R}^n$, then there is a convex body $K \dot{+} L$ such that

$$S(K \dot{+} L, \cdot) = S(K, \cdot) + S(L, \cdot),$$

where $S(K, \cdot)$ denotes the surface area measure of K . The operation $\dot{+}$ is called the Blaschke sum (see e.g. [13]).

Received by the editors April 4, 2011 and, in revised form, June 4, 2011 and June 26, 2011.

2010 *Mathematics Subject Classification*. Primary 52A40, 53A15.

Key words and phrases. Blaschke-Minkowski homomorphism, radial Blaschke-Minkowski homomorphism, polar Blaschke-Minkowski homomorphisms, radial Blaschke sum, radial Minkowski sum.

This research is supported by the National Natural Science Foundation of China (10971205).

If K is a convex body that contains the origin in its interior, the polar body K^* of K is defined by

$$K^* := \{x \in \mathbb{R}^n | x \cdot y \leq 1, y \in K\}.$$

1. DUAL MIXED VOLUMES

The radial Minkowski linear combination, $\lambda_1 K_1 \tilde{+} \cdots \tilde{+} \lambda_r K_r$, is defined by

$$(1.1) \quad \lambda_1 K_1 \tilde{+} \cdots \tilde{+} \lambda_r K_r = \{\lambda_1 x_1 \tilde{+} \cdots \tilde{+} \lambda_r x_r : x_i \in K_i, i = 1, \dots, r\},$$

for $K_1, \dots, K_r \in \mathcal{S}^n$ and $\lambda_1, \dots, \lambda_r \in \mathbb{R}$.

It has the following important property (see [13]):

$$(1.2) \quad \rho(\lambda K \tilde{+} \mu L, \cdot) = \lambda \rho(K, \cdot) + \mu \rho(L, \cdot),$$

for $K, L \in \mathcal{S}^n$ and $\lambda, \mu \geq 0$.

For $K_1, \dots, K_r \in \mathcal{S}^n$ and $\lambda_1, \dots, \lambda_r \geq 0$, the volume of the radial Minkowski linear combination $\lambda_1 K_1 \tilde{+} \cdots \tilde{+} \lambda_r K_r$ is a homogeneous polynomial of degree n in the λ_i ,

$$(1.3) \quad V(\lambda_1 K_1 \tilde{+} \cdots \tilde{+} \lambda_r K_r) = \sum \tilde{V}(K_{i_1}, \dots, K_{i_n}) \lambda_{i_1} \cdots \lambda_{i_n},$$

where the sum is taken over all n -tuples $(i_1, \dots, i_n)$ whose entries are positive integers not exceeding r . If we require the coefficients of the polynomial in (1.3) to be symmetric in their argument, then they are uniquely determined. The coefficient $\tilde{V}(K_{i_1}, \dots, K_{i_n})$ is nonnegative and depends only on the bodies $K_{i_1}, \dots, K_{i_n}$. It is called the dual mixed volume of $K_{i_1}, \dots, K_{i_n}$.

If $K_1, \dots, K_n \in \mathcal{S}^n$, the dual mixed volume $\tilde{V}(K_1, \dots, K_n)$ can be represented in the form (see [14])

$$(1.4) \quad \tilde{V}(K_1, \dots, K_n) = \frac{1}{n} \int_{S^{n-1}} \rho(K_1, u) \cdots \rho(K_n, u) dS(u).$$

If $K_1 = \cdots = K_{n-i} = K$, $K_{n-i+1} = \cdots = K_n = L$, the dual mixed volume is written as $\tilde{V}_i(K, L)$. If $L = B$, the dual mixed volume $\tilde{V}_i(K, L) = \tilde{V}_i(K, B)$ is written as $\tilde{W}_i(K)$.

For $K, L \in \mathcal{S}^n$, the i -th dual mixed volume of K and L , $\tilde{V}_i(K, L)$, can be extended to all $i \in \mathbb{R}$ by

$$(1.5) \quad \tilde{V}_i(K, L) = \frac{1}{n} \int_{S^{n-1}} \rho(K, u)^{n-i} \rho(L, u)^i dS(u), \quad i \in \mathbb{R}.$$

Thus, if $K \in \mathcal{S}^n$ and $i \in \mathbb{R}$, then (see [14])

$$(1.6) \quad \tilde{W}_i(K) = \frac{1}{n} \int_{S^{n-1}} \rho(K, u)^{n-i} dS(u), \quad i \in \mathbb{R}.$$

If K and L are star bodies in $\mathbb{R}^n$, $s \neq 0$ and $\lambda, \mu \geq 0$, then $\lambda \cdot K \tilde{+}_s \mu \cdot L$ is the star body whose radial function is given by

$$(1.7) \quad \rho(\lambda \cdot K \tilde{+}_s \mu \cdot L, \cdot)^s = \lambda \rho(K, \cdot)^s + \mu \rho(L, \cdot)^s.$$

The addition $\tilde{+}_s$ is called the L_s radial sum.

The L_s dual Brunn-Minkowski inequality states: If $K, L \in \mathcal{S}^n$, then

$$V(K \tilde{+}_s L)^{-s/n} \geq V(K)^{-s/n} + V(L)^{-s/n},$$

with equality if and only if K and L are dilates.

2. RADIAL BLASCHKE-MINKOWSKI HOMOMORPHISMS

Definition 2.1 ([16]). A map $\Psi : \mathcal{S}^n \rightarrow \mathcal{S}^n$ is called a radial Blaschke-Minkowski homomorphism if it satisfies the following conditions:

- (a) Ψ is continuous.
- (b) For all $K, L \in \mathcal{S}^n$,

$$\Psi(K \hat{+} L) = \Psi(K) \tilde{+} \Psi(L),$$

where $\hat{+}$ denotes the L_{n-1} radial sum of K and L .

- (c) For all $K, L \in \mathcal{S}^n$ and every $\vartheta \in SO(n)$,

$$\Psi(\vartheta K) = \vartheta \Psi(K),$$

where $SO(n)$ is the group of rotations in n dimensions.

In 2006, Schuster [16] established the following Brunn-Minkowski inequality for radial Blaschke-Minkowski homomorphisms of star bodies.

If K and L are star bodies in $\mathbb{R}^n$, then

$$(2.1) \quad V(\Psi(K \tilde{+} L))^{1/n(n-1)} \leq V(\Psi K)^{1/n(n-1)} + V(\Psi L)^{1/n(n-1)},$$

with equality if and only if K and L are dilates.

In fact a more general version of the Brunn-Minkowski inequality for radial Blaschke-Minkowski homomorphisms holds (see [16]): If K and L are star bodies in $\mathbb{R}^n$ and $0 \leq i \leq n-1$, $0 \leq j < n-2$, then

$$(2.2) \quad \tilde{W}_i(\Psi_j(K \tilde{+} L))^{1/(n-i)(n-j-1)} \leq \tilde{W}_i(\Psi_j K)^{1/(n-i)(n-j-1)} + \tilde{W}_i(\Psi_j L)^{1/(n-i)(n-j-1)},$$

with equality if and only if K and L are dilates. Here Ψ_j denotes the mixed radial Blaschke-Minkowski homomorphism defined by:

Theorem 2.2 (see [16]). Let $\Psi : \mathcal{S}^n \rightarrow \mathcal{S}^n$ be a radial Blaschke-Minkowski homomorphism. There is a continuous operator $\Psi : \underbrace{\mathcal{S}^n \times \cdots \times \mathcal{S}^n}_{n-1} \rightarrow \mathcal{S}^n$, symmetric in

its arguments such that, for $K_1, \dots, K_m \in \mathcal{S}^n$ and $\lambda_1, \dots, \lambda_m \leq 0$,

$$\Psi(\lambda_1 K_1 \tilde{+} \cdots \tilde{+} \lambda_m K_m) = \sum_{i_1, \dots, i_{n-1}} \lambda_{i_1} \cdots \lambda_{i_{n-1}} \Psi(K_{i_1}, \dots, K_{i_{n-1}}).$$

Clearly, Theorem 2.2 generalizes the notion of radial Blaschke-Minkowski homomorphisms. We call $\Psi : \mathcal{S}^n \times \cdots \times \mathcal{S}^n \rightarrow \mathcal{S}^n$ the mixed radial Blaschke-Minkowski homomorphism induced by Ψ . The mixed radial Blaschke-Minkowski homomorphisms were first studied in more detail in [17]–[18]. If $K_1 = \cdots = K_{n-i-1} = K$, $K_{n-i} = \cdots = K_{n-1} = B$, we write $\Psi_i K$ for $\Psi(\underbrace{K, \dots, K}_{n-i-1}, \underbrace{B, \dots, B}_i)$ and call

$\Psi_i K$ the mixed Blaschke-Minkowski homomorphism of order i of K . We write $\Psi_i(K, L)$ for $\Psi(\underbrace{K, \dots, K}_{n-i-1}, \underbrace{L, \dots, L}_i)$.

The aim of this paper is to establish the following new Brunn-Minkowski inequality for mixed radial Blaschke-Minkowski homomorphisms.

Theorem 2.3. *If $K, L \in \mathcal{S}^n$ and $i, j \in \mathbb{R}, s \in \mathbb{N}$ satisfy $i \leq n-1 \leq j \leq n, 0 \leq s \leq n-1$, then*

$$(2.3) \quad \left(\frac{\tilde{W}_i(\Psi_s(K \dot{+}_s L))}{\tilde{W}_j(\Psi_s(K \dot{+}_s L))} \right)^{1/(j-i)} \leq \left(\frac{\tilde{W}_i(\Psi_s K)}{\tilde{W}_j(\Psi_s K)} \right)^{1/(j-i)} + \left(\frac{\tilde{W}_i(\Psi_s L)}{\tilde{W}_j(\Psi_s L)} \right)^{1/(j-i)},$$

with equality if and only if K and L are dilates.

3. POLAR BLASCHKE-MINKOWSKI HOMOMORPHISMS

Definition 3.1 (see [16]). A map $\Phi : \mathcal{K}^n \rightarrow \mathcal{K}^n$ is called a Blaschke-Minkowski homomorphism if it satisfies the following conditions:

- (a) Φ is continuous.
- (b) For all $K, L \in \mathcal{K}^n$,

$$\Phi(K \dot{+} L) = \Phi(K) + \Phi(L).$$

- (c) For all $K, L \in \mathcal{K}^n$ and every $\vartheta \in SO(n)$,

$$\Phi(\vartheta K) = \vartheta \Phi(K).$$

In [16] it was shown that the polar body $(\Phi K)^*$ is well defined for every Blaschke-Minkowski homomorphism Φ and $K \in \mathcal{K}^n$. In the following we simply write $\Phi^* K$ rather than $(\Phi K)^*$.

In 2006, Schuster [16] also established the following Brunn-Minkowski inequality for polars of *even* Blaschke-Minkowski homomorphisms Φ of convex bodies.

If K and L are convex bodies in $\mathbb{R}^n$, then

$$(3.1) \quad V(\Phi^*(K + L))^{-1/n(n-1)} \geq V(\Phi^* K)^{1/(n(n-1))} + V(\Phi^* L)^{1/n(n-1)},$$

with equality if and only if K and L are homothetic.

In fact a more general version of the Brunn-Minkowski inequality for polars of even Blaschke-Minkowski homomorphisms holds (see [16]): If K and L are convex bodies in $\mathbb{R}^n$ and $0 \leq j \leq n-3$, then

$$(3.2) \quad V(\Phi_j^*(K + L))^{-1/n(n-j-1)} \geq V(\Phi_j^* K)^{1/(n(n-j-1))} + V(\Phi_j^* L)^{1/n(n-j-1)},$$

with equality if and only if K and L are homothetic. Here, $\Phi_j K$ denotes the mixed Blaschke-Minkowski homomorphism induced by Φ defined by:

Theorem 3.2 (see [16]). *There is a continuous operator $\Phi : \underbrace{\mathcal{K}^n \times \cdots \times \mathcal{K}^n}_{n-1}$, symmetric in its arguments such that, for $K_1, \dots, K_r$ and $\lambda_1, \dots, \lambda_r \geq 0$,*

$$\Phi(\lambda_1 K_1 + \cdots + \lambda_r K_r) = \sum_{i_1, \dots, i_{n-1}} \lambda_{i_1} \cdots \lambda_{i_{n-1}} \Phi(K_{i_1}, \dots, K_{i_{n-1}}).$$

Clearly, Theorem 3.2 generalizes the notion of Blaschke-Minkowski homomorphisms. We call $\Phi : \mathcal{K}^n \times \cdots \times \mathcal{K}^n \rightarrow \mathcal{K}^n$ the mixed Blaschke-Minkowski homomorphism induced by Φ . Mixed Blaschke-Minkowski homomorphisms were first studied in more detail in [18]. If $K_1 = \cdots = K_{n-i-1} = K, K_{n-i} = \cdots = K_{n-1} = B$, we write $\Phi_i K$ for $\Phi(\underbrace{K, \dots, K}_{n-i-1}, \underbrace{B, \dots, B}_i)$ and call $\Phi_i K$ the mixed Blaschke-Minkowski

homomorphism of order i . We write $\Phi_i(K, L)$ for $\Phi(\underbrace{K, \dots, K}_{n-i-1}, \underbrace{L, \dots, L}_i)$ and write $\Phi_0 K$ as ΦK .

Blaschke-Minkowski homomorphisms are an important notion in the theory of convex body-valued valuations (see, e.g., [5]–[6], [9]–[10], [15], [19], [22] and [1]–[2], [7]–[8], [11]–[12], [20]–[21]). They are natural duals to radial Blaschke-Minkowski homomorphisms which are important examples of star body-valued valuations.

Another aim of this paper is to establish the following Brunn-Minkowski inequality for polars of even Blaschke-Minkowski homomorphisms.

Theorem 3.3. *Let K, L be convex bodies in $\mathbb{R}^n$ and $i, j \in \mathbb{R}$ satisfy $i \geq n + 1 \geq j \geq n$. Then*

$$(3.3) \quad \left(\frac{\tilde{W}_i(\Phi^*(K \ddot{+} L))}{\tilde{W}_j(\Phi^*(K \ddot{+} L))} \right)^{1/(i-j)} \leq \left(\frac{\tilde{W}_i(\Phi^*K)}{\tilde{W}_j(\Phi^*K)} \right)^{1/(i-j)} + \left(\frac{\tilde{W}_i(\Phi^*L)}{\tilde{W}_j(\Phi^*L)} \right)^{1/(i-j)},$$

with equality if and only if K and L are homothetic.

4. BRUNN-MINKOWSKI TYPE INEQUALITIES FOR RADIAL AND POLAR BLASCHKE-MINKOWSKI HOMOMORPHISMS

An extension of Beckenbach's inequality (see [3], p. 27) was obtained by Dresher [4] by means of moment-space techniques:

Lemma 4.1 (The Beckenbach-Dresher inequality). *If $p \geq 1 \geq r \geq 0$, $f, g \geq 0$, and ϕ is a distribution function, then*

$$(4.1) \quad \left(\frac{\int (f+g)^p d\phi}{\int (f+g)^r d\phi} \right)^{1/(p-r)} \leq \left(\frac{\int f^p d\phi}{\int f^r d\phi} \right)^{1/(p-r)} + \left(\frac{\int g^p d\phi}{\int g^r d\phi} \right)^{1/(p-r)},$$

with equality if and only if the functions f and g are proportional.

Lemma 4.2 (see [16]). *If $\Phi : \mathcal{K}^n \rightarrow \mathcal{K}^n$ is a Blaschke-Minkowski homomorphism, then there is a function $g \in \mathcal{C}(S^{n-1}, \hat{e})$ such that*

$$(4.2) \quad h(\Phi K, \cdot) = S_{n-1}(K, \cdot)^{n-1} * g,$$

where $\mathcal{C}(S^{n-1}, \hat{e})$ denotes the set of continuous zonal functions on S^{n-1} .

As a consequence of Lemma 4.2, we have for the mixed Blaschke-Minkowski homomorphism induced by Φ ,

$$(4.3) \quad h(\Phi(K_1, \dots, K_{n-1}), \cdot) = S(K_1, \dots, K_{n-1}; \cdot) * g,$$

where $S(K_1, \dots, K_{n-1}; \cdot)$ is the mixed surface area measure of $K_1, \dots, K_{n-1}$.

Let $\Phi : \underbrace{\mathcal{K}^n \times \dots \times \mathcal{K}^n}_{n-1} \rightarrow \mathcal{K}^n$ be a mixed Blaschke-Minkowski homomorphism.

If $K_1, \dots, K_{n-1} \in \mathcal{K}^n$, then (see [16])

$$(4.4) \quad \rho(\Phi^*(K_1, \dots, K_{n-1}), \cdot)^{-1} = h(\Phi^*(K_1, \dots, K_{n-1}), \cdot).$$

Lemma 4.3 (see [16]). *A map $\Psi : \mathcal{S}^n \rightarrow \mathcal{S}^n$ is a radial Blaschke-Minkowski homomorphism if and only if there is a measure $\mu \in \mathcal{M}_+(S^{n-1}, \hat{e})$ such that*

$$(4.5) \quad \rho(\Psi K, \cdot) = \rho(K, \cdot)^{n-1} * \mu,$$

where $\mathcal{M}_+(S^{n-1}, \hat{e})$ denotes the set of nonnegative zonal measures on S^{n-1} .

For the mixed radial Blaschke-Minkowski homomorphism induced by Ψ , we have

$$(4.6) \quad \rho(\Psi(K_1, \dots, K_{n-1}), \cdot) = \rho(K_1, \cdot) \cdots \rho(K_{n-1}, \cdot) * \mu.$$

We are now in a position to prove Theorem 2.3. The following statement is just a slight reformulation of Theorem 2.3.

Theorem 4.4. *If $K, L \in \mathcal{S}^n$ and $p, r \in \mathbb{R}, j \in \mathbb{N}$ satisfy $0 \leq r \leq 1 \leq p$, $0 \leq j \leq n-1$, then*

$$(4.7) \quad \left(\frac{\tilde{W}_{n-p}(\Psi_j(K \tilde{+}_j L))}{\tilde{W}_{n-r}(\Psi_j(K \tilde{+}_j L))} \right)^{1/(p-r)} \leq \left(\frac{\tilde{W}_{n-p}(\Psi_j K)}{\tilde{W}_{n-r}(\Psi_j K)} \right)^{1/(p-r)} + \left(\frac{\tilde{W}_{n-p}(\Psi_j L)}{\tilde{W}_{n-r}(\Psi_j L)} \right)^{1/(p-r)},$$

with equality if and only if K and L are dilates.

Proof. From (1.7), we have

$$\rho(K \tilde{+}_s L, \cdot)^s * \mu = \rho(K, \cdot)^s * \mu + \rho(L, \cdot)^s * \mu, \quad s \neq 0,$$

where μ is defined in Lemma 4.3. Hence, from (1.2) and (4.6), we obtain

$$\rho(\Psi_s(K \tilde{+}_s L), \cdot) = \rho(\Psi_s K, \cdot) + \rho(\Psi_s L, \cdot) = \rho(\Psi_s K \tilde{+} \Psi_s L, \cdot).$$

Namely,

$$\Psi_s(K \tilde{+}_s L) = \Psi_s(K) \tilde{+} \Psi_s(L).$$

Therefore, from (1.2) and (1.6), we have

$$(4.8) \quad \begin{aligned} \tilde{W}_{n-p}(\Psi_j(K \tilde{+}_j L)) &= \frac{1}{n} \int_{S^{n-1}} \rho(\Psi_j(K \tilde{+}_j L), u)^p dS(u) \\ &= \frac{1}{n} \int_{S^{n-1}} \rho(\Psi_j K \tilde{+} \Psi_j L, u)^p dS(u) = \frac{1}{n} \int_{S^{n-1}} (\rho(\Psi_j K, u) + \rho(\Psi_j L, u))^p dS(u) \end{aligned}$$

and

$$(4.9) \quad \tilde{W}_{n-r}(\Psi_j(K \tilde{+}_j L)) = \frac{1}{n} \int_{S^{n-1}} (\rho(\Psi_j K, u) + \rho(\Psi_j L, u))^r dS(u).$$

From (4.8) and (4.9) and in view of Lemma 4.1, we obtain

$$\begin{aligned} \left(\frac{\tilde{W}_{n-p}(\Psi_j(K \tilde{+}_j L))}{\tilde{W}_{n-r}(\Psi_j(K \tilde{+}_j L))} \right)^{\frac{1}{p-r}} &= \left(\frac{\int_{S^{n-1}} (\rho(\Psi_j K, u) + \rho(\Psi_j L, u))^p dS(u)}{\int_{S^{n-1}} (\rho(\Psi_j K, u) + \rho(\Psi_j L, u))^r dS(u)} \right)^{\frac{1}{p-r}} \\ &\leq \left(\frac{\int_{S^{n-1}} \rho(\Psi_j K, u)^p dS(u)}{\int_{S^{n-1}} \rho(\Psi_j K, u)^r dS(u)} \right)^{\frac{1}{p-r}} + \left(\frac{\int_{S^{n-1}} \rho(\Psi_j L, u)^p dS(u)}{\int_{S^{n-1}} \rho(\Psi_j L, u)^r dS(u)} \right)^{\frac{1}{p-r}} \\ &= \left(\frac{\tilde{W}_{n-p}(\Psi_j K)}{\tilde{W}_{n-r}(\Psi_j K)} \right)^{\frac{1}{p-r}} + \left(\frac{\tilde{W}_{n-p}(\Psi_j L)}{\tilde{W}_{n-r}(\Psi_j L)} \right)^{\frac{1}{p-r}}. \end{aligned}$$

Equality holds if and only if the functions $\rho(\Psi_j K, u)$ and $\rho(\Psi_j L, u)$ are proportional. Namely, $\rho(\Psi_j K, u) = \lambda \rho(\Psi_j L, u)$. From (4.5), we obtain $\rho(\Psi_j K, u) = \rho(\Psi_j(\lambda^{1/(n-j-1)} L), u)$. Hence, equality holds if and only if K and L are dilates.

Let $p = n - i$ and $r = n - j$. Since $0 \leq r \leq 1 \leq p$, we have

$$r \leq 1 \leq p \Rightarrow i \leq n - 1 \leq j, \quad 0 \leq r \Rightarrow j \leq n.$$

Therefore,

$$(4.10) \quad i \leq n-1 \leq j \leq n.$$

Taking for $p = n-i$ and $r = n-j$ in (4.7) and using (4.10), we see that (4.7) changes to the inequality in Theorem 2.3.

Taking for $p = n-i, j = 0$ and $r = 1$ in (4.7), (4.7) changes to the following inequality:

$$(4.11) \quad \left(\frac{\tilde{W}_i(\Psi(K \tilde{+} L))}{\tilde{W}_{n-1}(\Psi(K \tilde{+} L))} \right)^{1/(n-i-1)} \leq \left(\frac{\tilde{W}_i(\Psi K)}{\tilde{W}_{n-1}(\Psi K)} \right)^{1/(n-i-1)} + \left(\frac{\tilde{W}_i(\Psi L)}{\tilde{W}_{n-1}(\Psi L)} \right)^{1/(n-i-1)},$$

with equality if and only if K and L are dilates.

For $K \in \mathcal{S}^n$, there is a unique star body $\mathbf{I}K$ whose radial function satisfies for $u \in S^{n-1}$,

$$\rho(\mathbf{I}K, u) = v(K \cap E_u).$$

It is called the *intersection bodies* of K . The volume of intersection bodies is given by

$$V(\mathbf{I}K) = \frac{1}{n} \int_{S^{n-1}} v(K \cap E_u)^n dS(u).$$

The mixed intersection body of $K_1, \dots, K_{n-1} \in \mathcal{S}^n$, $\mathbf{I}(K_1, \dots, K_{n-1})$, is defined by

$$\rho(\mathbf{I}(K_1, \dots, K_{n-1}), u) = \tilde{v}(K_1 \cap E_u, \dots, K_{n-1} \cap E_u),$$

where $\tilde{v}$ is $(n-1)$ -dimensional dual mixed volume.

If $K_1 = \dots = K_{n-i-1} = K, K_{n-i} = \dots = K_{n-1} = L$, then $\mathbf{I}(K_1, \dots, K_{n-1})$ is written as $\mathbf{I}_i(K, L)$. If $L = B$, then $\mathbf{I}_i(K, L)$ is written as $\mathbf{I}_i K$ and called the i th intersection body of K . For $\mathbf{I}_0 K$ we simply write $\mathbf{I}K$. $\square$

If $\Psi : \underbrace{\mathcal{S}^n \times \dots \times \mathcal{S}^n}_{n-1} \rightarrow \mathcal{S}^n$ is the mixed intersection operator $\mathbf{I} : \underbrace{\mathcal{S}^n \times \dots \times \mathcal{S}^n}_{n-1} \rightarrow \mathcal{S}^n$ in (4.7), we obtain

Corollary 4.5. *If $K, L \in \mathcal{S}^n$ and $p, r \in \mathbb{R}, j \in \mathbb{N}$ satisfy $0 \leq r \leq 1 \leq p, 0 \leq j \leq n-1$, then*

$$(4.12) \quad \left(\frac{\tilde{W}_{n-p}(\mathbf{I}_j(K \tilde{+} L))}{\tilde{W}_{n-r}(\mathbf{I}_j(K \tilde{+} L))} \right)^{1/(p-r)} \leq \left(\frac{\tilde{W}_{n-p}(\mathbf{I}_j K)}{\tilde{W}_{n-r}(\mathbf{I}_j K)} \right)^{1/(p-r)} + \left(\frac{\tilde{W}_{n-p}(\mathbf{I}_j L)}{\tilde{W}_{n-r}(\mathbf{I}_j L)} \right)^{1/(p-r)},$$

with equality if and only if K and L are dilates.

The following statement is just a slight reformulation of Theorem 3.3.

Theorem 4.6. *If $K, L \in \mathcal{K}^n$ and $p, r \in \mathbb{R}$ satisfy $p \leq -1 \leq r \leq 0$, then*

$$(4.13) \quad \left(\frac{\tilde{W}_{n-p}(\Phi^*(K \ddot{+} L))}{\tilde{W}_{n-r}(\Phi^*(K \ddot{+} L))} \right)^{1/(r-p)} \leq \left(\frac{\tilde{W}_{n-p}(\Phi^* K)}{\tilde{W}_{n-r}(\Phi^* K)} \right)^{1/(r-p)} + \left(\frac{\tilde{W}_{n-p}(\Phi^* L)}{\tilde{W}_{n-r}(\Phi^* L)} \right)^{1/(r-p)},$$

with equality if and only if K and L are homothetic.

Proof. From (1.6), (4.4) and in view of Definition 3.1, we have

$$\begin{aligned}
 \tilde{W}_{n-p}(\Phi^*(K \ddot{+} L)) &= \frac{1}{n} \int_{S^{n-1}} \rho(\Phi^*(K \ddot{+} L), u)^p dS(u) \\
 &= \frac{1}{n} \int_{S^{n-1}} h(\Phi(K \ddot{+} L), u)^{-p} dS(u) = \frac{1}{n} \int_{S^{n-1}} h(\Phi K + \Phi L, u)^{-p} dS(u) \\
 (4.14) \quad &= \frac{1}{n} \int_{S^{n-1}} (h(\Phi K, u) + h(\Phi L, u))^{-p} dS(u)
 \end{aligned}$$

and

$$(4.15) \quad \tilde{W}_{n-r}(\Phi^*(K \ddot{+} L)) = \frac{1}{n} \int_{S^{n-1}} (h(\Phi K, u) + h(\Phi L, u))^{-r} dS(u).$$

From (4.14), (4.15), and Lemma 4.1, we obtain

$$\begin{aligned}
 \left(\frac{\tilde{W}_{n-p}(\Phi^*(K \ddot{+} L))}{\tilde{W}_{n-r}(\Phi^*(K \ddot{+} L))} \right)^{1/(r-p)} &= \left(\frac{\int_{S^{n-1}} (h(\Phi K, u) + h(\Phi L, u))^{-p} dS(u)}{\int_{S^{n-1}} (h(\Phi K, u) + h(\Phi L, u))^{-r} dS(u)} \right)^{1/(r-p)} \\
 &\leq \left(\frac{\int_{S^{n-1}} h(\Phi K, u)^{-p} dS(u)}{\int_{S^{n-1}} h(\Phi K, u)^{-r} dS(u)} \right)^{1/(r-p)} + \left(\frac{\int_{S^{n-1}} h(\Phi L, u)^{-p} dS(u)}{\int_{S^{n-1}} h(\Phi L, u)^{-r} dS(u)} \right)^{1/(r-p)} \\
 &= \left(\frac{\int_{S^{n-1}} \rho(\Phi^* K, u)^p dS(u)}{\int_{S^{n-1}} \rho(\Phi^* K, u)^r dS(u)} \right)^{1/(r-p)} + \left(\frac{\int_{S^{n-1}} \rho(\Phi^* L, u)^p dS(u)}{\int_{S^{n-1}} \rho(\Phi^* L, u)^r dS(u)} \right)^{1/(r-p)} \\
 &= \left(\frac{\tilde{W}_{n-p}(\Phi^* K)}{\tilde{W}_{n-r}(\Phi^* K)} \right)^{1/(r-p)} + \left(\frac{\tilde{W}_{n-p}(\Phi^* L)}{\tilde{W}_{n-r}(\Phi^* L)} \right)^{1/(r-p)}.
 \end{aligned}$$

Equality holds if and only if the functions $h(\Phi^* K, u)$ and $h(\Phi^* L, u)$ are proportional, namely, $h(\Phi^* K, u) = \lambda h(\Phi^* L, u)$, and from (4.3), we obtain $h(\Phi^* K, u) = \lambda h(\Phi^*(\lambda^{-1/(n-1)} L), u)$. Hence, equality holds if and only if K and L are homothetic.

Let $p = n - i$ and $r = n - j$. Since $p \leq -1 \leq r \leq 0$, we have

$$(4.16) \quad p \leq -1 \leq r \leq 0 \Rightarrow i \geq n + 1 \geq j \geq n.$$

Taking for $p = n - i$ and $r = n - j$ in (4.13) and using (4.16), we see that (4.13) changes to the inequality in Theorem 3.3.

Taking $p = -n, r = -1$ and $s = 0$ in (4.13), we have

$$(4.17) \quad \left(\frac{\tilde{W}_{2n}(\Phi^*(K \ddot{+} L))}{\tilde{W}_{n+1}(\Phi^*(K \ddot{+} L))} \right)^{1/(n-1)} \leq \left(\frac{\tilde{W}_{2n}(\Phi^* K)}{\tilde{W}_{n+1}(\Phi^* K)} \right)^{1/(n-1)} + \left(\frac{\tilde{W}_{2n}(\Phi^* L)}{\tilde{W}_{n+1}(\Phi^* L)} \right)^{1/(n-1)},$$

with equality if and only if K and L are homothetic.

If $K_1, \dots, K_r \in \mathcal{K}^n$ and $\lambda_1, \dots, \lambda_r \geq 0$, then the projection body of the Minkowski linear combination $\lambda_1 K_1 + \dots + \lambda_r K_r \in \mathcal{K}^n$ can be written as a symmetric homogeneous polynomial of degree $(n - 1)$ in the λ_i (see [15]):

$$(4.18) \quad \Pi(\lambda_1 K_1 + \dots + \lambda_r K_r) = \sum \lambda_{i_1} \dots \lambda_{i_{n-1}} \Pi_{i_1 \dots i_{n-1}},$$

where the sum is a Minkowski sum taken over all $(n - 1)$ -tuples $(i_1, \dots, i_{n-1})$ of positive integers not exceeding r . The body $\Pi_{i_1 \dots i_{n-1}}$ depends only on the bodies $K_{i_1}, \dots, K_{i_{n-1}}$ and is uniquely determined by (4.18). It is called *the mixed projection bodies* of $K_{i_1}, \dots, K_{i_{n-1}}$, and is written as $\Pi(K_{i_1}, \dots, K_{i_{n-1}})$.

If $K_1 = \cdots = K_{n-1-i} = K$ and $K_{n-i} = \cdots = K_{n-1} = L$, then $\Pi(K_{i_1}, \dots, K_{i_{n-1}})$ will be written as $\Pi_i(K, L)$. If $L = B$, then $\Pi_i(K, L)$ is denoted by $\Pi_i K$ and when $i = 0$, $\Pi_i K$ is denoted by ΠK , where ΠK is the projection body of K . $\square$

If $\Phi : \underbrace{\mathcal{K}^n \times \cdots \times \mathcal{K}^n}_{n-1} \rightarrow \mathcal{K}^n$ is the mixed projection operator $\Pi : \underbrace{\mathcal{K}^n \times \cdots \times \mathcal{K}^n}_{n-1} \rightarrow \mathcal{K}^n$ in (4.13), we obtain

Corollary 4.7. *If $K, L \in \mathcal{K}^n$ and $p, r \in \mathbb{R}$ satisfy $p \leq -1 \leq r \leq 0$, then*

$$(4.19) \quad \left(\frac{\tilde{W}_{n-p}(\Pi^*(K \ddot{+} L))}{\tilde{W}_{n-r}(\Pi^*(K \ddot{+} L))} \right)^{1/(r-p)} \leq \left(\frac{\tilde{W}_{n-p}(\Pi^* K)}{\tilde{W}_{n-r}(\Pi^* K)} \right)^{1/(r-p)} + \left(\frac{\tilde{W}_{n-p}(\Pi^* L)}{\tilde{W}_{n-r}(\Pi^* L)} \right)^{1/(r-p)},$$

with equality if and only if K and L are homothetic.

We finally remark that inequalities for the intersection operator $\mathbf{I}$ were also established in [23], [27], for the L_p -intersection operator $\mathbf{I}_p$ in [24] and for the polar projection body operator Π^* in [25]–[26].

ACKNOWLEDGEMENTS

The author is very grateful to the referee for many very valuable suggestions and comments. The author also express his grateful thanks to Prof. Gang-song Leng for his help.

REFERENCES

- [1] J. Abardia, A. Bernig, Projection bodies in complex vector spaces, *Adv. Math.*, **227**(2011), 830–846.
- [2] S. Alesker, A. Bernig, F.E. Schuster, Harmonic analysis of translation invariant valuations, *Geom. Funct. Anal.*, **21** (2011), 751–773. MR2827009
- [3] E. F. Beckenbach, R. Bellman, Inequalities, Springer-Verlag, Berlin-Göttingen, Heidelberg, 1961. MR0158038 (28:1266)
- [4] M. Dresher, Moment space and inequalities, *Duke. Math. J.*, **20**(1953), 261–271. MR0055389 (14:1067d)
- [5] C. Haberl, Star body valued valuations, *Indiana Univ. Math. J.*, **58**(2009), 2253–2276. MR2583498 (2011b:52018)
- [6] C. Haberl, F.E. Schuster, General L_p affine isoperimetric inequalities, *J. Diff. Geom.*, **83**(2009), 1–26. MR2545028 (2010j:52026)
- [7] C. Haberl, L_p intersection bodies, *Adv. Math.*, **217**(2008), 2599–2624. MR2397461 (2009a:52001)
- [8] C. Haberl, M. Ludwig, A characterization of L_p intersection bodies, *Int. Math. Res. Not.*, Art ID 10548, 29 pages. MR2250020 (2007k:52007)
- [9] M. Kiderlen, Blaschke- and Minkowski-endomorphisms of convex bodies, *Trans. Amer. Math. Soc.*, **358**(2006), 5539–5564. MR2238926 (2007e:52006)
- [10] M. Ludwig, Minkowski valuations, *Trans. Amer. Math. Soc.*, **357**(10)(2005), 4191–4213. MR2159706 (2006f:52005)
- [11] M. Ludwig, General affine surface areas, *Adv. Math.*, **224**(2010), 2346–2360. MR2652209 (2011h:52006)
- [12] M. Ludwig, Minkowski areas and valuations, *J. Diff. Geom.*, **86**(2010), 133–161. MR2772547
- [13] E. Lutwak, Intersection bodies and dual mixed volumes, *Adv. Math.*, **71**(1988), 232–261. MR963487 (90a:52023)
- [14] E. Lutwak, Dual mixed volumes, *Pacific J. Math.*, **58**(1975), 531–538. MR0380631 (52:1528)
- [15] E. Lutwak, Mixed projection inequalities, *Trans. Amer. Math. Soc.*, **287**(1)(1985), 91–105. MR766208 (86c:52015)
- [16] F. E. Schuster, Volume inequalities and additive maps of convex bodies, *Mathematika*, **53**(2006), 211–234. MR2343256 (2008i:52011)

- [17] F. E. Schuster, Valuations and Busemann-Petty type problems, *Adv. Math.*, **219**(2008), 344-368. MR2435426 (2009f:52018)
- [18] F. E. Schuster, Convolutions and multiplier transformations of convex bodies, *Trans. Amer. Math. Soc.*, **359**(2007), 5567-5591. MR2327043 (2009e:52008)
- [19] F. E. Schuster, Crofton measures and Minkowski valuations, *Duke Math. J.*, **154**(2010), 1-30. MR2668553 (2011g:52026)
- [20] F. E. Schuster, T. Wannerer, $GL(n)$ contravariant Minkowski valuations, *Trans. Amer. Math. Soc.*, **364**(2012), 815-826. MR2846354
- [21] T. Wannerer, $GL(n)$ equivariant Minkowski valuations, *Indiana Univ. Math. J.*, in press. <http://www.iuj.indiana.edu/IUMJ/forthcoming.php>
- [22] C. Zhao, On Blaschke-Minkowski homomorphisms, *Geom. Dedicata*, **149**(2010), 373-378. MR2737698
- [23] C. Zhao, On intersection and mixed intersection bodies, *Geom. Dedicata*, **141**(2009), 109-122. MR2520066 (2010k:52010)
- [24] C. Zhao, L_p -mixed intersection bodies, *Sci. China*, **51**(2008), 2172-2188. MR2462021 (2009k:52023)
- [25] C. Zhao, L_p -dual Quermassintegrals sums, *Sci. China*, **50**(2007), 1347-1360. MR2370622 (2009e:52019)
- [26] C. Zhao, G. Leng, On polars of mixed projection bodies, *J. Math. Anal. Appl.*, **316**(2006), 664-678. MR2207338 (2006m:52008)
- [27] C. Zhao, G. Leng, Brunn-Minkowski inequality for mixed intersection bodies, *J. Math. Anal. Appl.*, **301**(2005), 115-123. MR2105924 (2005i:52016)

DEPARTMENT OF MATHEMATICS, CHINA JILIANG UNIVERSITY, HANGZHOU 310018, PEOPLE'S REPUBLIC OF CHINA

E-mail address: chjzhao@yahoo.com.cn

E-mail address: chjzhao@163.com