CRITERIA FOR TOEPLITZ OPERATORS ON THE SPHERE

JINGBO XIA

(Communicated by Marius Junge)

ABSTRACT. Let $H^2(S)$ be the Hardy space on the unit sphere S in $\mathbf{C}^n$. We show that a set of inner functions Λ is sufficient for the purpose of determining which $A \in \mathcal{B}(H^2(S))$ is a Toeplitz operator if and only if the multiplication operators $\{M_u : u \in \Lambda\}$ on $L^2(S, d\sigma)$ generate the von Neumann algebra $\{M_f : f \in L^\infty(S, d\sigma)\}$.

1. INTRODUCTION

Throughout the paper, we denote the unit sphere $\{z \in \mathbf{C}^n : |z| = 1\}$ in $\mathbf{C}^n$ by S . Let σ be the positive, regular Borel measure on S which is invariant under the orthogonal group $O(2n)$, i.e., the group of isometries on $\mathbf{C}^n \cong \mathbf{R}^{2n}$ which fix 0. Furthermore we normalize σ such that $\sigma(S) = 1$. Recall that the Hardy space $H^2(S)$ is the closure in $L^2(S, d\sigma)$ of polynomials in the coordinate variables $z_1, \dots, z_n$. Let

$$P : L^2(S, d\sigma) \rightarrow H^2(S)$$

be the orthogonal projection. For each $f \in L^\infty(S, d\sigma)$, we have the Toeplitz operator T_f on $H^2(S)$ defined by the formula

$$T_f = PM_f|_{H^2(S)}.$$

That is, T_f is the compression of the multiplication operator M_f to the subspace $H^2(S)$.

Toeplitz operators on various reproducing-kernel Hilbert spaces have been extensively studied in the literature. This paper concerns one of the most elementary questions in the theory, namely, how does one characterize a Toeplitz operator on $H^2(S)$?

In the case where the complex dimension n equals 1, i.e., in the unit circle case, there is a very simple answer due to Brown and Halmos. In [2], Brown and Halmos showed that if A is a bounded operator on the Hardy space H^2 of the unit circle $\mathbf{T} = \{z \in \mathbf{C} : |z| = 1\}$, then A is a Toeplitz operator if and only if it satisfies the equation

$$(1.1) \quad T_{\bar{z}}AT_z = A.$$

This criterion for Toeplitz operators was later generalized to an arbitrary complex dimension n . In [3], Davie and Jewell showed that, for whatever n , a bounded

Received by the editors December 18, 2010 and, in revised form, July 10, 2011.
 2010 *Mathematics Subject Classification*. Primary 46L10, 47B35.

operator A on $H^2(S)$ is a Toeplitz operator if and only if it satisfies the equation

$$(1.2) \quad \sum_{j=1}^n T_{\bar{z}_j} A T_{z_j} = A.$$

By any reasonable standard, (1.2) is a satisfactory generalization of (1.1) to the high-dimensional case. But the fact that (1.2) is a satisfactory generalization of (1.1) also causes one to neglect the other side of the story, namely, when $n \geq 2$, (1.2) is really a *different* kind of test for Toeplitz operators. The substantive difference between (1.1) and (1.2) is the simple fact that when $n = 1$, z is an inner function on the unit disc; in contrast, if $n \geq 2$, then the functions $z_1, \dots, z_n$ are far from being inner.

Let B denote the open unit ball $\{z \in \mathbf{C}^n : |z| < 1\}$ in $\mathbf{C}^n$. Recall that an analytic function u on B is said to be *inner* if

$$\lim_{r \uparrow 1} |u(r\zeta)| = 1 \quad \text{for } \sigma\text{-a.e. } \zeta \in S.$$

As usual, we identify the function u on B with its boundary value on S . In this paper, we consider the problem of characterizing Toeplitz operators in terms of inner functions, which is not addressed by (1.2) in the case $n \geq 2$. When [3] was published in 1977, it was not yet known that there are non-constant inner functions on B in the case $n \geq 2$. Therefore (1.2) was the best that could be managed in terms of characterizing Toeplitz operators at the time. Later, Aleksandrov [1] and Løw [6] showed that there are non-constant inner functions on B for every $n \geq 2$. This makes it possible to consider the problem of characterizing Toeplitz operators in terms of inner functions.

Note that if u is an inner function, then for every $f \in L^\infty(S, d\sigma)$ we have

$$T_{\bar{u}} T_f T_u = T_{\bar{u}fu} = T_f.$$

Thus, in order for an operator A on $H^2(S)$ to be a Toeplitz operator, it is necessary that

$$T_{\bar{u}} A T_u = A \quad \text{for every inner function } u \text{ on } B.$$

Our question is the following. Suppose that Λ is a non-empty set of inner functions on B . If A is a bounded operator on $H^2(S)$ and if it satisfies the condition

$$T_{\bar{u}} A T_u = A \quad \text{for every } u \in \Lambda,$$

can we conclude that A is a Toeplitz operator on $H^2(S)$? Note that even in the case $n = 1$, this question goes beyond the Brown-Halmos criterion for Toeplitz operators.

We will characterize those sets Λ which yield the answer “yes” to the above question. Interestingly, this characterization involves von Neumann algebras and a rare use of the double-commutant relation of Murray and von Neumann. We will then deal with specific Λ ’s which yield the answer “yes” and specific Λ ’s which yield the answer “no” to the above question.

To conclude the introduction, let us summarize these specific results.

(1) Let $\text{Aut}(B)$ denote the group of biholomorphic bijections on the ball B . Let u be any non-constant inner function on B . Then the set $\{u \circ \psi : \psi \in \text{Aut}(B)\}$ is an example of Λ that yields the answer “yes” to the above question. That is, if A is a bounded operator on $H^2(S)$ such that $T_{\overline{u \circ \psi}} A T_{u \circ \psi} = A$ for every $\psi \in \text{Aut}(B)$, then A is a Toeplitz operator.

(2) Suppose that $n \geq 2$. Then for each singleton set $\Lambda = \{u\}$, the answer to the above question is always “no”. In other words, for every inner function u on B , there is a bounded operator Y on $H^2(S)$ such that $T_{\bar{u}}YT_u = Y$ and yet Y is not a Toeplitz operator.

(3) Suppose that $n = 1$. Let u be an inner function on the unit disc. If u does not have the form

$$e^{i\theta} \frac{a - z}{1 - \bar{a}z},$$

where $\theta \in \mathbf{R}$ and $|a| < 1$, then there is a bounded operator Y on H^2 such that $T_{\bar{u}}YT_u = Y$ and yet Y is not a Toeplitz operator. These comments put the Brown-Halmos criterion in the proper perspective.

2. TRANSFORMATIONS ON THE UNIT BALL

For each $a \in B \setminus \{0\}$, we have the Möbius transformation

$$\varphi_a(z) = \frac{1}{1 - \langle z, a \rangle} \left\{ a - \frac{\langle z, a \rangle}{|a|^2} a - (1 - |a|^2)^{1/2} \left(z - \frac{\langle z, a \rangle}{|a|^2} a \right) \right\}, \quad z \in B.$$

Each φ_a is an involution, i.e., $\varphi_a \circ \varphi_a = \text{id}$ [7, Theorem 2.2.2]. We also define $\varphi_0(z) = -z$ on B . By Theorems 3.3.8 and 2.2.2 in [7], the formula

$$(2.1) \quad (U_a f)(z) = \frac{(1 - |a|^2)^{n/2}}{(1 - \langle z, a \rangle)^n} f(\varphi_a(z)), \quad f \in L^2(S, d\sigma),$$

defines a unitary operator with the property $[U_a, P] = 0$.

Let $\mathcal{U} = \mathcal{U}(n)$ denote the collection of unitary transformations on $\mathbf{C}^n$. For each $V \in \mathcal{U}$, define the operator $W_V : L^2(S, d\sigma) \rightarrow L^2(S, d\sigma)$ by the formula

$$(2.2) \quad (W_V g)(z) = g(Vz),$$

$g \in L^2(S, d\sigma)$. By the invariance of σ , W_V is a unitary operator on $L^2(S, d\sigma)$.

Let $\text{Aut}(B)$ denote the group of biholomorphic bijections on B . If $\psi \in \text{Aut}(B)$ and if $a \in B$ is such that $\psi(a) = 0$, then

$$(2.3) \quad \psi = V\varphi_a$$

for some $V \in \mathcal{U}$ [7, Theorem 2.2.5]. For such a ψ , set

$$(2.4) \quad R_\psi = U_a W_V.$$

Then R_ψ is a unitary operator on $L^2(S, d\sigma)$ which has the properties that

$$R_\psi M_f R_\psi^* = M_{f \circ \psi}$$

for every $f \in L^\infty(S, d\sigma)$ and that $[R_\psi, P] = 0$. Consequently, $H^2(S)$ is a reducing subspace for R_ψ , and if we regard R_ψ as a unitary operator on $H^2(S)$, then

$$R_\psi T_f R_\psi^* = T_{f \circ \psi}$$

for every $f \in L^\infty(S, d\sigma)$.

Let $H^\infty(S)$ be the collection of bounded analytic functions on B . As usual, each $h \in H^\infty(S)$ is identified with its boundary value on S . Our first proposition is based on ideas in Section VI of the paper [4] by Feldman and Rochberg, where the main interest was Hankel operators with conjugate analytic symbols.

Proposition 2.1. *Let $h \in H^\infty(S)$. If h is not a constant then, on the Hilbert space $L^2(S, d\sigma)$, the von Neumann algebra generated by*

$$\{M_{h \circ \psi} : \psi \in \text{Aut}(B)\}$$

equals $\{M_f : f \in L^\infty(S, d\sigma)\}$.

Proof. Let $\partial_1, \dots, \partial_n$ denote the differentiations with respect to the complex variables $z_1, \dots, z_n$ respectively. We first show that if h is not a constant, then there exist $\psi_1, \dots, \psi_n \in \text{Aut}(B)$ such that

$$(2.5) \quad \partial_j(h \circ \psi_j)(0) \neq 0, \quad j = 1, \dots, n.$$

For any analytic function g on B , write $\text{grad}(g) = (\partial_1 g, \dots, \partial_n g)$. If h is not a constant, then there exists an $a \in B$ such that $\text{grad}(h)(a) \neq 0$, that is, $\text{grad}(h)(\varphi_a(0)) \neq 0$. Since the derivative $\varphi'_a(0)$ is an invertible $n \times n$ matrix [7, Theorem 2.2.2(ii)], by the chain rule, we have $\text{grad}(h \circ \varphi_a)(0) \neq 0$. That is, there is at least one $\nu \in \{1, \dots, n\}$ such that

$$\partial_\nu(h \circ \varphi_a)(0) \neq 0.$$

By simple transpositions of coordinates, we see that there are $V_1, \dots, V_n \in \mathcal{U}$ such that $\partial_j(h \circ \varphi_a \circ V_j)(0) \neq 0$ for $j = 1, \dots, n$. Thus (2.5) holds for $\psi_j = \varphi_a \circ V_j = V_j \varphi_{V_j^* a}$, $j = 1, \dots, n$.

Let $\mathbf{T}^n$ denote the n -dimensional torus $\{(\tau_1, \dots, \tau_n) : |\tau_1| = \dots = |\tau_n| = 1\}$. Let dm_n be the Lebesgue measure on $\mathbf{T}^n$ with the normalization $m_n(\mathbf{T}^n) = 1$. Now, for each pair of $j \in \{1, \dots, n\}$ and $\tau = (\tau_1, \dots, \tau_n) \in \mathbf{T}^n$, define the function

$$\eta_\tau^{(j)}(z_1, \dots, z_n) = (h \circ \psi_j)(\tau_1 z_1, \dots, \tau_n z_n).$$

Then, of course, we still have $\eta_\tau^{(j)} \in \{h \circ \psi : \psi \in \text{Aut}(B)\}$. Using a power-series expansion, it is straightforward to verify that for each $j \in \{1, \dots, n\}$,

$$\int \bar{\tau}_j M_{\eta_\tau^{(j)}} dm_n(\tau) = \partial_j(h \circ \psi_j)(0) M_{z_j}.$$

By (2.5), this means that the von Neumann algebra generated by $\{M_{h \circ \psi} : \psi \in \text{Aut}(B)\}$ contains $M_{z_1}, \dots, M_{z_n}$. Since $M_{z_1}, \dots, M_{z_n}$ generate the von Neumann algebra $\{M_f : f \in L^\infty(S, d\sigma)\}$, this completes the proof. $\square$

3. MAIN RESULTS

To better state our results, let us introduce the following terminology:

Definition 3.1. Let Λ be a non-empty set of inner functions. We say that the set Λ is *Toeplitz-determining* if it has the property that for $A \in \mathcal{B}(H^2(S))$, the condition that $T_{\bar{u}} A T_u = A$ for every $u \in \Lambda$ implies that A is a Toeplitz operator on $H^2(S)$.

The following is our main result:

Theorem 3.2. *Let Λ be a non-empty set of inner functions. Let $\mathcal{N}(\Lambda)$ denote the von Neumann algebra generated by*

$$\{M_u : u \in \Lambda\}$$

on the Hilbert space $L^2(S, d\sigma)$. Then the set Λ is Toeplitz-determining if and only if $\mathcal{N}(\Lambda) = \{M_f : f \in L^\infty(S, d\sigma)\}$.

Proof. First, assuming that $\mathcal{N}(\Lambda) = \{M_f : f \in L^\infty(S, d\sigma)\}$, we will show that Λ is Toeplitz-determining. The set Λ is, of course, a subset of $H^2(S)$. Since $H^2(S)$ is separable and since separability is a hereditary property for metric spaces, there is a countable subset Λ_0 of Λ which is dense in Λ with respect to the norm topology of $H^2(S)$. We can always list Λ_0 as

$$\Lambda_0 = \{u_1, u_2, \dots, u_k, \dots\}$$

if we allow the possibility $u_j = u_k$ for distinct j and k .

Now suppose that A is a bounded operator on $H^2(S)$ such that

$$(3.1) \quad T_{\bar{u}}AT_u = A \quad \text{for every } u \in \Lambda.$$

To show that $A = T_\varphi$ for some $\varphi \in L^\infty(S, d\sigma)$, we follow the ideas in the proof of [3, Lemma 2.5]. Define the operator

$$\tilde{A} = A \oplus 0$$

on $L^2(S, d\sigma)$, where the direct sum corresponds to the space decomposition

$$L^2(S, d\sigma) = H^2(S) \oplus \{H^2(S)\}^\perp.$$

For each natural number k , define the operator

$$L_k = \frac{1}{k^k} \sum_{1 \leq i_1, \dots, i_k \leq k} M_{\bar{u}_k^{i_k}} \cdots M_{\bar{u}_1^{i_1}} \tilde{A} M_{u_1^{i_1}} \cdots M_{u_k^{i_k}}.$$

It is easy to see that if $1 \leq j \leq k$, then

$$(3.2) \quad \|M_{\bar{u}_j} L_k M_{u_j} - L_k\| \leq 2 \frac{k^{k-1}}{k^k} \|\tilde{A}\| = \frac{2}{k} \|A\|.$$

Since $\|L_k\| \leq \|\tilde{A}\| = \|A\|$ for every $k \geq 1$, there is a strictly increasing sequence of natural numbers $k(1) < k(2) < \cdots < k(\ell) < \cdots$ such that the limit

$$L = \lim_{\ell \rightarrow \infty} L_{k(\ell)}$$

exists in the weak operator topology. Clearly, it follows from (3.2) that $M_{\bar{u}_j} L M_{u_j} - L = 0$ for every $j \geq 1$. Since $|u_j| = 1$ a.e. on S , this means that

$$L M_{u_j} = M_{u_j} L \quad \text{for every } j \geq 1.$$

Since Λ_0 is dense in Λ with respect to the L^2 -norm, the above implies that L commutes with $\{M_u : u \in \Lambda\}$. Since each M_u , $u \in \Lambda$, is a unitary operator, it follows that L also commutes with $\{M_{\bar{u}} : u \in \Lambda\}$. The assumption $\mathcal{N}(\Lambda) = \{M_f : f \in L^\infty(S, d\sigma)\}$ then leads to the conclusion that L commutes with $\{M_f : f \in L^\infty(S, d\sigma)\}$. Hence there is a $\varphi \in L^\infty(S, d\sigma)$ such that $L = M_\varphi$.

Thus to complete the proof that A is a Toeplitz operator, it suffices to show that the compression of L to the subspace $H^2(S)$ equals A . Note that for $h, g \in H^2(S)$ and natural numbers $1 \leq i_1, \dots, i_k \leq k$, we have

$$\begin{aligned} \langle M_{\bar{u}_k^{i_k}} \cdots M_{\bar{u}_1^{i_1}} \tilde{A} M_{u_1^{i_1}} \cdots M_{u_k^{i_k}} h, g \rangle &= \langle \tilde{A} M_{u_1^{i_1}} \cdots M_{u_k^{i_k}} h, M_{u_1^{i_1}} \cdots M_{u_k^{i_k}} g \rangle \\ &= \langle AT_{u_1^{i_1}} \cdots T_{u_k^{i_k}} h, T_{u_1^{i_1}} \cdots T_{u_k^{i_k}} g \rangle \\ &= \langle T_{u_k^{i_k}} \cdots T_{u_1^{i_1}} AT_{u_1^{i_1}} \cdots T_{u_k^{i_k}} h, g \rangle = \langle Ah, g \rangle, \end{aligned}$$

where the last $=$ follows from repeated applications of (3.1). Thus the compression of each L_k to $H^2(S)$ equals A . Hence $A = PL|_{H^2(S)}$ as promised. This proves the "if" part of the theorem.

To prove the “only if” part, let us now assume that $\mathcal{N}(\Lambda) \neq \{M_f : f \in L^\infty(S, d\sigma)\}$. We will find a bounded operator Y on $H^2(S)$ such that $T_{\bar{u}}YT_u = Y$ for every $u \in \Lambda$ and such that $Y \notin \{T_f : f \in L^\infty(S, d\sigma)\}$.

By the double-commutant relation, the assumption that $\mathcal{N}(\Lambda) \neq \{M_f : f \in L^\infty(S, d\sigma)\}$ implies that the commutant of $\mathcal{N}(\Lambda)$ is strictly larger than the commutant of $\{M_f : f \in L^\infty(S, d\sigma)\}$. That is, there is a bounded operator Z which commutes with $\mathcal{N}(\Lambda)$ but which does not commute with $\{M_f : f \in L^\infty(S, d\sigma)\}$.

Now take any non-constant inner function v constructed by Aleksandrov [1] or Løw [6]. By Proposition 2.1, the unitary operators $\{M_{v \circ \psi} : \psi \in \text{Aut}(B)\}$ generate the von Neumann algebra $\{M_f : f \in L^\infty(S, d\sigma)\}$. Since Z does not commute with $\{M_f : f \in L^\infty(S, d\sigma)\}$, it follows that there is an inner function $w \in \{v \circ \psi : \psi \in \text{Aut}(B)\}$ such that

$$M_{\bar{w}}ZM_w \neq Z.$$

We follow the usual multi-index notation [7, page 3]. For each $\alpha \in \mathbf{Z}_+^n$, define the function $\epsilon_\alpha(z) = z^\alpha$ on S . Then, of course, the linear span of $\{\epsilon_\alpha \bar{\epsilon}_\beta : \alpha, \beta \in \mathbf{Z}_+^n\}$ is dense in $L^2(S, d\sigma)$. Hence there are $a, b, c, d \in \mathbf{Z}_+^n$ such that

$$\langle M_{\bar{w}}ZM_w \epsilon_a \bar{\epsilon}_b, \epsilon_c \bar{\epsilon}_d \rangle \neq \langle Z \epsilon_a \bar{\epsilon}_b, \epsilon_c \bar{\epsilon}_d \rangle.$$

Now define the operator $X = M_{\epsilon_d}ZM_{\bar{\epsilon}_b}$. Then the above gives us

$$(3.3) \quad \langle M_{\bar{w}}XM_w \epsilon_a, \epsilon_c \rangle \neq \langle X \epsilon_a, \epsilon_c \rangle.$$

Let Y be the compression of X to the subspace $H^2(S)$. That is, $Y = PX|_{H^2(S)}$. Since $\epsilon_a, \epsilon_c \in H^2(S)$ and since $(1 - P)M_wP = 0$, (3.3) tells us that $T_{\bar{w}}YT_w \neq Y$. Since w is an inner function, this means that Y is not a Toeplitz operator.

On the other hand, since Z is in the commutant of $\mathcal{N}(\Lambda)$, we have $M_{\bar{u}}ZM_u = Z$ for every $u \in \Lambda$. Since $X = M_{\epsilon_d}ZM_{\bar{\epsilon}_b}$, it follows that $M_{\bar{u}}XM_u = X$ for every $u \in \Lambda$. Compressing to the subspace $H^2(S)$, we see that $T_{\bar{u}}YT_u = Y$ for every $u \in \Lambda$. Thus we have produced the promised Y . This completes the proof. $\square$

As an immediate consequence of Theorem 3.2 and Proposition 2.1, we have

Corollary 3.3. *Let u be a non-constant inner function. Then a bounded operator A on $H^2(S)$ is a Toeplitz operator if and only if*

$$T_{\overline{u \circ \psi}}AT_{u \circ \psi} = A$$

for every $\psi \in \text{Aut}(B)$.

Recall that the unitary operator R_ψ defined by (2.4) has the property that $R_\psi^*T_fR_\psi = T_{f \circ \psi^{-1}}$, $f \in L^\infty(S, d\sigma)$. In other words, an operator A is a Toeplitz operator if and only if $R_\psi^*AR_\psi$ is a Toeplitz operator. In light of this, let us restate Corollary 3.3 as

Corollary 3.4. *Let u be a non-constant inner function. Then a bounded operator A on $H^2(S)$ is a Toeplitz operator if and only if*

$$T_{\bar{u}}R_\psi^*AR_\psiT_u = R_\psi^*AR_\psi$$

for every $\psi \in \text{Aut}(B)$.

4. SET OF A SINGLETON

Let us now consider the case where Λ is a set of a single inner function. In this case, the story is actually simpler in complex dimensions $n \geq 2$.

Proposition 4.1. *Suppose that $n \geq 2$. Then for each inner function u , there is a bounded operator Y on $H^2(S)$ such that $T_{\bar{u}}YT_u = Y$ and such that Y is not a Toeplitz operator.*

Proof. Let an inner function u be given and let $\mathcal{N}(u)$ denote the von Neumann algebra generated by the single unitary operator M_u . To prove the proposition, according to Theorem 3.2, it suffices to show that $\mathcal{N}(u) \neq \{M_f : f \in L^\infty(S, d\sigma)\}$.

Let $\mathcal{P}$ denote the linear span of all u^j and $\bar{u}^k$, $j, k = 0, 1, 2, \dots$. Note that since $u\bar{u} = 1$, $\mathcal{N}(u)$ is just the weak closure of $\{M_\xi : \xi \in \mathcal{P}\}$. Because of the assumption $n \geq 2$, we have the function

$$q(z_1, \dots, z_n) = z_1 \bar{z}_2$$

on S . To complete the proof, it suffices to show that M_q is not in the weak closure of $\{M_\xi : \xi \in \mathcal{P}\}$. Assuming the contrary, there would be a sequence $\{\xi_i\} \subset \mathcal{P}$ such that $\langle M_{\xi_i} 1, q \rangle \rightarrow \langle M_q 1, q \rangle$ as $i \rightarrow \infty$. In other words, we would have

$$(4.1) \quad \lim_{i \rightarrow \infty} \langle \xi_i, q \rangle = \langle q, q \rangle = \|q\|^2 > 0$$

for some sequence $\{\xi_i\} \subset \mathcal{P}$. But for each integer $j \geq 0$, the analyticity of u^j gives us

$$\langle u^j, q \rangle = \langle z_2 u^j, z_1 \rangle = 0.$$

Similarly, for each integer $k \geq 0$, we have

$$\langle \bar{u}^k, q \rangle = \langle z_2, z_1 u^k \rangle = 0.$$

Thus, in the Hilbert space $L^2(S, d\sigma)$, q is orthogonal to the set $\mathcal{P}$. This clearly contradicts (4.1). This contradiction shows that M_q is not in the weak closure of $\{M_\xi : \xi \in \mathcal{P}\}$. $\square$

Proposition 4.2. *Suppose that $n = 1$. If u is an inner function on the unit disc and if u does not have the form*

$$(4.2) \quad e^{i\theta} \frac{a - z}{1 - \bar{a}z}, \quad \text{where } \theta \in \mathbf{R} \text{ and } |a| < 1,$$

then there is a bounded operator Y on H^2 such that $T_{\bar{u}}YT_u = Y$ and such that Y is not a Toeplitz operator.

Proof. For the given u , again let $\mathcal{N}(u)$ denote the von Neumann algebra generated by the single unitary operator M_u . To prove the proposition, according to Theorem 3.2, it suffices to show that if u does not have the form (4.2), then $\mathcal{N}(u) \neq \{M_f : f \in L^\infty\}$.

We only need, of course, to consider the case where u is not a constant. Then

$$u = bs,$$

where b is a Blaschke product or a constant, and s is a so-called singular inner function or a constant. Suppose that u does not have the form (4.2). Then either b has at least two zeros (counting multiplicity) or s is a non-trivial singular inner function. For such a u it is well known (and easy to verify) that the dimension of

the subspace $H^2 \ominus uH^2$ is at least 2. Hence there is a $q \in H^2 \ominus uH^2$ with $\|q\| \neq 0$ which is orthogonal to the one-dimensional subspace $\mathbf{C}$, that is,

$$(4.3) \quad q \in H^2, \quad q \perp uH^2, \quad \text{and} \quad q \perp \mathbf{C}.$$

Since $\|q\| \neq 0$, there exists a $p \in L^\infty$ such that $\langle p, q \rangle \neq 0$.

As in the previous proof, let $\mathcal{P}$ denote the linear span of all u^j and $\bar{u}^k$, $j, k = 0, 1, 2, \dots$. Again, $\mathcal{N}(u)$ is just the weak closure of $\{M_\xi : \xi \in \mathcal{P}\}$. To complete the proof, it suffices to show that M_p is not in the weak closure of $\{M_\xi : \xi \in \mathcal{P}\}$. Assuming the contrary, there would be a sequence $\{\xi_i\} \subset \mathcal{P}$ such that $\langle M_{\xi_i} 1, q \rangle \rightarrow \langle M_p 1, q \rangle$ as $i \rightarrow \infty$. In other words, we would have

$$(4.4) \quad \lim_{i \rightarrow \infty} \langle \xi_i, q \rangle = \langle p, q \rangle \neq 0$$

for some sequence $\{\xi_i\} \subset \mathcal{P}$. But clearly, (4.3) implies $q \perp \mathcal{P}$, which contradicts (4.4). This contradiction shows that M_p is not in the weak closure of $\{M_\xi : \xi \in \mathcal{P}\}$. $\square$

REFERENCES

1. A. Aleksandrov, *The existence of inner functions in a ball* (Russian), Mat. Sb. (N.S.) **118(160)** (1982), 147-163, 287; (English) Math. USSR Sbornik **46** (1983), 143-159. MR658785 (83i:32002)
2. A. Brown and P. Halmos, *Algebraic properties of Toeplitz operators*, J. Reine Angew. Math. **213** (1963/64), 89-102. MR0160136 (28:3350)
3. A. Davie and N. Jewell, *Toeplitz operators in several complex variables*, J. Funct. Anal. **26** (1977), 356-368. MR0461195 (57:1180)
4. M. Feldman and R. Rochberg, *Singular value estimates for commutators and Hankel operators on the unit ball and the Heisenberg group*, Analysis and partial differential equations, 121-159, Lecture Notes in Pure and Appl. Math., **122**, Dekker, New York, 1990. MR1044786 (91e:47024)
5. R. Kadison and J. Ringrose, *Fundamentals of the theory of operator algebras*. Vol. I. *Elementary theory*. Pure and Applied Mathematics, **100**. Academic Press, Inc., New York, 1983. MR719020 (85j:46099)
6. E. Löw, *A construction of inner functions on the unit ball in $\mathbf{C}^p$* , Invent. Math. **67** (1982), 223-229. MR665154 (84j:32008b)
7. W. Rudin, *Function theory in the unit ball of $\mathbf{C}^n$* , Springer-Verlag, New York-Berlin, 1980. MR601594 (82i:32002)
8. W. Rudin, *Inner functions in the unit ball of $\mathbf{C}^n$* , J. Funct. Anal. **50** (1983), 100-126. MR690001 (84i:32007)

DEPARTMENT OF MATHEMATICS, STATE UNIVERSITY OF NEW YORK AT BUFFALO, BUFFALO, NEW YORK 14260

E-mail address: jxia@acsu.buffalo.edu