COMMENT ON A RESULT BY ALPIN, CHIEN, AND YEH

A. MELMAN

(Communicated by Walter Van Assche)

ABSTRACT. Theorem 3 in Alpin, Yu. A.; Chien, Mao-Ting; and Yeh, Lina, *The numerical radius and bounds for zeros of a polynomial*, Proc. Amer. Math. Soc., 131 (2003), no. 3, 725–730, is incorrect. In this note, we identify the error and present a correctly stated theorem.

1. INTRODUCTION

This note concerns the paper [1] by Alpin, Chien, and Yeh. One of their results, namely, Theorem 3, is erroneous. We provide a counterexample and then identify the flaw in the argument of the proof before stating an amended theorem. Most of our notation will be the same as in [1].

Let

$$(1.1) \quad \begin{pmatrix} 0 & 0 & \cdots & 0 & -a_0 \\ 1 & 0 & \cdots & 0 & -a_1 \\ 0 & 1 & \cdots & 0 & -a_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & -a_{n-1} \end{pmatrix}$$

be the companion matrix of the monic polynomial

$$(1.2) \quad p(t) = t^n + a_{n-1}t^{n-1} + a_{n-2}t^{n-2} + \cdots + a_1t + a_0 ,$$

and define for a complex $n \times n$ matrix $A = (a_{ij})$:

$$(1.3) \quad R'_i(A) = \sum_{\substack{j=1 \\ j \neq i}}^n |a_{ij}| \quad \text{and} \quad g_i(A) = \frac{1}{2} (R'_i(A) + R'_i(A^*)) , \quad 1 \leq i \leq n ,$$

where A^* is the Hermitian conjugate of A . We denote the spectrum of A by $\sigma(A)$ and its numerical range by $W(A)$, i.e.,

$$W(A) = \{z^*Az : z \in \mathbb{C}^n, |z| = 1\} .$$

The set of points $z \in \mathbb{C}$ satisfying $|z - a||z - b| = \rho$ for $a, b \in \mathbb{C}$ and $\rho \geq 0$ is known as an oval of Cassini.

We now state Theorem 3 in [1].

Theorem 1.1 ([1, Theorem 3]). *Let p be the polynomial (1.2) and $p(z) = 0$, and let A be the matrix defined by (1.1). Let $\alpha = \max_{1 \leq i < n} g_i(A)$. Then*

$$|z| \leq |a_{n-1}|/2 + \left(|a_{n-1}|^2/4 + (\alpha^2 g_n^2(A) + |a_{n-1}|^2)^{1/2} \right)^{1/2} .$$

Received by the editors May 16, 2011 and, in revised form, July 16, 2011.

2010 *Mathematics Subject Classification*. Primary 15A60, 26C10.

©2012 American Mathematical Society
 Reverts to public domain 28 years from publication

The following claim presents a counterexample to this theorem.

Claim 1.2. The upper bound from Theorem 1 is not an upper bound for the modulus of any zero of $q(t) = t^9 + 0.01t^8 + 0.2$.

Proof. For the companion matrix A of $q(t)$, we have

$$\begin{aligned} R'_1(A) &= 0.2, \quad R'_2(A) = R'_3(A) = \cdots = R'_8(A) = 1, \quad R'_9(A) = 1, \\ R'_1(A^*) &= 1, \quad R'_2(A^*) = R'_3(A^*) = \cdots = R'_8(A^*) = 1, \quad R'_9(A^*) = 0.2, \\ g_1(A) &= 0.6, \quad g_2(A) = g_3(A) = \cdots = g_8(A) = 1, \quad g_9(A) = 0.6. \end{aligned}$$

Therefore, $\alpha = \max_{1 \leq i < 9} g_i(A) = 1$, and the upper bound in Theorem 1 is given by

$$0.01/2 + \left(0.01^2/4 + (1^2 \cdot 0.6^2 + 0.01^2)^{1/2}\right)^{1/2} = 0.7797.$$

However, any zero z of $q(t)$ satisfies $0.8352 \leq |z| \leq 0.8374$, with the entire range being larger than 0.7797. $\square$

2. CORRECTED THEOREM

The error in the proof of Theorem 3 in [1] was caused by the following incorrect inequality:

$$(2.1) \quad \alpha \leq |a_{n-1}|/2 + \left(|a_{n-1}|^2/4 + (\alpha^2 g_n^2(A) + |a_{n-1}|^2)^{1/2}\right)^{1/2}.$$

To see why this inequality cannot be correct in general, assume, e.g., that for the coefficients of p in (1.2): $a_1 = a_2 = \cdots = a_{n-2} = 0$, $a_{n-1} = \beta$, and $a_0 = \epsilon$ with $\beta, \epsilon > 0$ and $\beta, \epsilon \ll 1$. Then $\alpha = 1$ and $g_n = (1 + \epsilon)/2$. For β sufficiently small, inequality (2.1) becomes

$$\begin{aligned} 1 &\leq \frac{\beta}{2} + \left(\frac{\beta^2}{4} + \left(\frac{(1+\epsilon)^2}{4} + \beta^2\right)^{1/2}\right)^{1/2} \\ &= \frac{\beta}{2} + \left(\frac{1+\epsilon}{2} + \mathcal{O}(\beta^2)\right)^{1/2} \\ &= \left(\frac{1+\epsilon}{2}\right)^{1/2} + \mathcal{O}(\beta). \end{aligned}$$

Clearly, as $\beta, \epsilon \rightarrow 0$, this inequality cannot hold. The following theorem corrects Theorem 3 in [1].

Theorem 2.1. Let p be the polynomial (1.2) and $p(z) = 0$, and let A be the matrix defined by (1.1). Let α_1 and α_2 be the largest and (possibly equal) second largest of the numbers $\{g_i(A)\}_{i=1}^{n-1}$. Then

$$|z| \leq \max \left\{ \sqrt{\alpha_1 \alpha_2}, \frac{|a_{n-1}|}{2} + \left(\frac{|a_{n-1}|^2}{4} + (\alpha_1^2 g_n^2(A) + |a_{n-1}|^2)^{1/2}\right)^{1/2} \right\}.$$

Proof. The proof follows, just as in Theorem 3 in [1], by applying Theorem 2 in [1] to the companion matrix A of p . The closed interiors of the ovals of Cassini defined in that theorem are then the collection of points $z \in \mathbb{C}$ for which

$$(2.2) \quad |z| \leq (g_i(A)g_j(A))^{1/2}, \quad 1 \leq i \neq j < n,$$

and those for which

$$(2.3) \quad |z||z + a_{n-1}| \leq (g_i^2(A)g_n^2(A) + |a_{n-1}|^2)^{1/2}, \quad 1 \leq i < n.$$

The points defined by (2.2) are all contained in the disk defined by $|z| \leq (\alpha_1\alpha_2)^{1/2}$, and the points defined by (2.3) are all contained in the disk defined by

$$|z| \leq \frac{|a_{n-1}|}{2} + \left(\frac{|a_{n-1}|^2}{4} + (\alpha_1^2 g_n^2(A) + |a_{n-1}|^2)^{1/2} \right)^{1/2}.$$

This, together with the fact that $\sigma(A) \subset W(A)$, completes the proof. $\square$

For our counterexample with $q(t) = t^9 + 0.01t^8 + 0.2$, we obtain $\alpha_1 = \alpha_2 = 1$, so that Theorem 2 states, correctly, that for any zero z of q : $|z| \leq 1$.

REFERENCES

1. Alpin, Yu. A., Chien, Mao-Ting, and Yeh, Lina. *The numerical radius and bounds for zeros of a polynomial*. Proc. Amer. Math. Soc., 131 (2003), no. 3, 725–730. MR1937409 (2003h:26021)

DEPARTMENT OF APPLIED MATHEMATICS, SCHOOL OF ENGINEERING, SANTA CLARA UNIVERSITY, SANTA CLARA, CALIFORNIA 95053

E-mail address: `amelman@scu.edu`