AN EMBEDDING THEOREM

N. A. CHERNYAVSKAYA AND L. A. SHUSTER

(Communicated by Michael Hitrik)

ABSTRACT. We consider a weighted space $W_1^{(2)}(\mathbb{R}, q)$ of Sobolev type:

$$W_1^{(2)}(\mathbb{R}, q) = \left\{ y \in AC_{\text{loc}}^{(1)}(\mathbb{R}) : \|y''\|_{L_1(\mathbb{R})} + \|qy\|_{L_1(\mathbb{R})} < \infty \right\},$$

where $0 \leq q \in L_1^{\text{loc}}(\mathbb{R})$ and

$$\|y\|_{W_1^{(2)}(\mathbb{R}, q)} = \|y''\|_{L_1(\mathbb{R})} + \|qy\|_{L_1(\mathbb{R})}.$$

We obtain a precise condition which guarantees the embedding

$$W_1^{(2)}(\mathbb{R}, q) \hookrightarrow L_p(\mathbb{R}), \quad p \in [1, \infty).$$

1. INTRODUCTION

In the present paper, we consider a weighted function space $W_1^{(2)}(\mathbb{R}, q)$ of Sobolev type:

$$(1.1) \quad W_1^{(2)}(\mathbb{R}, q) = \left\{ y \in AC_{\text{loc}}^{(1)}(\mathbb{R}) : \|y''\|_{L_1(\mathbb{R})} + \|qy\|_{L_1(\mathbb{R})} < \infty \right\}.$$

Here and in the sequel, $\|f\|_p = \|f\|_{L_p(\mathbb{R})}$, $p \in [1, \infty)$, $AC_{\text{loc}}^{(1)}(\mathbb{R})$, is the set of functions absolutely continuous together with the derivative on every finite interval.

In this paper we always require that

$$(1.2) \quad 0 \leq q \in L_1^{\text{loc}}(\mathbb{R})$$

and usually do not mention this requirement in the statements.

Our general goal is to find conditions under which there is an embedding

$$W_1^{(2)}(\mathbb{R}, q) \hookrightarrow L_p(\mathbb{R}).$$

To be more precise, we give the following definitions.

Definition 1.1 ([6]). Let $p \in [1, \infty)$. We say that the space $W_1^{(2)}(\mathbb{R}, q)$ is embedded into the space $L_p(\mathbb{R})$ (and write $W_1^{(2)}(\mathbb{R}, q) \hookrightarrow L_p(\mathbb{R})$) if $W_1^{(2)}(\mathbb{R}, q) \subseteq L_p(\mathbb{R})$ and

$$(1.3) \quad \|y\|_p \leq c(p) \{ \|y''\|_1 + \|qy\|_1 \}, \quad \forall y \in W_1^{(2)}(\mathbb{R}, q).$$

Our general convention is that the letters $c, c(\cdot)$ stand for absolute positive constants which are not essential for exposition and may differ even within a single chain of calculations.

We now state our main result.

Received by the editors December 1, 2011.

2010 *Mathematics Subject Classification*. Primary 46E35; Secondary 34B24.

Key words and phrases. Embedding theorem.

Theorem 1.2. *Let $p \in [1, \infty)$. Then the embedding $W_1^{(2)}(\mathbb{R}, q) \hookrightarrow L_p(\mathbb{R})$ takes place if and only if*

$$(1.4) \quad \exists a > 0 : \quad q_0(a) \stackrel{\text{def}}{=} \inf_{x \in \mathbb{R}} \int_{x-a}^{x+a} q(t) dt > 0.$$

Note that for $p = 1$ this assertion was proved in [4] and will not be reconsidered here (see §2).

Example 1.3. Let

$$(1.5) \quad q(x) = 1 + \cos(|x|^\theta), \quad x \in \mathbb{R}, \quad \theta \in (0, \infty).$$

In [3] it was shown that in case (1.5) condition (1.4) holds if and only if $\theta \geq 1$. Together with Theorem 1.2, this implies that in case (1.5) the embedding $W_1^{(2)}(\mathbb{R}, q) \hookrightarrow L_p(\mathbb{R})$, $p \in [1, \infty)$, takes place if and only if $\theta \geq 1$.

2. PRELIMINARIES

In this section, we present some facts needed for the proofs (see §3).

Lemma 2.1 ([1, 2]). *Suppose conditions (1.2) and (2.1) hold:*

$$(2.1) \quad \int_{-\infty}^x q(t) dt > 0, \quad \int_x^{\infty} q(t) dt > 0, \quad \forall x \in \mathbb{R}.$$

Then there exists a fundamental system of solutions (FSS) $\{u, v\}$ of the equation

$$(2.2) \quad z''(x) = q(x)z(x), \quad x \in \mathbb{R},$$

which has the following properties:

$$(2.3) \quad u(x) > 0, \quad v(x) > 0, \quad u'(x) \leq 0, \quad v'(x) \geq 0, \quad \forall x \in \mathbb{R},$$

$$(2.4) \quad v'(x)u(x) - u'(x)v(x) = 1, \quad \forall x \in \mathbb{R},$$

$$(2.5) \quad \lim_{x \rightarrow -\infty} \frac{v(x)}{u(x)} = \lim_{x \rightarrow \infty} \frac{u(x)}{v(x)} = 0.$$

The solutions $\{u, v\}$ and the function ρ ,

$$(2.6) \quad \rho(x) = u(x)v(x), \quad x \in \mathbb{R},$$

satisfy the relations

$$(2.7) \quad \int_{-\infty}^0 \frac{dt}{\rho(t)} = \int_0^{\infty} \frac{dt}{\rho(t)} = \infty,$$

$$(2.8) \quad |\rho'(x)| < 1, \quad \forall x \in \mathbb{R},$$

$$(2.9) \quad \frac{v'(x)}{v(x)} = \frac{1 + \rho'(x)}{2\rho(x)}, \quad \frac{u'(x)}{u(x)} = -\frac{1 - \rho'(x)}{2\rho(x)}, \quad x \in \mathbb{R}.$$

In addition, if $G(x, t)$ is the Green function,

$$(2.10) \quad G(x, t) = \begin{cases} u(x)v(t), & x \geq t, \\ u(t)v(x), & x \leq t, \end{cases}$$

corresponding to the equation

$$(2.11) \quad -y''(x) + q(x)y(x) = f(x), \quad x \in \mathbb{R},$$

then we have the inequality

$$(2.12) \quad \sup_{t \in \mathbb{R}} \Phi(t) \leq 1, \quad \Phi(t) = \int_{-\infty}^{\infty} q(\xi) G(t, \xi) d\xi, \quad t \in \mathbb{R}.$$

Finally, for every $p \in [1, \infty)$, equation (2.2) has no solutions $z \in L_p(\mathbb{R})$ apart from $z \equiv 0$.

Lemma 2.2 ([1]). *Suppose conditions (1.2) and (2.1) hold. For a given $x \in \mathbb{R}$, consider the equation in $d \geq 0$:*

$$(2.13) \quad d \int_{x-d}^{x+d} q(t) dt = 2.$$

Equation (2.13) has a unique finite positive solution for every $x \in \mathbb{R}$. Denote this solution by $d(x)$. The function $d(x)$ is continuous for $x \in \mathbb{R}$.

Note that the function d was introduced by M. Otelbaev (see [6]).

Theorem 2.3 ([1]). *Suppose conditions (1.2) and (2.1) hold. Then we have the inequalities*

$$(2.14) \quad \frac{d(x)}{4} \leq \rho(x) \leq \frac{3}{2} d(x), \quad x \in \mathbb{R}.$$

Estimates of type (2.14) were first obtained (under some additional requirements to the function q) by Otelbaev (see [7]) and are therefore called Otelbaev's inequalities (see also [6]).

Consider equation (2.11) with $f \in L_p(\mathbb{R})$, $p \in [1, \infty)$. Below by a solution of (2.11) we mean any function $y \in AC_{\text{loc}}^{(1)}(\mathbb{R})$ satisfying equality (2.11) almost everywhere on $\mathbb{R}$.

Definition 2.4 ([3]). We say that equation (2.11) is correctly solvable in the space $L_p(\mathbb{R})$, $p \in [1, \infty)$, if assertions I–II hold:

- I) for any function $f \in L_p(\mathbb{R})$, there is a unique solution $y \in L_p(\mathbb{R})$ of (2.11);
- II) the solution $y \in L_p(\mathbb{R})$ of (2.11) satisfies the estimate

$$(2.15) \quad \|y\|_p \leq c(p) \|f\|_p, \quad \forall f \in L_p(\mathbb{R}).$$

Theorem 2.5 ([3]). *Let $p \in [1, \infty)$. Equation (2.11) is correctly solvable in $L_p(\mathbb{R})$ if and only if one of the following two equivalent conditions, A, B, holds:*

- A) *One has inequalities (2.1) and $d_0 < \infty$. Here*

$$(2.16) \quad d_0 = \sup_{x \in \mathbb{R}} d(x).$$

- B) *There is $a \in (0, \infty)$ such that $q_0(a) > 0$:*

$$(2.17) \quad q_0(a) = \inf_{x \in \mathbb{R}} \int_{x-a}^{x+a} q(t) dt.$$

In addition, if equation (2.11) is correctly solvable in $L_p(\mathbb{R})$, $p \in [1, \infty)$, then its solution $y \in L_p(\mathbb{R})$ is of the form

$$(2.18) \quad y(x) = (Gf)(x) \stackrel{\text{def}}{=} \int_{-\infty}^{\infty} G(x, t) f(t) dt, \quad x \in \mathbb{R}.$$

Theorem 2.6 ([4]). *The embedding $W_1^{(2)}(\mathbb{R}, q) \hookrightarrow L_1(\mathbb{R})$ takes place if and only if equation (2.11) is correctly solvable in $L_1(\mathbb{R})$.*

Theorem 2.7 ([5, Ch. XI, §6]). *Let $K(x, t)$ be a continuous positive function in variables $x, t \in \mathbb{R}$, and let K be the integral operator*

$$(2.19) \quad (Kf)(x) = \int_{-\infty}^{\infty} K(x, t)f(t)dt, \quad \forall f \in L_1(\mathbb{R}).$$

Then for every $p \in (1, \infty)$, we have the equality

$$(2.20) \quad \|K\|_{L_1(\mathbb{R}) \rightarrow L_p(\mathbb{R})} = \sup_{t \in \mathbb{R}} \left(\int_{-\infty}^{\infty} K(x, t)^p dx \right)^{1/p}.$$

3. PROOFS

Proof of Theorem 1.2. Necessity.

Lemma 3.1. *Let $W_1^{(2)}(\mathbb{R}, q) \hookrightarrow L_p(\mathbb{R})$, $p \in [1, \infty)$. Then relations (2.1) hold.*

Proof. Assume the contrary, say,

$$\int_{x_0}^{\infty} q(t)dt = 0,$$

for some $x_0 \in \mathbb{R}$. Let $\varphi \in C^\infty(\mathbb{R})$ be such that $\text{supp } \varphi = [x_0, \infty)$, $0 \leq \varphi \leq 1$, for $x \in \mathbb{R}$ and $\varphi(x) \equiv 1$ for $x \geq x_0 + 1$. Clearly, $\varphi \in AC_{\text{loc}}^{(1)}(\mathbb{R})$ and, in addition,

$$\begin{aligned} \int_{-\infty}^{\infty} |q(t)\varphi(t)|dt &= \int_{x_0}^{\infty} q(t)\varphi(t)dt = 0, \\ 0 &< \int_{-\infty}^{\infty} |\varphi''(t)|dt = \int_{x_0}^{x_0+1} |\varphi''(t)|dt = c < \infty. \end{aligned}$$

Hence $\varphi \in W_1^{(2)}(\mathbb{R}, q)$, and since $W_1^{(2)}(\mathbb{R}, q) \hookrightarrow L_p(\mathbb{R})$, we have

$$\begin{aligned} \infty &> c \left\{ \int_{-\infty}^{\infty} |\varphi''(t)|dt + \int_{-\infty}^{\infty} |q(t)\varphi(t)|dt \right\} \geq \left(\int_{-\infty}^{\infty} |\varphi(t)|^p dt \right) \geq \left(\int_{x_0+1}^{\infty} 1dt \right)^{1/p} \\ &= \infty. \end{aligned}$$

Contradiction. □

Below, conditions (1.2) and (2.1) are assumed to be valid and do not appear in the statements.

Lemma 3.2. *For a given $x \in \mathbb{R}$, consider the equation in $s \geq 0$:*

$$(3.1) \quad \int_{x-s}^{x+s} \frac{dt}{d(t)} = 2.$$

Equation (3.1) has a unique finite positive solution for every $x \in \mathbb{R}$. Denote it by $s(x)$. Then for every $t \in [x - s(x), x + s(x)]$ and $x \in \mathbb{R}$, we have the inequalities

$$(3.2) \quad c^{-1} \leq \frac{u(t)}{u(x)}, \quad \frac{v(t)}{v(x)} \leq c,$$

$$(3.3) \quad c^{-1}d(x) \leq s(x) \leq cd(x).$$

Proof. Consider the function

$$(3.4) \quad F(s) = \int_{x-s}^{x+s} \frac{d\xi}{d(\xi)}, \quad s \geq 0.$$

Clearly, $F(0) = 0$, the function $F(s)$ is monotone increasing and continuous on $[0, \infty)$, and, in addition, in view of (2.7) and (2.14), we have

$$F(s) = \int_{x-s}^{x+s} \frac{dt}{d(t)} \geq \frac{1}{4} \int_{x-s}^{x+s} \frac{dt}{\rho(t)} \rightarrow \infty \quad \text{as } s \rightarrow \infty.$$

Hence, for a given $x \in \mathbb{R}$, equation (3.1) has a unique finite positive solution $s(x)$. Further, from (2.8), (2.9) and (2.14), it follows that

$$\frac{v'(\xi)}{v(\xi)} = \frac{1 + \rho'(\xi)}{2\rho(\xi)} \leq \frac{1}{\rho(\xi)} \leq \frac{4}{d(\xi)}, \quad \xi \in \mathbb{R}.$$

For $t \in [x, x + s(x)]$ this implies that

$$\ln \frac{v(t)}{v(x)} \leq 4 \int_x^t \frac{d\xi}{d(\xi)} \leq 4 \int_{x-s(x)}^{x+s(x)} \frac{d\xi}{d(\xi)} = 8.$$

According to (2.3), this gives

$$(3.5) \quad c^{-1} \leq 1 \leq \frac{v(t)}{v(x)} \leq c \quad \text{for } t \in [x, x + s(x)], \quad x \in \mathbb{R}.$$

Similarly, if $t \in [x - s(x), x]$, then

$$(3.6) \quad \begin{aligned} \ln \frac{v(x)}{v(t)} &\leq 4 \int_t^x \frac{d\xi}{d(\xi)} \leq 4 \int_{x-s(x)}^{x+s(x)} \frac{d\xi}{d(\xi)} = 8 \\ \Rightarrow c^{-1} &\leq 1 \leq \frac{v(x)}{v(t)} \leq c \quad \text{for } t \in [x - s(x), x], \quad x \in \mathbb{R}. \end{aligned}$$

From (3.5) and (3.6) we obtain (3.2) for v (inequalities (3.2) for u are proved in a similar way). Further, by (3.1), (3.2) and (2.14), we get

$$\begin{aligned} 2 &= \int_{x-s(x)}^{x+s(x)} \frac{d\xi}{d(\xi)} = \int_{x-s(x)}^{x+s(x)} \frac{\rho(\xi)}{d(\xi)} \cdot \frac{u(x)}{u(\xi)} \cdot \frac{v(x)}{v(\xi)} \cdot \frac{d(x)}{\rho(x)} \cdot \frac{d\xi}{d(x)} \geq \frac{1}{c} \cdot \frac{s(x)}{d(x)}, \\ 2 &= \int_{x-s(x)}^{x+s(x)} \frac{d\xi}{d(\xi)} = \int_{x-s(x)}^{x+s(x)} \frac{\rho(\xi)}{d(\xi)} \cdot \frac{u(x)}{u(\xi)} \cdot \frac{v(x)}{v(\xi)} \cdot \frac{d(x)}{\rho(x)} \cdot \frac{d\xi}{d(x)} \leq c \frac{s(x)}{d(x)}. \end{aligned}$$

□

Lemma 3.3. Suppose conditions (1.2) and (2.1) hold. Set

$$(3.7) \quad f_x(t) = \begin{cases} 1, & \text{if } t \in [x - s(x), x + s(x)], \\ 0, & \text{if } t \notin [x - s(x), x + s(x)], \end{cases}$$

$$(3.8) \quad y_x(t) = \int_{-\infty}^{\infty} G(t, \xi) f_x(\xi) d\xi, \quad t \in \mathbb{R} \quad (\text{see (2.10)}).$$

Then we have the relations

$$(3.9) \quad -(y_x(t))'' + q(t)y_x(t) = f(t), \quad t \in \mathbb{R},$$

$$(3.10) \quad \|(y_x)''\|_1 + \|qy_x\|_1 \leq cs(x), \quad x \in \mathbb{R}.$$

Proof. From (2.10) and (3.8) it follows that

$$(3.11) \quad y_x(t) = u(t) \int_{-\infty}^t v(\xi) f_x(\xi) d\xi + v(t) \int_t^{\infty} u(\xi) f_x(\xi) d\xi.$$

According to (3.7), integration in (3.11) is along segments which are contained in the finite segment $[x - s(x), x + s(x)]$, and therefore the function y_x is well-defined. Equality (3.9) can be checked directly using Lemma 2.1. Further, from Fubini's Theorem and (2.12), it follows that

$$(3.12) \quad \begin{aligned} \|qy\|_1 &= \int_{-\infty}^{\infty} q(t) \left| \int_{-\infty}^{\infty} G(t, \xi) f_x(\xi) d\xi \right| dt = \int_{-\infty}^{\infty} q(t) \int_{-\infty}^{\infty} G(t, \xi) f_x(\xi) d\xi dt \\ &= \int_{-\infty}^{\infty} f_x(\xi) \left[\int_{-\infty}^{\infty} q(t) G(t, \xi) dt \right] d\xi \leq \int_{-\infty}^{\infty} f_x(\xi) d\xi = 2s(x). \end{aligned}$$

Estimate (3.10) follows from (3.12), (3.7), (3.9) and the triangle inequality for norms. $\square$

Corollary 3.4. *Suppose conditions (1.2) and (2.1) hold. Let $x \in \mathbb{R}$, and let $y_x(t)$, $t \in \mathbb{R}$, be the function defined in (3.8). Then we have the inequalities*

$$(3.13) \quad y_x(t) \geq c^{-1}s^2(x) \quad \text{if} \quad |t - x| \leq s(x),$$

$$(3.14) \quad \|y_x\|_p \geq c^{-1}s(x)^{2+1/p}, \quad x \in \mathbb{R}, \quad p \in [1, \infty).$$

Proof. Now we use (3.11), (3.2), (3.3) and (2.14):

$$\begin{aligned} y_x(t) &= u(t) \int_{x-s(x)}^t v(\xi) d\xi + v(t) \int_t^{x+s(x)} u(\xi) d\xi \\ &= \rho(x) \left\{ \frac{u(t)}{u(x)} \int_{x-s(x)}^t \frac{v(\xi)}{v(x)} d\xi + \frac{v(t)}{v(x)} \int_t^{x+s(x)} \frac{u(\xi)}{u(x)} d\xi \right\} \\ &\geq \rho(x)s(x)c^{-1} \geq c^{-1}d(x)s(x) \geq c^{-1}s^2(x) \Rightarrow (3.13). \end{aligned}$$

Estimate (3.14) follows from (3.13):

$$\|y_x\|_p \geq \left[\int_{x-s(x)}^{x+s(x)} |y_x(t)|^p dt \right]^{1/p} \geq c^{-1}s(x)^{2+1/p}.$$

$\square$

Let us now go to the proof of the theorem. Let $p \in (1, \infty)$, and let $W_1^{(2)}(\mathbb{R}, q) \hookrightarrow L_p(\mathbb{R})$. According to the facts proven above, we have (2.1), the functions d , s and y_x (see (3.8)) are defined, and $y_x \in W_1^{(2)}(\mathbb{R}, q)$ in view of (3.10). This implies that

$$cs(x) \geq \|(y_x)''\|_1 + \|qy_x\|_1 \geq c^{-1}\|y_x\|_p \geq c^{-1}s(x)^{2+1/p}.$$

Hence $s(x) \leq c < \infty$, $x \in \mathbb{R}$, and therefore $d_0 < \infty$ (see (3.3), (2.16)). Then $q_0(a) > 0$ for some $a \in (0, \infty)$ (see Theorem 2.5).

Proof of Theorem 1.2. Sufficiency.

Lemma 3.5. *Suppose conditions (1.2) and (2.1) hold. Then for every $x \in \mathbb{R}$, we have the equality*

$$(3.15) \quad 4\sqrt{\rho(x)} = \int_{-\infty}^{\infty} \frac{G(x, \xi) d\xi}{\rho(\xi)^{3/2}}.$$

Proof. The following elementary relations follow from Lemma 2.1:

$$\begin{aligned}
 \left(\sqrt{\frac{v(\xi)}{u(\xi)}} \right)' &= \frac{1}{2} \sqrt{\frac{u(\xi)}{v(\xi)}} \frac{v'(\xi)u(\xi) - u'(\xi)v(\xi)}{u^2(\xi)} \\
 &= \frac{1}{2\sqrt{v(\xi)}} \frac{1}{u(\xi)^{3/2}} = \frac{1}{2} \frac{v(\xi)}{\rho(\xi)^{3/2}}, \quad \xi \in \mathbb{R} \\
 \Rightarrow \sqrt{\frac{v(x)}{u(x)}} - \sqrt{\frac{v(t)}{u(t)}} &= \frac{1}{2} \int_t^x \frac{v(\xi)d\xi}{\rho(\xi)^{3/2}}, \quad t \leq x \\
 \Rightarrow \sqrt{\frac{v(x)}{u(x)}} &= \sqrt{\frac{v(x)}{u(x)}} - \lim_{t \rightarrow -\infty} \sqrt{\frac{v(t)}{u(t)}} = \frac{1}{2} \int_{-\infty}^x \frac{v(\xi)d\xi}{\rho(\xi)^{3/2}} \\
 \Rightarrow \sqrt{\rho(x)} &= \frac{1}{2} u(x) \int_{-\infty}^x \frac{v(\xi)d\xi}{\rho(\xi)^{3/2}} = \frac{1}{2} \int_{-\infty}^x \frac{G(x, \xi)d\xi}{\rho(\xi)^{3/2}}, \quad x \in \mathbb{R}.
 \end{aligned}$$

Similarly, one can check the equality

$$\sqrt{\rho(x)} = \frac{1}{2} \int_x^\infty \frac{G(x, \xi)d\xi}{\rho(\xi)^{3/2}}, \quad x \in \mathbb{R}.$$

The latter two relations imply (3.15). $\square$

Let us introduce the notation

$$(3.16) \quad H_p = \sup_{x \in \mathbb{R}} \int_{-\infty}^\infty G(x, t)^p dt, \quad p \in [1, \infty).$$

Lemma 3.6. *Suppose conditions (1.2) and (2.1) hold. Then we have the inequalities (see (2.16))*

$$(3.17) \quad c^{-1}d_0^2 \leq H_1 \leq cd_0^2.$$

Proof. Let $x \in \mathbb{R}$. Now we use Lemmas 2.2, 3.2, and the estimate (3.13):

$$\begin{aligned}
 \int_{-\infty}^\infty G(x, t)dt &\geq \int_{x-s(x)}^{x+s(x)} G(x, t)dt = u(x) \int_{x-s(x)}^x v(t)dt + v(x) \int_x^{x+s(x)} u(t)dt \\
 &\geq c^{-1}s^2(x) \geq c^{-1}d^2(x) \quad \Rightarrow \quad H_1 \geq c^{-1}d_0^2.
 \end{aligned}$$

In particular, if $d_0 = \infty$, then from the facts proven above it follows that $H_1 = \infty$, and therefore in this case the upper estimate in (3.17) also holds. Now let $d_0 < \infty$. In the following relations, we use (3.15) and (2.14):

$$\begin{aligned}
 4\sqrt{d_0} &\geq 4\sqrt{d(x)} \geq c^{-1}\sqrt{\rho(x)} = \frac{c^{-1}}{4} \int_{-\infty}^\infty \frac{G(x, \xi)d\xi}{\rho(\xi)^{3/2}} \\
 &\geq c^{-1} \int_{-\infty}^\infty \frac{G(x, \xi)d\xi}{d(\xi)^{3/2}} \geq \frac{c^{-1}}{d_0^{3/2}} \int_{-\infty}^\infty G(x, \xi)d\xi, \quad x \in \mathbb{R}.
 \end{aligned}$$

Hence $H_1 \leq cd_0^2$, as required. $\square$

Corollary 3.7. *Suppose conditions (1.2) and (2.1) hold, and let $p \in (1, \infty)$. Then we have the inequalities*

$$(3.18) \quad c^{-1}(p)d_0^{p+1} \leq H_p \leq c(p)d_0^{p+1}.$$

Proof. The lower estimate in (3.18) is proved exactly as the lower estimate in (3.17). Here, if $d_0 = \infty$, then also $H_p = \infty$, and then the upper estimate in (3.18) also holds. Now let $d_0 < \infty$. The following relations are based on Lemma 2.1, (2.14) and (3.17):

$$\begin{aligned} \int_{-\infty}^{\infty} G(x, t)^p dt &= u^p(x) \int_{-\infty}^x v^p(t) dt + v^p(x) \int_x^{\infty} u(t)^p dt \\ &\leq u^p(x) v^{p-1}(x) \int_{-\infty}^x v(t) dt + v(x)^p u(x)^{p-1} \int_x^{\infty} u(t) dt \\ &= \rho(x)^{p-1} \int_{-\infty}^{\infty} G(x, t) dt \leq c_p d(x)^{p-1} d_0^2 \leq c(p) d_0^{p+1}. \end{aligned}$$

This gives $H_p \leq c(p) d_0^{p+1}$. $\square$

Let us now go to the proof of the theorem. Let $q_0(a) > 0$ for some $a \in (0, \infty)$, and let $y \in W_1^{(2)}(\mathbb{R}, q)$. Denote

$$(3.19) \quad f(x) = -y''(x) + q(x)y(x), \quad x \in \mathbb{R}.$$

Then, clearly, $f \in L_1(\mathbb{R})$ since

$$(3.20) \quad \|f\|_1 \leq \|y''\|_1 + \|qy\|_1 < \infty.$$

Besides, since $W_1^{(2)}(\mathbb{R}, q) \hookrightarrow L_1(\mathbb{R})$ (see Theorems 2.5 and 2.6), in view of the embedding, we have

$$(3.21) \quad \|y\|_1 \leq c\{\|y''\|_1 + \|qy\|_1\} < \infty.$$

Let us now consider the equation

$$(3.22) \quad -\tilde{y}'' + q(x)\tilde{y}(x) = f(x), \quad x \in \mathbb{R},$$

where the function f is defined according to (3.19). By Theorem 2.5, equation (2.11) is correctly solvable in the space $L_1(\mathbb{R})$, and therefore (3.22) has a unique solution $\tilde{y} \in L_1(\mathbb{R})$, and

$$(3.23) \quad \|\tilde{y}\|_1 \leq c\|f\|_1 \leq c\{\|y''\|_1 + \|qy\|_1\} < \infty.$$

From (3.22) and (3.19) it follows that the function

$$z(x) = \tilde{y}(x) - y(x), \quad x \in \mathbb{R},$$

is a solution of (2.2), and $z \in L_1(\mathbb{R})$ in view of (3.21) and (3.23). Hence $z \equiv 0$ by Lemma 2.1 and $y \equiv \tilde{y}$. Then from (2.18) we get

$$(3.24) \quad y(x) \equiv \tilde{y}(x) = \int_{-\infty}^{\infty} G(x, t) f(t) dt, \quad x \in \mathbb{R},$$

where the function f is defined by (3.19). Since $f \in L_1(\mathbb{R})$ (see (3.20)) and, by Theorem 2.7, Corollary 3.7 and Theorem 2.5, the operator $G : L_1(\mathbb{R}) \rightarrow L_p(\mathbb{R})$ is bounded, we obtain from (3.24) that

$$\|y\|_p = \|Gf\|_p \leq \|G\|_{L_1(\mathbb{R}) \rightarrow L_p(\mathbb{R})} \cdot \|f\|_1 \leq c(p) d_0^{1+1/p} \{\|y''\|_1 + \|qy\|_1\},$$

as required. $\square$

REFERENCES

- [1] N. Chernyavskaya and L. Shuster, *Estimates for the Green function of a general Sturm-Liouville operator and their applications*, Proc. Amer. Math. Soc. **127** (1999), 1413-1426. MR1625725 (99h:34040)
- [2] N. Chernyavskaya and L. Shuster, *Regularity of the inversion problem for the Sturm-Liouville equation in the spaces L_p* , Methods and Applications of Analysis **7** (1) (2000), 65-84. MR1796006 (2001i:34010)
- [3] N. Chernyavskaya and L. Shuster, *A criterion for correct solvability of the Sturm-Liouville equation in the space $L_p(R)$* , Proc. Amer. Math. Soc. **130** (2001), 1043-1054. MR1873778 (2002j:34040)
- [4] N. A. Chernyavskaya and L. Shuster, *An embedding theorem for a weighted space of Sobolev type and correct solvability of the Sturm-Liouville equation*, Czechoslovak Math. J. **62** (2012), no. 3, 709-716. MR2984630
- [5] L.W. Kantorovich and G.P. Akilov, *Functional Analysis*, Nauka, Moscow, 1977. MR0511615 (58:23465)
- [6] K. Mynbaev and M. Otelbaev, *Weighted Function Spaces and the Spectrum of Differential Operators*, Nauka, Moscow, 1988. MR950172 (89h:46036)
- [7] M. Otelbaev, *A criterion for the resolvent of a Sturm-Liouville operator to be a kernel*, Math. Notes **25** (1979), 296-297 (translation of Mat. Zametki). MR534299 (80k:34037)

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, BEN-GURION UNIVERSITY OF THE NEGEV, P.O. BOX 653, BEER-SHEVA, 84105, ISRAEL

DEPARTMENT OF MATHEMATICS, BAR-ILAN UNIVERSITY, 52900 RAMAT GAN, ISRAEL
E-mail address: `miriam@macs.biu.ac.il`