ON THE HAUSDORFF DIMENSION OF JULIA SETS OF SOME REAL POLYNOMIALS

GENADI LEVIN AND MICHEL ZINSMEISTER

(Communicated by Mario Bonk)

ABSTRACT. We show that the supremum for c real of the Hausdorff dimension of the Julia set of the polynomial $z \mapsto z^d + c$ (d is an even natural number) is greater than $2d/(d+1)$.

1. INTRODUCTION AND STATEMENT OF THE RESULT

The Julia set of a non-linear polynomial $P : \mathbf{C} \rightarrow \mathbf{C}$ is the set of points having no neighborhood on which the family of iterates (P^n) is normal. It is a compact non-empty set which is, except for very special polynomials P , a fractal set. For $c \in \mathbf{C}$ we denote by f_c the polynomial

$$f_c(z) = z^d + c,$$

where $d \geq 2$ is an even integer number, which we fix. Denote by J_c the Julia set of f_c .

In the quadratic case $d = 2$, the values of c for which the Hausdorff dimension (HD) of J_c is big has attracted a lot of attention. It has been known since the pioneering work of Douady-Hubbard [3] that the Hausdorff dimension of the Julia set is less than 2 for every hyperbolic polynomial. Thus $HD(J_c) < 2$ outside ∂M and the (hypothetic) non-hyperbolic components of (the interior) M . We recall that M stands for the Mandelbrot set, that is, the compact subset of $c \in \mathbf{C}$ such that J_c is connected. Shishikura [9] was the first to find quadratic Julia sets with Hausdorff dimension 2. He indeed proved that this property holds on a dense $\mathcal{G}_\delta$ subset of ∂M or even on a dense $\mathcal{G}_\delta$ subset on the boundary of every hyperbolic component of M . More recently, Buff and Chéritat [12] have found quadratic Julia sets with positive Lebesgue measure (see also <http://annals.math.princeton.edu/articles/3682>).

Both Shishikura's and Buff-Chéritat's results are based on the phenomenon of parabolic implosion which was discovered and studied by Douady-Hubbard [3]. It should be pointed out that Buff-Chéritat's result is very involved and that we will make no use of it. There is no doubt that if they exist, values of $c \in \mathbf{R}$ such that the Julia set J_c has positive measure must be as hard to find as those of Buff-Chéritat's. The aim of the present note is much more modest. Its starting point is the second author's re-reading of Shishikura's result [10]: it states that if one implodes a polynomial with a parabolic cycle having q petals, then the dimension

Received by the editors November 23, 2010 and, in revised form, October 18, 2011 and January 4, 2012.

2010 *Mathematics Subject Classification*. Primary 37F10.

The authors were supported in part by the IMPAN-BC European Community Centre of Excellence and by the Marie Curie European network CODY.

of its Julia set automatically gets bigger than $2q/(q+1)$. Shishikura's result follows from this by a Baire argument.

Very little is known about Hausdorff dimension of J_c for real c . In particular, it is not known if, for a given degree d ,

$$\sup\{HD(J_c), c \in \mathbf{R}\} = 2.$$

Possible candidates for high dimension are of course (just look at them!) infinitely renormalizable polynomials, but the analysis seems to be very delicate, and at least no result concerning dimension 2 has been proven so far (for results in the opposite direction, see [1]). It is unknown for example if the Julia set of the Feigenbaum polynomial has dimension 2 or not (see [4]-[5] for the Julia set of the Feigenbaum universal map). The only known result about this set is a very general result of Zdunik [14]: it has dimension bigger than 1.

If one tries to use the same ideas as in [9] for real polynomials f_c , one immediately faces the problem that if f_c , for c real, has a parabolic cycle that may be imploded along the real axis, then the number of petals is 1 and this does not imply more than Zdunik's general result. The only trick of this paper is to make use of a virtual doubling of petals when the critical point is mapped to a parabolic point (by Lavaurs map). It was inspired by the Douady et al. paper [13] and implies the following theorem, which is the main result of this work.

Theorem 1. *Let $f_c(z) = z^d + c$, d even. Let N be the set of parameters $c \in \mathbf{R}$, such that f_c has a parabolic cycle of period at least 2 and multiplier 1. Then there exists an open set Y of $\mathbf{R}$ whose closure contains N such that J_c is connected and*

$$HD(J_c) > \frac{2d}{d+1},$$

for every $c \in Y$.

Comment 1. In fact, we prove a stronger statement: hyperbolic dimension [9] of J_c is bigger than $2d/(d+1)$.

Comment 2. By [11] ([6], [8] for $d=2$), the set of real c such that f_c is hyperbolic is dense in $\mathbf{R}$. In particular, hyperbolic parameters c are dense in Y .

2. PROOF OF THEOREM 1

We fix an even integer $d \geq 2$ and consider the family $f_c(x) = x^d + c$, for real c . Then the Julia set J_c is connected if and only if $c \in [a, b]$, where $a < 0$ is such that $f_a^2(0)$ is a fixed point and $b > 0$ is such that f_b has a fixed point with multiplier 1. It is sufficient to prove that, given $c_0 \in (a, b)$ such that f_{c_0} has a neutral cycle of period $k > 1$ with multiplier 1, that is, with one petal, there is an open set Z accumulating at c_0 for which $HD(J_c) > 2d/(d+1)$ for $c \in Z$.

We begin with three pictures. The first two illustrate how to choose the parameter (for $d=2$) and how the corresponding Julia set looks. The third has kindly been drawn for us by the referee and shows the case $d=4$.

Since $c_0 \in (a, b)$, the set J_c is connected for c in a small neighborhood U of c_0 . It is known that f_c has an attracting periodic orbit of period k for $c \in U$ on the left side of c_0 . Since $k > 1$, the corresponding filled-in Julia set K_{c_0} is such that its interior has a component Δ containing 0 and has infinitely many preimages of Δ . The boundary $\partial\Delta$ contains a parabolic point α of period k . In particular, there is a sequence of preimages of Δ which intersect the real line and accumulate at α .

FIGURE 1. Choice of the parameter

FIGURE 2. The corresponding Julia set

FIGURE 3. The case $d = 4$

For definiteness, one can assume that $\alpha > 0$. Then $F = f_{c_0}^k$ has the following local form:

$$(1) \quad F(z) = z + a(z - \alpha)^2 + b(z - \alpha)^3 + \dots,$$

where $a > 0$. This implies that a parabolic implosion phenomenon occurs as $\epsilon \rightarrow 0$, $\epsilon > 0$, for the maps $f_{c_0+\epsilon}$.

At this point we digress somewhat and briefly describe the theory of parabolic implosion.

Let $\delta > 0$ be very small and $D_{\pm}$ be the disks centered at $\alpha \pm \delta$ with radius δ . The map F sends D_- into itself, while $D_+ \subset F(D_+)$. This defines, after identification of z with $F(z)$ at the boundary, two cylinders $U_- = D_- \setminus F(D_-)$ and $U_+ = F(D_+) \setminus D_+$. The fact that $U_{\pm}$ are actual cylinders is best seen in the approximate Fatou coordinate $I : z \mapsto -1/(a(z - \alpha))$ which sends α to ∞ and conjugates F to a map which is asymptotically the translation by 1 at ∞ :

$$(2) \quad F_{\infty}(w) = I \circ F \circ I^{-1}(w) = w + 1 + \frac{A}{w} + O\left(\frac{1}{|w|^2}\right),$$

where $A = 1 - b/a^2$. The real number A is a conformal invariant. In the case of a real polynomial F which has a parabolic fixed point α with multiplier 1 and with a single critical point in its immediate basin Δ , it is known [9] that $A > 0$.

By the Riemann mapping theorem these two cylinders may be uniformized by “straight” cylinders. In other words, there exists $\varphi_{\pm}$ mapping the cylinders $U_{\pm}$ to vertical strips $V_{\pm}$ of width 1 conjugating F to the translation by 1. For further use we notice that, by symmetry, c_0 being real, we may assume that $\varphi_{\pm}(\bar{z}) = \overline{\varphi_{\pm}(z)}$.

FIGURE 4. Fatou coordinates

We also notice that these maps are unique up to post-composition by a real translation. These two maps are called respectively repelling (+) and attracting (-) Fatou coordinates. We normalize them as follows. For every $\kappa > 0$ and $R > 0$, consider two sectors $\Sigma_-(\kappa, R) = \{w : \operatorname{Re}(w) > R - \kappa|\operatorname{Im}(w)|\}$, $\Sigma_+(\kappa, R) = \{w : \operatorname{Re}(w) < -R + \kappa|\operatorname{Im}(w)|\}$. Then for any $\kappa > 0$ there is a big enough $R(\kappa)$ such that if we introduce two sectors $\Sigma_\pm(\kappa) = \Sigma_\pm(\kappa, R(\kappa))$, then $\varphi_\pm = \Phi_\pm \circ I$, where

$$(3) \quad \Phi_\pm(w) = w - A \log_\pm(w) + C_\pm + o(1)$$

as $w \rightarrow \infty$ within $\Sigma_\pm(\kappa)$ respectively. We specify the constants $C_\pm$ and the log-branches in such a way that $\Phi_\pm$ are real for real w . Namely, we choose $C_- = 0$ and $\log_-$ to be the standard log-branch in the slit plane $\mathbf{C} \setminus \{\mathbf{x} \leq \mathbf{0}\}$. In turn, let $C_+ = iA\pi$ and $\log_+$ be a branch of $\log$ in $\mathbf{C} \setminus \{\mathbf{x} \geq \mathbf{0}\}$ so that $\log_+(w) = \log|w| + i\pi$ for $w < 0$.

Now, we extend $\varphi_\pm$ in the following way. Since

$$(4) \quad \varphi_-(F(z)) = T_1(\varphi_-(z)),$$

where T_σ denotes the translation by σ , and since every orbit converging to α passes through U_- exactly once, φ_- extends uniquely to Δ to a holomorphic function still satisfying (4).

It is seen from (4) that the map $\varphi_- : \Delta \rightarrow \mathbf{C}$ is a branched covering, with the critical points at 0 and all its preimages in Δ by F^n , $n > 0$, and the critical values at the real numbers $\varphi_-(0) - n$, $n \geq 0$. In particular, there exists a simply-connected domain $\Omega_- \subset \Delta \cap \mathbf{H}^+$, where $\mathbf{H}^+$ is the upper half-plane, such that $\varphi_- : \Omega_- \rightarrow \mathbf{H}^+$

is a holomorphic homeomorphism. Moreover, the intersection of the boundary of Ω_- with $\mathbf{R}$ is the interval $(0, \alpha)$.

Concerning the repelling Fatou coordinate it is best to consider $\psi_+ = \varphi_-^{-1} : V_+ \rightarrow U_+$. The functional relation is now

$$(5) \quad \psi_+(T_1(z)) = F(\psi_+(z)),$$

and we can extend ψ_+ to an entire function by putting, for $n \in \mathbf{Z}$, $\psi_+(T_n(z)) = F^n(\psi_+(z))$ for $z \in V_+$. There exists a simply-connected domain $\Omega_+ \subset \mathbf{H}^+$ such that $\psi_+ : \Omega_+ \rightarrow \mathbf{H}^+$ is a homeomorphism.

Now let σ be a real number. We define the Lavaurs map g_σ on the component Δ of the interior of the filled-in Julia set of f_{c_0} by

$$g_\sigma = \psi_+ \circ T_\sigma \circ \varphi_-.$$

The raison d'être of this definition is the following theorem due to Douady and Lavaurs [2]:

Theorem 2. *There exists a sequence of positive ϵ_n converging to 0 and a sequence of positive integers N_n such that*

$$(6) \quad g_\sigma(z) = \lim_{n \rightarrow \infty} f_{c_0 + \epsilon_n}^{kN_n}(z)$$

uniformly on compact sets of Δ .

Using (3) with the constants $C_\pm$ and the log-branches specified above, it is easy to get that, for every κ , if w tends to ∞ in $\Sigma(\kappa) := \Sigma_-(\kappa) \cap \Sigma_+(\kappa) \cap \mathbf{H}^+$, then $g_\sigma(w) := I \circ g_\sigma \circ I^{-1}(w) = w + (\sigma - iA\pi) + O(\frac{1}{|w|})$, where A is real and positive. Therefore, for every real σ and every $\kappa > \kappa(\sigma)$, the inverse map g_σ^{-1} leaves the sector $\Sigma(\kappa)$ invariant and $g_\sigma^{-n} \rightarrow \infty$ as $n \rightarrow \infty$. Coming back to the z -plane, we conclude that the branch $G = I^{-1} \circ g_\sigma^{-1} \circ I$ of g_σ^{-1} leaves the set $S(\kappa) = I^{-1}(\Sigma(\kappa))$ invariant and $G^n(z) \rightarrow \alpha$ as $n \rightarrow \infty$, for $z \in S(\kappa)$ and every $\kappa > \kappa(\sigma)$. We have, for $w \in \Sigma(\kappa)$,

$$(7) \quad G_\infty(w) := I \circ G \circ I^{-1}(w) = w + (-\sigma + iA\pi) + O(\frac{1}{|w|}).$$

Now, from the definition of $g_\sigma(z)$ and the global properties of the maps φ_- and ψ_+ , there follows the existence of a simply-connected domain $\Omega_0 \subset \Omega_-$, which is mapped by g_σ homeomorphically onto $\mathbf{H}^+$ and such that $\alpha \in \bar{\Omega}_0$. Moreover, from the above description, $S(\kappa) \subset \Omega_0$, for every $\kappa > \kappa(\sigma)$. Therefore, the branch G of g_σ^{-1} which is defined above extends to a global univalent branch $G : \mathbf{H}^+ \rightarrow \Omega_0$ of g_σ^{-1} . Since $\Omega_0 \subset \mathbf{H}^+$, the iterates $G^n : \mathbf{H}^+ \rightarrow \Omega_0$, $n > 0$, converge uniformly on compact sets in $\mathbf{H}^+$ to a unique fixed point in $\bar{\Omega}_0$, which must be α .

Let us consider the continuous map $\sigma \mapsto g_\sigma(0) = \psi_+(\varphi_-(0) + \sigma)$: if σ runs in the interval $I = \{\varphi_+(x) - \varphi_-(0) : x \in U_+ \cap \mathbf{R}\}$, then $g_\sigma(0)$ runs over $U_+ \cap \mathbf{R}$. It is thus clear, and this is the key point in the proof, that we can choose σ in such a way that $g_\sigma(0)$ is a preimage of α : there is $j \geq 1$ such that $f_{c_0}^j \circ g_\sigma(0) = \alpha$. Since 0 is a critical point for g_σ , taking the inverse image by $f_{c_0}^j \circ g_\sigma$ has the same effect as multiplying the number of petals by d , and we may state:

Lemma 2.1. *There exists an infinite iterated function system defined on a small compact neighborhood B_0 of zero and generated by some holomorphic branches of $f_{c_0}^{-1}$ and g_σ^{-1} such that its limit set has Hausdorff dimension bigger than $2d/(d+1)$.*

Comment 3. In fact, the limit set is a subset of a so-called Julia-Lavaurs set denoted by $J_{c_0, \sigma}$. It is defined as follows. The map g_σ can be extended in a natural way from Δ to the interior of the filled-in Fatou set of f_{c_0} : if $f_{c_0}^k(z) \in \Delta$, set $g_\sigma(z) = g_\sigma \circ f_{c_0}^k(z)$. Then $J_{c_0, \sigma}$ is simply the closure of the set of points z for which there exists $m \in \mathbf{N}$ such that $g_\sigma^m(z)$ is defined and belongs to $J(f_0)$.

This lemma together with the above discussion implies the theorem. Indeed, by a general property of iterated function systems [7], there exists its finite subsystem with the Hausdorff dimension of its limit set bigger than $2d/(d+1)$. On the other hand, the finite iterated function system persists for $f_{c_0+\epsilon}$ by (6). To be more precise, if $\{I_j : B_0 \rightarrow X_j, 1 \leq j \leq j_0\}$ is this finite iterated function system, then each I_j can be extended to a univalent map to a fixed neighborhood Y of B_0 as $I_j : Y \rightarrow Y_j$. Consider the inverse univalent map $I_j^{-1} : Y_j \rightarrow Y$. Since the convergence in (6) is uniform on compacts in Δ , for every ϵ_n small enough there is some integer $N_j > 0$ and a compact set $X_{j,n}$ so that $f_{c_0+\epsilon_n}^{N_j} : X_{j,n} \rightarrow B_0$ is univalent too. Now it is clear that, for every ϵ_n small enough, the non-escaping set K_n of the dynamical system which consists of finitely many maps $f_{c_0+\epsilon_n}^{N_j} : X_{j,n} \rightarrow B_0$, $1 \leq j \leq j_0$, has the Hausdorff dimension which is bigger than $2d/(d+1)$. On the other hand, K_n must lie in the Julia set of $f_{c_0+\epsilon_n}$ because some iterate of the map $f_{c_0+\epsilon_n}^{N_1 N_2 \dots N_{j_0}}$ leaves the set K_n invariant and is expanding on it.

Proof of Lemma 2.1. As the first step, let us fix a small enough closed ball B_0 around zero so that it does not contain points of the postcritical set of f_{c_0} . There exists its preimage B' by F^{-1} in $\Delta \cap \mathbf{H}^+$. Then we can apply to B' the maps G^n , $n > 0$. By the above, $B'_n = G^n(B')$ are pairwise disjoint, compactly contained in Δ , and $B'_n \rightarrow \alpha$ as $n \rightarrow \infty$. Now, for every $n \geq n_0$, so that B'_n lies in a small enough neighborhood U of α , we make “clones” of B'_n in $U \cap \Delta$, applying to it F^r , $r \in \mathbf{Z}$, where F^r for $r < 0$ is a well-defined in $U \cap \Delta$ branch which fixes α . We obtain the sets $B'_{n,r} = F^r(B'_n)$. In the second step, we consider the map $f_{c_0}^j \circ g_\sigma$ from a neighborhood V of 0 onto U . This map is a ramified cover with the only ramification point at 0 of order d . Let $U^* = U \setminus \{x \geq \alpha\}$ and $V^* = V \cap \{z : \text{Arg}(z) \in (0, 2\pi/d)\}$. Denote by h a branch of $(f_{c_0}^j \circ g_\sigma)^{-1}$ from U^* onto V^* . Let $B_{n,r} = h(B'_{n,r})$. We obtain a system of holomorphic maps $\Psi = \{\psi_{n,r} : B_0 \rightarrow B_{n,r}\}$, where $\psi_{n,r} = h \circ F^r \circ G^n \circ F^{-1}$, $r \in \mathbf{Z}$, $n \geq n_0$. If the neighborhood U is chosen small enough, the maps $\psi_{n,r}$ extend to univalent maps in a fixed neighborhood $\tilde{B}$ of B_0 into itself. In particular, the compact sets $B_{n,r}$ are pairwise disjoint and compactly contained in B_0 . Now, it is quite standard to check that Ψ forms a conformal infinite iterated function system in the sense of [7] (strictly speaking, in the hyperbolic metric of $\tilde{B}$, which is equivalent to the Euclidean one on B_0). Let us calculate the parameter $\theta = \inf\{t : \psi(t) < \infty\}$ of Ψ , where $\psi(t) = \sum_{(n,r)} \max_{z \in B_0} |\psi'_{n,r}(z)|^t$. The map $\psi_{n,r} = h \circ I^{-1} \circ F_\infty^r \circ G_\infty^n \circ I \circ F^{-1}$. Here F^{-1} is a univalent map of a neighborhood $\tilde{B}$ of B_0 into Δ . Now, routine and well-known calculations based on (2)-(7) show (see e.g. [10]) that, for all $r \in \mathbf{Z}$, $n \geq n_0$ and some C , which depends only on a compact set in $\mathbf{H}^+ \cap \Delta$, from which w is taken, $C^{-1}|r + (-\sigma + i\pi A)n| \leq |F_\infty^r \circ G_\infty^n(w)| \leq C|r + (-\sigma + i\pi A)n|$ and $C^{-1} \leq |(F_\infty^r \circ G_\infty^n)'(w)| \leq C$. On the other hand, the map h is a composition of a univalent map with an inverse branch of $z^{1/d}$. This gives us $C_1^{-1}|r + (-\sigma + i\pi A)n|^{-1-1/d} \leq |\psi'_{n,r}(z)| \leq C_1|r + (-\sigma + i\pi A)n|^{-1-1/d}$, for some

C_1 and every $z \in B_0$. It follows that the series for $\psi(t)$ converges if and only if $t > \theta = 2d/(d+1)$ and $\psi(\theta) = \infty$. Hence [7], the Hausdorff dimension of the limit set of Ψ is strictly bigger than $2d/(d+1)$. $\square$

ACKNOWLEDGMENT

This work was done during the first author's one-month visit at the University of Orléans in 2008.

REFERENCES

1. A. Avila and M. Lyubich, *Examples of Feigenbaum Julia sets with small Hausdorff dimension*, Dynamics on the Riemann Sphere, in honor of B. Branner **1** (2006), 71–87. MR2348955 (2008j:37102)
2. A. Douady, *Does a Julia set depend continuously on the polynomial?*, Proceedings of Symposia in Applied Mathematics **49** (1994), 91–135. MR1315535
3. A. Douady and J. Hubbard, *Etude dynamique des Polynômes complexes: Première partie*, Publications Mathématiques d'Orsay **84-02** (1984), 1–75. MR762431 (87f:58072a)
4. G. Levin and G. Świątek, *Hausdorff dimension of Julia sets of Feigenbaum type*, Communications in Mathematical Physics **258** (2005), 135–148. MR2166843 (2007b:37104)
5. ———, *Measure of the Julia set of the Feigenbaum map with high criticality*, Ergodic Theory and Dynamical Systems **30** (2009), 855–875. MR2643714 (2011f:37066)
6. J. Graczyk and G. Świątek, *Generic hyperbolicity in the logistic family*, Annals of Mathematics **146** (1997), 1–52. MR1469316 (99b:58079)
7. R. D. Mauldin and M. Urbanski, *Dimension and measure in infinite iterated function systems*, Proc. London Math. Soc. **73** (1996), 105–154. MR1387085 (97c:28020)
8. M. Lyubich, *Dynamics of quadratic polynomials*, Acta Mathematica **178** (2007), 185–297. MR1459261 (98e:58145)
9. M. Shishikura, *The Hausdorff dimension of the boundary of the Mandelbrot set and Julia sets*, Annals of Mathematics **147** (1998), 225–267. MR1626737 (2000f:37056)
10. M. Zinsmeister, *Fleur de Leau-Fatou et dimension de Hausdorff*, Comptes-rendus de l'Académie des Sciences **326** (1998), 1227–1232. MR1650223 (99j:58180)
11. O. Kozlovsky, W. Shen, and S. van Strien, *Density of hyperbolicity in dimension one*, Annals of Mathematics **166** (2007), 145–182. MR2342693 (2008j:37081)
12. X. Buff and A. Chéritat, *Ensembles de Julia quadratiques de mesure strictement positive*, Comptes-rendus de l'Académie des Sciences **341** (2005), 669–674. MR2183346 (2006k:37133)
13. X. Buff, R. Devaney, A. Douady, and P. Sentenac, *Baby Mandelbrot sets are born in cauliflowerers*, London Mathematical Society Lecture Notes Series **274** (2000), 19–36. MR1765083 (2001g:37062)
14. Anna Zdunik, *Parabolic orbifolds and the dimension of the maximal measure for rational maps*, Inventiones Mathematicae **99** (1990), 627–649. MR1032883 (90m:58120)

INSTITUTE OF MATHEMATICS, HEBREW UNIVERSITY OF JERUSALEM, JERUSALEM 91904, ISRAEL

MAPMO, UNIVERSITÉ D'ORLÉANS, BP 6759, 45067 ORLÉANS CEDEX 2, FRANCE