INVERTIBLE WEIGHTED SHIFT OPERATORS WHICH ARE m -ISOMETRIES

MUNEO CHÔ, SCHÔICHI ÔTA, AND KÔTARÔ TANAHASHI

(Communicated by Richard Rochberg)

ABSTRACT. For a bounded linear operator T on a complex Hilbert space $\mathcal{H}$, let $\Delta_{T,m} = \sum_{k=0}^m (-1)^k \binom{m}{k} T^{*m-k} T^{m-k}$ for $m \in \mathbb{N}$. T is called an m -isometry if $\Delta_{T,m} = 0$. In this paper, for every even number m , we give an example of invertible $(m+1)$ -isometry which is not an m -isometry. Next we show that if T is an m -isometry, then the operator $\Delta_{T,m-1}$ is not invertible.

1. INTRODUCTION

J. Agler and M. Stankus published excellent papers about m -isometric operators, [1], [2] and [3]. They showed that m -isometries have interesting spectral properties. For example, if T is an m -isometry, then the approximate point spectrum of T lies on the unit circle. In [10], Patel showed that Weyl's theorem holds for a 2-isometry. Applying Uchiyama and Tanahashi's result [11], in [8] we showed that if T is an m -isometry, then T has the single valued extension property. Let $\mathcal{H}$ be a complex Hilbert space and $B(\mathcal{H})$ be a set of all bounded linear operators on $\mathcal{H}$. Let $\binom{m}{k}$ be the binomial coefficient. For an operator $T \in B(\mathcal{H})$, let

$$\Delta_{T,m} = \sum_{k=0}^m (-1)^k \binom{m}{k} T^{*m-k} T^{m-k}.$$

T is said to be an m -isometry if $\Delta_{T,m} = 0$. Agler and Stankus proved that if T is an m -isometry, then $\Delta_{T,m-1} \geq 0$ (Proposition 1.5, [1]). It is easy to see that $T^* \Delta_{T,m} T - \Delta_{T,m} = \Delta_{T,m+1}$. Hence it holds that if T is an m -isometry, then T is an $(m+1)$ -isometry. In [7], T. Bermudez, A. Martinon and E. Negrin studied characterizations of weighted shift operators which are m -isometries. In [1], Agler and Stankus proved that if m is even and T is an invertible m -isometry, then T is an $(m-1)$ -isometry. In [4] A. Athavale proved that if S is a unilateral weighted shift

Received by the editors October 2, 2011 and, in revised form, January 31, 2012.

2010 *Mathematics Subject Classification*. Primary 47B37.

Key words and phrases. Hilbert space, m -isometry, bilateral weighted shift.

The first author's research was partially supported by Grant-in-Aid for Scientific Research, No. 20540192.

The second author's research was partially supported by Grant-in-Aid for Scientific Research, No. 20540178.

The third author's research was partially supported by Grant-in-Aid for Scientific Research, No. 20540184.

©2013 American Mathematical Society
Reverts to public domain 28 years from publication

$Se_n = w_n e_{n+1}$ with weights $w_n = \sqrt{1 + \frac{m}{n}}$ for $n = 1, 2, 3, \dots$, then S is an $(m+1)$ -isometry which is not an m -isometry. The operator S is not invertible. We have not seen invertible $(m+1)$ -isometries which are weighted shifts for m even. We give an example of an invertible $(m+1)$ -isometry which is not an m -isometry for every even number m . Next we prove that power bounded m -isometries are isometries, and if T is an m -isometry for a natural number $m \geq 2$, then the operator $\Delta_{T,m-1}$ is not invertible.

2. INVERTIBLE WEIGHTED SHIFT OF AN m -ISOMETRY

Agler and Stankus showed the following result.

Proposition A (Proposition 1.23, [1]). *If T is an invertible m -isometry and m is even, then T is an $(m-1)$ -isometry.*

Their proof is fine. We give another proof.

Proof. Since T is an m -isometry, it holds that

$$(2.1) \quad \Delta_{T,m-1} = \sum_{k=0}^{m-1} (-1)^k \binom{m-1}{k} T^{*m-1-k} T^{m-1-k} \geq 0.$$

It is easy to see that T^{-1} also is an m -isometry. Since T^{-1} is an m -isometry, it holds that

$$(2.2) \quad \Delta_{T^{-1},m-1} = \sum_{k=0}^{m-1} (-1)^k \binom{m-1}{k} (T^{-1})^{*m-1-k} (T^{-1})^{m-1-k} \geq 0.$$

Since $m-1$ is an odd number, by (2.2) we have

$$\begin{aligned} -\Delta_{T,m-1} &= \sum_{k=0}^{m-1} (-1)^k \binom{m-1}{k} T^{*k} T^k \\ &= \sum_{k=0}^{m-1} (-1)^k \binom{m-1}{k} T^{*m-1} (T^{-1})^{*m-1-k} (T^{-1})^{m-1-k} T^{m-1} \\ &= T^{*m-1} (\Delta_{T^{-1},m-1}) T^{m-1} \geq 0. \end{aligned}$$

Hence $\Delta_{T,m-1} \leq 0$. By (2.1) we have $\Delta_{T,m-1} = 0$, so the proof is complete. $\square$

For every even number m , we give an example of an invertible $(m+1)$ -isometry T which is not an m -isometry.

Theorem 1. *For any even number m , there exists an invertible $(m+1)$ -isometry T which is not an m -isometry.*

Proof. Let m be any even number and put $\psi(x) = x(x+1)\cdots(x+m-1)$. Then $\psi(x)$ is a polynomial of even degree m . Hence $\psi(n) \geq 0$ for $n = 0, \pm 1, \pm 2, \dots$. So, for any $k > 0$, we have $\psi(n) + k > 0$ for $n = 0, \pm 1, \pm 2, \dots$. Let

$$w_n = \sqrt{\frac{\psi(n+1) + k}{\psi(n) + k}} > 0 \quad \text{for } n = 0, \pm 1, \pm 2, \dots$$

Let $\{e_n\}_{n=-\infty}^{\infty}$ be an orthonormal basis of $\mathcal{H} = \ell^2$ and T be a bilateral weighted shift such that $Te_n = w_n e_{n+1}$. We define

$$\begin{aligned} I_{m,n} &= w_n^2 w_{n+1}^2 \cdots w_{n+m-1}^2 - \binom{m}{1} w_n^2 w_{n+1}^2 \cdots w_{n+m-2}^2 \\ &\quad + \binom{m}{2} w_n^2 w_{n+1}^2 \cdots w_{n+m-3}^2 + \cdots + (-1)^{m-1} \binom{m}{m-1} w_n^2 + (-1)^m. \end{aligned}$$

Note that T is an m -isometry if and only if $I_{m,n} = 0$ for $n = 0, \pm 1, \pm 2, \dots$. To show T is an $(m+1)$ -isometry, we will show that $I_{m+1,n} = 0$ for $n = 0, \pm 1, \pm 2, \dots$. We define a function $f(x)$ as

$$\begin{aligned} f(x) &= x^{n+m-1}(1-x)^{m+1} \\ &= x^{n+m-1} - \binom{m+1}{1} x^{n+m} + \binom{m+1}{2} x^{n+m+1} \\ &\quad + \cdots + (-1)^m \binom{m+1}{m} x^{n+2m-1} + (-1)^{m+1} x^{n+2m}. \end{aligned}$$

Note that $f^{(m)}(1) = 0$. By differentiating m times, we have

$$\begin{aligned} 0 &= f^{(m)}(1) \\ &= (n+m-1)(n+m-2) \cdots n \\ &\quad - \binom{m+1}{1} (n+m)(n+m-1) \cdots (n+1) \\ &\quad + \cdots + (-1)^m \binom{m+1}{m} (n+2m-1)(n+2m-2) \cdots (n+m) \\ &\quad + (-1)^{m+1} (n+2m)(n+2m-1) \cdots (n+m+1). \end{aligned}$$

Hence

$$\begin{aligned} &(-1)^{m+1} I_{m+1,n} (\psi(n) + k) \\ &= (n+m+1)(n+m+2) \cdots (n+2m) + k \\ &\quad - \binom{m+1}{1} \{(n+m)(n+m+1) \cdots (n+2m-1) + k\} \\ &\quad + \cdots + (-1)^m \binom{m+1}{m} \{(n+1)(n+2) \cdots (n+m) + k\} \\ &\quad + (-1)^{m+1} \{n(n+1)(n+2) \cdots (n+m-1) + k\} \\ &= f^{(m)}(1) + k(1-1)^{m+1} = 0. \end{aligned}$$

This implies that $I_{m+1,n} = 0$. Hence T is an $(m+1)$ -isometry.

To show that T is not an m -isometry, we will show that $I_{m,n} \neq 0$ for $n = 0, \pm 1, \pm 2, \dots$. We define a function $g(x)$ as

$$\begin{aligned} g(x) &= x^{n+m-1}(1-x)^m \\ &= x^{n+m-1} - \binom{m}{1} x^{n+m} + \binom{m}{2} x^{n+m+1} \\ &\quad + \cdots + (-1)^{m-1} \binom{m}{m-1} x^{n+2m-2} + (-1)^m x^{n+2m-1}. \end{aligned}$$

Note that $g^{(m)}(1) = m!(-1)^m$. By differentiating m times, we have

$$\begin{aligned} m!(-1)^m &= g^{(m)}(1) \\ &= (n+m-1)(n+m-2)\cdots n \\ &\quad - \binom{m}{1}(n+m)(n+m-1)\cdots(n+1) \\ &\quad + \cdots + (-1)^{m-1} \binom{m}{m-1}(n+2m-2)(n+2m-3)\cdots(n+m-1) \\ &\quad + (-1)^m(n+2m-1)(n+2m-2)\cdots(n+m). \end{aligned}$$

Hence

$$\begin{aligned} &(-1)^m I_{m,n}(\psi(n) + k) \\ &= (n+m)(n+m+1)\cdots(n+2m-1) + k \\ &\quad - \binom{m}{1}\{(n+m-1)(n+m)\cdots(n+2m-2) + k\} \\ &\quad + \cdots + (-1)^{m-1} \binom{m}{m-1}\{(n+1)(n+2)\cdots(n+m) + k\} \\ &\quad + (-1)^m\{n(n+1)(n+2)\cdots(n+m-1) + k\} \\ &= g^{(m)}(1) + k(1-1)^m = m!(-1)^m. \end{aligned}$$

This implies that $I_{m,n} = m!(\psi(n) + k)^{-1} \neq 0$. Hence T is not an m -isometry.

Finally, since $w_n \rightarrow 1$ ($n \rightarrow \pm\infty$), we have $\sigma(T) = \{\lambda : |\lambda| = 1\}$ by Proposition 2.6.8 (b) of [9]. Therefore T is invertible. The proof is complete. $\square$

Remark 1. This result provides an example of an $(m+1)$ -isometry T which is not an m -isometry if we take $Te_n = w_n e_{n+1}$ for $n = 1, 2, \dots$ where

$$w_n = \sqrt{\frac{(n+1)(n+2)\cdots(n+m) + k}{n(n+1)\cdots(n+m-1) + k}}$$

with $0 \leq k$. If $k = 0$, then

$$w_n = \sqrt{\frac{(n+1)(n+2)\cdots(n+m) + 0}{n(n+1)\cdots(n+m-1) + 0}} = \sqrt{\frac{n+m}{n}}.$$

This is the result of A. Athavale, Proposition 8 of [4].

3. SOME PROPERTIES OF m -ISOMETRIES

An operator $T \in B(\mathcal{H})$ is said to be power bounded if there exists a positive number M such that $\|T^n\| \leq M$ for every $n \in \mathbb{N}$.

Theorem 2. *A power bounded m -isometry is an isometry.*

To prove this result, we will use Berberian's method (cf. [6], [12]).

Proposition (Lemma 2.7, [12]). *Let $\mathcal{H}$ be a complex Hilbert space. Then there exist a Hilbert space $\mathcal{H}^\circ \supset \mathcal{H}$ and a unital linear map $\circ : B(\mathcal{H}) \rightarrow B(\mathcal{H}^\circ)$ such that*

- (i) $(ST)^\circ = S^\circ T^\circ, (T^*)^\circ = (T^\circ)^*, \|T\| = \|T^\circ\|,$
- (ii) $S \leq T \implies S^\circ \leq T^\circ,$
- (iii) $\sigma(T) = \sigma(T^\circ), \sigma_a(T) = \sigma_a(T^\circ) = \sigma_p(T^\circ).$

We prepare notation and a lemma. For a unit vector $x \in \mathcal{H}$, assume that $(T^*T - I)x = a_1x$. Let $a_n = \langle \Delta_{T,n}x, x \rangle$ and $b_n = \|T^n x\|^2$ for $n = 1, 2, 3, \dots$. Then we have the following lemma.

Lemma. *For an operator $T \in B(\mathcal{H})$ with the above notation, it holds that*

$$(3.1) \quad b_n = a_n + \binom{n}{n-1} a_{n-1} + \binom{n}{n-2} a_{n-2} + \dots + na_1 + 1.$$

Proof. Note that

$$b_1 = \|Tx\|^2 = \langle T^*Tx, x \rangle = \langle a_1x + x, x \rangle = a_1 + 1.$$

Since

$$\begin{aligned} T^{*2}T^2 &= (T^{*2}T^2 - 2T^*T + I) + 2(T^*T - I) + I \\ &= \Delta_{T,2} + 2\Delta_{T,1} + I, \end{aligned}$$

we have

$$\begin{aligned} b_2 &= \langle T^{*2}T^2x, x \rangle \\ &= \langle \Delta_{T,2}x, x \rangle + 2\langle \Delta_{T,1}x, x \rangle + \langle x, x \rangle \\ &= a_2 + 2a_1 + 1. \end{aligned}$$

Hence (3.1) holds for $n = 1, 2$. Note that

$$\begin{aligned} t^n &= (t-1+1)^n \\ &= (t-1)^n + \binom{n}{1}(t-1)^{n-1} + \binom{n}{2}(t-1)^{n-2} + \dots + n(t-1) + 1. \end{aligned}$$

Hence

$$T^{*n}T^n = \Delta_{T,n} + \binom{n}{1}\Delta_{T,n-1} + \binom{n}{2}\Delta_{T,n-2} + \dots + n\Delta_{T,n-1} + I$$

and

$$\begin{aligned} b_n &= \langle T^{*n}T^n x, x \rangle \\ &= a_n + \binom{n}{1}a_{n-1} + \binom{n}{2}a_{n-2} + \dots + na_1 + 1 \\ &= a_n + \binom{n}{n-1}a_{n-1} + \binom{n}{n-2}a_{n-2} + \dots + na_1 + 1. \end{aligned}$$

This completes the proof. $\square$

Proof of Theorem 2. Let T be a power bounded m -isometry. Since $T^*T - I$ is self-adjoint, it suffices to show that

$$\sigma(T^*T - I) = \sigma_a(T^*T - I) = \{0\}.$$

Assume that there exists a non-zero real number $a \in \sigma(T^*T - I)$. Since a belongs to the approximate point spectrum of $T^*T - I$, by Berberian's method we consider

an extension $\mathcal{H}^\circ$ of $\mathcal{H}$ and the mapping $S \rightarrow S^\circ$ of $B(\mathcal{H})$ into $B(\mathcal{H}^\circ)$. Then a is an eigen-value of $T^{*\circ}T^\circ - I^\circ$ and T° is also a power bounded m -isometry. For simplification, we denote T° by T . Since a is an eigen-value of $T^*T - I$, there exists a unit vector x such that $(T^*T - I)x = ax$. Hence

$$\langle (T^*T - I)x, x \rangle = \langle ax, x \rangle = a.$$

Let $a_1 = a$ and $a_n = \langle \Delta_{T,n}x, x \rangle$. Then since T is an m -isometry, we have $a_m = a_{m+1} = \cdots = a_n = 0$ for $n > m$. By the Lemma, we have

$$\|T^n x\|^2 = \binom{n}{m-1} a_{m-1} + \binom{n}{m-2} a_{m-2} + \cdots + na_1 + 1.$$

Hence

$$\frac{\|T^n x\|^2}{\binom{n}{m-1}} = a_{m-1} + \frac{1}{\binom{n}{m-1}} \left\{ \binom{n}{m-2} a_{m-2} + \cdots + na_1 + 1 \right\}.$$

Since T is power bounded, we have $a_{m-1} = 0$ by $n \rightarrow \infty$. Repeating this, we have

$$a_{m-1} = a_{m-2} = \cdots = a_1 = a = 0.$$

This is a contradiction, so the proof is complete. $\square$

Patel (Corollary 2.8, [10]) proved that if T is a 2-isometry, then $1 \in \sigma(T^*T)$. That is, if T is a 2-isometry, then $0 \in \sigma(\Delta_{T,1})$. We now generalize this result as follows.

Theorem 3. *If T is an m -isometry for $m \geq 2$, then $0 \in \sigma(\Delta_{T,m-1})$.*

Proof. For simplification, we denote $\Delta_{T,m-1}$ by Δ . Assume that $0 \notin \sigma(\Delta)$. Since $\Delta_{T,m} = T^*\Delta T - \Delta = 0$, it holds that $T^*\Delta T = \Delta$. Let $S = \Delta^{1/2}T\Delta^{-1/2}$. Then

$$S^*S = (\Delta^{-1/2}T^*\Delta^{1/2})(\Delta^{1/2}T\Delta^{-1/2}) = I.$$

Hence S is an isometry and $T = \Delta^{-1/2}S\Delta^{1/2}$. So T is similar to the isometry S . Since S is power bounded, T is an isometry by Theorem 3. Hence $\Delta_{T,1} = 0$, and so $\Delta_{T,m-1} = \Delta = 0$. This contradicts our assumption that $0 \notin \sigma(\Delta)$, so the proof is complete. $\square$

Remark 2. Since Weyl's theorem holds for an isometry, Weyl's theorem holds for a power bounded m -isometry by Theorem 3 (cf. [5], [10]).

Finally we show the following result. For a 2-isometry, Patel proved it (Theorem 2.1, [10]).

Theorem 4. *A power of an m -isometry is again an m -isometry.*

Proof. Let T be an m -isometry, i.e.,

$$\Delta_{T,m} = T^{*m}T^m - mT^{*m-1}T^{m-1} + \cdots + (-1)^m I = 0.$$

We prove T^k is also an m -isometry. Write

$$\begin{aligned} (t^k - 1)^m &= (t - 1)^m(t^{k-1} + t^{k-2} + \cdots + t + 1)^m \\ &= (t^m - mt^{m-1} + \cdots + (-1)^m)(a_{m(k-1)}t^{m(k-1)} + \cdots + a_1t + a_0). \end{aligned}$$

Hence we have

$$\Delta_{T^k, m} = \sum_{j=0}^{m(k-1)} a_j T^{*j} (\Delta_{T^k, m}) T^j = 0.$$

So the proof is complete. $\square$

ACKNOWLEDGMENT

The authors would like to express their sincere thanks to the referee for a careful reading and much kind advice. The proofs became very simple because of the referee's suggestions.

REFERENCES

- [1] J. Agler and M. Stankus, *m-Isometric transformations of Hilbert space I*, Integr. Equat. Oper. Theory, **21** (1995), 387-429. MR1321694 (96j:47008a)
- [2] J. Agler and M. Stankus, *m-Isometric transformations of Hilbert space II*, Integr. Equat. Oper. Theory, **23** (1995), 1-48. MR1346617 (96j:47008b)
- [3] J. Agler and M. Stankus, *m-Isometric transformations of Hilbert space III*, Integr. Equat. Oper. Theory, **24** (1996), 379-421. MR1382018 (96j:47008c)
- [4] A. Athavale, *Some operator theoretic calculus for positive definite kernels*, Proc. Amer. Math. Soc., **112**(3) (1991), 701-708. MR1068114 (92j:47050)
- [5] J.V. Baxley, *Some general conditions implying Weyl's theorem*, Rev. Roum. Math. Pures Appl., **16** (1971), 1163-1166. MR0305107 (46:4237)
- [6] S.K. Berberian, *Approximate proper vectors*, Proc. Amer. Math. Soc., **13** (1962), 111-114. MR0133690 (24:A3516)
- [7] T. Bermudez, A. Martinon and E. Negrin, *Weighted shift operators which are m-isometry*, Integr. Equ. Oper. Theory, **68** (2010), 301-312. MR2735438 (2012a:47074)
- [8] M. Chō, S. Ōta, K. Tanahashi, and A. Uchiyama, *Spectral properties of m-isometric operators*, Funct. Anal. Approx. Comput., **4**:2 (2012), 33-39.
- [9] J.B. Conway, *The Theory of Subnormal Operators*, Math. Surveys and Monographs **36**, Amer. Math. Soc., 1991. MR1112128 (92h:47026)
- [10] S.M. Patel, *2-Isometric operators*, Glasnik Mat., **37** (2002), 143-147. MR1918101 (2003f:47003)
- [11] A. Uchiyama and K. Tanahashi, *Bishop's property (β) for paranormal operators*, Operators and Matrices, **3** (2009), 517-524. MR2597677 (2011b:47047)
- [12] D. Xia, *Spectral theory of hyponormal operators*, Birkhäuser Verlag, Boston, 1983. MR806959 (87j:47036)

DEPARTMENT OF MATHEMATICS, KANAGAWA UNIVERSITY, YOKOHAMA 221-8686, JAPAN
E-mail address: `chiyom01@kanagawa-u.ac.jp`

DEPARTMENT OF CONTENT AND CREATIVE DESIGN, KYUSHU UNIVERSITY, FUKUOKA 815-8540, JAPAN
E-mail address: `ota@design.kyushu-u.ac.jp`

DEPARTMENT OF MATHEMATICS, TOHOKU PHARMACEUTICAL UNIVERSITY, SENDAI 981-8558, JAPAN
E-mail address: `tanahasi@tohoku-pharm.ac.jp`