EXPANSIONS OF QUADRATIC MAPS IN PRIME FIELDS

MEI-CHU CHANG

(Communicated by Bryna Kra)

ABSTRACT. Let $f(x) = ax^2 + bx + c \in \mathbb{Z}[x]$ be a quadratic polynomial with $a \not\equiv 0 \pmod{p}$. Take $z \in \mathbb{F}_p$ and let $\mathcal{O}_z = \{f_i(z)\}_{i \in \mathbb{Z}^+}$ be the orbit of z under f , where $f_i(z) = f(f_{i-1}(z))$ and $f_0(z) = z$. For $M < |\mathcal{O}_z|$, we study the diameter of the partial orbit $\mathcal{O}_M = \{z, f(z), f_2(z), \dots, f_{M-1}(z)\}$ and prove that there exists $c_1 > 0$ such that

$$\text{diam } \mathcal{O}_M \gtrsim \min \left\{ Mp^{c_1}, \frac{1}{\log p} M^{\frac{4}{5}} p^{\frac{1}{5}}, M^{\frac{1}{13} \log \log M} \right\}.$$

For a complete orbit $\mathcal{C}$, we prove that

$$\text{diam } \mathcal{C} \gtrsim \min \{ p^{5c_1}, e^{T/4} \},$$

where T is the period of the orbit.

INTRODUCTION

This paper belongs to the general theme of dynamical systems over finite fields. Let p be a prime and $\mathbb{F}_p$ the finite field of p elements, represented by the set $\{0, 1, \dots, p-1\}$. Let $f \in \mathbb{F}_p[x]$ be a polynomial, which we view as a transformation of $\mathbb{F}_p$. Thus if $z \in \mathbb{F}_p$ is some element, we consider its orbit

$$(0.1) \quad z_0 = z, \quad z_{n+1} = f(z_n), \quad n = 0, 1, \dots,$$

which eventually becomes periodic. The *period* $T_z = T$ is the smallest integer satisfying

$$(0.2) \quad \{z_n : n = 0, 1, \dots, T-1\} = \{z_n : n \in \mathbb{N}\}.$$

We are interested in the metrical properties of orbits and partial orbits. More precisely, for $M < T_z$, we define

$$(0.3) \quad \text{diam } \mathcal{O}_M = \max_{0 \leq n < M} |z_n - z|.$$

Following the papers [GS] and [CGOS], we study the expansion properties of f , in the sense of establishing lower bounds on $\text{diam } \mathcal{O}_M$. Obviously, if $M \leq T$, then $\text{diam } \mathcal{O}_M \geq M$. But, assuming that f is nonlinear and $M = o(p)$, one reasonably expects that the diameter of the partial orbit is much larger. Results along these lines were obtained in [GS] under the additional assumption that $M > p^{\frac{1}{2} + \epsilon}$. In this situation, Weil's theorem on exponential sums permits proving equidistribution of the partial orbit. For $M \leq p^{1/2}$, Weil's theorem becomes inapplicable and

Received by the editors October 10, 2011 and, in revised form, March 2, 2012.

2010 *Mathematics Subject Classification*. Primary 11B50, 37A45, 11B75; Secondary 11T23, 37F10, 11G99.

Key words and phrases. Dynamical system, orbits, additive combinatorics, exponential sums. The author's research was partially financed by the National Science Foundation.

lower bounds on $\text{diam } \mathcal{O}_M$ based on Vinogradov's theorem were established in [CGOS]. Our paper is a contribution of this line of research. We restrict ourselves to quadratic polynomials, though certainly the methods can be generalized. (See [CCGHSZ] for a generalization of Proposition 2 and Theorem 2 to higher degree polynomials and rational maps.)

Our first result is the following.

Theorem 1. *There is a constant $c_1 > 0$ such that if $f(x) = ax^2 + bx + c \in \mathbb{Z}[x]$ with $(a, p) = 1$, then with the above notation, for any $z \in \mathbb{F}_p$ and $M \leq T_z$,*

$$(0.4) \quad \text{diam } \mathcal{O}_M \gtrsim^1 \min \left\{ Mp^{c_1}, \frac{1}{\log p} M^{\frac{4}{5}} p^{\frac{1}{5}}, M^{\frac{1}{13}} \log \log M \right\}.$$

In view of Theorem 2, one could at least expect that $\text{diam } \mathcal{O}_M \gtrsim \min(p^c, e^{cM})$ as is the case when $M = T_z$.

In the proof, we distinguish the cases $\text{diam } \mathcal{O}_M > p^{c_0}$ and $\text{diam } \mathcal{O}_M \leq p^{c_0}$, where $c_0 > 0$ is a suitable constant. First, we again exploit exponential sum techniques (though, from the analytical side, our approach differs from [CGOS] and exploits a specific multilinear setup of the problem). More precisely, Proposition 1 in §1 states that (for $M \leq T$ large enough)

$$(0.5) \quad \text{diam } \mathcal{O}_M \gtrsim \frac{1}{\log p} \min \left(M^{\frac{5}{4}}, M^{\frac{4}{5}} p^{\frac{1}{5}} \right).$$

(Note that (0.5) is a clear improvement over Theorem 8 from [CGOS] for the case $d = 2$.)

When $\text{diam } \mathcal{O}_M \leq p^{c_0}$, a different approach becomes available as explained in Proposition 2. In this situation, we are able to replace the $(\text{mod } p)$ iteration by a similar problem in the field $\mathbb{C}$ of complex numbers, for an appropriate quadratic polynomial $F(z) \in \mathbb{Q}[z]$. Elementary arithmetic permits us to then prove that $\log \text{diam } \mathcal{O}_M$ is at least as large as $\frac{1}{13} \log M \log \log M$.

Interestingly, assuming $\mathcal{C}$ is a complete periodic cycle and $\text{diam } \mathcal{C} < p^{c_0}$, the transfer argument from Proposition 2 enables us to invoke bounds on the number of rational preperiodic points of a quadratic map, for instance the results from R. Benedetto [B]. The conclusion is the following.

Theorem 2. *There is a constant $c_0 > 0$ such that if $f(x) = ax^2 + bx + c \in \mathbb{Z}[x]$ with $(a, p) = 1$ and $\mathcal{C} \subset \mathbb{F}_p$ is a periodic cycle for f of length T , then*

$$(0.6) \quad \text{diam } \mathcal{C} \gtrsim \min\{p^{c_0}, e^{T/4}\}.$$

1. DIAMETER OF PARTIAL ORBITS

Let $f(x) = ax^2 + bx + c \in \mathbb{Z}[x]$, where $a \not\equiv 0 \pmod{p}$. Fix $x_0 \in \mathbb{F}_p$ and denote the *orbit* of x_0 by

$$\mathcal{O}_{x_0} = \{f_j(x_0)\}_{j \in \mathbb{Z}^+},$$

where $f_j(x_0) = f(f_{j-1}(x_0))$ and $f_0(x_0) = x_0$. The *period* of the orbit of x_0 under f is denoted $T = T_{x_0} = |\mathcal{O}_{x_0}|$. For $A \subset \mathbb{F}_p$, we denote the *diameter* of A by

$$\text{diam } A = \max_{x, y \in A} p \left\| \frac{x - y}{p} \right\|,$$

¹ $h \lesssim g$ if there exist constants C, M such that $|h(x)| \leq Cg(x)$ for all $x > M$.

where $\|a\|$ is the distance from a to the nearest integer. We are interested in the expansion of part of an orbit.

Proposition 1. *For $1 \ll M < T$, consider a partial orbit*

$$\mathcal{O}_M = \{x_0, f(x_0), f_2(x_0), \dots, f_{M-1}(x_0)\}.$$

Then

$$(1.1) \quad \text{diam } \mathcal{O}_M \gtrsim \frac{1}{\sqrt{\log p}} \min(M^{5/4}, M^{4/5} p^{1/5}).$$

Proof. Let $M_1 = \text{diam } \mathcal{O}_M$. Take $I \subset \mathbb{F}_p$ with $|I| = M_1$ and $\mathcal{O}_M \subset I$; then

$$(1.2) \quad |f(I) \cap I| \geq M - 1.$$

We will express (1.2) using exponential sums.

Let $0 \leq \varphi \leq 1$ be a smooth function on $\mathbb{F}_p$ such that $\varphi = 1$ on I and $\text{supp } \varphi \subset \tilde{I}$, where $\tilde{I}$ is an interval with the same center and double the length of I . Equation (1.2) implies that

$$\sum_{x \in I} \varphi(f(x)) \geq M$$

and expanding φ in Fourier gives

$$\varphi(x) = \sum_{\xi \in \mathbb{F}_p} \widehat{\varphi}(\xi) e_p(x\xi), \quad \text{with } \widehat{\varphi}(\xi) = \frac{1}{p} \sum_{x \in \mathbb{F}_p} \varphi(x) e_p(-x\xi).$$

Combining these gives

$$(1.3) \quad \sum_{\xi \in \mathbb{F}_p} |\widehat{\varphi}(\xi)| \left| \sum_{x \in I} e_p(\xi f(x)) \right| \gtrsim M.$$

We will estimate $\sum_{x \in I} e_p(\xi f(x))$ using van der Corput-Weyl.

Take $M_0 = O(M)$, e.g. $M_0 = \frac{1}{100}M$. Then

$$(1.4) \quad \begin{aligned} \left| \sum_{x \in I} e_p(\xi f(x)) \right| &\leq \frac{1}{M_0} \sum_{0 \leq y < M_0} \left| \sum_{x \in I} e_p(\xi(a(x+y)^2 + b(x+y))) \right| + O(M_0) \\ &\leq \frac{1}{\sqrt{M_0}} \left[\sum_{0 \leq y < M_0} \left| \sum_{x \in I} e_p(\xi(ax^2 + 2axy + bx)) \right|^2 \right]^{1/2} + O(M_0) \\ &= \frac{1}{\sqrt{M_0}} \left| \sum_{\substack{0 \leq y < M_0 \\ x_1, x_2 \in I}} e_p(\xi(x_1 - x_2)(a(x_1 + x_2) + 2ay + b)) \right|^{1/2} + O(M_0). \end{aligned}$$

(The second inequality is by Cauchy-Schwarz.)

Take φ sufficiently smooth so as to ensure that

$$(1.5) \quad \sum_{\xi \in \mathbb{F}_p} |\widehat{\varphi}(\xi)| = O(1).$$

Equations (1.3) and (1.4) imply

$$\sum_{\xi \in \mathbb{F}_p} |\widehat{\varphi}(\xi)| \left| \sum_{\substack{0 \leq y < M_0 \\ x_1, x_2 \in I}} e_p(\xi(x_1 - x_2)(a(x_1 + x_2) + 2ay + b)) \right|^{1/2} \gtrsim M^{3/2}.$$

Hence by Cauchy-Schwarz and (1.5),

$$(1.6) \quad \sum_{\xi \in \mathbb{F}_p} |\widehat{\varphi}(\xi)| \left| \sum_{\substack{0 \leq y < M_0 \\ x_1, x_2 \in I}} e_p(\xi(x_1 - x_2)(a(x_1 + x_2) + 2ay + b)) \right| \gtrsim M^3.$$

Fix $x_1 + x_2 = s \leq 2M_1$; then

$$(1.7) \quad \sum_{\xi \in \mathbb{F}_p} |\widehat{\varphi}(\xi)| \left| \sum_{\substack{0 \leq y < M_0 \\ x \in I}} e_p(\xi(2x - s)(as + 2ay + b)) \right| \gtrsim \frac{M^3}{2M_1}.$$

Next, for $z \in \mathbb{F}_p$, denote

$$(1.8) \quad \eta(z) = |\{(x, y) \in I \times [1, M_0] : (2x - s)(2ay + b + as) \equiv z \pmod{p}\}|,$$

and write the left hand side of (1.7) as

$$(1.9) \quad \begin{aligned} & \sum_{\xi \in \mathbb{F}_p} |\widehat{\varphi}(\xi)| \left| \sum_{z \in \mathbb{F}_p} \eta(z) e_p(\xi z) \right| \\ & \leq \left(\sum_{\xi \in \mathbb{F}_p} |\widehat{\varphi}(\xi)|^2 \right)^{1/2} \left(\sum_{\xi \in \mathbb{F}_p} \left| \sum_{z \in \mathbb{F}_p} \eta(z) e_p(\xi z) \right|^2 \right)^{1/2} \\ & = \left(\frac{1}{p} \sum_{x \in \mathbb{F}_p} |\varphi(x)|^2 \right)^{1/2} \sqrt{p} \left(\sum_{z \in \mathbb{F}_p} \eta(z)^2 \right)^{1/2} \\ & < 2M_1^{1/2} \left(\sum_{z \in \mathbb{F}_p} \eta(z)^2 \right)^{1/2}, \end{aligned}$$

by Cauchy-Schwarz and Parseval.

Recall that $(a, p) = 1$. Let $I' = I - \frac{s}{2}$, $I'' = [1, M_0] + \frac{b+as}{2a} \subset \mathbb{F}_p$ so that

$$\sum_{z \in \mathbb{F}_p} \eta(z)^2 = E(I', I''),$$

the multiplicative energy of I' and I'' .

It is well-known that

$$(1.10) \quad \begin{aligned} E(I', I'') & \leq \log p \max \left\{ |I'| |I''|, \frac{|I'|^2 |I''|^2}{p} \right\} \\ & \leq \log p \max \left\{ M_1 M, \frac{M_1^2 M^2}{p} \right\}. \end{aligned}$$

Thus, by (1.7), (1.9) and (1.10),

$$(1.11) \quad \frac{M^3}{M_1} \lesssim M_1^{1/2} (\log p)^{1/2} \max \left\{ M_1^{1/2} M^{1/2}, \frac{M_1 M}{p^{1/2}} \right\}.$$

Distinguish the cases $M_1 M \leq p$ and $M_1 M > p$, and then (1.11) implies

$$(1.12) \quad M_1 \gtrsim \min \left\{ (\log p)^{-1/4} M^{5/4}, (\log p)^{-1/5} M^{4/5} p^{1/5} \right\}. \quad \square$$

2. PARTIAL ORBITS OF SMALL DIAMETERS

For $M < p^{c_0}$, one obtains the following stronger result. (Notation is as in Proposition 1.)

Proposition 2. *There exists $c_0 > 0$ such that*

$$(2.1) \quad \text{diam } \mathcal{O}_M > \min \left(p^{c_0}, M^{\frac{1}{13} \log \log M} \right).$$

Consequently,

$$(2.2) \quad \text{diam } \mathcal{O}_M \gtrsim \min \left\{ Mp^{\frac{c_0}{5}}, \frac{1}{\log p} M^{\frac{4}{5}} p^{\frac{1}{5}}, M^{\frac{1}{13} \log \log M} \right\}.$$

Proof. Let $\mathcal{O}_M = \{x_0, x_1, \dots, x_{M-1}\}$ with $x_j = f(x_{j-1})$ as before, and let $\text{diam } \mathcal{O}_M = M_1$. Since $|x_j - x_0| \leq M_1$, we can write $x_j = x_0 + z_j$ with $z_j \in [-M_1, M_1]$. Thus, a, b, c, x_0 satisfy the $M - 1$ equations

$$a(x_0 + z_j)^2 + b(x_0 + z_j) + c \equiv x_0 + z_{j+1} \pmod{p}, \quad j = 0, \dots, M - 2,$$

and the $\mathbb{F}_p$ -variety

$$\mathcal{V}_p = \bigcap_{j=0}^{M-2} [(r + z_j)^2 + v(r + z_j) + w = u(r + z_{j+1}) \pmod{p}]$$

in the variables $(u, v, w, r) \in \mathbb{F}_p^4$ is therefore nonempty. Note that the coefficients of the $M - 1$ defining polynomials in $\mathbb{Z}[u, v, w, r]$ are $O(M_1^2)$.

Assume

$$(2.3) \quad M_1 < p^{c_0}$$

with $c_0 > 0$ small enough. Elimination theory² implies that $\mathcal{V}_p \neq \emptyset$ is a $\mathbb{C}$ -variety. Hence there are $U, V, W, R \in \mathbb{C}$ such that for all j ,

$$(R + z_j)^2 + V(R + z_j) + W = U(R + z_{j+1}), \quad j = 0, \dots, M - 2.$$

Obviously, $U \neq 0$, since $z_1, \dots, z_{M-2}$ are distinct. We therefore have a quadratic polynomial

$$(2.4) \quad F(z) := \frac{1}{U}(R + z)^2 + \frac{V}{U}(R + z) + \frac{W}{U} - R =: Az^2 + Bz + C,$$

satisfying

$$(2.5) \quad F(z_j) = z_{j+1} \quad \text{in } \mathbb{C}, \quad \text{for } j = 0, \dots, M - 2.$$

Since $z_0 = 0$, (2.4) and (2.5) imply $C = z_1 \in \mathbb{Z} \cap [-M_1, M_1]$, and the equations

$$\begin{aligned} z_1^2 A + z_1 B &= z_2 - z_1, \\ z_2^2 A + z_2 B &= z_3 - z_1 \end{aligned}$$

imply $A, B \in \mathbb{Q}$ with $A = \frac{a}{d}$, $B = \frac{b}{d}$, and $a, b, d \in \mathbb{Z}$ being $O(M_1^3)$. Equation (2.5) becomes

$$(2.6) \quad z_{j+1} = \frac{a}{d} z_j^2 + \frac{b}{d} z_j + C.$$

²See [C], where a similar elimination procedure was used in a combinatorial problem. In particular, see [C], Lemma 2.14 and its proof.

Hence

$$\frac{a}{d}z_{j+1} + \frac{b}{2d} = \left(\frac{a}{d}z_j + \frac{b}{2d}\right)^2 + C\frac{a}{d} - \frac{b^2}{4d^2} + \frac{b}{2d}.$$

Putting

$$y_j = \frac{a}{d}z_j + \frac{b}{2d} \in \frac{1}{2d}\mathbb{Z}, \quad j = 0, \dots, M-1,$$

and

$$\frac{r}{s} = C\frac{a}{d} - \frac{b^2}{4d^2} + \frac{b}{2d} \quad \text{with } s > 0, (r, s) = 1, |r|, s = O(M_1^6),$$

gives

$$(2.7) \quad y_{j+1} = y_j^2 + \frac{r}{s}, \quad j = 0, \dots, M-2.$$

Next, write $y_j = \alpha_j/\beta_j$, where $\beta_j|2d$ and $(\alpha_j, \beta_j) = 1$; thus (2.7) gives

$$(2.8) \quad \frac{\alpha_{j+1}}{\beta_{j+1}} = \frac{\alpha_j^2}{\beta_j^2} + \frac{r}{s}, \quad j = 0, \dots, M-2.$$

Note also that

$$(2.9) \quad |\alpha_j| = O(M_1^4).$$

Write the prime factorizations

$$s = \prod_p p^{v(p)} \quad \text{and} \quad \beta_j = \prod_p p^{v_j(p)}, \quad j = 0, \dots, M-1.$$

Claim. $2v_j(p) \leq v(p)$, for $j < M - O(\log \log M_1)$.

Proof. We may assume $v_j(p) > 0$.

Case 1. $2v_j(p) > v_{j+1}(p)$.

Fact 2.1 (which will be stated at the end of this section) and (2.8) imply that $v(p) = 2v_j(p)$.

Case 2. $2v_j(p) \leq v_{j+1}(p)$.

Again, we separate two cases.

Case 2.1. $2v_{j+1}(p) > v_{j+2}(p)$. Reasoning as in Case 1, we have

$$v(p) = 2v_{j+1}(p) \geq 2^2v_j(p) > 2v_j(p).$$

Case 2.2. $2v_{j+1}(p) \leq v_{j+2}(p)$. Therefore, $v_{j+2}(p) \geq 2^2v_j(p)$. We repeat the argument for Case 2 with $j = j+1$. Continuing this process, after τ steps, we obtain either $v(p) \geq 2v_j(p)$ or

$$(2.10) \quad v_{j+\tau}(p) \geq 2^\tau v_j(p),$$

when necessarily $\tau \lesssim \log v_{j+\tau}(p) \lesssim \log \log \beta_{j+\tau} \lesssim \log \log d \lesssim \log \log M_1$. Since $j + \tau \leq M$, the claim is proved. $\square$

It follows from the claim that $\beta_j^2|s$ for $j < M - O(\log \log M_1)$. Back to (2.8), if $v(p) > 2v_j(p)$ for some $j < M - O(\log \log M_1)$, then $v_{j+1}(p) = v(p)$. This contradicts the fact that $\beta_{j+1}^2|s$. So we conclude that

$$(2.11) \quad \beta_j^2 = s =: s_1^2 \quad \text{for } j < M - O(\log \log M_1).$$

Hence

$$(2.12) \quad \alpha_{j+1} = \frac{\alpha_j^2}{s_1} + \frac{r}{s_1},$$

which implies

$$(2.13) \quad \alpha_j^2 + r \equiv 0 \pmod{s_1}.$$

Let $s_1 = \prod p_i^{v_i}$. Then α_j satisfies (2.13) if and only if α_j satisfies $\alpha_j^2 + r \equiv 0 \pmod{p_i^{v_i}}$ for all i . Since $-r$ is a quadratic residue modulo p^v if and only if it is a quadratic residue modulo p for odd prime p , we have

$$(2.14) \quad \left| \{\pi_{s_1}(\alpha_j)\}_j \right| \leq 2 \cdot 2^{\omega(s_1)} < e^{\frac{\log s_1}{\log \log s_1}} < e^{\frac{4 \log M_1}{\log \log M_1}}.$$

Here $\pi_{s_1}(\alpha_j)$ is the projection of α_j in $\mathbb{Z}_{s_1}$.

To show $M_1 > M^{\frac{1}{13} \log \log M}$, we assume

$$(2.15) \quad \log M_1 < \frac{1}{13} \log M \log \log M.$$

Then (2.14) implies there exists $\xi \in \mathbb{Z}_{s_1}$ such that

$$(2.16) \quad |\mathcal{J}| = \left| \left\{ 0 \leq j \leq \frac{M}{2} : \pi_{s_1}(\alpha_j) = \xi \right\} \right| > M^{1/2}.$$

Thus

$$\alpha_{j_1} - \alpha_{j_2} \in s_1 \mathbb{Z}, \quad \text{for } j_1, j_2 \in \mathcal{J},$$

and

$$|\alpha_{j_1} - \alpha_{j_2}| \geq s_1, \quad \text{for } j_1 \neq j_2 \in \mathcal{J}.$$

In particular, there exists $j \in \mathcal{J}$ such that

$$(2.17) \quad |\alpha_j| \geq \frac{M^{1/2}}{8} s_1 \quad \text{and} \quad \left| |\alpha_j| - |r|^{1/2} \right| > \frac{M^{1/2}}{8} s_1.$$

Claim. Either $|\alpha_j| > 10|r|^{1/2}$ or $|\alpha_{j+1}| > 10|r|^{1/2}$.

Proof. Assume

$$(2.18) \quad |\alpha_j|, |\alpha_{j+1}| < 10|r|^{1/2}.$$

Hence, $|r|^{1/2} \gtrsim M^{1/2} s_1$ by (2.17). From (2.12), (2.17) and (2.18),

$$\begin{aligned} 10|r|^{1/2} s_1 &> |\alpha_{j+1}| s_1 = |\alpha_j^2 + r| \\ &\geq (|\alpha_j| + |r|^{1/2})(|\alpha_j| - |r|^{1/2}) \\ &\geq |r|^{1/2} \cdot \frac{M^{1/2}}{8} s_1, \end{aligned}$$

a contradiction, proving the claim. □

Thus, there exists $j < M/2$ such that either

$$(2.19) \quad |\alpha_j| > 10s_1 \quad \text{and} \quad |\alpha_j| > 10|r|^{1/2}$$

or

$$(2.20) \quad |\alpha_j| > 10s_1 \quad \text{and} \quad |\alpha_{j+1}| > 10|r|^{1/2}.$$

Clearly, (2.19) implies (2.20). Indeed, by (2.12),

$$|\alpha_{j+1}| \geq \frac{1}{s_1} |\alpha_j^2 - |r|| \geq \frac{99}{100s_1} \alpha_j^2 > 2|\alpha_j|.$$

Iteration shows that

$$|\alpha_{j+\frac{M}{3}}| > 2^{\frac{M}{3}} |\alpha_j| > 2^{\frac{M}{3}},$$

contradicting (2.9). This proves (2.1).

Combining Proposition 1 and (2.1), we have (2.2). $\square$

Fact 2.1. Let $\frac{a_1}{d_1}, \frac{a_2}{d_2}, \frac{a_3}{d_3} \in \mathbb{Q}$ be rational numbers in lowest terms, and $p^{v_p(d_i)} || d_i$. If $\frac{a_1}{d_1} + \frac{a_2}{d_2} + \frac{a_3}{d_3} = 0$ and $v_p(d_1) \geq v_p(d_2) \geq v_p(d_3)$, then $v_p(d_1) = v_p(d_2)$.

3. FULL CYCLES

In this section, we will prove Theorem 2.

Assume $M_1 = \text{diam } \mathcal{C} < p^{c_0}$ with c_0 as in Proposition 2. The proof of Proposition 2 gives a quadratic polynomial (cf. (2.7))

$$(3.1) \quad F(z) = z^2 + \frac{r}{s} \quad \text{with } r, s \in \mathbb{Z}, |s| = O(M_1^6)$$

and a rational F -cycle $\{y_j\}_{0 \leq j < T}$, i.e.

$$y_{j+1} = F(y_j) \quad \text{for } 0 \leq j \leq T-2$$

and $F(y_{T-1}) = y_0$.

We now invoke a result of R. Benedetto [B], which gives quantitative bounds on the number of preperiodic points of a polynomial f in a number field (z is *preperiodic* if the set $\{z, f(z), f(f(z)), \dots\}$ is finite). According to Theorem 7.1 in [B], the number of preperiodic points of F in $\mathbb{Q}$ is bounded by

$$(3.2) \quad (2\sigma + 1) [\log_2(2\sigma + 1) + \log_2(\log_2(2\sigma + 1) - 1) + 2]$$

with σ the number of primes where F has bad reduction. Hence $\sigma \leq \omega(s) \leq \frac{\log M_1}{\log \log M_1}$ and (3.2) implies

$$(3.3) \quad T < 4 \log M_1 = 4 \log \text{diam } \mathcal{C}.$$

ACKNOWLEDGEMENTS

The author would like to thank the referee for many helpful comments and the mathematics department of the University of California at Berkeley for its hospitality.

REFERENCES

- [B] Robert L. Benedetto, *Preperiodic points of polynomials over global fields*, J. Reine Angew. Math. **608** (2007), 123–153, DOI 10.1515/CRELLE.2007.055. MR2339471 (2008j:11071)
- [CCGHSZ] M.-C. Chang, J. Cilleruelo, M. Garaev, J. Hernandez, I. Shparlinski, A. Zumalacarregui, *Points on curves in small boxes and applications* (preprint).
- [C] M. Chang, *Factorization in generalized arithmetic progressions and applications to the Erdős-Szemerédi sum-product problems*, Geom. Funct. Anal. **13** (2003), no. 4, 720–736, DOI 10.1007/s00039-003-0428-5. MR2006555 (2004g:11007)
- [CGOS] Javier Cilleruelo, Mubbariz Z. Garaev, Alina Ostafe, and Igor E. Shparlinski, *On the concentration of points of polynomial maps and applications*, Math. Z. **272** (2012), no. 3-4, 825–837, DOI 10.1007/s00209-011-0959-7. MR2995141
- [GS] Jaime Gutierrez and Igor E. Shparlinski, *Expansion of orbits of some dynamical systems over finite fields*, Bull. Aust. Math. Soc. **82** (2010), no. 2, 232–239, DOI 10.1017/S0004972709001270. MR2685147 (2011j:11153)