SYMPLECTIC 4-MANIFOLDS WITH FIXED POINT FREE CIRCLE ACTIONS

JONATHAN BOWDEN

(Communicated by Daniel Ruberman)

ABSTRACT. We show that recent results of Friedl-Vidussi and Chen imply that a symplectic 4-manifold admits a fixed point free circle action if and only if it admits a symplectic structure that is invariant under the action and we give a complete description of the symplectic cone in this case. This then completes the topological characterisation of symplectic 4-manifolds that admit non-trivial circle actions.

1. INTRODUCTION

Recently Friedl and Vidussi [7] solved the long standing Taubes Conjecture, which classifies which 4-manifolds of the form $M \times S^1$ admit symplectic forms. Moreover, they determined exactly which cohomology classes can be represented by symplectic forms. Using recent results of D. Wise, [13] they have extended their results to the case of non-trivial S^1 -bundles in [9]. In this note we observe that their results as well of those of Chen, who obtained partial results in the fixed point free case in [3], imply the analogue of ([9], Theorem 1.3) for all fixed point free circle actions.

Before stating our main result we fix some notation and terminology. Let $X \xrightarrow{p} M$ be an orientable 4-manifold with a fixed point free circle action and quotient space $M = X/S^1$. The quotient space is an orbifold whose underlying topological space $|M|$ is a manifold since all the stabilisers of the S^1 -action are necessarily cyclic and the singular locus consists of a collection of branching circles (cf. [1], [6]).

We let M_{reg} denote the complement of an open tubular neighbourhood of the singular locus of M and $X_{reg} = p^{-1}(M_{reg})$, which is an honest S^1 -bundle so that the pushforward map p_* is well-defined for cohomology classes in $H^*(X_{reg}, \mathbb{R})$. The manifold M_{reg} has toroidal boundary and thus one may define the Thurston norm on $H^1(M_{reg}, \mathbb{R})$ in the usual fashion. Finally for $\psi \in H^2(X, \mathbb{R})$ we let ψ_{reg} denote the restriction of ψ to X_{reg} .

Theorem 1. *Let $X \xrightarrow{p} M$ be an oriented manifold admitting a fixed point free S^1 -action with quotient space M and let $\psi \in H^2(X, \mathbb{R})$. Then the following are equivalent:*

- (1) ψ can be represented by a symplectic form,
- (2) ψ can be represented by an S^1 -invariant symplectic form,
- (3) $\psi^2 > 0$ and $p_*\psi_{reg} \in H^1(M_{reg}, \mathbb{R})$ lies in the open cone over a fibered face of the Thurston norm ball and is the restriction of a class in $H^1(|M|, \mathbb{R})$.

Received by the editors May 11, 2012 and, in revised form, September 1, 2012.

2010 *Mathematics Subject Classification.* Primary 57R17; Secondary 57N10, 57N13.

©2014 American Mathematical Society
 Reverts to public domain 28 years from publication

Note that if a class $\phi \in H^1(|M|, \mathbb{R})$ is integral, then ϕ can be represented by a fibration over S^1 that is transverse to the singular locus of $|M|$ if and only if its restriction to M_{reg} , which we denote by ϕ_{reg} , is fibered. Recall that a fibration of a manifold with boundary is required to be transverse to the boundary. Furthermore, since ϕ_{reg} is the restriction of a class in $H^1(|M|, \mathbb{R})$, it automatically vanishes on the meridian classes in ∂M_{reg} so that if ϕ_{reg} is fibered, then the induced fibration on the boundary is necessarily meridional. Thus the fibration dual to ϕ_{reg} extends to $|M|$ in the desired way by filling in discs near the singular locus. In particular, part (3) of Theorem 1 implies that the underlying manifold $|M|$ is fibered and we obtain a positive answer to the following conjecture, which implies [3], Conjecture 1.7, as a special case.

Conjecture 1 (Generalised Taubes Conjecture). *Let X be a symplectic 4-manifold that admits a non-trivial fixed point free circle action with quotient orbifold M . Then the (possibly empty) singular locus L of M is a meridionally fibered link.*

Furthermore, as noted in [3], p. 6, Theorem 1 completes the characterisation of which symplectic manifolds admit non-trivial S^1 -actions. For Baldridge, [1] showed that if a non-trivial S^1 -action on a symplectic 4-manifold has fixed points, then X is rational or ruled and thus admits an S^1 -invariant symplectic form for *some* non-trivial S^1 -action. In view of this we obtain the following corollary.

Corollary 1. *Let X be a symplectic 4-manifold that admits a non-trivial S^1 -action. Then either the action is fixed point free and the quotient space fibers over S^1 or X is rational or ruled. In either case, X admits a non-trivial symplectic S^1 -action.*

2. PROOF OF THEOREM 1

The proof of Theorem 1 is based on the following lemma, which provides a generalisation of [4], Theorem 5.2, to include irrational classes. For the proof we assume a certain familiarity with the basic properties of the Thurston norm (cf. [12]).

Lemma 1. *Let $\overline{M}$ be a 3-manifold with an orientation preserving smooth action of a finite group G and quotient orbifold $M = \overline{M}/G$. An element $\overline{\phi}$ in the invariant subspace $H^1(\overline{M}, \mathbb{R})^G$ admits a non-degenerate de Rham representative if and only if it admits a non-degenerate de Rham representative that is G -invariant.*

In particular, the restriction of the associated class $\phi \in H^1(|M|, \mathbb{R}) \cong H^1(\overline{M}, \mathbb{R})^G$ to M_{reg} lies in the open cone over a fibered face of the Thurston norm ball.

Proof. We first assume that $\overline{\phi}$ is rational. Since nothing changes after multiplying with positive constants, we may assume that $\overline{\phi}$ is in fact integral. In this case the first claim is just a restatement of [4], Theorem 5.2, which can be applied in complete generality in view of [11], Theorem 8.1. Note that the assumption $H^1(\overline{M}, \mathbb{Q})^G = \mathbb{Q}$ in [4], Theorem 5.2, can be replaced by the fact that the fibration is given by a fibered class $\overline{\phi}$ that is G -invariant. Moreover, the proof in [4] actually gives a fibration that is transverse to the branching locus in $\overline{M}$. The quotient map π induces an isomorphism $H^1(\overline{M}, \mathbb{R})^G \cong H^1(|M|, \mathbb{R})$ so that there is a unique class ϕ with $\overline{\phi} = \pi^*\phi$ and the fibration dual to $\overline{\phi}$ descends to a fibration of $|M|$ dual to ϕ . Finally since the fibration is transverse to the singular locus it follows that the restriction of ϕ to M_{reg} is fibered.

We next assume that $\bar{\phi}$ is irrational and let ϕ be the unique class with $\bar{\phi} = \pi^*\phi$. We let ι_{reg} denote the natural inclusion $M_{reg} \hookrightarrow |M|$ and set $V = \text{Im}(\iota_{reg}^*)$. By the previous case all rational classes in V that are sufficiently close to $\phi_{reg} = \iota_{reg}^*\phi$ are fibered. If ϕ_{reg} itself did not lie in the open cone over a fibered face of the Thurston unit ball, then it must lie in the closed cone over the boundary of a fibered face by the assumption that it can be approximated by fibered elements. Since the Thurston unit ball is rational, the intersection of the closed cone containing ϕ_{reg} with V must contain non-fibered rational points arbitrarily close to ϕ_{reg} , which gives a contradiction. Thus ϕ_{reg} admits a non-degenerate de Rham representative η_{reg} . Since η_{reg} can be approximated by rational classes that are fibered and restrict themselves to meridional fibrations on the boundary of M_{reg} , the foliation induced by η_{reg} on the boundary is also meridional.

We let $(z, \theta) \in D^2 \times S^1$ denote coordinates on a tubular neighbourhood of a component of the branching locus of $|M|$. After applying a suitable isotopy we may assume that η_{reg} has the form $f(\theta)d\theta$ near $\partial D^2 \times S^1$. It follows that η_{reg} extends to a non-degenerate closed form η which is transverse to the branching locus of $|M|$. The pullback $\bar{\eta} = \pi^*\eta$ then gives the desired non-degenerate G -equivariant representative of $\bar{\phi}$. $\square$

Proof of Theorem 1. The implication (2) $\implies$ (1) is trivial.

(1) $\implies$ (3): Let (X, ω) be a symplectic manifold with a fixed point free S^1 -action and quotient space M . By [3], Proposition 1.8, there is a manifold $\bar{M}$ and a smooth action by a finite group so that $M = \bar{M}/G$. Furthermore, we have the following commutative diagram:

$$\begin{array}{ccc} \pi^*X = \bar{X} & \xrightarrow{\bar{p}} & \bar{M} \\ \bar{\pi} \downarrow & & \downarrow \pi \\ X & \xrightarrow{p} & M, \end{array}$$

where π is the quotient map, $\bar{\pi}$ is an unramified covering and the induced S^1 -action on $\bar{X}$ is free. Moreover, the group G acts naturally on $\bar{X}$ as the group of deck transformations of $\bar{\pi}$.

Thus $\bar{\omega} = \bar{\pi}^*\omega$ is a symplectic form and by [9], Theorem 1.4, its image under the pushforward map $\bar{p}_*(\bar{\omega}) \in H^1(\bar{M}, \mathbb{R})$ lies in the open cone over a fibered face of the Thurston norm ball. Since $\bar{\omega}$ is G -invariant and the action on $\bar{X}$ is fiber preserving, the class $\bar{\phi} = \bar{p}_*(\bar{\omega})$ is also G -invariant. We let ϕ be the unique class such that $\pi^*\phi = \bar{\phi}$. By Lemma 1 the restriction of ϕ_{reg} to M_{reg} lies in the open cone over a fibered face. Finally the naturality of the transfer homomorphism implies that the restriction of ϕ_{reg} agrees with $p_*\omega_{reg}$.

(3) $\implies$ (2): By assumption $\phi_{reg} = p_*\psi_{reg}$ lies in the open cone over a fibered face of the Thurston norm ball and ϕ_{reg} is the restriction of a class $\phi \in H^1(|M|, \mathbb{R})$. In particular, $|M|$ fibers over S^1 . We first note that M is a *very good* orbifold so that it is a quotient of a manifold $\bar{M}$ by a smooth action of a finite group G . For this it suffices to rule out bad 2-suborbifolds by [2], Corollary 3.28. However, a bad 2-suborbifold is topologically a sphere that is essential in $H_2(|M|, \mathbb{Z})$ and as in the proof of [3], Lemma 2.3, this implies that $b_2^+(X) = b_2(|M|) - 1$. Thus $|M| = S^2 \times S^1$ and $b_2^+(X) = 0$, contradicting the assumption that $\psi^2 > 0$.

Thus since M is very good we can proceed as in the proof of the previous implication. In particular, M is a quotient of a manifold $\overline{M}$ by a smooth action of a finite group G , the total space has a finite covering $\overline{X}$ which is a genuine S^1 -bundle and these bundles fit into a pullback diagram as above. Since a degree one cohomology class on $\overline{M}$ is determined by its restriction to the complement of the branching locus, we deduce that $\overline{\phi} = \pi^*\phi$ and $\overline{p}_*(\pi^*\psi)$ agree as cohomology classes. We then note that the construction of S^1 -invariant forms in [5] and its extension to irrational classes ([8], Theorem 1.1) can be done G -equivariantly.

First choose a G -invariant representative γ of $e(\overline{X})$, which can be obtained as the curvature of a G -equivariant angular form. By [8], Lemma 2.1, we may write $\gamma = \overline{\phi} \wedge \beta$. After averaging over G this equation still holds, so β can be assumed to be G -equivariant. Let η be a G -invariant angular 1-form so that $d\eta = \overline{p}^*\gamma$ and let $\overline{\Omega} \in H^2(\overline{M}, \mathbb{R})$ be the unique class such that the following holds in cohomology:

$$\pi^*\psi - \eta \wedge \overline{p}^*\overline{\phi} = \overline{p}^*\overline{\Omega}.$$

Such an $\overline{\Omega}$ exists in view of the Gysin sequence since the left hand lies in the kernel of $\overline{p}_*$, and since the left hand side is G -equivariant so is $\overline{\Omega}$. The fact that $\pi^*\psi^2 > 0$ implies that $\overline{p}^*\overline{\phi} \wedge \overline{\Omega} > 0$. Thus by [8], Lemma 2.2, there is a non-vanishing 2-form representing the class $\overline{\Omega}$ so that $\overline{\phi} \wedge \overline{\Omega} > 0$; again after averaging we may assume that $\overline{\Omega}$ is G -invariant. Thus the S^1 -invariant form

$$\overline{\omega}_{inv} = \eta \wedge \overline{p}^*\overline{\phi} + \overline{p}^*\overline{\Omega}$$

represents $\pi^*\psi$ and descends to an S^1 -invariant form ω_{inv} on X which is cohomologous to ψ . $\square$

Remark 1. A vital step in the proof of Theorem 1 was Chen's observation that the base orbifold of a symplectic manifold X admitting a fixed point free S^1 -action is a quotient of a manifold by a finite group action. The main technical point in the proof of [3], Proposition 1.8, is to rule out bad 2-orbifolds in the base. This is achieved by results relating the Seiberg-Witten invariants of the base orbifold to those of the underlying manifold.

We sketch a different proof which uses more standard Seiberg-Witten vanishing results. For background on the Seiberg-Witten invariants we refer to [10] and the references therein. First observe that a bad 2-suborbifold Σ in the quotient orbifold $M = X/S^1$ can intersect at most 2 singular curves L_1, L_2 , each in at most one point. Taking a neighbourhood N of $|\Sigma| \cup L_1$ gives a topological splitting of the base $|M| = (S^2 \times S^1) \# M'$ so that the preimage of the splitting sphere in $|M|$ induces a splitting $X = X_1 \cup_S X_2$, where S is either $S^2 \times S^1$ or S^3 depending on whether L_2 is empty or not. Moreover, as in the proof of [3], Lemma 2.3, we must have $b_1(M') > 0$ by the assumption that $b_2^+(X) > 0$. If S is a 3-sphere, then $b_2^+(X_2) > 0$, and by taking the covering $\overline{X}$ of X induced by the natural surjection

$$\pi_1(X) \rightarrow \pi_1(S^2 \times S^1) \rightarrow \mathbb{Z}_n$$

we obtain a splitting of $\overline{X} = \overline{X}_1 \cup_{S^3} \overline{X}_2$, where $b_2^+(\overline{X}_1), b_2^+(\overline{X}_2) \geq 1$. It follows that the Seiberg-Witten invariants of $\overline{X}$ are trivial.

If $S = S^2 \times S^1$, then we take the covering $\overline{X}$ of X induced by a surjection

$$\pi_1(X) \rightarrow \pi_1(S^2 \times S^1) * \pi_1(|M'|) \rightarrow \mathbb{Z}_n \times \mathbb{Z}_n.$$

The embedded 2-sphere $S^2 \times \{pt\}$ in S then becomes essential in the covering and $b_1(\overline{X})$, and hence $b_2^+(\overline{X})$, may be assumed to be arbitrarily large. Furthermore, the sphere $S^2 \times \{pt\}$ has trivial self-intersection and consequently the Seiberg-Witten invariants of $\overline{X}$ are trivial. Thus in both cases we obtain a contradiction to the non-vanishing results of Taubes for the Seiberg-Witten invariants of a symplectic 4-manifold.

ACKNOWLEDGMENTS

The author thanks P. Suárez-Serrato for his stimulating questions and S. Friedl for helpful comments. The hospitality of the Max-Planck-Institut für Mathematik in Bonn, where this research was carried out, is also gratefully acknowledged.

REFERENCES

- [1] Scott Baldridge, *Seiberg-Witten vanishing theorem for S^1 -manifolds with fixed points*, Pacific J. Math. **217** (2004), no. 1, 1–10, DOI 10.2140/pjm.2004.217.1. MR2105762 (2005j:57040)
- [2] Michel Boileau, Sylvain Maillot, and Joan Porti, *Three-dimensional orbifolds and their geometric structures* (English, with English and French summaries), Panoramas et Synthèses [Panoramas and Synthèses], vol. 15, Société Mathématique de France, Paris, 2003. MR2060653 (2005b:57030)
- [3] W. Chen, *Seifert fibered four-manifolds with nonzero Seiberg-Witten invariant*, Preprint: arXiv:1103.5681v3, 2011.
- [4] Allan L. Edmonds and Charles Livingston, *Group actions on fibered three-manifolds*, Comment. Math. Helv. **58** (1983), no. 4, 529–542, DOI 10.1007/BF02564651. MR728451 (85k:57037)
- [5] Marisa Fernández, Alfred Gray, and John W. Morgan, *Compact symplectic manifolds with free circle actions, and Massey products*, Michigan Math. J. **38** (1991), no. 2, 271–283, DOI 10.1307/mmj/1029004333. MR1098863 (92e:57049)
- [6] Ronald Fintushel, *Classification of circle actions on 4-manifolds*, Trans. Amer. Math. Soc. **242** (1978), 377–390, DOI 10.2307/1997745. MR496815 (81e:57036)
- [7] S. Friedl and S. Vidussi, *Twisted Alexander polynomials detect fibered 3-manifolds*, Ann. Math. (2) **173** (2011), no. 3, 1587–1643, DOI 10.4007/annals.2011.173.3.8. MR2800721 (2012f:57025)
- [8] Stefan Friedl and Stefano Vidussi, *Twisted Alexander polynomials and fibered 3-manifolds*, Low-dimensional and symplectic topology, Proc. Sympos. Pure Math., vol. 82, Amer. Math. Soc., Providence, RI, 2011, pp. 111–130. MR2768657 (2012j:57028)
- [9] S. Friedl and S. Vidussi, *A vanishing theorem for twisted Alexander polynomials with applications to symplectic 4-manifolds*, J. Eur. Math. Soc. (JEMS) **15** (2013), no. 6, 2127–2041. MR3120739
- [10] Robert E. Gompf and András I. Stipsicz, *4-manifolds and Kirby calculus*, Graduate Studies in Mathematics, vol. 20, American Mathematical Society, Providence, RI, 1999. MR1707327 (2000h:57038)
- [11] William H. Meeks III and Peter Scott, *Finite group actions on 3-manifolds*, Invent. Math. **86** (1986), no. 2, 287–346, DOI 10.1007/BF01389073. MR856847 (88b:57039)
- [12] William P. Thurston, *A norm for the homology of 3-manifolds*, Mem. Amer. Math. Soc. **59** (1986), no. 339, i–vi and 99–130. MR823443 (88h:57014)
- [13] D. Wise, *The structure of groups with a quasiconvex hierarchy*, 187 pages, Preprint: available on the webpage for the NSF-CBMS conference.

MATHEMATISCHES INSTITUT, UNIVERSITÄT AUGSBURG, UNIVERSITÄTSSTRASSE 14, 86159 AUGSBURG, GERMANY

E-mail address: jonathan.bowden@math.uni-augsburg.de