HERMITIAN OPERATORS ON BANACH ALGEBRAS OF LIPSCHITZ FUNCTIONS

FERNANDA BOTELHO, JAMES JAMISON, A. JIMÉNEZ-VARGAS,
 AND MOISÉS VILLEGAS-VALLECILLOS

(Communicated by Thomas Schlumprecht)

ABSTRACT. For compact metric spaces (X, d) , we show that the Lipschitz spaces $\text{Lip}(X, d)$ and the little Lipschitz spaces $\text{lip}(X, d^\alpha)$ with $0 < \alpha < 1$, equipped with the sum norm, support only trivial hermitian operators, that is, real multiples of the identity operator.

1. INTRODUCTION

Let A be a complex Banach algebra with unity I and let A^* be its dual space. Given $a \in A$, recall that the algebraic numerical range $V(a)$ is given by

$$V(a) = \{F(a) : F \in A^*, \|F\| = F(I) = 1\}.$$

An element $a \in A$ is said to be hermitian if $V(a) \subset \mathbb{R}$. It is known that $a \in A$ is hermitian if and only if $\|\exp(ita)\| = 1$ for all $t \in \mathbb{R}$; see [3].

Let E be a complex Banach space and $\mathcal{B}(E)$ the Banach algebra of all bounded linear operators on E equipped with the operator norm. It is well-known that an operator $T \in \mathcal{B}(E)$ is hermitian if and only if $\exp(itT)$ is an isometry for each $t \in \mathbb{R}$; see [6, Theorem 5.2.6]. The set of hermitian operators on E is a real subspace of $\mathcal{B}(E)$ which contains all operators of the form λI , where λ is a real number. A hermitian operator is said to be trivial if it is a real multiple of the identity operator.

Some important Banach spaces only support trivial hermitian operators, as for example, the Bergman spaces $L_a^p(\Delta)$ ($1 \leq p < \infty$, $p \neq 2$) [9, Corollary 5.4] and the Hardy spaces $H^p(\Delta)$ ($1 \leq p < \infty$, $p \neq 2$) [1]. Also, the hermitian operators on several spaces of scalar-valued continuous functions defined on the interval $[0, 1]$ are known to be just real scalar multiples of the identity. Such spaces include the space of continuously differentiable functions $C^1[0, 1]$; the space of absolutely continuous functions $AC[0, 1]$; and the spaces of Lipschitz functions: $\text{Lip}[0, 1]$ and $\text{lip } \alpha$, $0 < \alpha < 1$. We recall that $\text{lip } \alpha$ consists of all period 1 functions on $\mathbb{R}$ satisfying $|f(x) - f(y)| = o(|x - y|^\alpha)$ uniformly as $|x - y| \rightarrow 0$; cf. [2, Theorem 3.1].

Received by the editors February 6, 2012 and, in revised form, September 5, 2012; September 13, 2012; and October 9, 2012.

2010 *Mathematics Subject Classification*. Primary 46E15, 47B15, 47B38.

Key words and phrases. Spaces of Lipschitz functions, hermitian operator, derivation, bi-circular projection.

The third and fourth authors were partially supported by Junta de Andalucía grants FQM-3737 and FQM-194 and by MICINN project MTM 2010-17687.

©2014 American Mathematical Society
 Reverts to public domain 28 years from publication

In this paper we investigate the hermitian operators on spaces of Lipschitz functions defined on a compact metric space. More precisely, for a compact metric space (X, d) and a positive real parameter $\alpha \in (0, 1]$, we consider the space of all α -Lipschitz functions $f: X \rightarrow \mathbb{C}$ such that

$$p_\alpha(f) := \sup \left\{ \frac{|f(x) - f(y)|}{d(x, y)^\alpha} : x, y \in X, x \neq y \right\} < \infty,$$

and also the subspace of all α -Lipschitz functions $f: X \rightarrow \mathbb{C}$ satisfying the additional local flatness condition:

$$\lim_{d(x, y) \rightarrow 0} \frac{|f(x) - f(y)|}{d(x, y)^\alpha} = 0.$$

These two spaces with the standard operations of addition, multiplication and scalar multiplication are complex algebras, and when equipped with the norm

$$\|f\|_\alpha = p_\alpha(f) + \|f\|_\infty$$

become Banach algebras. These two algebras are denoted by $\text{Lip}(X, d^\alpha)$ and $\text{lip}(X, d^\alpha)$, respectively.

It is important to observe that $\text{Lip}(X, d^\alpha)$ and $\text{lip}(X, d^\alpha)$ are unital semi-simple commutative complex Banach algebras, and $\text{lip}(X, d^\alpha)$ is a closed subalgebra of $\text{Lip}(X, d^\alpha)$. Notice that $\text{lip}(X, d)$ may contain only constant functions, for example $\text{lip}[0, 1]$ with the usual metric. When $X = [0, 1]$ or $X = \mathbb{T}$ with the usual metrics, $\text{Lip}(X, d^\alpha)$ and $\text{lip}(X, d^\alpha)$ are among the classical algebras considered by de Leeuw in [4]. These algebras were first studied by Sherbert in [14, 15].

In [2], it was shown that the hermitian operators on the Lipschitz spaces $\text{Lip}[0, 1]$ and $\text{lip } \alpha$, $0 < \alpha < 1$, are real multiples of the identity operator. In this paper we prove that the same property holds for the spaces $\text{Lip}(X, d)$ and the spaces $\text{lip}(X, d^\alpha)$ with $0 < \alpha < 1$, for (X, d) a compact metric space. This generalizes the aforementioned result.

We also mention the natural connection between hermitian operators and the class of bi-circular projections. A projection P on a complex Banach space is bi-circular if $e^{is}P + e^{it}(I - P)$ is an isometry for all $s, t \in \mathbb{R}$. This type of projection was introduced by Stachó and Zalar in [17]. Jamison [10] showed that these projections are exactly the hermitian projections. Our result implies that the only bi-circular projections on $\text{Lip}(X, d)$ and $\text{lip}(X, d^\alpha)$ with $0 < \alpha < 1$ are the trivial projections, 0 and I .

2. PRELIMINARIES

In this section we give a representation for all surjective linear isometries on $\text{Lip}(X, d)$ or $\text{lip}(X, d^\alpha)$ ($0 < \alpha < 1$) that fix the constant function everywhere equal to 1. Then we characterize the hermitian elements of $\text{Lip}(X, d)$ and $\text{lip}(X, d^\alpha)$ ($0 < \alpha < 1$). The last result provides a useful description of the continuous linear functionals on both spaces.

Throughout this paper (X, d) is a compact metric space, 1_X denotes the constant function equal to 1 on X , I_X represents the identity function on X and I is the identity operator on $\text{Lip}(X, d)$ or $\text{lip}(X, d^\alpha)$, $0 < \alpha < 1$. For each $x \in X$, δ_x stands for the evaluation functional at the point x defined on $\text{Lip}(X, d)$ or $\text{lip}(X, d^\alpha)$, $0 < \alpha < 1$.

Our approach requires that the surjective linear isometries on the spaces $\text{Lip}(X, d)$ and $\text{lip}(X, d^\alpha)$ ($0 < \alpha < 1$) have a suitable representation. Rao and Roy [13] proved that any surjective linear isometry of $\text{Lip}[0, 1]$ can be expressed as a weighted composition operator $f \mapsto \tau f \circ \varphi$ ($f \in \text{Lip}[0, 1]$) where τ is a scalar of modulus 1 and φ is a surjective isometry of $[0, 1]$. They asked whether every isometry on the Banach spaces $\text{Lip}(X, d)$ and $\text{lip}(X, d^\alpha)$ ($0 < \alpha < 1$) are induced by the isometries of the metric space X . Next we derive a characterization of surjective linear isometries on these spaces that fix 1_X . This characterization follows from a theorem due to Jarosz in [11], a theorem in [8] (page 144) and a result by Sherbert in [14].

Theorem 2.1. *Let X be a compact metric space. Then $T: \text{Lip}(X, d) \rightarrow \text{Lip}(X, d)$ is a surjective linear isometry such that $T(1_X) = 1_X$ if and only if there exists a surjective isometry $\varphi: X \rightarrow X$ such that T is of the form $T(f) = f \circ \varphi$ for all $f \in \text{Lip}(X, d)$. The same characterization holds for a surjective linear isometry T on $\text{lip}(X, d^\alpha)$ ($0 < \alpha < 1$) such that $T(1_X) = 1_X$.*

Proof. It is straightforward to check that an operator T of the form described in the theorem is a surjective isometry. Then we just prove the reversed implication.

Let $A(X)$ represent either $\text{Lip}(X, d)$ or $\text{lip}(X, d^\alpha)$ with $0 < \alpha < 1$ and let $C(X)$ be the algebra of continuous complex-valued functions on X . We first observe that $A(X)$ is a regular subspace of $C(X)$ and the sum norm is a p -norm for the norm on $\mathbb{R}^2$ given by $p(s, t) = |s| + |t|$. Let us recall (see [11]) that given a compact Hausdorff space X , a complex linear subspace A of $C(X)$ that contains the function 1_X , is said to be regular if for any $\varepsilon > 0$, any $x_0 \in \text{Ch}A$ and any open neighborhood U of x_0 , there is an $f \in A$ with $\|f\|_\infty \leq 1 + \varepsilon$, $f(x_0) = 1$, and $|f(x)| < \varepsilon$ for $x \in X \setminus U$. $\text{Ch}A$ denotes the set of extreme points F of the unit ball of $(A, \|\cdot\|_\infty)^*$ such that $F(1_X) = 1$, and we identify $\text{Ch}A$ with a subset of X . Suppose that T is a surjective linear isometry on $A(X)$ such that $T(1_X) = 1_X$. An application of the main theorem in [11] to $A(X)$ yields that T is a surjective isometry on $(A(X), \|\cdot\|_\infty)$.

Next we quote a theorem from Hoffman's book [8, p. 44]: Let X be a compact Hausdorff space and let B be a complex linear subalgebra of $C(X)$ that contains the function 1_X . Suppose that S is a linear map of B onto B such that $\|S(f)\|_\infty = \|f\|_\infty$ for all $f \in B$. If $S(1_X) = 1_X$, then S is multiplicative.

Therefore T is an automorphism of $A(X)$. By Sherbert's theorem [14, Corollary 5.2], every automorphism T of $\text{Lip}(X, d)$ that carries 1_X into 1_X is of the form $T(f) = f \circ \varphi$, where $\varphi: X \rightarrow X$ is a homeomorphism. Similarly, we can prove that this is also true for those automorphisms of $\text{lip}(X, d^\alpha)$ ($0 < \alpha < 1$) that fix 1_X .

We now show that φ is an isometry of X . Observe that given any $\alpha \in (0, 1]$, we have $p_\alpha(T(f)) = p_\alpha(f)$ for all $f \in A(X)$ since T is an isometry for both norms $\|\cdot\|_\alpha$ and $\|\cdot\|_\infty$.

For the case $A(X) = \text{Lip}(X, d)$, fix $y \in X$ and define $f_y: X \rightarrow \mathbb{R}$ by $f_y(z) = d(z, \varphi(y))$ for all $z \in X$. Clearly, $f_y \in \text{Lip}(X, d)$ and $p_1(f_y) \leq 1$. For any $x, y \in X$, we have

$$\begin{aligned} d(\varphi(x), \varphi(y)) &= |f_y(\varphi(x)) - f_y(\varphi(y))| \\ &= |T(f_y)(x) - T(f_y)(y)| \\ &\leq p_1(T(f_y))d(x, y) \\ &\leq d(x, y). \end{aligned}$$

For the case $A(X) = \text{lip}(X, d^\alpha)$ ($0 < \alpha < 1$), fix $x, y \in X$, $x \neq y$, choose $\beta \in (\alpha, 1)$ and define

$$f_{xy}(z) = \frac{d(z, \varphi(y))^\beta - d(z, \varphi(x))^\beta}{2d(\varphi(x), \varphi(y))^{\beta-\alpha}}, \quad \forall z \in X.$$

It is not hard to check that $f_{xy} \in \text{lip}(X, d^\alpha)$ and $p_\alpha(f_{xy}) = 1$ (see, for example, [12, p. 62]). An easy calculation gives

$$\begin{aligned} d(\varphi(x), \varphi(y))^\alpha &= |f_{xy}(\varphi(x)) - f_{xy}(\varphi(y))| \\ &= |T(f_{xy})(x) - T(f_{xy})(y)| \\ &\leq p_\alpha(T(f_{xy}))d(x, y)^\alpha \\ &= d(x, y)^\alpha. \end{aligned}$$

In either case we have $d(\varphi(x), \varphi(y)) \leq d(x, y)$ for all $x, y \in X$.

Since T^{-1} is also a surjective linear isometry on $A(X)$ such that $T^{-1}(1_X) = 1_X$, the same argument used above implies the existence of a homeomorphism $\phi: X \rightarrow X$ such that $T^{-1}(f) = f \circ \phi$ for all $f \in A(X)$. Therefore $d(\phi(x), \phi(y)) \leq d(x, y)$ for all $x, y \in X$. Given $x \in X$, we have

$$f(\varphi^{-1}(x)) = T(T^{-1}(f))(\varphi^{-1}(x)) = T^{-1}(f)(x) = f(\phi(x))$$

for all $f \in A(X)$. Since $A(X)$ separates the points of X , this implies that $\varphi^{-1} = \phi$. Consequently, φ is a surjective isometry. This completes the proof of the theorem. $\square$

We will next characterize the hermitian elements of the spaces $\text{Lip}(X, d)$ and $\text{lip}(X, d^\alpha)$, $0 < \alpha < 1$.

Lemma 2.2. *Let (X, d) be a compact metric space and $h \in \text{Lip}(X, d)$ (or $\text{lip}(X, d^\alpha)$, $0 < \alpha < 1$). Then h is a hermitian element in $\text{Lip}(X, d)$ (or $\text{lip}(X, d^\alpha)$) if and only if h is a real constant function.*

Proof. Assume that h is hermitian in $\text{Lip}(X, d)$. Then $F(h) \in V(h) \subset \mathbb{R}$ for all $F \in \text{Lip}(X, d)^*$ such that $\|F\| = F(1_X) = 1$. In particular, $h(x) = \delta_x(h) \in \mathbb{R}$ for all $x \in X$, and so h is real-valued. Using that $|e^a - e^b| \leq |a - b| \exp(\max\{|a|, |b|\})$ for all $a, b \in \mathbb{C}$, we deduce that $\exp(ih)$ is a function in $\text{Lip}(X, d)$. We also have that, for each $t \in \mathbb{R}$, $\|\exp(i th)\|_1 = 1$. Since $\|\exp(i th)\|_\infty = 1$, it follows that $p_1(\exp(i th)) = 0$. Hence $\exp(i th)$ is a constant function on X for all $t \in \mathbb{R}$ which implies that h is constant.

Conversely, if h is a real constant function, then h is a real multiple of 1_X . Therefore h is hermitian in $\text{Lip}(X, d)$. The same proof works for $\text{lip}(X, d^\alpha)$, $0 < \alpha < 1$. $\square$

Following an idea of de Leeuw [4], we embed the Banach spaces $\text{Lip}(X, d)$ and $\text{lip}(X, d^\alpha)$ ($0 < \alpha < 1$) isometrically into some suitable spaces of complex-valued continuous functions.

Let X be a compact metric space and let $\tilde{X}$ be the set $\{(x, y) \in X^2: x \neq y\}$. It is easy to check that $\tilde{X}$ is completely regular; we denote by $\beta\tilde{X}$ the Stone-Ćech compactification of $\tilde{X}$. Let $C(X \cup \beta\tilde{X})$ denote the Banach space of all complex-valued continuous functions on $X \cup \beta\tilde{X}$, under the norm

$$\|f\| = \|f|_X\|_\infty + \left\|f|_{\beta\tilde{X}}\right\|_\infty \quad (f \in C(X \cup \beta\tilde{X})),$$

and let $C_0(X \cup \tilde{X})$ denote the Banach space of all complex-valued continuous functions on $X \cup \tilde{X}$ vanishing at infinity, endowed with the norm

$$\|f\| = \|f|_X\|_\infty + \|f|_{\tilde{X}}\|_\infty \quad (f \in C_0(X \cup \tilde{X})).$$

We now recall that the Riesz representation theorem states that the map $\mu \mapsto F_\mu$, given by

$$F_\mu(f) = \int_{X \cup \beta\tilde{X}} f \, d\mu \quad (f \in C(X \cup \beta\tilde{X})),$$

defines an isometric isomorphism from the Banach space $\mathcal{M}(X \cup \beta\tilde{X})$ of all complex-valued regular Borel measures on $X \cup \beta\tilde{X}$ equipped with the norm of total variation:

$$\|\mu\| = |\mu|(X \cup \beta\tilde{X}) \quad (\mu \in \mathcal{M}(X \cup \beta\tilde{X}))$$

onto the dual space of $(C(X \cup \beta\tilde{X}), \|\cdot\|_\infty)$. Similarly, the map $\nu \mapsto G_\nu$ defined by

$$G_\nu(f) = \int_{X \cup \tilde{X}} f \, d\nu \quad (f \in C_0(X \cup \tilde{X}))$$

is an isometric isomorphism from the Banach space $\mathcal{M}(X \cup \tilde{X})$ with the norm

$$\|\nu\| = |\nu|(X \cup \tilde{X}) \quad (\nu \in \mathcal{M}(X \cup \tilde{X}))$$

onto the dual space of $(C_0(X \cup \tilde{X}), \|\cdot\|_\infty)$.

For each $f \in \text{Lip}(X, d)$ or $f \in \text{lip}(X, d^\alpha)$, $0 < \alpha < 1$, we set $\tilde{f}: \tilde{X} \rightarrow \mathbb{C}$ to be the map given by

$$\tilde{f}(x, y) = \frac{f(x) - f(y)}{d(x, y)^\alpha}, \quad \forall (x, y) \in \tilde{X},$$

where $\alpha = 1$ when $f \in \text{Lip}(X, d)$. It is easy to show that $\tilde{f}$ is continuous on $\tilde{X}$ and $\|\tilde{f}\|_\infty = p_\alpha(f)$ ($0 < \alpha \leq 1$). Hence there exists a unique continuous function $\beta\tilde{f}$ on $\beta\tilde{X}$ such that $(\beta\tilde{f})|_{\tilde{X}} = \tilde{f}$ and $\|\beta\tilde{f}\|_\infty = \|\tilde{f}\|_\infty$. Furthermore, if $f \in \text{lip}(X, d^\alpha)$, then $\tilde{f}$ vanishes at infinity on $\tilde{X}$. The maps $\Phi: \text{Lip}(X, d) \rightarrow C(X \cup \beta\tilde{X})$ and $\Psi: \text{lip}(X, d^\alpha) \rightarrow C_0(X \cup \tilde{X})$, defined by

$$(1) \quad \Phi(f)(w) = \begin{cases} f(w) & \text{if } w \in X, \\ \beta\tilde{f}(w) & \text{if } w \in \beta\tilde{X}, \end{cases}$$

and

$$(2) \quad \Psi(f)(w) = \begin{cases} f(w) & \text{if } w \in X, \\ \tilde{f}(w) & \text{if } w \in \tilde{X}, \end{cases}$$

are isometric linear embeddings from $\text{Lip}(X, d)$ with the norm $\|\cdot\|_1$ into $C(X \cup \beta\tilde{X})$, and from $\text{lip}(X, d^\alpha)$ with the norm $\|\cdot\|_\alpha$ into $C_0(X \cup \tilde{X})$, respectively.

The Hahn–Banach theorem and the Riesz representation theorem yield the following lemma.

Lemma 2.3. *Let (X, d) be a compact metric space.*

- (1) For each $F \in \text{Lip}(X, d)^*$, there exists $\mu \in \mathcal{M}(X \cup \beta\tilde{X})$ with $\|F\| \leq \|\mu\|$ satisfying

$$F(f) = \int_{X \cup \beta\tilde{X}} \Phi(f)(w) d\mu(w), \quad \forall f \in \text{Lip}(X, d).$$

- (2) Let $\alpha \in (0, 1)$. For each $G \in \text{lip}(X, d^\alpha)^*$, there exists $\nu \in \mathcal{M}(X \cup \tilde{X})$ with $\|G\| \leq \|\nu\|$ such that

$$G(f) = \int_{X \cup \tilde{X}} \Psi(f)(w) d\nu(w), \quad \forall f \in \text{lip}(X, d^\alpha).$$

Proof. Let $F \in \text{Lip}(X, d)^*$. The functional $T: \Phi(\text{Lip}(X, d)) \rightarrow \mathbb{C}$, defined by $T(\Phi(f)) = F(f)$ for all $f \in \text{Lip}(X, d)$, is linear, continuous and $\|T\| = \|F\|$. By the Hahn–Banach theorem, there exists a linear continuous functional $\tilde{T}: C(X \cup \beta\tilde{X}) \rightarrow \mathbb{C}$ such that $\tilde{T}(\Phi(f)) = T(\Phi(f))$ for all $f \in \text{Lip}(X, d)$ and $\|\tilde{T}\| = \|T\|$.

Since $\|g\| \leq 2\|g\|_\infty$ for all $g \in C(X \cup \beta\tilde{X})$, it follows that the linear functional $\tilde{T}$ is continuous on the space $C(X \cup \beta\tilde{X})$ equipped with the norm $\|\cdot\|_\infty$. We denote by $\|\cdot\|_\infty^*$ the norm on the dual Banach space of $(C(X \cup \beta\tilde{X}), \|\cdot\|_\infty)$. By the Riesz representation theorem, there exists $\mu \in \mathcal{M}(X \cup \beta\tilde{X})$ with $\|\tilde{T}\|_\infty^* = \|\mu\|$ satisfying

$$\tilde{T}(g) = \int_{X \cup \beta\tilde{X}} g(w) d\mu(w), \quad \forall g \in C(X \cup \beta\tilde{X}).$$

Since $\|g\|_\infty \leq \|g\|$ for all $g \in C(X \cup \beta\tilde{X})$, we have $\|\tilde{T}\| \leq \|\tilde{T}\|_\infty^*$, and so $\|F\| \leq \|\mu\|$. Moreover,

$$F(f) = T(\Phi(f)) = \tilde{T}(\Phi(f)) = \int_{X \cup \beta\tilde{X}} \Phi(f)(w) d\mu(w)$$

for all $f \in \text{Lip}(X, d)$, as we wanted. Similarly, we prove statement (2). $\square$

Such a μ is called a representing measure for F (analogously, ν for G). We should note that a representing measure for F or G is not always determined uniquely.

3. THE MAIN RESULT

In this section we describe all the hermitian operators on $\text{Lip}(X, d)$ or $\text{lip}(X, d^\alpha)$ with $0 < \alpha < 1$. We proceed with the statement and proof of our main result.

Theorem 3.1. *Let (X, d) be a compact metric space. A bounded linear operator $T: \text{Lip}(X, d) \rightarrow \text{Lip}(X, d)$ is hermitian if and only if T is a real multiple of the identity operator on $\text{Lip}(X, d)$. An analogous assertion holds for $T: \text{lip}(X, d^\alpha) \rightarrow \text{lip}(X, d^\alpha)$ with $0 < \alpha < 1$.*

Before proving this theorem we set notation and prove some preliminary lemmas. Let $A(X)$ denote either $\text{Lip}(X, d)$ or $\text{lip}(X, d^\alpha)$, $0 < \alpha < 1$. Recall that $\alpha = 1$ in the case $A(X) = \text{Lip}(X, d)$.

Lemma 3.2. *If $T: A(X) \rightarrow A(X)$ is a hermitian bounded linear operator, then the following statements hold:*

- (i) *There exists $\lambda \in \mathbb{R}$ such that $T(1_X) = \lambda 1_X$.*
- (ii) *For each $t \in \mathbb{R}$, $\exp(it(T - \lambda I))$ is a surjective linear isometry on $A(X)$ fixing 1_X .*

(iii) For each $t \in \mathbb{R}$, there exists a surjective isometry φ_t on X such that

$$\exp(it(T - \lambda I))(f)(x) = f(\varphi_t(x)), \quad \forall f \in A(X), \forall x \in X.$$

(iv) $\{\varphi_t\}_{t \in \mathbb{R}}$ is a one-parameter group of surjective isometries on X such that, for each $x \in X$, the map $t \mapsto \varphi_t(x)$ from $\mathbb{R}$ to X is continuous.

(v) For every $f \in A(X)$,

$$(3) \quad \lim_{t \rightarrow 0} (f \circ \varphi_t - f)(x) = 0, \quad \forall x \in X,$$

and

$$(4) \quad \lim_{t \rightarrow 0} \frac{(f \circ \varphi_t - f)(x) - (f \circ \varphi_t - f)(y)}{d(x, y)^\alpha} = 0, \quad \forall (x, y) \in \tilde{X}.$$

Proof. (i) For each $F \in A(X)^*$ with $\|F\| = F(1_X) = 1$, define $\Phi_F: \mathcal{B}(A(X)) \rightarrow \mathbb{C}$ by

$$\Phi_F(S) = F(S(1_X)), \quad \forall S \in \mathcal{B}(A(X)).$$

It is easy to check that Φ_F is a linear functional on $\mathcal{B}(A(X))$, and since

$$|\Phi_F(S)| = |F(S(1_X))| \leq \|F\| \|S(1_X)\|_\alpha \leq \|S\| \|1_X\|_\alpha = \|S\|,$$

for all $S \in \mathcal{B}(A(X))$, then Φ_F is continuous and $\|\Phi_F\| \leq 1$. Moreover, $\Phi_F(I) = F(1_X) = 1$; hence $\|\Phi_F\| \geq |\Phi_F(I)| = 1$ and thus $\|\Phi_F\| = \Phi_F(I) = 1$.

Since $T \in \mathcal{B}(A(X))$ is hermitian, it follows that $F(T(1_X)) = \Phi_F(T) \in V(T) \subset \mathbb{R}$ for all $F \in A(X)^*$ such that $\|F\| = F(1_X) = 1$. This means that $T(1_X)$ is a hermitian element in $A(X)$. Then, according to Lemma 2.2, there exists $\lambda \in \mathbb{R}$ such that $T(1_X) = \lambda 1_X$.

(ii) By (i), we have $(T - \lambda I)(1_X) = 0$ and so $\exp(it(T - \lambda I))(1_X) = 1_X$ for all $t \in \mathbb{R}$. Indeed, since

$$\exp(it(T - \lambda I)) = I + \sum_{n=1}^{\infty} i^n t^n (T - \lambda I)^n,$$

it follows that

$$\exp(it(T - \lambda I))(1_X) = 1_X + \sum_{n=1}^{\infty} i^n t^n (T - \lambda I)^n(1_X) = 1_X.$$

Since T and λI are hermitian operators in $\mathcal{B}(A(X))$, it is easily seen that so is $T - \lambda I$. Indeed, using the fact that $\exp(it(T - \lambda I)) = \exp(itT)\exp(-it\lambda I)$ for all $t \in \mathbb{R}$, we have

$$\begin{aligned} 1 &= \|1_X\|_\alpha = \|\exp(it(T - \lambda I))(1_X)\|_\alpha \leq \|\exp(it(T - \lambda I))\| \\ &\leq \|\exp(itT)\| \|\exp(-it\lambda I)\| = 1. \end{aligned}$$

Therefore, for each $t \in \mathbb{R}$, $\exp(it(T - \lambda I))$ is a linear isometry from $A(X)$ onto itself, fixing 1_X .

(iii) In view of (ii), by applying Theorem 2.1, for each $t \in \mathbb{R}$ there exists a surjective isometry φ_t on X such that

$$(5) \quad \exp(it(T - \lambda I))(f)(x) = f(\varphi_t(x)), \quad \forall f \in A(X), \forall x \in X.$$

(iv) Using the fact that $A(X)$ separates the points of X , it is easily derived that $\varphi_{(s+t)} = \varphi_s \circ \varphi_t$ for all $s, t \in \mathbb{R}$ and $\varphi_0 = I_X$. More precisely, given $f \in A(X)$ and $x \in X$, we have

$$\begin{aligned} f(\varphi_{s+t}(x)) &= \exp(i(s+t)(T - \lambda I))(f)(x) \\ &= \exp(i(t+s)(T - \lambda I))(f)(x) \\ &= \exp(it(T - \lambda I))\exp(is(T - \lambda I))(f)(x) \\ &= \exp(it(T - \lambda I))(\exp(is(T - \lambda I))(f))(x) \\ &= \exp(is(T - \lambda I))(f)(\varphi_t(x)) \\ &= f(\varphi_s(\varphi_t(x))) \\ &= f(\varphi_s \circ \varphi_t(x)) \end{aligned}$$

and

$$f(\varphi_0(x)) = \exp(i0(T - \lambda I))(f)(x) = \exp(0)(f)(x) = I(f)(x) = f(x).$$

We next prove that for each $x \in X$, the map $t \mapsto \varphi_t(x)$ from $\mathbb{R}$ to X is continuous. Note first that $\delta: X \rightarrow A(X)^*$ defined by $\delta(x) = \delta_x$ is a Lipschitz bijection from (X, d^α) onto $\delta(X)$. Indeed, δ is injective since $A(X)$ separates points; and given $x, y \in X$, we have

$$|(\delta(x) - \delta(y))(f)| = |f(x) - f(y)| \leq \|f\|_\alpha d(x, y)^\alpha$$

for all $f \in A(X)$. Hence $\|\delta(x) - \delta(y)\| \leq d(x, y)^\alpha$. Since X is compact, we deduce that $\delta^{-1}: \delta(X) \rightarrow X$ is continuous. Notice that $\delta^{-1}(\delta_x) = x$ for all $x \in X$.

Fix $x \in X$. The maps $t \mapsto \exp(it(T - \lambda I))$ from $\mathbb{R}$ to $\mathcal{B}(A(X))$, $U \mapsto U^*$ from $\mathcal{B}(A(X))$ to $\mathcal{B}(A(X)^*)$ and $S \mapsto S(\delta(x))$ from $\mathcal{B}(A(X)^*)$ to $A(X)^*$ are clearly continuous. From (5), we deduce that

$$(6) \quad (\exp(it(T - \lambda I)))^*(\delta(x)) = \delta(\varphi_t(x)) \quad (t \in \mathbb{R}, x \in X).$$

Since

$$\varphi_t(x) = \delta^{-1}(\exp(it(T - \lambda I))^*(\delta(x))) \quad (t \in \mathbb{R}, x \in X),$$

we conclude that $t \mapsto \varphi_t(x)$ from $\mathbb{R}$ to X is continuous.

(v) Let $f \in A(X)$. Given $x \in X$, we have $\lim_{t \rightarrow 0}(f \circ \varphi_t)(x) = (f \circ \varphi_0)(x) = f(x)$ by (iv), and thus $\lim_{t \rightarrow 0}(f \circ \varphi_t - f)(x) = 0$. Using this, for $(x, y) \in \tilde{X}$, we deduce that

$$\lim_{t \rightarrow 0} \frac{(f \circ \varphi_t - f)(x) - (f \circ \varphi_t - f)(y)}{d(x, y)^\alpha} = 0.$$

□

We recall that for $f \in A(X)$ the map $\tilde{f}: \tilde{X} \rightarrow \mathbb{C}$ is defined to be

$$\tilde{f}(x, y) = (f(x) - f(y))/d(x, y)^\alpha.$$

We recall that $\beta\tilde{X}$ represents the Stone-Ćech compactification of $\tilde{X}$. This entails that every bounded, continuous and scalar-valued map defined on $\tilde{X}$ has a unique continuous extension to $\beta\tilde{X}$.

Lemma 3.3. *If $f \in A(X)$, then*

$$(7) \quad \lim_{t \rightarrow 0} \beta\tilde{f}_t(w) = 0, \quad \forall w \in \beta\tilde{X},$$

where, for each $t \in \mathbb{R}$, f_t denotes the function $f \circ \varphi_t - f$.

Proof. We define $g: [-1, 1] \times \tilde{X} \rightarrow \mathbb{C}$ by

$$g(t, (x, y)) = \frac{(f \circ \varphi_t - f)(x) - (f \circ \varphi_t - f)(y)}{d(x, y)^\alpha}.$$

The function g is continuous and bounded. In fact, we have

$$|g(t, (x, y))| \leq p_\alpha(f) + p_\alpha(f) = 2p_\alpha(f)$$

for all $t \in [-1, 1]$ and $(x, y) \in \tilde{X}$. For the continuity of g , define $\sigma: [-1, 1] \times \tilde{X} \rightarrow A(X)^*$ by

$$\sigma(t, (x, y)) = \frac{\delta(\varphi_t(x)) - \delta(x) - (\delta(\varphi_t(y)) - \delta(y))}{d(x, y)^\alpha}$$

and notice that

$$g(t, (x, y)) = \sigma(t, (x, y))(f) \quad (t \in [-1, 1], (x, y) \in \tilde{X}).$$

Taking into account the equality (6), for any $t, s \in [-1, 1]$ and $x, y \in X$, we have

$$\begin{aligned} \|\delta(\varphi_t(x)) - \delta(\varphi_s(y))\| &= \|(\exp(it(T - \lambda I)))^*(\delta(x)) - (\exp(is(T - \lambda I)))^*(\delta(y))\| \\ &\leq \|(\exp(it(T - \lambda I)))^*\| \|\delta(x) - \delta(y)\| \\ &\quad + \|(\exp(it(T - \lambda I)))^* - (\exp(is(T - \lambda I)))^*\| \|\delta(y)\| \\ &\leq d(x, y) + \|\exp(it(T - \lambda I)) - \exp(is(T - \lambda I))\| \|\delta(y)\|. \end{aligned}$$

Let us recall now that if A and B are bounded commuting operators on a Banach algebra, then

$$\|\exp(iA) - \exp(iB)\| \leq \|A - B\| \exp(\max\{\|A\|, \|B\|\}).$$

Applying this formula to $A = t(T - \lambda I)$ and $B = s(T - \lambda I)$, we obtain

$$(8) \quad \|\exp(it(T - \lambda I)) - \exp(is(T - \lambda I))\| \leq |t - s| k,$$

where $k = \|T - \lambda I\| \exp(\|T - \lambda I\|)$ is a constant, and so

$$\|\delta(\varphi_t(x)) - \delta(\varphi_s(y))\| \leq d(x, y)^\alpha + k |t - s|.$$

Therefore, for every $t, s \in [-1, 1]$ and $x, y \in X$, we have

$$\|\delta(\varphi_t(x)) - \delta(x) - (\delta(\varphi_s(y)) - \delta(y))\| \leq 2d(x, y)^\alpha + k |t - s|.$$

Hence the mapping $(t, (x, y)) \mapsto \delta(\varphi_t(x)) - \delta(x)$, defined on $[-1, 1] \times \tilde{X}$ and with values in $A(X)^*$, is continuous. Since $(x, y) \mapsto d(x, y)^\alpha$ from $\tilde{X}$ to $\mathbb{R}$ is continuous, it follows that σ is continuous. Hence, given $\varepsilon > 0$ and $(t_0, (x_0, y_0)) \in [-1, 1] \times \tilde{X}$, there is a neighborhood V of $(t_0, (x_0, y_0))$ such that if $(t, (x, y)) \in V$, then $\|\sigma(t, (x, y)) - \sigma(t_0, (x_0, y_0))\| < \varepsilon/(1 + \|f\|_\alpha)$. Therefore, for every $(t, (x, y)) \in V$, we have

$$|g(t, (x, y)) - g(t_0, (x_0, y_0))| < \frac{\varepsilon}{1 + \|f\|_\alpha} \|f\|_\alpha < \varepsilon,$$

and this proves that g is continuous.

We now check that the mapping $t \mapsto g(t, \cdot)$ from $[-1, 1]$ into $C(\widetilde{X})$ is continuous. Indeed, by using (6) and (8) we have

$$\begin{aligned} & |g(t, (x, y)) - g(s, (x, y))| \\ &= \left| \frac{\delta(\varphi_t(x)) - \delta(\varphi_t(y)) - (\delta(\varphi_s(x)) - \delta(\varphi_s(y)))}{d(x, y)^\alpha} (f) \right| \\ &\leq \left\| \frac{((\exp(it(T - \lambda I)))^* - (\exp(is(T - \lambda I)))^*)(\delta(x) - \delta(y))}{d(x, y)^\alpha} \right\| \|f\|_\alpha \\ &\leq k \|f\|_\alpha |t - s| \end{aligned}$$

for all $t, s \in [-1, 1]$ and $(x, y) \in \widetilde{X}$, and so $\|g(t, \cdot) - g(s, \cdot)\|_\infty \leq k \|f\|_\alpha |t - s|$.

Then g has a continuous extension h from $[-1, 1] \times \beta\widetilde{X}$ to $\mathbb{C}$ by [7, Lemma 2]. Hence $h(0, (x, y)) = g(0, (x, y)) = 0$ for all $(x, y) \in \widetilde{X}$. Now fix $w \in \beta\widetilde{X}$ and take a net $\{(x_i, y_i)\}$ in $\widetilde{X}$ converging to w . The continuity of h yields $h(0, w) = \lim_i h(0, (x_i, y_i)) = 0$. Using the continuity of $\beta\widetilde{f}_t$, with $t \in [-1, 1]$, and the continuity of h , we have

$$\begin{aligned} \lim_{t \rightarrow 0} \beta\widetilde{f}_t(w) &= \lim_{t \rightarrow 0} \lim_i \beta\widetilde{f}_t(x_i, y_i) = \lim_{t \rightarrow 0} \lim_i \widetilde{f}_t(x_i, y_i) = \lim_{t \rightarrow 0} \lim_i g(t, (x_i, y_i)) \\ &= \lim_{t \rightarrow 0} \lim_i h(t, (x_i, y_i)) = \lim_{t \rightarrow 0} h(t, w) = h(0, w) = 0. \end{aligned}$$

This completes the proof. $\square$

We now proceed with the details for the proof of our main theorem.

Proof of Theorem 3.1. Assume that $T: A(X) \rightarrow A(X)$ is a hermitian bounded linear operator. In view of (3), (4) and (7), the maps defined in (1) and (2) satisfy

$$(9) \quad \lim_{t \rightarrow 0} \Phi(f \circ \varphi_t - f)(w) = 0, \quad \forall w \in X \cup \beta\widetilde{X},$$

and

$$(10) \quad \lim_{t \rightarrow 0} \Psi(f \circ \varphi_t - f)(w) = 0, \quad \forall w \in X \cup \widetilde{X}.$$

Note also that

$$(11) \quad |\Phi(f \circ \varphi_t - f)(w)| \leq \|\Phi(f \circ \varphi_t - f)\| = \|f \circ \varphi_t - f\|_1 \leq 2 \|f\|_1, \\ \forall t \in \mathbb{R}, \forall w \in X \cup \beta\widetilde{X},$$

and

$$(12) \quad |\Psi(f \circ \varphi_t - f)(w)| \leq \|\Psi(f \circ \varphi_t - f)\| = \|f \circ \varphi_t - f\|_\alpha \leq 2 \|f\|_\alpha, \\ \forall t \in \mathbb{R}, \forall w \in X \cup \widetilde{X}.$$

Now, for each $t \in \mathbb{R}$, define $T_t: A(X) \rightarrow A(X)$ by

$$(13) \quad T_t(f)(x) = f(\varphi_t(x)), \quad \forall f \in A(X), \forall x \in X.$$

Each T_t is an isometric algebra automorphism of $A(X)$. Moreover, it is immediate that $T_{(s+t)} = T_s \circ T_t$ for all $s, t \in \mathbb{R}$ and $T_0 = I$. Indeed, for any $f \in A(X)$, we have

$$T_{(s+t)}(f) = f \circ \varphi_{(s+t)} = f \circ \varphi_{(t+s)} = f \circ \varphi_t \circ \varphi_s = T_s(f \circ \varphi_t) = T_s(T_t(f)) = T_s \circ T_t(f)$$

and

$$T_0(f) = f \circ \varphi_0 = f \circ I_X = f.$$

Thus $\{T_t: t \in \mathbb{R}\}$ is a one-parameter group of isometric algebra automorphisms of $A(X)$. Indeed, notice that $T_{(-t)} = (T_t)^{-1}$ for every $t \in \mathbb{R}$ since

$$I = T_0 = T_{(t+(-t))} = T_t \circ T_{(-t)}, \quad I = T_0 = T_{((-t)+t)} = T_{(-t)} \circ T_t \quad (t \in \mathbb{R}).$$

Next we will prove that $\{T_t: t \in \mathbb{R}\}$ is strongly continuous; that is, the maps $t \mapsto T_t(f)$ from $\mathbb{R}$ to $A(X)$ are continuous for each $f \in A(X)$ [5, Chapter I, 1.1 Definition]. According to [5, Chapter I, 1.6 Theorem], it is sufficient to show that $\{T_t: t \in \mathbb{R}\}$ is weakly continuous; that is, the maps $t \mapsto F(T_t(f))$ from $\mathbb{R}$ to $\mathbb{C}$ are continuous for every $f \in A(X)$ and $F \in A(X)^*$. To prove this, notice that we only need to see that the aforementioned maps are continuous at 0, and it is easily checked.

Let $f \in \text{Lip}(X, d)$ and $F \in \text{Lip}(X, d)^*$. By Lemma 2.3 (1), there exists $\mu \in \mathcal{M}(X \cup \beta\tilde{X})$ such that

$$F(g) = \int_{X \cup \beta\tilde{X}} \Phi(g)(w) d\mu(w) \quad (g \in \text{Lip}(X, d)).$$

In particular, we have

$$F(T_t(f) - f) = \int_{X \cup \beta\tilde{X}} \Phi(T_t(f) - f)(w) d\mu(w) \quad (t \in \mathbb{R}).$$

Taking into account (9) and (11), we can apply Lebesgue's bounded convergence theorem and get

$$\begin{aligned} \lim_{t \rightarrow 0} F(T_t(f) - f) &= \lim_{t \rightarrow 0} \int_{X \cup \beta\tilde{X}} \Phi(T_t(f) - f)(w) d\mu(w) \\ &= \lim_{t \rightarrow 0} \int_{X \cup \beta\tilde{X}} \Phi(f \circ \varphi_t - f)(w) d\mu(w) \\ &= \int_{X \cup \beta\tilde{X}} \lim_{t \rightarrow 0} \Phi(f \circ \varphi_t - f)(w) d\mu(w) = 0. \end{aligned}$$

Similarly, for any $f \in \text{lip}(X, d^\alpha)$ and $G \in \text{lip}(X, d^\alpha)^*$, we obtain

$$\lim_{t \rightarrow 0} G(T_t(f) - f) = 0$$

by using Lemma 2.3 (2) and the equalities (10) and (12).

Let $A: D(A) \subset A(X) \rightarrow A(X)$ be the generator of $\{T_t: t \in \mathbb{R}\}$. That is, A is the linear operator

$$A(f) = \lim_{t \rightarrow 0} \frac{T_t(f) - f}{t}$$

defined for every f in its domain

$$D(A) = \left\{ f \in A(X) : \exists \lim_{t \rightarrow 0} \frac{T_t(f) - f}{t} \right\}$$

(see [5, Chapter II, 1.2 Definition]). By the equalities (5) and (13), we have

$$T_t(f) = \exp(it(T - \lambda I))(f), \quad \forall f \in A(X).$$

It follows that

$$\lim_{t \rightarrow 0} \frac{T_t(f) - f}{t} = i(T - \lambda I)(f), \quad \forall f \in A(X)$$

[5, Chapter I, 2.11 Proposition]; hence $D(A) = A(X)$ and

$$A(f) = i(T - \lambda I)(f), \quad \forall f \in A(X).$$

Since $\{T_t : t \in \mathbb{R}\}$ is a group of algebra homomorphisms on $A(X)$, then A is a derivation on $A(X)$; that is, $A(fg) = A(f)g + fA(g)$ for all $f, g \in A(X)$. Indeed,

$$\begin{aligned} A(fg) &= \lim_{t \rightarrow 0} \frac{T_t(fg) - fg}{t} \\ &= \lim_{t \rightarrow 0} \frac{T_t(f)T_t(g) - fg}{t} \\ &= \lim_{t \rightarrow 0} \frac{(T_t(f) - f)T_t(g) + f(T_t(g) - g)}{t} \\ &= \lim_{t \rightarrow 0} \left(\frac{T_t(f) - f}{t} T_t(g) + f \frac{T_t(g) - g}{t} \right) \\ &= A(f)T_0(g) + fA(g) \\ &= A(f)g + fA(g). \end{aligned}$$

Since all linear derivations on a complex commutative semi-simple Banach algebra are trivial [16, 18], it follows that A is zero, and so $T = \lambda I$, as desired. The converse implication follows immediately, and so the proof of the theorem is finished. $\square$

ACKNOWLEDGEMENT

The authors wish to thank the referee for making several suggestions which improved this paper.

REFERENCES

- [1] E. Berkson and H. Porta, *Hermitian operators and one-parameter groups of isometries in Hardy spaces*, Trans. Amer. Math. Soc. **185** (1973), 331–344. MR0338833
- [2] Earl Berkson and Ahmed Sourour, *The Hermitian operators on some Banach spaces*, Studia Math. **52** (1974), 33–41. MR0355668 (50 #8142)
- [3] Frank F. Bonsall and John Duncan, *Complete normed algebras*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 80, Springer-Verlag, New York, 1973. MR0423029 (54 #11013)
- [4] K. de Leeuw, *Banach spaces of Lipschitz functions*, Studia Math. **21** (1961/1962), 55–66. MR0140927 (25 #4341)
- [5] Klaus-Jochen Engel and Rainer Nagel, *A short course on operator semigroups*, Universitext, Springer, New York, 2006. MR2229872 (2007e:47001)
- [6] Richard J. Fleming and James E. Jamison, *Isometries on Banach spaces: function spaces*, Chapman & Hall/CRC Monographs and Surveys in Pure and Applied Mathematics, vol. 129, Chapman & Hall/CRC, Boca Raton, FL, 2003. MR1957004 (2004j:46030)
- [7] Irving Glicksberg, *Stone-Čech compactifications of products*, Trans. Amer. Math. Soc. **90** (1959), 369–382. MR0105667 (21 #4405)
- [8] Kenneth Hoffman, *Banach spaces of analytic functions*, Prentice-Hall Series in Modern Analysis, Prentice-Hall Inc., Englewood Cliffs, N.J., 1962. MR0133008 (24 #A2844)
- [9] William Hornor and James E. Jamison, *Isometries of some Banach spaces of analytic functions*, Integral Equations Operator Theory **41** (2001), no. 4, 410–425, DOI 10.1007/BF01202102. MR1857800 (2002h:46035)
- [10] James E. Jamison, *Bicircular projections on some Banach spaces*, Linear Algebra Appl. **420** (2007), no. 1, 29–33, DOI 10.1016/j.laa.2006.05.009. MR2277626 (2007m:47038)
- [11] Krzysztof Jarosz, *Isometries in semisimple, commutative Banach algebras*, Proc. Amer. Math. Soc. **94** (1985), no. 1, 65–71, DOI 10.2307/2044953. MR781058 (86d:46044)
- [12] E. Mayer-Wolf, *Isometries between Banach spaces of Lipschitz functions*, Israel J. Math. **38** (1981), no. 1-2, 58–74, DOI 10.1007/BF02761849. MR599476 (82e:46036)
- [13] N. V. Rao and A. K. Roy, *Linear isometries of some function spaces*, Pacific J. Math. **38** (1971), 177–192. MR0308763 (46 #7877)
- [14] Donald R. Sherbert, *Banach algebras of Lipschitz functions*, Pacific J. Math. **13** (1963), 1387–1399. MR0156214 (27 #6143)

- [15] Donald R. Sherbert, *The structure of ideals and point derivations in Banach algebras of Lipschitz functions*, Trans. Amer. Math. Soc. **111** (1964), 240–272. MR0161177 (28 #4385)
- [16] I. M. Singer and J. Wermer, *Derivations on commutative normed algebras*, Math. Ann. **129** (1955), 260–264. MR0070061 (16,1125c)
- [17] L. L. Stachó and B. Zalar, *Bicircular projections on some matrix and operator spaces*, Linear Algebra Appl. **384** (2004), 9–20, DOI 10.1016/j.laa.2003.11.014. MR2055340 (2005a:46146)
- [18] Marc P. Thomas, *The image of a derivation is contained in the radical*, Ann. of Math. (2) **128** (1988), no. 3, 435–460, DOI 10.2307/1971432. MR970607 (90d:46075)

DEPARTMENT OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF MEMPHIS, MEMPHIS, TENNESSEE 38152

E-mail address: mbotelho@memphis.edu

DEPARTMENT OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF MEMPHIS, MEMPHIS, TENNESSEE 38152

E-mail address: jjamison@memphis.edu

DEPARTAMENTO DE ÁLGEBRA Y ANÁLISIS MATEMÁTICO, UNIVERSIDAD DE ALMERÍA, 04120 ALMERÍA, SPAIN

E-mail address: ajimenez@ual.es

CAMPUS UNIVERSITARIO DE PUERTO REAL, FACULTAD DE CIENCIAS, NUESTRA UNIVERSIDAD, 11510 PUERTO REAL, CÁDIZ, SPAIN

E-mail address: moises.villegas@uca.es