AN EXTENSION OF MARKOV'S THEOREM

A. BRANQUINHO, U. FIDALGO PRIETO, AND A. FOULQUIÉ MORENO

(Communicated by Walter Van Assche)

ABSTRACT. We give a general sufficient condition for the uniform convergence of sequences of type II Hermite-Padé approximants associated with Nikishin systems of functions.

1. INTRODUCTION

Let $\Delta \subset \mathbb{R}$ be a compact interval and $\mathcal{M}(\Delta)$ the set of finite Borel measures with constant sign whose support $S(\mu)$ is a subset of Δ such that Δ is the smallest interval which contains $S(\mu)$; we write $\text{Co}(S(\mu)) = \Delta$. Given $\mu \in \mathcal{M}(\Delta)$, the associated Markov function is defined by

$$\hat{\mu}(z) = \int \frac{d\mu(x)}{z - x} \in \mathcal{H}(\overline{\mathbb{C}} \setminus S(\mu)),$$

which is holomorphic in $\overline{\mathbb{C}} \setminus S(\mu)$.

Fix a measure $\sigma \in \mathcal{M}(\Delta)$ and a system of m weights $\mathbf{r} = (\rho_1, \dots, \rho_m)$ with respect to σ ; that is, each $\rho_k \in L_1(\sigma)$ and has constant sign. Consider the system of measures $\mathbf{s} = (s_1, \dots, s_m)$, where $ds_j = \rho_j d\sigma$, and the corresponding system of Markov functions $\hat{\mathbf{s}} = (\hat{s}_1, \dots, \hat{s}_m)$. Take a multi-index $\mathbf{n} = (n_1, \dots, n_m) \in \mathbb{Z}_+^m$, where $\mathbb{Z}_+ = \{0, 1, 2, \dots\}$. There exist polynomials $Q_{\mathbf{n}}$ and $P_{\mathbf{n},j}$, $j = 1, \dots, m$, such that

$$(1.1) \quad \begin{aligned} i) \quad & \deg Q_{\mathbf{n}} \leq |\mathbf{n}| = n_1 + \dots + n_m, \quad Q_{\mathbf{n}} \not\equiv 0, \\ ii) \quad & (Q_{\mathbf{n}} \hat{s}_j - P_{\mathbf{n},j})(z) = \mathcal{O}(1/z^{n_j+1}), \quad z \rightarrow \infty, \quad j = 1, \dots, m. \end{aligned}$$

In the sequel we assume that $Q_{\mathbf{n}}$ is monic.

For each $j = 1, \dots, m$, $Q_{\mathbf{n}}$ annihilates the terms corresponding to the powers between -1 and $-n_j$ of the Laurent expansion of $Q_{\mathbf{n}} \hat{s}_j$, whereas $P_{\mathbf{n},j}$ represents the polynomial part of $Q_{\mathbf{n}} \hat{s}_j$. Hence, $Q_{\mathbf{n}}$ determines univocally $P_{\mathbf{n},j}$ and, consequently, the rational fraction $P_{\mathbf{n},j}/Q_{\mathbf{n}}$.

Received by the editors October 16, 2012 and, in revised form, October 17, 2012.

2010 *Mathematics Subject Classification.* Primary 30E10.

This work was supported by the Center for Mathematics of the University of Coimbra, Center for Research and Development in Mathematics and Applications (University of Aveiro), Acção Integrada Luso-Espanhola n. E-4/10, and Fundação para a Ciência e Tecnologia, through European program COMPETE/FEDER and grant SFRH/BPD/62947/2009. The second author also acknowledges support from Ministerio de Economía y Competitividad, under grant MTM2009-12740-C03-01.

The authors are indebted to Professor G. López Lagomasino, who carefully read the manuscript and whose suggestions improved it.

The vector rational fractions $\mathbf{R}_n = (P_{n,1}/Q_n, \dots, P_{n,m}/Q_n)$ are called the type II Hermite-Padé approximant corresponding to the system $\widehat{\mathbf{s}}$ and the multi-index $\mathbf{n}$.

When $m = 1$, $\mathbf{R}_n = P_{n,1}/Q_n = P_n/Q_n$, $\mathbf{n} = n$, is the n th diagonal Padé approximant of $\widehat{s}_1 = \widehat{s}$. It is well known (for example, see Chapter II in [15]) that in this case Q_n is the n th monic orthogonal polynomial with respect to the measure s . Usually, monic orthogonal polynomials are defined for positive measures, however, the definition is trivially extended to measures with constant sign. Q_n has n simple zeros in the interior of $\text{Co}(S(s))$ (see [16, Lemma 1.1.3]).

In [13], A.A. Markov proved that given an arbitrary measure $s \in \mathcal{M}(\Delta)$ the sequence $\{P_n/Q_n\}_{n \in \mathbb{Z}_+}$ converges uniformly to $\widehat{s}$ on every compact subset contained in the domain $\overline{\mathbb{C}} \setminus \Delta$. We write

$$\frac{P_n}{Q_n} \underset{n \rightarrow \infty}{\rightrightarrows} \widehat{s} \quad \text{on} \quad \overline{\mathbb{C}} \setminus \Delta.$$

In the present paper, we extend Markov's Theorem to the context of type II Hermite-Padé approximation.

The first drawback in extending Markov's Theorem to the context of Hermite-Padé approximation is that in the vector case, in general, Q_n is not uniquely determined by (1.1). However, in [10] it is shown that uniqueness takes place for the so-called Nikishin systems of measures which we introduce below. In this case, Q_n also has $|\mathbf{n}|$ simple zeros in the interior of Δ .

Nikishin systems of measures were introduced by E.M. Nikishin in his famous article [14]. Take two compact intervals Δ_α and Δ_β of the real line such that $\Delta_\alpha \cap \Delta_\beta = \emptyset$ and two measures $\sigma_\alpha \in \mathcal{M}(\Delta_\alpha)$ and $\sigma_\beta \in \mathcal{M}(\Delta_\beta)$. We define a third measure $\langle \sigma_\alpha, \sigma_\beta \rangle$ whose differential expression is

$$d\langle \sigma_\alpha, \sigma_\beta \rangle(x) = \int \frac{d\sigma_\beta(t)}{x-t} d\sigma_\alpha(x) = \widehat{\sigma}_\beta(x) d\sigma_\alpha(x).$$

Observe that $\langle \sigma_\alpha, \sigma_\beta \rangle \in \mathcal{M}(\Delta_\alpha)$.

Now, take m compact intervals $\Delta_1, \dots, \Delta_m$ with the property that for each $j = 1, \dots, m-1$, $\Delta_j \cap \Delta_{j+1} = \emptyset$. Let $(\sigma_1, \dots, \sigma_m)$ be a system of measures such that $\sigma_j \in \mathcal{M}(\Delta_j)$, $j = 1, \dots, m$. The system of measures $(s_1, \dots, s_m)$ given by

$$s_1 = \sigma_1, \quad s_2 = \langle \sigma_1, \sigma_2 \rangle, \quad s_3 = \langle \sigma_1, \langle \sigma_2, \sigma_3 \rangle \rangle = \langle \sigma_1, \sigma_2, \sigma_3 \rangle, \dots, s_m = \langle \sigma_1, \dots, \sigma_m \rangle,$$

is the so-called Nikishin system of measures generated by $(\sigma_1, \dots, \sigma_m)$. For short, we write $(s_1, \dots, s_m) = \mathcal{N}(\sigma_1, \dots, \sigma_m)$, whereas $\widehat{\mathbf{s}} = (\widehat{s}_1, \dots, \widehat{s}_m) = \widehat{\mathcal{N}}(\sigma_1, \dots, \sigma_m)$ is the corresponding Nikishin system of functions. Nikishin systems have received a great deal of attention in the recent past and have found numerous applications; see for example [1], [2], [3], [4], [6], [7] [8], [11], [12] and [17].

In order to state our main result we need to review some concepts. Given two disjoint compact sets K_1 and K_2 of $\mathbb{R}$, $\text{dist}(K_1, K_2)$ denotes the distance between K_1 and K_2 , i.e. $\text{dist}(K_1, K_2) = \min\{|x_1 - x_2| : (x_1, x_2) \in K_1 \times K_2\}$, whereas $\text{diam}(K_1) = \max\{|x_1 - x_2| : x_1, x_2 \in K_1\}$ denotes the diameter of K_1 .

The main result of this paper is the following theorem.

Theorem 1.1. *Let $\{\mathbf{R}_n = (P_{n,1}/Q_n, \dots, P_{n,m}/Q_n)\}_{\mathbf{n} \in \Lambda}$ be the sequence of type II Hermite-Padé approximants corresponding to a sequence of distinct multi-indices $\Lambda \subset \mathbb{Z}_+^m$ and a system $(\widehat{s}_1, \dots, \widehat{s}_m) = \widehat{\mathcal{N}}(\sigma_1, \dots, \sigma_m)$. Assume*

$$\text{diam}(\Delta_k) < \text{dist}(\Delta_1, \Delta_2).$$

Then, for each compact set $K \subset \overline{\mathbb{C}} \setminus \Delta_1$

$$\limsup_{\mathbf{n} \in \Lambda} \left\| \widehat{s}_j - \frac{P_{\mathbf{n},j}}{Q_{\mathbf{n}}} \right\|_K^{1/(|\mathbf{n}|+n_j)} \leq \|\phi_\infty\|_K < 1, \quad j = 1, \dots, m,$$

where $\|\cdot\|_K$ denotes the sup-norm on K and ϕ_∞ denotes the conformal representation of $\overline{\mathbb{C}} \setminus \Delta_1$ onto the open unit disk such that $\phi_\infty(\infty) = 0$ and $\phi'_\infty(\infty) > 0$.

Notice that the sequence of multi-indices may be completely arbitrary. In Markov's Theorem, there is no assumption on the measure. This is also true in our case whenever $\text{diam}(\Delta_k) < \text{dist}(\Delta_1, \Delta_2)$, $k = 1, 2$. We have imposed no restrictions on the measures $\sigma_3, \dots, \sigma_m$ at all. Another extension of Markov's Theorem was given in [10, Corollary 1.1] without any assumption on the measures, but the indices are required to satisfy $n_j \geq |\mathbf{n}|/m - c|\mathbf{n}|^\kappa$, $j = 1, \dots, m$, for $c > 0$ and $\kappa < 1$. We believe that a complete analogue of Markov's Theorem should hold.

The following result extends [9, Corollary 2] to a larger class of multi-indices.

Theorem 1.2. *Let $\Lambda \subset \mathbb{Z}_+^m$ be a sequence of multi-indices such that either there exists $k \in \{2, \dots, m\}$ such that for every $\mathbf{n} = (n_1, \dots, n_m) \in \Lambda$, $n_k = \max\{n_1 + 1, n_2, \dots, n_m\}$, or $n_1 = \max\{n_1, n_2 - 1, \dots, n_m - 1\}$ (in which case we take $k = 1$). Then, for each compact set $K \subset \overline{\mathbb{C}} \setminus \Delta_1$,*

$$(1.2) \quad \limsup_{\mathbf{n} \in \Lambda} \left\| \widehat{s}_k - \frac{P_{\mathbf{n},k}}{Q_{\mathbf{n}}} \right\|_K^{1/2|\mathbf{n}|} \leq \kappa(K) < 1,$$

where

$$\kappa(K) = \sup\{\|\phi_t\|_K : t \in \Delta_2 \cup \{\infty\}\}$$

and ϕ_t denotes the conformal representation of $\overline{\mathbb{C}} \setminus \Delta_1$ onto the open unit disk such that $\phi_t(t) = 0$ and $\phi'_t(t) > 0$.

In the first three sections we give some preliminary results which are necessary for the proof of the theorems above. Section 2 includes some properties of multiple orthogonal polynomials corresponding to Nikishin systems of measures. In Section 3 we study properties of Fourier series of functions expanded in terms of orthogonal polynomials with respect to varying measures. Theorem 1.2 is proved in Section 4 as a first step to the proof of Theorem 1.1 which is completed in Section 5.

2. MULTIPLE ORTHOGONALITY IN NIKISHIN SYSTEMS

Let $\mathbf{s} = (s_1, \dots, s_m) = \mathcal{N}(\sigma_1, \dots, \sigma_m)$ and $\mathbf{n} = (n_1, \dots, n_m)$ be given. It is well known and easy to verify that the conditions (1.1) imply

$$(2.1) \quad 0 = \int x^\nu Q_{\mathbf{n}}(x) ds_j(x), \quad \nu = 0, \dots, n_j - 1, \quad j = 1, \dots, m.$$

For each $j = 1, \dots, m$, let h be an arbitrary polynomial such that $\deg h \leq n_j$. Then

$$(2.2) \quad 0 = \int \frac{h(z) - h(x)}{z - x} Q_{\mathbf{n}}(x) ds_j(x),$$

hence

$$\int \frac{Q_{\mathbf{n}}(x)}{z - x} ds_j(x) = \frac{1}{h(z)} \int \frac{h(x) Q_{\mathbf{n}}(x)}{z - x} ds_j(x) = \mathcal{O}\left(\frac{1}{z^{n_j+1}}\right) \quad \text{as } z \rightarrow \infty.$$

Define

$$P(z) = \int \frac{Q_{\mathbf{n}}(z) - Q_{\mathbf{n}}(x)}{z - x} ds_j(x).$$

Thus

$$(Q_{\mathbf{n}}\widehat{s}_j - P)(z) = \int \frac{Q_{\mathbf{n}}(x)}{z - x} ds_j(x) = \mathcal{O}\left(\frac{1}{z^{n_j+1}}\right) \quad \text{as } z \rightarrow \infty.$$

From (1.1) we see that

$$P(z) - P_{\mathbf{n},j}(z) = \mathcal{O}\left(\frac{1}{z^{n_j+1}}\right) \in \mathcal{H}(\overline{\mathbb{C}}), \quad z \rightarrow \infty.$$

Consequently,

$$(2.3) \quad P_{\mathbf{n},j}(z) \equiv \int \frac{Q_{\mathbf{n}}(z) - Q_{\mathbf{n}}(x)}{z - x} ds_j(x), \quad (Q_{\mathbf{n}}\widehat{s}_j - P_{\mathbf{n},j})(z) = \int \frac{Q_{\mathbf{n}}(x)}{z - x} ds_j(x).$$

From [10] we know that the conditions in (2.1) imply that $Q_{\mathbf{n}}$ has $|\mathbf{n}|$ simple zeros which lie in the interior of Δ_1 . Let $x_{\mathbf{n},1} < \dots < x_{\mathbf{n},|\mathbf{n}|}$ be the zeros of $Q_{\mathbf{n}}$. Decomposing into simple fractions, we get

$$(2.4) \quad \frac{P_{\mathbf{n},j}(z)}{Q_{\mathbf{n}}(z)} = \sum_{i=1}^{|\mathbf{n}|} \frac{\lambda_{i,j,\mathbf{n}}}{z - x_{\mathbf{n},i}}, \quad j = 1, \dots, m.$$

The coefficients $\lambda_{i,j,\mathbf{n}}$, $i = 1, \dots, |\mathbf{n}|$ and $j = 1, \dots, m$, were called Nikishin-Christoffel coefficients in [9, Definition 2]. Taking into account the equality in (2.3), we have that

$$(2.5) \quad \lambda_{i,j,\mathbf{n}} = \lim_{z \rightarrow x_{\mathbf{n},i}} (z - x_{\mathbf{n},i}) \frac{P_{\mathbf{n},j}(z)}{Q_{\mathbf{n}}(z)} = \int \frac{Q_{\mathbf{n}}(x) ds_j(x)}{Q'_{\mathbf{n}}(x_{\mathbf{n},i})(x - x_{\mathbf{n},i})}.$$

For each $j = 1, \dots, m$,

$$(2.6) \quad \left| \sum_{i=1}^{|\mathbf{n}|} \lambda_{i,j,\mathbf{n}} \right| = \left| \sum_{i=1}^{|\mathbf{n}|} \int \frac{Q_{\mathbf{n}}(x) ds_j(x)}{Q'_{\mathbf{n}}(x_{\mathbf{n},i})(x - x_{\mathbf{n},i})} \right|$$

$$= \left| \int \sum_{i=1}^{|\mathbf{n}|} \frac{Q_{\mathbf{n}}(x)}{Q'_{\mathbf{n}}(x_{\mathbf{n},i})(x - x_{\mathbf{n},i})} ds_j(x) \right| = \left| \int ds_j(x) \right| = \|s_j\| < +\infty,$$

where $\|s\|$ represents the total variation of the measure s . In this chain of equalities we have used the fact that $\mathcal{P}(x) = \sum_{i=1}^{|\mathbf{n}|} Q_{\mathbf{n}}(x) / (Q'_{\mathbf{n}}(x_{\mathbf{n},i})(x_{\mathbf{n},i} - x))$ is the polynomial of degree $\leq |\mathbf{n}| - 1$ which interpolates the constant function 1 at the zeros of $Q_{\mathbf{n}}$. Thus $\mathcal{P} \equiv 1$.

From [10, Lemma 3.2] one can state the following result. (We wish to point out that the measure denoted here with τ are products of those in [10].)

Lemma 2.1. *Let $(\widehat{s}_{2,2}, \dots, \widehat{s}_{2,m}) = \widehat{\mathcal{N}}(\sigma_2, \dots, \sigma_m)$. There is a system of $m - 1$ measures $(\tau_{2,1}^k, \dots, \tau_{2,k-1}^k, \tau_{2,k+1}^k, \dots, \tau_{2,m}^k)$ where $\text{Co}(S(\tau_{2,j}^k)) \subset \Delta_2$, $j = 1, \dots, k - 1, k + 1, \dots, m$, such that*

$$(2.7) \quad \frac{1}{\widehat{s}_{2,k}(z)} = \ell_{2,k}(z) + \widehat{\tau}_{2,1}^k(z),$$

where $\ell_{2,k}$ denotes a polynomial with degree one, and

$$(2.8) \quad \frac{\widehat{s}_{2,j}(z)}{\widehat{s}_{2,k}(z)} - \frac{|s_{2,j}|}{|s_{2,k}|} = \widehat{\tau}_{2,j}^k(z), \quad j = 2, \dots, k-1, k+1, \dots, m.$$

Theorem 1.4 in [10] refers to so-called mixed type multiple orthogonal polynomials of two Nikishin systems. When reduced to type II multiple orthogonal polynomials of a Nikishin system it may be restated in the following form.

Lemma 2.2. *Let $(s_1, \dots, s_m) = \mathcal{N}(\sigma_1, \dots, \sigma_m)$ and $\mathbf{n} = (n_1, \dots, n_m) \in \mathbb{Z}_+^m$ be given. Set $k = 1$ if $n_1 + 1 = M = \max\{n_1 + 1, n_2 \dots n_m\}$; otherwise k is equal to the subscript of the first component of $\mathbf{n}$ such that $M = n_k$. Then, there exists a permutation λ of $\{1, \dots, m\}$ which reorders the components of $\mathbf{n}$ such that $n_{\lambda(1)} + \delta_{\lambda(1),1} \geq n_{\lambda(2)} \geq \dots \geq n_{\lambda(m)}$ with $n_k = n_{\lambda(1)}$ and $\delta_{\lambda(1),1}$ denoting the known Kronecker delta function, and an associated Nikishin system $\widetilde{\mathbf{s}} = (r_1, \dots, r_m) = \mathcal{N}(\rho_1, \dots, \rho_m)$, where $s_k = r_1 = \rho_1$ and $\text{Co}(S(\rho_j)) \subset \Delta_j$, $j = 1, \dots, m$, such that if $\widetilde{\mathbf{n}} = (n_{\lambda(1)}, \dots, n_{\lambda(m)})$, the pairs $(\mathbf{s}, \mathbf{n})$ and $(\widetilde{\mathbf{s}}, \widetilde{\mathbf{n}})$ have the same type II multiple orthogonal polynomial. That is, $Q_{\mathbf{n}}$ satisfies (2.1) and*

$$(2.9) \quad 0 = \int x^\nu Q_{\mathbf{n}}(x) \widehat{r}_{2,j}(x) ds_k(x), \quad \nu = 0, \dots, n_{\lambda(j)} - 1, \quad j = 1, \dots, m,$$

where $r_{2,j} = \langle \rho_2, \dots, \rho_j \rangle$, $j = 2, \dots, m$, and $\widehat{r}_{2,1} \equiv 1$.

Type II multiple orthogonal polynomials of Nikishin systems with respect to decreasing multi-indices satisfy other orthogonality relations. In particular, from Propositions 2 and 3 in [11] (see also relations (5)-(7) in [2]), we have

Lemma 2.3. *Let $\mathbf{s} = (s_1, \dots, s_m) = \mathcal{N}(\sigma_1, \dots, \sigma_m)$ and $\mathbf{n} = (n_1, \dots, n_m)$ be given. Let $k \in \{1, \dots, m\}$ be as in Lemma 2.2. Then, there exist two monic polynomials $Q_{\mathbf{n},2}$, $\deg Q_{\mathbf{n},2} = |\mathbf{n}| - n_k$, and $Q_{\mathbf{n},3} = |\mathbf{n}| - n_k - n_{\lambda(2)}$, whose zeros are simple and lie in the interior of Δ_2 and Δ_3 , respectively, such that*

$$(2.10) \quad \left(\frac{Q_{\mathbf{n}} \widehat{s}_k - P_{\mathbf{n},k}}{Q_{\mathbf{n},2}} \right) (z) = \mathcal{O} \left(\frac{1}{z^{|\mathbf{n}|+1}} \right) \in \mathcal{H}(\overline{\mathbb{C}} \setminus S(\sigma_1)),$$

$$(2.11) \quad 0 = \int x^\nu Q_{\mathbf{n}}(x) \frac{ds_k(x)}{Q_{\mathbf{n},2}(x)}, \quad \nu = 0, \dots, |\mathbf{n}| - 1,$$

and

$$(2.12) \quad 0 = \int t^\nu Q_{\mathbf{n},2}(t) \int \frac{Q_{\mathbf{n}}^2(x)}{t-x} \frac{ds_k(x)}{Q_{\mathbf{n},2}(x)} \frac{d\rho_2(t)}{Q_{\mathbf{n}}(t)Q_{\mathbf{n},3}(t)}, \quad \nu = 0, \dots, |\mathbf{n}| - n_k - 1.$$

(Here, ρ_2 is the measure coming from Lemma 2.2.)

Formulas (2.11) and (2.12) state that $Q_{\mathbf{n}}$ and $Q_{\mathbf{n},2}$ are the $|\mathbf{n}|$ th and $(|\mathbf{n}| - n_k)$ th monic orthogonal polynomials with respect to the varying measures

$$(2.13) \quad \frac{ds_k}{Q_{\mathbf{n},2}} \quad \text{and} \quad \int \frac{Q_{\mathbf{n}}^2(x)}{t-x} \frac{ds_k(x)}{Q_{\mathbf{n},2}(x)} \frac{d\rho_2(t)}{Q_{\mathbf{n}}(t)Q_{\mathbf{n},3}(t)}, \quad \text{respectively.}$$

There are other full orthogonality relations with respect to varying measures satisfied deeper in the system, but we will not need them.

3. VARYING MEASURES AND ASSOCIATED FOURIER SERIES

Let $\text{sign} : \mathbb{R} \setminus \{0\} \rightarrow \{-1, 1\}$ denote the sign function. Analogously, $\text{sign}(\mu)$ will denote the sign of a given measure $\mu \in \mathcal{M}(\Delta)$. Notice that $\text{sign}(\mu) \cdot \mu$ is a positive measure. Given a measurable function $f : \Delta \rightarrow \mathbb{R}$,

$$\|f\|_{2,\mu} = \sqrt{\text{sign}(\mu) \int f^2(x) d\mu(x)}$$

denotes the L_2 norm with respect to μ . If $\|f\|_{2,\mu} < +\infty$ we write $f \in L_2(\mu)$.

Let $\{q_{\mu,n}\}_{n \in \mathbb{Z}_+}$ be the family of monic orthogonal polynomials with respect to μ . For each $n \in \mathbb{Z}_+$ let $p_{\mu,n}(z) \equiv q_{\mu,n}/\|q_{\mu,n}\|_{2,\mu}$ denote the n th orthonormal polynomial with respect to the measure μ . That is,

$$\int p_{\mu,n}(x) p_{\mu,k}(x) d\mu(x) = \delta_{n,k} = \begin{cases} 1 & \text{if } n = k \\ 0 & \text{if } n \neq k \end{cases}, \quad (n, k) \in \mathbb{Z}_+^2.$$

Fix $n \in \mathbb{Z}_+$. For each polynomial h of degree $\leq n$ we have the identity

$$0 = \int \frac{h(z) - h(x)}{z - x} p_{\mu,n}(x) d\mu(x);$$

thus

$$(3.1) \quad \int \frac{p_{\mu,n}(x) d\mu(x)}{z - x} = \frac{1}{p_{\mu,n}(z)} \int \frac{p_{\mu,n}^2(x) d\mu(x)}{z - x}.$$

From (2.11) we see that $q_{\mu,|\mathbf{n}|} \equiv Q_{\mathbf{n}}$ when $d\mu = ds_k/Q_{\mathbf{n},2}$ and $p_{\mu,|\mathbf{n}|} \equiv Q_{\mathbf{n}}/\|Q_{\mathbf{n}}\|_{2,\mu}$.

Lemma 3.1. *Let $\{d\mu_n\}_{n \in \mathbb{Z}_+} \subset \mathcal{M}(\Delta)$ be given. Then for each $t \in \mathbb{C} \setminus \Delta$ we have that*

$$(3.2) \quad \left| \frac{q_{\mu_n,n}(x)}{q_{\mu_n,n}(t)} \right|^{1/n} \leq \frac{\text{diam}(\Delta)}{\text{dist}(t, \Delta)}, \quad n \in \mathbb{Z}_+,$$

uniformly in $\{x \in \Delta\}$.

Proof. Fix $n \in \mathbb{Z}_+$. Since $q_{\mu_n,n}$ has its n zeros in the interior of Δ , then

$$\left| \frac{q_{\mu_n,n}(x)}{q_{\mu_n,n}(z)} \right| \leq \left(\frac{\text{diam}(\Delta)}{\text{dist}(K, \Delta)} \right)^n.$$

This proves immediately (3.2). □

Fix two integers $n, \nu \in \mathbb{Z}_+$ and a function $f \in L_2(\mu_\nu)$. The sum

$$(3.3) \quad S_{f,n,\mu_\nu}(z) = \sum_{i=0}^n \gamma_{i,\nu} p_{\mu_\nu,i}(z),$$

where

$$\gamma_{i,\nu} = \text{sign}(\mu_\nu) \int f(x) p_{\mu_\nu,i}(x) d\mu_\nu(x), \quad i = 0, \dots, n,$$

defines the n th partial sum of the Fourier series corresponding to f in terms of the orthonormal system $\{p_{\mu_n,i}\}_{i \in \mathbb{Z}_+}$.

Substituting in (3.3) the well known Christoffel-Darboux identity (Theorem 4.5 in [5], page 23), we obtain

$$(3.4) \quad S_{f,n,\mu_\nu}(z) = a_{\mu_\nu,n+1} \int \frac{p_{\mu_\nu,n+1}(z)p_{\mu_\nu,n}(x) - p_{\mu_\nu,n+1}(x)p_{\mu_\nu,n}(z)}{z-x} f(x) d\mu_\nu(x),$$

where

$$a_{\mu_\nu,n+1} = \int x p_{\mu_\nu,n+1}(x) p_{\mu_\nu,n}(x) d\mu_\nu(x).$$

Notice that $\text{sign}(a_{\mu_n,n+1}) = \text{sign}(\mu_n)$. For an arbitrary polynomial $\mathcal{P}$ of degree $\leq n$, $S_{\mathcal{P},n,\mu_n} \equiv \mathcal{P}$.

Proposition 3.1. *Let $\{\mu_n\}_{n \in \mathbb{Z}_+} \subset \mathcal{M}(\Delta)$ be given. Fix $t \in \mathbb{C} \setminus \Delta$ such that $\text{dist}(t, \Delta) > \text{diam}(\Delta)$. Then*

$$(3.5) \quad S_{1/(z-t),n,\mu_n} \rightrightarrows \frac{1}{z-t}, \quad \text{for } z \in \Delta.$$

Proof. Fix $N \in \mathbb{Z}_+$. We start by proving

$$(3.6) \quad S_{1/(z-t),n,\mu_N} \rightrightarrows \frac{1}{z-t}, \quad \text{for } z \in \Delta.$$

For two nonnegative integers $n > n'$ we analyze the difference,

$$(3.7) \quad \varepsilon_{N,n,n'} = |S_{1/(z-t),n',\mu_N} - S_{1/(z-t),n,\mu_N}| = \left| \sum_{i=n'+1}^n \gamma_{i,N} p_{\mu_N,i}(z) \right|,$$

where $\gamma_{i,N} = \int p_{\mu_N,i}(x)/(x-t) d\mu_N(x)$. So

$$\varepsilon_{N,n,n'} = \left| \sum_{i=n'+1}^n p_{\mu_N,i}(z) \int \frac{p_{\mu_N,i}(x) d\mu_N(x)}{\rho_N(x) (x-t)} \right|.$$

Taking into account the identity given in (3.1) we have that

$$\begin{aligned} \varepsilon_{N,n,n'} &= \left| \sum_{i=n'+1}^n \frac{p_{\mu_N,i}(z)}{p_{\mu_N,i}(t)} \int \frac{p_{\mu_N,i}^2(x) d\mu_N(x)}{x-t} \right| \\ &\leq \sum_{i=n'+1}^n \left| \frac{p_{\mu_N,i}(z)}{p_{\mu_N,i}(t)} \right| \left| \int \frac{p_{\mu_N,i}^2(x) d\mu_N(x)}{x-t} \right| \leq \sum_{i=n'+1}^n \left| \frac{p_{\mu_N,i}(z)}{p_{\mu_N,i}(t)} \right| \frac{\left| \int p_{\mu_N,i}^2(x) d\mu_N(x) \right|}{\text{dist}(t, \Delta)}. \end{aligned}$$

Hence we obtain that

$$\varepsilon_{N,n,n'} \leq \frac{1}{\text{dist}(t, \Delta)} \sum_{i=n'+1}^n \left| \frac{p_{\mu_N,i}(z)}{p_{\mu_N,i}(t)} \right|.$$

Lemma 3.1 implies that there exists a nonnegative integer N' such that for every pair (n, n') , with $n \geq n' \geq N'$,

$$\varepsilon_{N,n,n'} \leq \varepsilon_{N,n,n'} \leq \frac{1}{\text{dist}(t, \Delta)} \sum_{i=n'+1}^n M^i \rightarrow 0 \quad \text{as } n, n' \rightarrow \infty,$$

where $M = \text{diam}(\Delta)/\text{dist}(\Delta, t) < 1$. This proves (3.6).

So, for each $n \in \mathbb{Z}_+$ fixed we can write

$$\frac{1}{z-x} = \sum_{i=0}^{\infty} p_{\mu_n,i}(z) \int \frac{p_{\mu_n,i}(x)}{x-t} d\mu_n(x) = \sum_{i=0}^{\infty} \frac{p_{\mu_n,i}(z)}{p_{\mu_n,i}(t)} \int \frac{p_{\mu_n,i}^2(x) d\mu_n(x)}{x-t}.$$

Then

$$\varepsilon_{n,n,\infty} = \left| S_{1/(z-t),n,\mu_n} - \frac{1}{z-t} \right| = \left| \sum_{i=n+1}^{\infty} \frac{p_{\mu_n,i}(z)}{p_{\mu_n,i}(t)} \int \frac{p_{\mu_n,i}^2(x) d\mu_n(x)}{x-t} \right|.$$

Taking again into account Lemma 3.1 we see that there exists a nonnegative integer N' such that for all $n \geq N'$,

$$\varepsilon_{n,n,\infty} \leq \frac{1}{\text{dist}(t, \Delta)} \sum_{i=n}^{\infty} M^i \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

This proves (3.5) and completes the proof of Proposition 3.1. $\square$

Recall the definition of Nikishin-Christoffel coefficients introduced in Section 2.

Proposition 3.2. *Let $\mathbf{n} = (n_1, \dots, n_m) \in \mathbb{Z}_+^m$ and $(s_1, \dots, s_m) = \mathcal{N}(\sigma_1, \dots, \sigma_m)$ be given. Set $k = 1$ if $n_1 + 1 = M = \max\{n_1 + 1, n_2, \dots, n_m\}$; otherwise k is equal to the subscript of the first component of $\mathbf{n}$ such that $M = n_k$. For each $n \in \mathbb{Z}_+$, denote $d\mu_{\mathbf{n}} = ds_k/Q_{\mathbf{n},2}$. Then, for each $j = 1, \dots, m$, the Nikishin-Christoffel coefficients can be written as follows:*

$$(3.8) \quad \lambda_{i,j,\mathbf{n}} = \frac{\|Q_{\mathbf{n}}\|_{2,\mu_{\mathbf{n}}} S_{Q_{\mathbf{n},2}\widehat{s}_{2,j}/\widehat{s}_{2,k},|\mathbf{n}|-1,\mu_{\mathbf{n}}}(x_{\mathbf{n},i})}{a_{\mu_{\mathbf{n}},|\mathbf{n}|} Q'_{\mathbf{n}}(x_{\mathbf{n},i}) p_{\mu_{\mathbf{n}},|\mathbf{n}|-1}(x_{\mathbf{n},i})}, \quad i = 1, \dots, |\mathbf{n}|.$$

When $j = k$, the Nikishin-Christoffel coefficients acquire the following form:

$$(3.9) \quad \lambda_{i,k,\mathbf{n}} = \frac{\|Q_{\mathbf{n}}\|_{2,\mu_{\mathbf{n}}}(x_{\mathbf{n},i})}{a_{\mu_{\mathbf{n}},|\mathbf{n}|} Q'_{\mathbf{n}}(x_{\mathbf{n},i}) p_{\mu_{\mathbf{n}},|\mathbf{n}|-1}(x_{\mathbf{n},i})}, \quad i = 1, \dots, |\mathbf{n}|.$$

Thus

$$(3.10) \quad \text{sign}(\lambda_{i,k,\mathbf{n}}) = \text{sign}(s_k), \quad i = 1, \dots, |\mathbf{n}|.$$

In particular,

$$(3.11) \quad \sum_{i=1}^{|\mathbf{n}|} |\lambda_{i,k,\mathbf{n}}| = \|s_k\| < +\infty.$$

Proof. Let us rewrite (2.5) for each $j = 1, \dots, m$ and each $i = 1, \dots, |\mathbf{n}|$ as

$$\begin{aligned} \lambda_{i,j,\mathbf{n}} &= \int \frac{Q_{\mathbf{n}}(x) ds_j(x)}{Q'_{\mathbf{n}}(x_{\mathbf{n},i})(x - x_{\mathbf{n},i})} = \int \frac{Q_{\mathbf{n}}(x)}{Q'_{\mathbf{n}}(x_{\mathbf{n},i})(x - x_{\mathbf{n},i})} \frac{\widehat{s}_{2,j}(x)}{\widehat{s}_{2,k}(x)} Q_{\mathbf{n},2}(x) \frac{ds_k(x)}{Q_{\mathbf{n},2}(x)} \\ &= \frac{\|Q_{\mathbf{n}}\|_{2,\mu_{\mathbf{n}}}}{a_{\mu_{\mathbf{n}},|\mathbf{n}|} Q'_{\mathbf{n}}(x_{\mathbf{n},i}) p_{\mu_{\mathbf{n}},|\mathbf{n}|-1}(x_{\mathbf{n},i})} \\ &\quad \times a_{\mu_{\mathbf{n}},|\mathbf{n}|} \int \frac{p_{\mu_{\mathbf{n}},|\mathbf{n}|}(x) p_{\mu_{\mathbf{n}},|\mathbf{n}|-1}(x_{\mathbf{n},i})}{x - x_{\mathbf{n},i}} \frac{\widehat{s}_{2,j}(x)}{\widehat{s}_{2,k}(x)} Q_{\mathbf{n},2}(x) \frac{ds_k(x)}{Q_{\mathbf{n},2}(x)}. \end{aligned}$$

Using the formula given in (3.4) it follows that

$$\lambda_{i,j,\mathbf{n}} = \frac{\|Q_{\mathbf{n}}\|_{2,\mu_{\mathbf{n}}} S_{Q_{\mathbf{n},2}\widehat{s}_{2,j}/\widehat{s}_{2,k},|\mathbf{n}|-1,\mu_{\mathbf{n}}}(x_{\mathbf{n},i})}{a_{\mu_{\mathbf{n}},|\mathbf{n}|-1} Q'_{\mathbf{n}}(x_{\mathbf{n},i}) p_{\mu_{\mathbf{n}},|\mathbf{n}|-1}(x_{\mathbf{n},i})}.$$

When $j = k$, since $\widehat{s}_{2,j}/\widehat{s}_{2,k} \equiv 1$ and $\deg Q_{\mathbf{n},2} = |\mathbf{n}| - n_k$,

$$\lambda_{i,k,\mathbf{n}} = \frac{\|Q_{\mathbf{n}}\|_{2,\mu_{\mathbf{n}}} Q_{\mathbf{n},2}(x_{\mathbf{n},i})}{a_{\mu_{\mathbf{n}},|\mathbf{n}|} Q'_{\mathbf{n}}(x_{\mathbf{n},i}) p_{\mu_{\mathbf{n}},|\mathbf{n}|-1}(x_{\mathbf{n},i})}.$$

So (3.8) and (3.9) have been proved. It is well known (see [5, Theorem 5.3]) that the zeros two two consecutive elements of a family of orthogonal polynomials interlace;

then $Q'_{\mathbf{n}}(x_{\mathbf{n},i})p_{\mu_{\mathbf{n}},|\mathbf{n}|-1}(x_{\mathbf{n},i})$ must be positive. Hence for each $i = 1, \dots, |\mathbf{n}|$ the equalities (3.9) imply

$$\begin{aligned}\operatorname{sign}(\lambda_{i,k,\mathbf{n}}) &= \operatorname{sign}(a_{\mu_{\mathbf{n}},|\mathbf{n}|})\operatorname{sign}(Q_{\mathbf{n},2}) \\ &= \operatorname{sign}(s_k)\operatorname{sign}(Q_{\mathbf{n},2})\operatorname{sign}(Q_{\mathbf{n},2}) = \operatorname{sign}(s_k).\end{aligned}$$

Combining (2.6) and (3.10) we obtain (3.11). $\square$

4. PROOF OF THEOREM 1.2

We proceed as in the proof of (34) in [9, Corollary 2]. Fix $\mathbf{n} \in \mathbf{A}$. Taking into account (3.11), from (2.4) we have that for each compact set $K \subset \overline{\mathbb{C}} \setminus \Delta_1$,

$$\left\| \frac{P_{\mathbf{n},k}}{Q_{\mathbf{n}}} \right\|_K \leq \frac{\|s_k\|}{\operatorname{dist}(K, \Delta_1)}.$$

Therefore, the family of functions $\{\widehat{s}_k - P_{\mathbf{n},k}/Q_{\mathbf{n}}\}_{\mathbf{n} \in \mathbf{A}}$ is uniformly bounded on each compact $K \subset \overline{\mathbb{C}} \setminus \Delta_1$ by $2\|s_k\|/\operatorname{dist}(K, \Delta_1)$.

Let $t_{\mathbf{n},1} < \dots < t_{\mathbf{n},|\mathbf{n}|-n_k}$ denote the zeros of $Q_{\mathbf{n},2}$. From Lemma 2.3 we know that $\{t_{\mathbf{n},1}, \dots, t_{\mathbf{n},|\mathbf{n}|-n_k}\} \subset \Delta_2$ and the zeros of $Q_{\mathbf{n}}$ lie in Δ_1 , and

$$\left(\frac{\widehat{s}_k - \frac{P_{\mathbf{n},k}}{Q_{\mathbf{n}}}}{Q_{\mathbf{n},2}} \right) (z) = \mathcal{O} \left(\frac{1}{z^{2|\mathbf{n}|+1}} \right), \quad z \rightarrow \infty.$$

So

$$\frac{\widehat{s}_k - \frac{P_{\mathbf{n},k}}{Q_{\mathbf{n}}}}{\phi_{\infty}^{|\mathbf{n}|+n_k+1} \prod_{i=1}^{|\mathbf{n}|-n_k} \phi_{t_{\mathbf{n},i}}} \in \mathcal{H}(\overline{\mathbb{C}} \setminus \Delta_1).$$

Take $\rho \in (0, 1)$ such that $\gamma_{\rho} = \{z : |\phi_{\infty}(z)| = \rho\}$ satisfies the fact that $\Delta_2 \subset \operatorname{Ext}(\gamma_{\rho})$, where $\operatorname{Ext}(\gamma_{\rho})$ denotes the unbounded connected component of the complement of γ_{ρ} . We have then

$$\left\| \frac{\widehat{s}_k - \frac{P_{\mathbf{n},k}}{Q_{\mathbf{n}}}}{\phi_{\infty}^{|\mathbf{n}|+n_k+1} \prod_{i=1}^{|\mathbf{n}|-n_k} \phi_{t_{\mathbf{n},i}}} \right\|_{\gamma_{\rho}} \leq \frac{2\|s_k\|}{\operatorname{dist}(\gamma_{\rho}, \Delta_1) \psi^{2|\mathbf{n}|+1}(\gamma_{\rho})},$$

where

$$\psi(\gamma_{\rho}) = \inf\{|\phi_t(z)| : z \in \gamma_{\rho}, t \in \Delta_2 \cup \{\infty\}\}.$$

Considered as a function of the two variables z and t , $\phi_t(z)$ is a continuous function in $\overline{\mathbb{C}}^2$. Since $\gamma_{\rho} \cap \Delta_2 = \emptyset$, then $\psi(\gamma_{\rho}) > 0$. Fix a compact $K \subset \overline{\mathbb{C}} \setminus \Delta_1$ and take ρ sufficient by close to 1 so that $K \subset \operatorname{Ext}(\gamma_{\rho})$. Since the function under the norm sign is analytic in $\overline{\mathbb{C}} \setminus \Delta_1$, from the maximum principle it follows that the same bound holds for all $z \in K$. Consequently,

$$\left\| \widehat{s}_k - \frac{P_{\mathbf{n},k}}{Q_{\mathbf{n}}} \right\|_K \leq \frac{2\|s_k\| \phi_{\infty}^{|\mathbf{n}|+n_k+1} \prod_{i=1}^{|\mathbf{n}|-n_k} \phi_{t_{\mathbf{n},i}}}{\operatorname{dist}(\gamma_{\rho}, \Delta_1) \psi^{2|\mathbf{n}|+1}(\gamma_{\rho})} \leq \frac{2\|s_k\|}{\operatorname{dist}(\gamma_{\rho}, \Delta_1)} \left(\frac{\kappa(K)}{\psi(\gamma_{\rho})} \right)^{2|\mathbf{n}|+1},$$

taking $\kappa(K)$ as in the statement of the theorem. Therefore,

$$\limsup_{|\mathbf{n}| \rightarrow \infty} \left\| \widehat{s}_k - \frac{P_{\mathbf{n},k}}{Q_{\mathbf{n}}} \right\|_K^{1/2|\mathbf{n}|} \leq \frac{\kappa(K)}{\psi(\gamma_{\rho})}.$$

So, the continuity of $|\phi_t(z)|$ in $\overline{\mathbb{C}}^2$ and the fact that $\lim_{\rho \rightarrow 1} \psi(\gamma_{\rho}) = 1$ prove (1.2). The fact that $\kappa(K) < 1$ is also a consequence of the continuity of $|\phi_t(z)|$ in $\overline{\mathbb{C}}^2$. $\square$

5. PROOF OF THEOREM 1.1

We will use the following auxiliary result.

Proposition 5.1. *Let $(s_1, \dots, s_m) = \mathcal{N}(\sigma_1, \dots, \sigma_m)$ and $\Lambda \subset \mathbb{Z}_+^m$ be given. Assume that $\text{diam}(\Delta_k) < \text{dist}(\Delta_1, \Delta_2)$, $k = 1, 2$. Then there exists $N \geq 0$ such that for each $\mathbf{n} \in \Lambda$, where $|\mathbf{n}| \geq N$, every coefficient $\lambda_{i,j,\mathbf{n}}$, $i = 1, \dots, |\mathbf{n}|$, $j = 1, \dots, m$, has the same sign as its corresponding measure s_j .*

Proof. Fix an arbitrary permutation λ of $\{1, \dots, m\}$. Define Λ_λ as the set of all $\mathbf{n} \in \Lambda$ such that there exists $\tilde{\mathbf{s}} = (r_1, \dots, r_m) = \mathcal{N}(\rho_1, \dots, \rho_m)$ for which $Q_{\mathbf{n}}$ is orthogonal with respect to $(\mathbf{s}, \mathbf{n})$ and $(\tilde{\mathbf{s}}, \tilde{\mathbf{n}})$ (recall that $\tilde{\mathbf{n}} = (n_{\lambda(1)}, \dots, n_{\lambda(m)})$) in such a way that $n_{\lambda(1)} + \delta_{\lambda(1),1} \geq n_{\lambda(2)} \geq \dots \geq n_{\lambda(m)}$. According to Lemma 2.2 we have that $\bigcup_\lambda \Lambda_\lambda = \Lambda$. Some of the sets Λ_λ may be empty or have a finite number of elements. Since the group of permutations of $\{1, \dots, m\}$ is finite, it is sufficient to prove that the result holds true for all λ such that Λ_λ has an infinite number of multi-indices. In the sequel we restrict our attention to such λ 's and fix one of them.

Fix $\mathbf{n} \in \Lambda_\lambda$. Let us denote the measures introduced in (2.13) as

$$(5.1) \quad d\mu_{\mathbf{n},1} = \frac{d\rho_1}{Q_{\mathbf{n},2}} = \frac{ds_k}{Q_{\mathbf{n},2}} \quad \text{and} \quad d\mu_{\mathbf{n},2}(t) = \int \frac{Q_{\mathbf{n}}^2(x)}{t-x} \frac{d\rho_1(x)}{Q_{\mathbf{n},2}(x)} \frac{d\rho_2(t)}{Q_{\mathbf{n}}(t)Q_{\mathbf{n},3}(t)}.$$

We say $k = \lambda(1)$. From identities (3.8) in Proposition 3.2 it is sufficient to show that for each $j = 1, \dots, k-1, k+1, \dots, m$ the sequence of functions $\{S_{Q_{\mathbf{n},2}\widehat{s}_{2,j}/\widehat{s}_{2,k},|\mathbf{n}|-1,\mu_{\mathbf{n},1}}\}_{\mathbf{n} \in \Lambda_\lambda}$ converges uniformly to $\widehat{s}_{2,j}/\widehat{s}_{2,k}$ on Δ_1 because this function has constant and constant sign and no zero on Δ_1 .

Denote

$$\mathcal{K}(z, x, |\mathbf{n}| - 1) = \frac{p_{\mu_{\mathbf{n},1},|\mathbf{n}|}(z)p_{\mu_{\mathbf{n},1},|\mathbf{n}|-1}(x) - p_{\mu_{\mathbf{n},1},|\mathbf{n}|}(x)p_{\mu_{\mathbf{n},1},|\mathbf{n}|-1}(z)}{z - x}.$$

Let us start by analyzing the case when $j = 1$. Taking into account the formula (3.4) and using the identity (2.7) in Lemma 2.1 we have that

$$\begin{aligned} & \left| \frac{S_{Q_{\mathbf{n},2}/\widehat{s}_{2,k},|\mathbf{n}|-1,\mu_{\mathbf{n},1}}(z)}{Q_{\mathbf{n},2}(z)} - \frac{1}{\widehat{s}_{2,k}(z)} \right| \\ &= \left| \frac{a_{\mu_{\mathbf{n},1},|\mathbf{n}|}}{Q_{\mathbf{n},2}(z)} \int \mathcal{K}(z, x, |\mathbf{n}| - 1) \left(\frac{Q_{\mathbf{n},2}(x)}{\widehat{s}_{2,k}(x)} - \frac{Q_{\mathbf{n},2}(z)}{\widehat{s}_{2,k}(z)} \right) d\mu_{\mathbf{n},1}(x) \right| \\ &= \left| \frac{a_{\mu_{\mathbf{n},1},|\mathbf{n}|}}{Q_{\mathbf{n},2}(z)} \int \mathcal{K}(z, x, |\mathbf{n}| - 1) (Q_{\mathbf{n},2}(x)\ell_{2,k}(x) - Q_{\mathbf{n},2}(z)\ell_{2,k}(z)) d\mu_{\mathbf{n},1}(x) \right. \\ & \quad \left. + \frac{a_{\mu_{\mathbf{n},1},|\mathbf{n}|}}{Q_{\mathbf{n},2}(z)} \int \mathcal{K}(z, x, |\mathbf{n}| - 1) (Q_{\mathbf{n},2}(x)\widehat{\tau}_{2,k}(x) - Q_{\mathbf{n},2}(z)\widehat{\tau}_{2,k}(z)) d\mu_{\mathbf{n},1}(x) \right|. \end{aligned}$$

Since $\deg Q_{\mathbf{n},2}\ell_{2,k} \leq |\mathbf{n}| - n_k + 1 < |\mathbf{n}| - 1$ ($n_k = \max\{n_1, \dots, n_m\}$), then

$$\begin{aligned} & \left| \frac{S_{Q_{\mathbf{n},2}/\widehat{s}_{2,k},|\mathbf{n}|-1,\mu_{\mathbf{n},1}}(z)}{Q_{\mathbf{n},2}(z)} - \frac{1}{\widehat{s}_{2,k}(z)} \right| = |\ell_{2,k}(z) - \ell_{2,k}(z)| \\ &+ \left| \frac{a_{\mu_{\mathbf{n},1},|\mathbf{n}|}}{Q_{\mathbf{n},2}(z)} \int \mathcal{K}(z, x, |\mathbf{n}| - 1) (Q_{\mathbf{n},2}(x)\widehat{\tau}_{2,k}(x) - Q_{\mathbf{n},2}(z)\widehat{\tau}_{2,k}(z)) d\mu_{\mathbf{n},1}(x) \right| \\ &= \left| \frac{a_{\mu_{\mathbf{n},1},|\mathbf{n}|}}{Q_{\mathbf{n},2}(z)} \int \mathcal{K}(z, x, |\mathbf{n}| - 1) (Q_{\mathbf{n},2}(x)\widehat{\tau}_{2,k}(x) - Q_{\mathbf{n},2}(z)\widehat{\tau}_{2,k}(z)) d\mu_{\mathbf{n},1}(x) \right|. \end{aligned}$$

Proceeding analogously as above, for $j = 2, \dots, k-1, k+1, \dots, m$, and taking into account (3.4) and (2.8), we obtain

$$\begin{aligned} & \left| \frac{S_{Q_{\mathbf{n},2}\widehat{s}_{2,j}/\widehat{s}_{2,k},|\mathbf{n}|-1,\mu_{\mathbf{n},1}}(z)}{Q_{\mathbf{n},2}(z)} - \frac{\widehat{s}_{2,j}(z)}{\widehat{s}_{2,k}(z)} \right| \\ &= \left| \frac{a_{\mu_{\mathbf{n},1},|\mathbf{n}|}}{Q_{\mathbf{n},2}(z)} \int \mathcal{K}(z, x, |\mathbf{n}| - 1) \left(\frac{Q_{\mathbf{n},2}(x)\widehat{s}_{2,j}(x)}{\widehat{s}_{2,k}(x)} - \frac{Q_{\mathbf{n},2}(z)\widehat{s}_{2,j}(z)}{\widehat{s}_{2,k}(z)} \right) d\mu_{\mathbf{n},1}(x) \right| \\ &= \left| \frac{a_{\mu_{\mathbf{n},1},|\mathbf{n}|}}{Q_{\mathbf{n},2}(z)} \int \mathcal{K}(z, x, |\mathbf{n}| - 1) (Q_{\mathbf{n},2}(x)\widehat{\tau}_{2,j}(x) - Q_{\mathbf{n},2}(z)\widehat{\tau}_{2,j}(z)) d\mu_{\mathbf{n},1}(x) \right|. \end{aligned}$$

Summarizing, for each $j = 1, \dots, k-1, k+1, \dots, m$, we need to analyze the expression

$$\left| \frac{a_{\mu_{\mathbf{n},1},|\mathbf{n}|}}{Q_{\mathbf{n},2}(z)} \int \mathcal{K}(z, x, |\mathbf{n}| - 1) (Q_{\mathbf{n},2}(x)\widehat{\tau}_{2,j}(x) - Q_{\mathbf{n},2}(z)\widehat{\tau}_{2,j}(z)) d\mu_{\mathbf{n},1}(x) \right|.$$

Using Fubini's Theorem we obtain the following chain of equalities:

$$\begin{aligned} & \left| \frac{a_{\mu_{\mathbf{n},1},|\mathbf{n}|}}{Q_{\mathbf{n},2}(z)} \int \mathcal{K}(z, x, |\mathbf{n}| - 1) (Q_{\mathbf{n},2}(x)\widehat{\tau}_{2,j}(x) - Q_{\mathbf{n},2}(z)\widehat{\tau}_{2,j}(z)) d\mu_{\mathbf{n},1}(x) \right| \\ &= \left| \int \frac{a_{\mu_{\mathbf{n},1},|\mathbf{n}|}}{Q_{\mathbf{n},2}(z)} \int \mathcal{K}(z, x, |\mathbf{n}| - 1) \left(\frac{Q_{\mathbf{n},2}(x)}{x-t} - \frac{Q_{\mathbf{n},2}(z)}{z-t} \right) d\mu_{\mathbf{n},1}(x) d\tau_{2,j}^k(t) \right| \\ &= \left| \int \frac{a_{\mu_{\mathbf{n},1},|\mathbf{n}|}}{Q_{\mathbf{n},2}(z)} \int \mathcal{K}(z, x, |\mathbf{n}| - 1) \left(\frac{Q_{\mathbf{n},2}(x) - Q_{\mathbf{n},2}(t)}{x-t} \right) d\mu_{\mathbf{n},1}(x) d\tau_{2,j}^k(t) \right. \\ &\quad \left. - \int \frac{a_{\mu_{\mathbf{n},1},|\mathbf{n}|}}{Q_{\mathbf{n},2}(z)} \int \mathcal{K}(z, x, |\mathbf{n}| - 1) \left(\frac{Q_{\mathbf{n},2}(z) - Q_{\mathbf{n},2}(t)}{z-t} \right) d\mu_{\mathbf{n},1}(x) d\tau_{2,j}^k(t) \right. \\ &\quad \left. + \int \frac{a_{\mu_{\mathbf{n},1},|\mathbf{n}|}}{Q_{\mathbf{n},2}(z)} \int \mathcal{K}(z, x, |\mathbf{n}| - 1) \left(\frac{Q_{\mathbf{n},2}(t)}{x-t} \right) d\mu_{\mathbf{n},1}(x) d\tau_{2,j}^k(t) \right. \\ &\quad \left. - \int \frac{Q_{\mathbf{n},2}(t)}{Q_{\mathbf{n},2}(z)} \int \mathcal{K}(z, x, |\mathbf{n}| - 1) \left(\frac{Q_{\mathbf{n},2}(t)}{z-t} \right) d\mu_{\mathbf{n},1}(x) d\tau_{2,j}^k(t) \right| \\ &= \left| \int \frac{Q_{\mathbf{n},2}(t)}{Q_{\mathbf{n},2}(z)} a_{\mu_{\mathbf{n},1},|\mathbf{n}|} \int \mathcal{K}(z, x, |\mathbf{n}| - 1) \left(\frac{1}{x-t} - \frac{1}{z-t} \right) d\mu_{\mathbf{n},1}(x) d\tau_{2,j}^k(t) \right| \\ &= \left| \int \frac{Q_{\mathbf{n},2}(t)}{Q_{\mathbf{n},2}(z)} \left(S_{1/(z-t),|\mathbf{n}|-1,\mu_{\mathbf{n},1}} - \frac{1}{z-t} \right) d\tau_{2,j}^k(t) \right| \\ &\leq \left\| \frac{Q_{\mathbf{n},2}(t)}{Q_{\mathbf{n},2}(z)} \right\|_{S(\sigma_2)} \left\| S_{1/(z-t),|\mathbf{n}|-1,\mu_{\mathbf{n},1}} - \frac{1}{z-t} \right\|_{\Delta_1} \|\tau_{2,j}^k\|. \end{aligned}$$

Combining the requirement $\text{diam}(\Delta_k) < \text{dist}(\Delta_1, \Delta_2)$, $k = 1, 2$, Lemma 3.1 and Proposition 3.1, we obtain that

$$\left\| \frac{Q_{\mathbf{n},2}(t)}{Q_{\mathbf{n},2}(z)} \right\|_{S(\sigma_2)} \rightarrow 0 \quad \text{and} \quad \left\| S_{1/(z-t),|\mathbf{n}|-1,\mu_{\mathbf{n},1}} - \frac{1}{z-t} \right\|_{\Delta_1} \rightarrow 0.$$

So this completes the proof. $\square$

Now we are ready to prove Theorem 1.1. As in Section 4, we take $\rho \in (0, 1)$ and $\gamma_\rho = \{z : |\phi_\infty(z)| = \rho\}$. For each $j = 1, \dots, k-1, k+1, \dots, m$ we have that

$$|\widehat{s}_j|_{\gamma_\rho} = \frac{|s_j|}{\text{dist}(\gamma_\rho, \Delta_1)} \quad \text{and} \quad \left\| \frac{P_j}{Q_{\mathbf{n}}} \right\|_{\gamma_\rho} = \left\| \sum_{i=1}^{|\mathbf{n}|} \frac{\lambda_{i,j,\mathbf{n}}}{z - x_{\mathbf{n},i}} \right\|_{\gamma_\rho} \leq \frac{|s_j|}{\text{dist}(\gamma_\rho, \Delta_1)}.$$

The second inequality can be deduced easily from Proposition 5.1. Combining the above inequalities we have that

$$\left\| \widehat{s}_j - \frac{P_{\mathbf{n},j}}{Q_{\mathbf{n}}} \right\|_{\gamma_\rho} \leq \frac{2|s_j|}{\text{dist}(\gamma_\rho, \Delta_1)\rho^{|\mathbf{n}|+n_j+1}}.$$

Let us fix a compact $K \subset \bar{\mathbb{C}} \setminus \Delta_1$ and take ρ sufficient close to 1. From the maximum principle it follows that the same bound holds for all $z \in K$. Consequently,

$$\left\| \widehat{s}_j - \frac{P_{\mathbf{n},j}}{Q_{\mathbf{n}}} \right\|_K \leq \frac{2|s_j| \|\phi_\infty\|_K^{|\mathbf{n}|+n_j+1}}{\text{dist}(\gamma_\rho, \Delta_1)\rho^{|\mathbf{n}|+n_j+1}}.$$

Therefore,

$$\limsup_{|\mathbf{n}| \rightarrow \infty} \left\| \widehat{s}_j - \frac{P_{\mathbf{n},j}}{Q_{\mathbf{n}}} \right\|_K^{1/(|\mathbf{n}|+n_j)} \leq \frac{\|\phi_\infty\|}{\rho},$$

and the result readily follows making $\rho \rightarrow 1$.

REFERENCES

- [1] A. I. Aptekarev, *Strong asymptotics of polynomials of simultaneous orthogonality for Nikishin systems* (Russian, with Russian summary), Mat. Sb. **190** (1999), no. 5, 3–44, DOI 10.1070/SM1999v190n05ABEH000401; English transl., Sb. Math. **190** (1999), no. 5-6, 631–669. MR1702555 (2001a:42023)
- [2] A. I. Aptekarev, G. Lopes Lagomasino, and I. A. Rocha, *Asymptotic behavior of the ratio of Hermite-Padé polynomials for Nikishin systems* (Russian, with Russian summary), Mat. Sb. **196** (2005), no. 8, 3–20, DOI 10.1070/SM2005v196n08ABEH002329; English transl., Sb. Math. **196** (2005), no. 7-8, 1089–1107. MR2188362 (2006h:41040)
- [3] Zh. Bustamante and G. Lopes Lagomasino, *Hermite-Padé approximations for Nikishin systems of analytic functions* (Russian, with Russian summary), Mat. Sb. **183** (1992), no. 11, 117–138, DOI 10.1070/SM1994v077n02ABEH003446; English transl., Russian Acad. Sci. Sb. Math. **77** (1994), no. 2, 367–384. MR1208217 (94d:41026)
- [4] Jorge Bustamante Gonzalez, *Asymptotics for Angelesco-Nikishin systems*, J. Approx. Theory **85** (1996), no. 1, 43–68, DOI 10.1006/jath.1996.0028. MR1382050 (97d:41017)
- [5] T. S. Chihara, *An introduction to orthogonal polynomials*, Mathematics and its Applications, Vol. 13, Gordon and Breach Science Publishers, New York, 1978. MR0481884 (58 #1979)
- [6] Kathy Driver and Herbert Stahl, *Normality in Nikishin systems*, Indag. Math. (N.S.) **5** (1994), no. 2, 161–187, DOI 10.1016/0019-3577(94)90023-X. MR1284561 (95h:30046)
- [7] Kathy Driver and Herbert Stahl, *Simultaneous rational approximants to Nikishin-systems. I*, Acta Sci. Math. (Szeged) **60** (1995), no. 1-2, 245–263. MR1348692 (96f:41016)
- [8] Kathy Driver and Herbert Stahl, *Simultaneous rational approximants to Nikishin systems. II*, Acta Sci. Math. (Szeged) **61** (1995), no. 1-4, 261–284. MR1377364 (96m:41014)
- [9] U. Fidalgo Prieto, J. Illán, and G. López Lagomasino, *Hermite-Padé approximation and simultaneous quadrature formulas*, J. Approx. Theory **126** (2004), no. 2, 171–197, DOI 10.1016/j.jat.2004.01.004. MR2045538 (2005c:41024)
- [10] U. Fidalgo Prieto and G. López Lagomasino, *Nikishin systems are perfect*, Constr. Approx. **34** (2011), no. 3, 297–356, DOI 10.1007/s00365-011-9139-6. MR2852293 (2012i:41012)
- [11] A. A. Gonchar, E. A. Rakhmanov, and V. N. Sorokin, *On Hermite-Padé approximants for systems of functions of Markov type* (Russian, with Russian summary), Mat. Sb. **188** (1997), no. 5, 33–58, DOI 10.1070/SM1997v188n05ABEH000225; English transl., Sb. Math. **188** (1997), no. 5, 671–696. MR1478629 (98h:41017)

- [12] Abey López García and Guillermo López Lagomasino, *Relative asymptotic of multiple orthogonal polynomials for Nikishin systems*, J. Approx. Theory **158** (2009), no. 2, 214–241, DOI 10.1016/j.jat.2008.09.002. MR2519070 (2010d:42043)
- [13] André Markoff, *Deux démonstrations de la convergence de certaines fractions continues* (French), Acta Math. **19** (1895), no. 1, 93–104, DOI 10.1007/BF02402872. MR1554864
- [14] E. M. Nikišin, *Simultaneous Padé approximants* (Russian), Mat. Sb. (N.S.) **113(155)** (1980), no. 4(12), 499–519, 637. MR602272 (82e:30053)
- [15] E. M. Nikishin and V. N. Sorokin, *Rational approximations and orthogonality*, Translations of Mathematical Monographs, vol. 92, American Mathematical Society, Providence, RI, 1991. Translated from the Russian by Ralph P. Boas. MR1130396 (92i:30037)
- [16] Herbert Stahl and Vilmos Totik, *General orthogonal polynomials*, Encyclopedia of Mathematics and its Applications, vol. 43, Cambridge University Press, Cambridge, 1992. MR1163828 (93d:42029)
- [17] Walter Van Assche, Jeffrey S. Geronimo, and Arno B. J. Kuijlaars, *Riemann-Hilbert problems for multiple orthogonal polynomials*, Special functions 2000: current perspective and future directions (Tempe, AZ), NATO Sci. Ser. II Math. Phys. Chem., vol. 30, Kluwer Acad. Publ., Dordrecht, 2001, pp. 23–59, DOI 10.1007/978-94-010-0818-1_2. MR2006283 (2004e:30065)

CMUC, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COIMBRA, LARGO D. DINIS, 3001-454 COIMBRA, PORTUGAL

E-mail address: `ajplb@mat.uc.pt`

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD CARLOS III DE MADRID, AVDA. UNIVERSIDAD 30, 28911 LEGANÉS, SPAIN

Current address: Department of Mathematics, Hume Hall 305, University of Mississippi, P. O. Box 1848, University, Mississippi 38677-1848

E-mail address: `ufidalgo@olemiss.edu`

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE DE AVEIRO, CAMPUS DE SANTIAGO 3810, AVEIRO, PORTUGAL

E-mail address: `foulquie@ua.pt`