A NOTE ON TRACE SCALING ACTIONS AND FUNDAMENTAL GROUPS OF C^* -ALGEBRAS

NORIO NAWATA

(Communicated by Marius Junge)

ABSTRACT. Using the Effros-Handelman-Shen theorem and Elliott’s classification theorem of AF algebras, we show that there exists a unital simple AF algebra A with unique trace such that $A \otimes \mathbb{K}$ admits no trace scaling action of the fundamental group of A .

1. INTRODUCTION

Let M be a factor of type II_1 with a normalized trace. Murray and von Neumann introduced the fundamental group $\mathcal{F}(M)$ of M in [14]. They showed that if M is hyperfinite, then $\mathcal{F}(M) = \mathbb{R}_+^\times$. Since then there have been many works on the computation of the fundamental groups. Voiculescu [24] showed that $\mathcal{F}(L(\mathbb{F}_\infty))$ of the group factor of the free group $\mathbb{F}_\infty$ contains the positive rationals and Radulescu proved that $\mathcal{F}(L(\mathbb{F}_\infty)) = \mathbb{R}_+^\times$ in [21]. Connes [3] showed that if G is an ICC group with property (T), then $\mathcal{F}(L(G))$ is a countable group. Popa showed that any countable subgroup of $\mathbb{R}_+^\times$ can be realized as the fundamental group of some factor of type II_1 in [18]. Furthermore, Popa and Vaes [19] exhibited a large family $\mathcal{S}$ of subgroups of $\mathbb{R}_+^\times$, containing $\mathbb{R}_+^\times$ itself, all of its countable subgroups, as well as uncountable subgroups with any Hausdorff dimension in $(0, 1)$, such that for each $G \in \mathcal{S}$ there exist many free ergodic measure preserving actions of $\mathbb{F}_\infty$ for which the associated II_1 factor M has the fundamental group equal to G . In our previous paper [16] (see also [15]), we introduced the fundamental group $\mathcal{F}(A)$ of a simple unital C^* -algebra A with a normalized trace τ based on the computation of Picard groups by Kodaka [11], [12] and [13]. The fundamental group $\mathcal{F}(A)$ is defined as the set of the numbers $\tau \otimes \text{Tr}(p)$ for some projection $p \in M_n(A)$ such that $pM_n(A)p$ is isomorphic to A . We computed fundamental groups of several C^* -algebras and showed that any countable subgroup of $\mathbb{R}_+^\times$ can be realized as the fundamental group of a separable simple unital C^* -algebra with unique trace [17].

The fundamental group of a II_1 factor M is equal to the set of trace scaling constants for automorphisms of $M \otimes B(\mathcal{H})$. We have a similar fact; that is, the fundamental group of a C^* -algebra A is equal to the set of trace scaling constants for automorphisms of $A \otimes \mathbb{K}$ [16] (see also [15]). It is of interest to know whether $A \otimes \mathbb{K}$ admits a trace scaling action of $\mathcal{F}(A)$. In the case where M is a factor of type II_1 , the existence of a trace scaling (continuous) $\mathbb{R}_+^\times$ -action on $M \otimes B(\mathcal{H})$ is equivalent

Received by the editors January 4, 2012 and, in revised form, December 12, 2012.

2010 *Mathematics Subject Classification.* Primary 46L40; Secondary 06F20.

Key words and phrases. Fundamental group, trace scaling action, dimension group, Effros-Handelman-Shen theorem.

The author is a Research Fellow of the Japan Society for the Promotion of Science.

to the existence of a type III₁ factor having a core isomorphic to $M \otimes B(\mathcal{H})$ by the continuous decomposition of type III₁ factors. (See [23] and [4].) Hence this question is important in the theory of von Neumann algebras. Radulescu showed that $L(\mathbb{F}_\infty) \otimes B(\mathcal{H})$ admits a trace scaling action of $\mathbb{R}_+^\times$ in [22]. Therefore there exists a type III₁ factor having a core isomorphic to $L(\mathbb{F}_\infty) \otimes B(\mathcal{H})$. Popa and Vaes [20] showed that there exists a II₁ factor M such that $\mathcal{F}(M) = \mathbb{R}_+^\times$ and $M \otimes B(\mathcal{H})$ admits no trace scaling (continuous) action of $\mathbb{R}_+^\times$.

In this paper we consider trace scaling actions on certain AF algebras. If A is a UHF algebra, then $A \otimes \mathbb{K}$ admits a trace scaling action of $\mathcal{F}(A)$. Using the Effros-Handelman-Shen theorem and Elliott's classification theorem of AF algebras, we show that there exists a unital simple AF algebra A with unique trace such that $A \otimes \mathbb{K}$ admits no trace scaling action of $\mathcal{F}(A)$. Note that there exist remarkable works of the classification of trace scaling automorphisms in [1], [8] and [9]. But we do not consider the classification of trace scaling actions in this paper.

2. EXAMPLES

We recall some definitions in [16]. Let A be a unital simple C*-algebra with a unique normalized trace τ and Tr the usual unnormalized trace on $M_n(\mathbb{C})$. Put

$$\mathcal{F}(A) := \{\tau \otimes \text{Tr}(p) \in \mathbb{R}_+^\times \mid p \text{ is a projection in } M_n(A) \text{ such that } pM_n(A)p \cong A\}.$$

Then $\mathcal{F}(A)$ is a multiplicative subgroup of $\mathbb{R}_+^\times$ by Theorem 3.1 in [16]. For an additive subgroup E of $\mathbb{R}$ containing 1, we define the positive inner multiplier group $IM_+(E)$ of E by

$$IM_+(E) = \{t \in \mathbb{R}_+^\times \mid t \in E, t^{-1} \in E, \text{ and } tE = E\}.$$

Then we have $\mathcal{F}(A) \subset IM_+(\tau_*(K_0(A)))$ by Proposition 3.7 in [16]. This obstruction enables us to compute fundamental groups easily. For $x \in (A \otimes \mathbb{K})_+$, set $\hat{\tau}(x) = \sup\{\tau \otimes \text{Tr}(y) : y \in \bigcup_n M_n(A), y \leq x\}$. Define $\mathcal{M}_\tau^+ = \{x \geq 0 : \hat{\tau}(x) < \infty\}$ and $\mathcal{M}_\tau = \text{span}\mathcal{M}_\tau^+$. Then $\hat{\tau}$ is a densely defined (with the domain $\mathcal{M}_\tau$) lower semicontinuous trace on $A \otimes \mathbb{K}$. Since the normalized trace on a unital C*-algebra A is unique, the lower semicontinuous densely defined trace on $A \otimes \mathbb{K}$ is unique up to constant multiple. It is clear that for any $\alpha \in \text{Aut}(A \otimes \mathbb{K})$, $\hat{\tau} \circ \alpha$ is a densely defined (with the domain $\alpha^{-1}(\mathcal{M}_\tau)$) lower semicontinuous trace on $A \otimes \mathbb{K}$. Therefore there exists a positive number λ such that $\hat{\tau} \circ \alpha = \lambda \hat{\tau}$, and hence $\alpha^{-1}(\mathcal{M}_\tau) = \mathcal{M}_\tau$. We define the set of trace scaling constants for automorphisms:

$$\mathfrak{S}(A) := \{\lambda \in \mathbb{R}_+^\times \mid \hat{\tau} \circ \alpha = \lambda \hat{\tau} \text{ for some } \alpha \in \text{Aut}(A \otimes \mathbb{K})\}.$$

Then $\mathcal{F}(A) = \mathfrak{S}(A)$ by Proposition 3.28 in [16]. Therefore it is of interest to know whether $A \otimes \mathbb{K}$ admits a trace scaling action of $\mathcal{F}(A)$.

It is clear that if the fundamental group of A is singly generated, $A \otimes \mathbb{K}$ admits a trace scaling action of $\mathcal{F}(A)$. See [16] and [17] for such examples. We shall show some examples of AF algebras A such that $A \otimes \mathbb{K}$ admits a trace scaling action of $\mathcal{F}(A)$.

Example 2.1. Consider a UHF algebra $M_{2^\infty 3^\infty}$. Then the fundamental group of $M_{2^\infty 3^\infty}$ is a multiplicative subgroup generated by 2 and 3. Hence $\mathcal{F}(M_{2^\infty 3^\infty})$ is isomorphic to $\mathbb{Z}^2$ as a group. Since $M_{2^\infty 3^\infty} \otimes \mathbb{K}$ is isomorphic to $M_{2^\infty} \otimes \mathbb{K} \otimes M_{3^\infty} \otimes \mathbb{K}$, there exists a trace scaling $\mathbb{Z}^2$ -action on $M_{2^\infty 3^\infty} \otimes \mathbb{K}$. In general, if A is a UHF algebra, then $\mathcal{F}(A)$ is a free abelian group (see [16]) and $A \otimes \mathbb{K}$ admits a trace scaling action of $\mathcal{F}(A)$ (see also [5]).

Example 2.2. Let A be a unital simple AF algebra such that $K_0(A) = \mathbb{Z} + \mathbb{Z}\sqrt{3}$, $K_0(A)_+ = (\mathbb{Z} + \mathbb{Z}\sqrt{3}) \cap \mathbb{R}_+$ and $[1]_0 = 1$. Then $\mathcal{F}(A) = \{(2 + \sqrt{3})^n : n \in \mathbb{Z}\}$ (see Proposition 3.17 and Corollary 3.18 in [16]). Consider $B = M_{5^\infty} \otimes A$. Then it is easily seen that $\tau_*(K_0(B)) = \mathbb{Z}[\frac{1}{5}] + \mathbb{Z}[\frac{1}{5}]\sqrt{3}$ and τ_* is an order isomorphism. We shall show that $IM_+(\tau_*(K_0(B)))$ is generated by 5 and $2 + \sqrt{3}$. Since $\tau_*(K_0(B))$ is a subring of $\mathbb{R}$, $IM_+(\tau_*(K_0(B)))$ is a group of positive invertible elements. Define a multiplicative map N of $\mathbb{Z}[\frac{1}{5}] + \mathbb{Z}[\frac{1}{5}]\sqrt{3}$ to $\mathbb{Z}[\frac{1}{5}]$ by $N(a + b\sqrt{3}) = a^2 - 3b^2$ for any $a, b \in \mathbb{Z}[\frac{1}{5}]$. If $a + b\sqrt{3}$ is an invertible element in $\mathbb{Z}[\frac{1}{5}] + \mathbb{Z}[\frac{1}{5}]\sqrt{3}$, then there exists an integer n such that $N(a + b\sqrt{3}) = \pm 5^n$. Elementary computations show that $x^2 - 3y^2 \equiv 0 \pmod{25}$ implies $x \equiv 0 \pmod{5}$ and $y \equiv 0 \pmod{5}$. It is easy to see that no integers x and y satisfy equations $x^2 - 3y^2 = 5$, $x^2 - 3y^2 = -5$ or $x^2 - 3y^2 = -1$. There exist integers x and y that satisfy the equation $x^2 - 3y^2 = 1$. Indeed, $x = 2, y = 1$ is such an example. (See, for example, [10] and [16].) Therefore, it can be easily checked that $IM_+(\tau_*(K_0(B)))$ is generated by 5 and $2 + \sqrt{3}$. Hence we see that $\mathcal{F}(B) = \{5^n(2 + \sqrt{3})^m : n, m \in \mathbb{Z}\}$ by Proposition 3.17 in [16] and $B \otimes \mathbb{K}$ admits a trace scaling action.

We shall show that there exists a unital simple AF algebra A with unique trace such that $A \otimes \mathbb{K}$ admits no trace scaling action of $\mathcal{F}(A)$. Define

$$E = \left\{ \left(\frac{j + k\sqrt{3}}{5^{6i}}, \begin{pmatrix} x \\ y \end{pmatrix} \right) \in \mathbb{R} \times \mathbb{Z}^2 \mid i, j, k, x, y \in \mathbb{Z}, x \equiv j \pmod{9}, y \equiv k \pmod{3} \right\},$$

$$E_+ = \left\{ \left(r, \begin{pmatrix} x \\ y \end{pmatrix} \right) \in E \mid r > 0 \right\} \cup \left\{ \left(0, \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right) \right\} \quad \text{and} \quad [u]_0 = \left(1, \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right).$$

Then there exists a simple AF algebra A with a unique normalized trace τ such that $(K_0(A), K_0(A)_+, [1_A]_0) = (E, E_+, u)$ by the Effros-Handelman-Shen theorem [6].

Lemma 2.3. *With notation as above the fundamental group of A is equal to the multiplicative group generated by 5 and $2 + \sqrt{3}$.*

Proof. Since $\tau_*(K_0(A))$ is equal to $\mathbb{Z}[\frac{1}{5}] + \mathbb{Z}[\frac{1}{5}]\sqrt{3}$, $\mathcal{F}(A)$ is a subgroup of $\{5^n(2 + \sqrt{3})^m : n, m \in \mathbb{Z}\}$ by an argument in Example 2.2. Define an additive homomorphism $\phi : E \rightarrow E$ by

$$\phi\left(\left(r, \begin{pmatrix} x \\ y \end{pmatrix}\right)\right) = \left(5r, \begin{pmatrix} 5 & 9 \\ 6 & 11 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}\right).$$

Computations show that ϕ is a well-defined order isomorphism of E with $\phi(u) = \left(5, \begin{pmatrix} 5 \\ 6 \end{pmatrix}\right)$. There exist a natural number n and a projection p in $M_n(A)$ such that $[p]_0 = \left(5, \begin{pmatrix} 5 \\ 6 \end{pmatrix}\right)$ and $\tau \otimes \text{Tr}(p) = 5$. Since we have $(K_0(pM_n(A)p), K_0(pM_n(A)p)_+, [p]_0) = (E, E_+, \left(5, \begin{pmatrix} 5 \\ 6 \end{pmatrix}\right))$, there exists an isomorphism $f : A \rightarrow pM_n(A)p$ with $f_* = \phi$ by Elliott's classification theorem of AF algebras [7]. Therefore $5 \in \mathcal{F}(A)$. Define an additive homomorphism $\psi : E \rightarrow E$ by

$$\psi\left(\left(r, \begin{pmatrix} x \\ y \end{pmatrix}\right)\right) = \left((2 + \sqrt{3})r, \begin{pmatrix} 2 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}\right).$$

Then we see that $2 + \sqrt{3} \in \mathcal{F}(A)$. Consequently $\mathcal{F}(A)$ is the multiplicative group generated by 5 and $2 + \sqrt{3}$. $\square$

We shall consider the order automorphisms of (E, E_+) .

Lemma 2.4. *Let ϕ be an order automorphism of (E, E_+) . Then there exist integers a, b, c, d and a positive invertible element λ in $\mathbb{Z}[\frac{1}{5}] + \mathbb{Z}[\frac{1}{5}]\sqrt{3}$ such that $ad - bc = \pm 1$ and*

$$\phi((r, \begin{pmatrix} x \\ y \end{pmatrix})) = (\lambda r, \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}).$$

Moreover if $\lambda = 5$, then

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \equiv \begin{pmatrix} 5 & 0 \\ 0 & 2 \end{pmatrix}, \begin{pmatrix} 5 & 0 \\ 3 & 2 \end{pmatrix}, \begin{pmatrix} 5 & 0 \\ 6 & 2 \end{pmatrix} \pmod{9},$$

and if $\lambda = 2 + \sqrt{3}$, then

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \equiv \begin{pmatrix} 2 & 3 \\ 1 & 2 \end{pmatrix}, \begin{pmatrix} 2 & 3 \\ 4 & 2 \end{pmatrix}, \begin{pmatrix} 2 & 3 \\ 7 & 2 \end{pmatrix} \pmod{9}.$$

Proof. We denote by $(\phi_1((r, \begin{pmatrix} x \\ y \end{pmatrix})), \phi_2((r, \begin{pmatrix} x \\ y \end{pmatrix})))$ the element $\phi((r, \begin{pmatrix} x \\ y \end{pmatrix}))$ for any $(r, \begin{pmatrix} x \\ y \end{pmatrix}) \in E$. Consider a subgroup F generated by $(0, \begin{pmatrix} 9 \\ 0 \end{pmatrix})$ and $(0, \begin{pmatrix} 0 \\ 3 \end{pmatrix})$. Then F is an ϕ -invariant subgroup because ϕ is an order isomorphism. Hence there exist integers m_1, m_2, m_3 and m_4 such that $m_1 m_4 - m_2 m_3 = \pm 1$ and $\phi_2((0, \begin{pmatrix} x \\ y \end{pmatrix})) = \begin{pmatrix} m_1 & 3m_2 \\ \frac{m_3}{3} & m_4 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$ for any $(0, \begin{pmatrix} x \\ y \end{pmatrix}) \in F$. Furthermore we see that there exists a positive invertible element λ in $\mathbb{Z}[\frac{1}{5}] + \mathbb{Z}[\frac{1}{5}]\sqrt{3}$ such that $\phi_1((r, \begin{pmatrix} x \\ y \end{pmatrix})) = \lambda r$. Since $5^{6i} \phi((\frac{9}{5^{6i}}, \begin{pmatrix} 0 \\ 0 \end{pmatrix})) = \phi((9, \begin{pmatrix} 0 \\ 0 \end{pmatrix}))$ for any $i \in \mathbb{Z}$, we see that $\phi_2((9, \begin{pmatrix} 0 \\ 0 \end{pmatrix})) = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$. This observation and easy computations show that $\phi((1, \begin{pmatrix} 1 \\ 0 \end{pmatrix})) = (\lambda, \begin{pmatrix} m_1 & 3m_2 \\ \frac{m_3}{3} & m_4 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix})$ and $\frac{m_3}{3} \in \mathbb{Z}$. In a similar way, we see that $\phi((\sqrt{3}, \begin{pmatrix} 0 \\ 1 \end{pmatrix})) = (\lambda\sqrt{3}, \begin{pmatrix} m_1 & 3m_2 \\ \frac{m_3}{3} & m_4 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix})$. It is easily seen that ϕ is determined by the values of $\phi((1, \begin{pmatrix} 1 \\ 0 \end{pmatrix}))$, $\phi((\sqrt{3}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}))$, $\phi((0, \begin{pmatrix} 9 \\ 0 \end{pmatrix}))$ and $\phi((0, \begin{pmatrix} 0 \\ 3 \end{pmatrix}))$. Therefore there exist integers a, b, c, d and a positive invertible element λ in $\mathbb{Z}[\frac{1}{5}] + \mathbb{Z}[\frac{1}{5}]\sqrt{3}$ such that $ad - bc = \pm 1$ and

$$\phi((r, \begin{pmatrix} x \\ y \end{pmatrix})) = (\lambda r, \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}).$$

Let $\lambda = 5$; then $a \equiv 5 \pmod{9}$, $b \equiv 0 \pmod{9}$, $c \equiv 0 \pmod{3}$ and $d \equiv 5 \pmod{3}$ by the definition of E . If $ad - bc = 1$, then $d \equiv 5^5 \pmod{9}$, $-b \equiv 0 \pmod{9}$, $-c \equiv 0 \pmod{3}$ and $a \equiv 5^5 \pmod{3}$ because ϕ is an isomorphism. Therefore computations show that

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \equiv \begin{pmatrix} 5 & 0 \\ 0 & 2 \end{pmatrix}, \begin{pmatrix} 5 & 0 \\ 3 & 2 \end{pmatrix}, \begin{pmatrix} 5 & 0 \\ 6 & 2 \end{pmatrix} \pmod{9}.$$

If $ad - bc = -1$, then $-d \equiv 5^5 \pmod{9}$, $b \equiv 0 \pmod{9}$, $c \equiv 0 \pmod{3}$ and $-a \equiv 5^5 \pmod{3}$. There does not exist an integer a such that $a \equiv 5 \pmod{9}$ and $-a \equiv 5^5 \pmod{3}$.

Therefore we reach a conclusion in the case $\lambda = 5$. In the case $\lambda = 2 + \sqrt{3}$, a similar argument as above proves the lemma. $\square$

Theorem 2.5. *There exists a unital simple AF algebra A with unique trace such that $A \otimes \mathbb{K}$ admits no trace scaling action of $\mathcal{F}(A)$.*

Proof. Let

$$E = \left\{ \left(\frac{j + k\sqrt{3}}{5^{6i}}, \begin{pmatrix} x \\ y \end{pmatrix} \right) \in \mathbb{R} \times \mathbb{Z}^2 \mid i, j, k, x, y \in \mathbb{Z}, x \equiv j \pmod{9}, y \equiv k \pmod{3} \right\},$$

$$E_+ = \left\{ (r, \begin{pmatrix} x \\ y \end{pmatrix}) \in E \mid r > 0 \right\} \cup \left\{ (0, \begin{pmatrix} 0 \\ 0 \end{pmatrix}) \right\} \quad \text{and} \quad [u]_0 = (1, \begin{pmatrix} 1 \\ 0 \end{pmatrix}).$$

Then there exists a simple AF algebra A with a unique normalized trace τ such that $(K_0(A), K_0(A)_+, [1_A]_0) = (E, E_+, u)$ by the Effros-Handelman-Shen theorem [6]. By Lemma 2.3, $\mathcal{F}(A) = \{5^n(2 + \sqrt{3})^m : n, m \in \mathbb{Z}\}$. Let α be an automorphism of $A \otimes \mathbb{K}$ such that $\hat{\tau} \circ \alpha = 5\hat{\tau}$ and β an automorphism of $A \otimes \mathbb{K}$ such that $\hat{\tau} \circ \beta = (2 + \sqrt{3})\hat{\tau}$. Then α_* and β_* are order isomorphisms of $(K_0(A), K_0(A)_+)$. Lemma 2.4 and computations show that $\alpha_* \circ \beta_* \neq \beta_* \circ \alpha_*$. Therefore $A \otimes \mathbb{K}$ admits no trace scaling action of $\mathcal{F}(A)$. $\square$

Remark 2.6. Let A be a unital simple C^* -algebra with a unique normalized trace τ . We denote by $\text{Pic}(A)$ the Picard group of A (see [2]). Assume that the normalized trace on A separates equivalence classes of projections. Then we have the following exact sequence [16] (see also [11]):

$$1 \longrightarrow \text{Out}(A) \xrightarrow{\rho_A} \text{Pic}(A) \xrightarrow{T} \mathcal{F}(A) \longrightarrow 1.$$

If $A \otimes \mathbb{K}$ admits a trace scaling action of $\mathcal{F}(A)$, then $\text{Pic}(A)$ is isomorphic to a semidirect product of $\text{Out}(A)$ with $\mathcal{F}(A)$. Example 2.1 and Example 2.2 are such examples. We do not know whether there exists a simple C^* -algebra A with a unique normalized trace τ such that the normalized trace on A separates equivalence classes of projections and $A \otimes \mathbb{K}$ admits no trace scaling action of $\mathcal{F}(A)$.

Remark 2.7. If A is a C^* -algebra in the proof of Theorem 2.5, then it can be checked that $\text{Out}(A)$ is not a normal subgroup of $\text{Pic}(A)$ by Lemma 2.4 in this paper, Proposition 1.5 in [11] and Elliott's classification theorem of AF algebras.

REFERENCES

- [1] Ola Bratteli and Akitaka Kishimoto, *Trace scaling automorphisms of certain stable AF algebras. II*, Q. J. Math. **51** (2000), no. 2, 131–154, DOI 10.1093/qjmath/51.2.131. MR1765786 (2001j:46092)
- [2] Lawrence G. Brown, Philip Green, and Marc A. Rieffel, *Stable isomorphism and strong Morita equivalence of C^* -algebras*, Pacific J. Math. **71** (1977), no. 2, 349–363. MR0463928 (57 #3866)
- [3] A. Connes, *A factor of type II_1 with countable fundamental group*, J. Operator Theory **4** (1980), no. 1, 151–153. MR587372 (81j:46099)
- [4] Alain Connes and Masamichi Takesaki, *The flow of weights on factors of type III*, Tôhoku Math. J. (2) **29** (1977), no. 4, 473–575, DOI 10.2748/tmj/1178240493. MR480760 (82a:46069a)
- [5] Giulio Della Rocca and Masamichi Takesaki, *The role of groupoids in classification theory: a new approach. The UHF algebra case*, Groupoids in analysis, geometry, and physics (Boulder, CO, 1999), Contemp. Math., vol. 282, Amer. Math. Soc., Providence, RI, 2001, pp. 47–65, DOI 10.1090/conm/282/04678. MR1855242 (2002i:46049)

- [6] Edward G. Effros, David E. Handelman, and Chao Liang Shen, *Dimension groups and their affine representations*, Amer. J. Math. **102** (1980), no. 2, 385–407, DOI 10.2307/2374244. MR564479 (83g:46061)
- [7] George A. Elliott, *On the classification of inductive limits of sequences of semisimple finite-dimensional algebras*, J. Algebra **38** (1976), no. 1, 29–44. MR0397420 (53 #1279)
- [8] George A. Elliott, David E. Evans, and Akitaka Kishimoto, *Outer conjugacy classes of trace scaling automorphisms of stable UHF algebras*, Math. Scand. **83** (1998), no. 1, 74–86. MR1662080 (2000h:46076)
- [9] David E. Evans and Akitaka Kishimoto, *Trace scaling automorphisms of certain stable AF algebras*, Hokkaido Math. J. **26** (1997), no. 1, 211–224. MR1432548 (98e:46081)
- [10] Michael J. Jacobson Jr. and Hugh C. Williams, *Solving the Pell equation*, CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, Springer, New York, 2009. MR2466979 (2009i:11003)
- [11] Kazunori Kodaka, *Full projections, equivalence bimodules and automorphisms of stable algebras of unital C^* -algebras*, J. Operator Theory **37** (1997), no. 2, 357–369. MR1452282 (98c:46149)
- [12] Kazunori Kodaka, *Picard groups of irrational rotation C^* -algebras*, J. London Math. Soc. (2) **56** (1997), no. 1, 179–188, DOI 10.1112/S0024610797005243. MR1462834 (98e:46071)
- [13] Kazunori Kodaka, *Projections inducing automorphisms of stable UHF-algebras*, Glasg. Math. J. **41** (1999), no. 3, 345–354, DOI 10.1017/S0017089599000336. MR1720450 (2000m:46124)
- [14] F. J. Murray and J. von Neumann, *On rings of operators. IV*, Ann. of Math. (2) **44** (1943), 716–808. MR0009096 (5,101a)
- [15] Norio Nawata, *Fundamental group of simple C^* -algebras with unique trace III*, Canad. J. Math. **64** (2012), no. 3, 573–587, DOI 10.4153/CJM-2011-052-0. MR2962317
- [16] Norio Nawata and Yasuo Watatani, *Fundamental group of simple C^* -algebras with unique trace*, Adv. Math. **225** (2010), no. 1, 307–318, DOI 10.1016/j.aim.2010.02.019. MR2669354 (2012h:46090)
- [17] Norio Nawata and Yasuo Watatani, *Fundamental group of simple C^* -algebras with unique trace II*, J. Funct. Anal. **260** (2011), no. 2, 428–435, DOI 10.1016/j.jfa.2010.09.014. MR2737407 (2012g:46080)
- [18] Sorin Popa, *Strong rigidity of II_1 factors arising from malleable actions of w -rigid groups. I*, Invent. Math. **165** (2006), no. 2, 369–408, DOI 10.1007/s00222-006-0501-4. MR2231961 (2007f:46058)
- [19] Sorin Popa and Stefaan Vaes, *Actions of $\mathbb{F}_\infty$ whose II_1 factors and orbit equivalence relations have prescribed fundamental group*, J. Amer. Math. Soc. **23** (2010), no. 2, 383–403, DOI 10.1090/S0894-0347-09-00644-4. MR2601038 (2011j:46102)
- [20] Sorin Popa and Stefaan Vaes, *On the fundamental group of II_1 factors and equivalence relations arising from group actions*, Quanta of maths, Clay Math. Proc., vol. 11, Amer. Math. Soc., Providence, RI, 2010, pp. 519–541. MR2732063 (2012j:37004)
- [21] Florin Rădulescu, *The fundamental group of the von Neumann algebra of a free group with infinitely many generators is $\mathbb{R}_+ \setminus 0$* , J. Amer. Math. Soc. **5** (1992), no. 3, 517–532, DOI 10.2307/2152703. MR1142260 (93a:46117)
- [22] Florin Rădulescu, *A one-parameter group of automorphisms of $\mathcal{L}(F_\infty) \otimes B(H)$ scaling the trace* (English, with French summary), C. R. Acad. Sci. Paris Sér. I Math. **314** (1992), no. 13, 1027–1032. MR1168529 (93i:46111)
- [23] Masamichi Takesaki, *Duality for crossed products and the structure of von Neumann algebras of type III*, Acta Math. **131** (1973), 249–310. MR0438149 (55 #11068)
- [24] Dan Voiculescu, *Circular and semicircular systems and free product factors*, Operator algebras, unitary representations, enveloping algebras, and invariant theory (Paris, 1989), Progr. Math., vol. 92, Birkhäuser Boston, Boston, MA, 1990, pp. 45–60. MR1103585 (92e:46124)

DEPARTMENT OF MATHEMATICS AND INFORMATICS, GRADUATE SCHOOL OF SCIENCE, CHIBA UNIVERSITY, 1-33 YAYOI-CHO, INAGE, CHIBA, 263-8522, JAPAN

Current address: Department of Arts and Sciences, Osaka Kyoiku University, 4-698-1 Asahi-gaoka, Kashiwara, Osaka, 582-8582, Japan

E-mail address: `nawata@cc.osaka-kyoiku.ac.jp`