PROPERTY (T) AND THE FURSTENBERG ENTROPY OF NONSINGULAR ACTIONS

LEWIS BOWEN, YAIR HARTMAN, AND OMER TAMUZ

(Communicated by Nimish Shah)

ABSTRACT. We establish a new characterization of property (T) in terms of the Furstenberg entropy of nonsingular actions. Given any generating measure μ on a countable group G , A. Nevo showed that a necessary condition for G to have property (T) is that the Furstenberg μ -entropy values of the ergodic, properly nonsingular G -actions are bounded away from zero. We show that this is also a sufficient condition.

1. INTRODUCTION

A measurable action on a probability space $G \curvearrowright (X, \eta)$ is called a *nonsingular action* if the measure η is quasi-invariant with respect to any $g \in G$; that is, if η and $g_*\eta$ are equivalent (i.e., mutually absolutely continuous) measures. We say that the action is *properly nonsingular* if η is not equivalent to a G -invariant probability measure.

A probability measure μ on a countable group G is *generating* if its support generates G as a semigroup.

Let $G \curvearrowright (X, \eta)$ be a nonsingular action and let μ be a probability measure on G . The *Furstenberg entropy* [5] or μ -*entropy* is defined by

$$h_\mu(X, \eta) = \sum_{g \in G} \mu(g) \int_X -\log \frac{dg_*^{-1}\eta}{d\eta}(x) \, d\eta(x).$$

Jensen's inequality implies that $h_\mu(X, \eta) \geq 0$ and that, for generating measures, equality holds if and only if η is an invariant measure. Furstenberg entropy is an important conjugacy invariant of nonsingular actions [11], and in particular of *stationary* actions; $G \curvearrowright (X, \eta)$ is μ -stationary if $\mu * \eta = \eta$.

The pair (G, μ) is said to have an *entropy gap* if there exists some constant $\epsilon = \epsilon(G, \mu) > 0$ such that the μ -entropy of any ergodic, properly nonsingular G -action on a probability space is at least ϵ .

A group G has *property (T)* if any unitary representation of G which has almost invariant vectors admits a nonzero invariant vector. The goal of this paper is to establish a similar characterization of property (T), using nonsingular actions instead of unitary representations, and thinking of those with small entropy values as being “almost invariant”.

Received by the editors June 30, 2014 and, in revised form, December 1, 2014.

2010 *Mathematics Subject Classification.* Primary 20F69, 37A40.

The first author was supported in part by NSF grant DMS-0968762, NSF CAREER Award DMS-0954606 and BSF grant 2008274.

The second author was supported by the European Research Council, grant 239885.

Theorem 1. *Let G be a countable group. Then G has property (T) if and only if for every (equivalently, for some) generating measure μ , (G, μ) has an entropy gap.*

For the case that G has property (T) this theorem follows from Nevo [10, Propositions 4.1, 4.2]. Note that the statements of these propositions omit the requirement that η is ergodic and properly nonsingular; the result is no longer true without either of these hypotheses. For completeness, we include in Appendix A a reproduction of Nevo's proof of this direction.

The idea of the proof of the other direction (i.e., that groups without property (T) have no entropy gap) is the following. We first consider the group $2_{fin}^{\mathbb{N}}$; the elements of this group are the finite subsets of $\mathbb{N}$ and the group operation is symmetric difference. For this group, we construct in Section 2 very simple nonsingular actions which show that $2_{fin}^{\mathbb{N}}$ has no entropy gap, for a family of natural measures on $2_{fin}^{\mathbb{N}}$.

Then, in Section 3, we observe that any countable group G without property (T) admits a nontrivial cocycle into $2_{fin}^{\mathbb{N}}$. Using these cocycles one can construct skew-product G -actions over $2_{fin}^{\mathbb{N}}$ spaces; the entropy formula for skew-product actions is established in Section 4. Finally, in Section 5 we prove Theorem 1, by showing that the $2_{fin}^{\mathbb{N}}$ -actions that we construct in Section 2, and which have arbitrarily low Furstenberg entropy, can be lifted to skew-product G -actions with arbitrarily low Furstenberg entropy.

1.1. Related literature.

1.1.1. *Characterizing property (T).* Recently, Ozawa [12] showed that property (T) can be characterized by the spectral gap of $\Delta_\mu \in \mathbb{R}[G]$, the Laplacian associated with a finitely supported, symmetric generating measure μ . This characterization is one of many which have been established since Kařdan's first definition of property (T) [9], some of which have been instrumental in proving property (T) for groups that were previously not known to have it. We refer the reader to Bekka, de La Harpe and Valette [1] for a complete discussion.

1.1.2. *Stationary actions.* A particularly interesting class of nonsingular actions are the stationary ones. Furstenberg entropy has been a useful tool in the study of stationary actions (e.g., [5, 8]), and the study of the set of entropy values realizable by properly nonsingular stationary actions has attracted some interest [2, 6, 11].

Nevo's theorem (Theorem A.1) implies that for every (G, μ) , where G has property (T), the Furstenberg entropy values of properly nonsingular, ergodic μ -stationary actions are bounded away from zero; in this case we say that (G, μ) has a *stationary entropy gap*. In [2] and [6] it is shown that some (G, μ) without property (T) have no stationary entropy gap. However, this is not a characterization of property (T): in a previous paper [3, Proposition 7.5] we show that there exist (G, μ) without property (T), but with a stationary entropy gap.

A question that therefore remains open is that of characterizing the pairs (G, μ) that have a stationary entropy gap. More narrowly, it is not even known that all amenable groups have no stationary entropy gap.

2. THE CASE OF $2_{fin}^{\mathbb{N}}$

Let $2_{fin}^{\mathbb{N}}$ denote the set of all finite subsets of the natural numbers $\mathbb{N} = \{1, 2, 3, \dots\}$. Endowed with the operation of symmetric difference, $2_{fin}^{\mathbb{N}}$ is an abelian group in which every element other than the identity is an involution. For $T \in 2_{fin}^{\mathbb{N}}$,

let $\max(T) = \max_{t \in T} t$. We also let $\max(\emptyset) = 0$. Because $2_{fin}^{\mathbb{N}}$ is an abelian group we will use additive notation when expressing group multiplication. Thus $T + S := T \triangle S = (T \setminus S) \cup (S \setminus T)$ for any $T, S \in 2_{fin}^{\mathbb{N}}$.

Let $2_{fin}^{\mathbb{N}}$ act on $2^{\mathbb{N}}$ by symmetric difference, identifying elements of $2^{\mathbb{N}}$ with subsets of $\mathbb{N}$. Let ω_p be the measure on $2^{\mathbb{N}}$ given by $\omega_p = \prod_{\mathbb{N}} B(p)$, where $B(p)$ is the Bernoulli p measure on $\{0, 1\}$.

Proposition 2.1. *Consider the action $2_{fin}^{\mathbb{N}} \curvearrowright (2^{\mathbb{N}}, \omega_p)$. Then, for any probability measure μ on $2_{fin}^{\mathbb{N}}$, the μ -entropy of $(2^{\mathbb{N}}, \omega_p)$ is*

$$h_{\mu}(2^{\mathbb{N}}, \omega_p) = \varphi(p) \sum_{T \in 2_{fin}^{\mathbb{N}}} \mu(T) |T| \leq \varphi(p) \sum_{T \in 2_{fin}^{\mathbb{N}}} \mu(T) \max(T),$$

where $\varphi(p) = \log \frac{p}{1-p} (2p - 1)$.

Note that $\lim_{p \rightarrow 1/2} \varphi(p) = 0$ and that for $p \neq 1/2$ it holds that $\varphi(p) > 0$.

Proof. For $T \in 2_{fin}^{\mathbb{N}}$, $T_*^{-1} \omega_p = T_* \omega_p = \prod_{n \in T} B(1-p) \prod_{n \notin T} B(p)$. Hence

$$\int_X -\log \frac{dT_* \omega}{d\omega}(x) d\omega(x) = \varphi(p) |T|,$$

where $\varphi(p) = \log \frac{p}{1-p} (2p - 1)$ is the Kullback-Leibler divergence between $B(p)$ and $B(1-p)$. Finally, it follows that for any measure μ on $2_{fin}^{\mathbb{N}}$,

$$h_{\mu}(2^{\mathbb{N}}, \omega_p) = \varphi(p) \sum_{T \in 2_{fin}^{\mathbb{N}}} \mu(T) |T| \leq \varphi(p) \sum_{T \in 2_{fin}^{\mathbb{N}}} \mu(T) \max(T).$$

□

It is easy to show that ω_p is ergodic and is not equivalent to an invariant probability measure (see Lemma 4.2; an invariant measure λ would have to satisfy $\lambda(\{x : x_n = 1\}) = 1/2$). Hence it follows from this proposition that $2_{fin}^{\mathbb{N}}$ does not have an entropy gap, for any μ with $\sum_{T \in 2_{fin}^{\mathbb{N}}} \mu(T) \max(T) < \infty$. In the remainder of this paper we lift this result to any group without property (T).

3. NONPROPERTY (T) AND COCYCLES

Before stating the next proposition, let us recall some standard definitions. Given a nonsingular action $G \curvearrowright (X, \eta)$ and a countable group Γ , a *cocycle* $c : G \times X \rightarrow \Gamma$ is a measurable map such that

$$c(gh, x) = c(g, hx) c(h, x)$$

for a.e. x and every $g, h \in G$. Two such cocycles $c_1, c_2 : G \times X \rightarrow \Gamma$ are *cohomologous* if there exists a measurable function $\beta : X \rightarrow \Gamma$ such that

$$c_1(g, x) = \beta(gx) c_2(g, x) \beta(x)^{-1}$$

for every $g \in G$ and a.e. x . A cocycle is a *coboundary* if it is cohomologous to the trivial cocycle (whose essential image is contained in the trivial element $\{e\}$).

Proposition 3.1. *Assume G does not have property (T). Let $\rho : G \rightarrow \mathbb{N}$ be a proper function (so $\rho^{-1}(n)$ is finite for every $n \in \mathbb{N}$). Then there exists an ergodic*

probability measure preserving action $G \curvearrowright (X, \eta)$ and a Borel cocycle $c : G \times X \rightarrow 2_{fin}^{\mathbb{N}}$ such that c is not cohomologous to a cocycle with a finite image. Moreover, for every element $g \in G$,

$$\int_X \max(c(g, x)) \, d\eta(x) \leq \rho(g).$$

Proof. This result follows from the proof of [7, Theorem 2.1] as we now explain. It is a well-known result of Connes and Weiss [4] that if G does not have property (T), then there exists an ergodic probability measure preserving action $G \curvearrowright (X, \eta)$ which is not strongly ergodic (see also [1, Theorem 6.3.4]). By [7, Lemma 2.4] there exist Borel sets $D_n \subset X$ such that the following hold.

- (1) D_n is asymptotically invariant: $\lim_{n \rightarrow \infty} \eta(gD_n \Delta D_n) = 0$ for every $g \in G$.
- (2) $\lim_{n \rightarrow \infty} \eta(D_n) = 1/2$.
- (3) For every $g \in G$ with $\rho(g) \leq n$, $\eta(D_n \Delta gD_n) < 2^{-n}$.

To be precise, the last statement follows from an easy modification of the proof of [7, Lemma 2.4]; see equation (2.10) there.

Define a map $\phi : X \rightarrow 2_{fin}^{\mathbb{N}}$ by $\phi(x) = \{n \in \mathbb{N} : x \in D_n\}$. We let $2_{fin}^{\mathbb{N}}$ act on $2^{\mathbb{N}}$ by symmetric difference.

Let $g \in G$. If $m \geq \rho(g)$, then

$$\eta(\{x \in X : \phi(gx) \notin 2_{fin}^{\mathbb{N}} \Delta \phi(x)\}) \leq \sum_{k=m}^{\infty} \eta(D_k \Delta gD_k) < 2^{-m+1}.$$

Since $m \geq \rho(g)$ is arbitrary, the left-hand side must equal zero. Therefore $\phi(gx) \in 2_{fin}^{\mathbb{N}} \Delta \phi(x)$ for η -a.e. $x \in X$.

Define the cocycle $c : G \times X \rightarrow 2_{fin}^{\mathbb{N}}$ by $c(g, x) = T$ if $\phi(gx) = T \Delta \phi(x)$. If $\rho(g) = m$, then

$$\begin{aligned} \int_X \max(c(g, x)) \, d\eta(x) &\leq m - 1 + \int_{\{x : \max(c(g, x)) \geq m\}} \max(c(g, x)) \, d\eta(x) \\ &\leq m - 1 + \sum_{n=m}^{\infty} \eta(\{x \in X : \max(c(g, x)) \geq n\}) \\ &\leq m - 1 + \sum_{n=m}^{\infty} \eta(D_n \Delta gD_n) \leq m - 1 + 2^{-m+1} \leq m. \end{aligned}$$

The cocycle c cannot be cohomologous to a cocycle with a finite image, since ϕ is nontrivial. To be precise, for $m \in \mathbb{N}$, we let $2^m < 2_{fin}^{\mathbb{N}}$ denote the subgroup consisting of all elements $T \in 2_{fin}^{\mathbb{N}}$ with support in $[m] := \{1, \dots, m\}$. Denote by $[m]^c = \{m+1, m+2, \dots\}$ the complement of $[m]$ in $\mathbb{N}$. To obtain a contradiction, suppose there is a Borel map $f : X \rightarrow 2_{fin}^{\mathbb{N}}$ and a cocycle $b : G \times X \rightarrow 2^m$ such that

$$c(g, x) = f(gx) + b(g, x) + f(x)$$

for a.e. $x \in X$ and every $g \in G$. Note that all elements of $2_{fin}^{\mathbb{N}}$ are involutions, and so $-f(x) = f(x)$.

Let $\psi(x) = f(x) \triangle \phi(x)$. Observe that

$$\begin{aligned}\psi(gx) &= f(gx) \triangle \phi(gx) = f(gx) \triangle c(g, x) \triangle \phi(x) \\ &= f(gx) \triangle f(gx) \triangle b(g, x) \triangle f(x) \triangle \phi(x) \\ &= b(g, x) \triangle \psi(x)\end{aligned}$$

for a.e. $x \in X$ and every $g \in G$.

Let $\tilde{\psi}(x) = \psi(x) \cap [m]^c$, $\tilde{f}(x) = f(x) \cap [m]^c$ and $\tilde{\phi}(x) = \phi(x) \cap [m]^c$. Then, since $b(g, x) \subseteq [m]$,

$$\tilde{\psi}(gx) = (b(g, x) \triangle \psi(x)) \cap [m]^c = \tilde{\psi}(x),$$

and $\tilde{\psi}$ is G -invariant. Because $G \curvearrowright (X, \eta)$ is ergodic, there is an element $R \in 2^{[m]^c}$ such that $\tilde{\psi}(x) = R$ for a.e. x . Thus

$$R = \tilde{\psi}(x) = \tilde{f}(x) \triangle \tilde{\phi}(x),$$

which implies $\tilde{\phi}(x) \triangle R = \tilde{f}(x) \in 2_{fin}^{\mathbb{N}}$ for a.e. $x \in X$.

Let $D'_n = D_n$ if $n \notin R$, and otherwise let $D'_n = X \setminus D_n$, the complement of D_n . Then, for a.e. $x \in X$, there are only a finite number of elements $j \in [m]^c$ such that $x \in D'_j$; these are precisely the elements in $\tilde{f}(x)$.

It follows that if we let $A_n = \{x \in X : \max(\tilde{f}(x)) \geq n\}$, then $\lim_n \eta(A_n) = 0$. But A_n contains D'_n , and so $\eta(A_n) \geq \eta(D'_n)$, in contradiction to the fact that $\lim_n \eta(D'_n) = 1/2$. $\square$

4. ENTROPY FOR SKEW-PRODUCTS

Lemma 4.1. *Let G be a countable group with a probability measure μ . Let $G \curvearrowright (X, \eta)$ be a probability measure preserving action, and let $c : G \times X \rightarrow \Gamma$ be a cocycle to a countable group Γ . Also, let $\Gamma \curvearrowright (W, \omega)$ be a nonsingular action on the probability space (W, ω) . Let $G \curvearrowright X \times W$ be the skew-product action*

$$g(x, w) = (gx, c(g, x)w).$$

Then

$$h_\mu(X \times W, \eta \times \omega) = \int_X h_{\mu_x}(W, \omega) \, d\eta(x),$$

where μ_x is the pushforward of μ under the map $g \mapsto c(g, x)$.

Proof.

$$\begin{aligned}h_\mu(X \times W, \eta \times \omega) &= \sum_{g \in G} \mu(g) \int_{X \times W} -\log \left(\frac{dg_*^{-1}(\eta \times \omega)}{d\eta \times \omega}(x, w) \right) \, d\eta \times \omega(x, w) \\ &= \sum_{g \in G} \mu(g) \int_{X \times W} -\log \left(\frac{dc(g, x)_*^{-1}\omega}{d\omega}(w) \right) \, d\eta \times \omega(x, w).\end{aligned}$$

Note that the integrand is the sum of an integrable function with a nonnegative function: if we denote $r(x, w) = \frac{dc(g, x)_*^{-1}\omega}{d\omega}(w)$, then r is in $L^1(X \times W, \eta \times \omega)$, and therefore the negative part of $-\log r$ is integrable. We can thus apply the Fubini-Tonelli Theorem and write

$$= \sum_{g \in G} \mu(g) \int_X \left(\int_W -\log \left(\frac{dc(g, x)_*^{-1}\omega}{d\omega}(w) \right) \, d\omega(w) \right) \, d\eta(x).$$

The innermost integral is always nonnegative, since it is a relative entropy. We can therefore exchange the order of integration, so that

$$\begin{aligned} &= \int_X \sum_{\gamma \in \Gamma} \mu_x(\gamma) \int_W -\log \left(\frac{d\gamma_*^{-1}\omega}{d\omega}(w) \right) d\omega(w) d\eta(x) \\ &= \int_X h_{\mu_x}(W, \omega) d\eta(x). \end{aligned}$$

□

Lemma 4.2. *Let $E_n = \{x \in 2^{\mathbb{N}} : x_n = 1\}$. Let ν be a Borel probability measure on $2^{\mathbb{N}}$ such that $\nu(E_n) \leq 1/2$ for infinitely many $n \in \mathbb{N}$, and let ω_p be the i.i.d. Bernoulli p measure on $2^{\mathbb{N}}$, with $p > 1/2$. Then ν is not absolutely continuous with respect to ω_p .*

Proof. Let $\{i_k\}_{k=1}^{\infty}$ be a sequence of indices such that $\nu(E_{i_k}) < 1/2$. For $x \in 2^{\mathbb{N}}$, denote

$$S_n(x) = \frac{1}{n} \sum_{k=1}^n 1_{E_{i_k}}(x),$$

and let

$$S(x) = \liminf_n S_n(x),$$

so that, clearly, $\omega_p(\{x \in 2^{\mathbb{N}} : S(x) \neq p\}) = 0$. But

$$\int_X S_n(x) d\nu(x) = \frac{1}{n} \sum_{k=1}^n \nu(E_{i_k}) \leq 1/2,$$

and so by Fatou's Lemma,

$$\int_X S(x) d\nu(x) \leq \liminf_n \frac{1}{n} \sum_{k=1}^n \nu(E_{i_k}) \leq 1/2 < p.$$

Hence $\nu(\{x \in 2^{\mathbb{N}} : S(x) \neq p\}) > 0$, and ν is not absolutely continuous with respect to ω_p . □

Lemma 4.3. *Let $G \curvearrowright (X, \eta)$ be an ergodic probability measure preserving action, let $c : G \times X \rightarrow 2_{fin}^{\mathbb{N}}$ be a cocycle, and let $2_{fin}^{\mathbb{N}}$ act on $2^{\mathbb{N}}$ by symmetric difference. Consider the skew-product action $G \curvearrowright X \times 2^{\mathbb{N}}$ given by $g(x, y) = (gx, c(g, x)y)$.*

Denote by ω_p the Bernoulli p i.i.d. measure on $2^{\mathbb{N}}$. For any $1/2 < p < 1$, if there exists a G -invariant probability measure that is absolutely continuous with respect to $\eta \times \omega_p$, then c is cohomologous to a cocycle with a finite image.

Proof. Let $\lambda \ll \eta \times \omega_p$ be a G -invariant probability measure. Note that λ projects to a G -invariant probability measure on X which is absolutely continuous with respect to η . Because η is ergodic, this projection must equal η . Hence

$$d\lambda(x, y) = d\eta(x) d\lambda_x(y),$$

where $\lambda = \int \delta_x \times \lambda_x d\eta(x)$ is the disintegration of λ over η (here δ_x is the Dirac measure concentrated on $\{x\}$). Also

$$\frac{d\lambda}{d\eta \times \omega_p}(x, y) = \frac{d\lambda_x}{d\omega_p}(y).$$

It follows that $\lambda_x \ll \omega_p$, for a.e. x . By the invariance of λ we have that, for every $g \in G$, $g_*\lambda = \lambda$ which implies

$$\lambda_{gx} = (g_*\lambda)_{gx} = c(g, x)_*\lambda_x$$

for a.e. x .

Let $E_n = \{x \in 2^{\mathbb{N}} : x_n = 1\}$, and let $f(x) = \{n \in \mathbb{N} : \lambda_x(E_n) = 1/2\}$. By Lemma 4.2 $f(x)$ is finite for a.e. $x \in X$, since $\lambda_x \ll \omega_p$ for a.e. $x \in X$. Then

$$\begin{aligned} f(gx) &= \{n \in \mathbb{N} : \lambda_{gx}(E_n) = 1/2\} \\ &= \{n \in \mathbb{N} : c(g, x)_*\lambda_x(E_n) = 1/2\} \\ &= f(x), \end{aligned}$$

since for any measure ν on $2^{\mathbb{N}}$ and $T \in 2_{fin}^{\mathbb{N}}$, $T_*\nu(E_n) = 1 - \nu(E_n)$ if $n \in T$ and $T_*\nu(E_n) = \nu(E_n)$ otherwise; in any case, $T_*\nu(E_n) = 1/2$ iff $\nu(E_n) = 1/2$.

By the ergodicity of η it follows that there exists a $T \in 2_{fin}^{\mathbb{N}}$ such that $f(x) = T$ for a.e. $x \in X$. Let $m = \max(T)$, and denote $\tilde{c}(g, x) = c(g, x) \setminus \{1, \dots, m\}$.

Let $\theta(x) = \{n > m : \lambda_x(E_n) < 1/2\}$. Then

$$\begin{aligned} \theta(gx) &= \{n > m : \lambda_{gx}(E_n) < 1/2\} \\ &= \{n > m : c(g, x)_*\lambda_x(E_n) < 1/2\} \\ &= \tilde{c}(g, x) \triangle \{n > m : \lambda_x(E_n) < 1/2\} \\ &= \tilde{c}(g, x) \triangle \theta(x), \end{aligned}$$

where the third equality is a consequence of the fact that $\lambda_x(E_n) \neq 1/2$ for every $n > m$ and a.e. $x \in X$. Note that $\theta(x)$ is finite, by another application of Lemma 4.2. Hence $\tilde{c}$ is a coboundary, and so c is cohomologous to the cocycle $c \triangle \tilde{c}$ whose image is in 2^m . $\square$

5. PROOF OF THE MAIN THEOREM

Proof of Theorem 1. Let μ be an arbitrary generating measure on G . We prove the theorem by showing that G has property (T) if and only if (G, μ) has an entropy gap.

By Theorem A.1 we may assume G does not have property (T). Let $\rho : G \rightarrow \mathbb{N}$ be a proper function such that

$$\sum_{g \in G} \mu(g) \rho(g) \leq 2.$$

By Proposition 3.1 there exists an ergodic probability measure preserving action $G \curvearrowright (X, \eta)$ and a cocycle $c : G \times X \rightarrow 2_{fin}^{\mathbb{N}}$ that is not cohomologous to a cocycle with a finite image. Moreover,

$$\int_X \max(c(g, x)) \, d\eta(x) \leq \rho(g)$$

for any $g \in G$. By Proposition 2.1 for every $\epsilon > 0$ there exists a Bernoulli probability measure $\omega_{p(\epsilon)}$ on $2^{\mathbb{N}}$ such that for any probability measure ν on $2_{fin}^{\mathbb{N}}$,

$$h_\nu(2^{\mathbb{N}}, \omega_{p(\epsilon)}) \leq \epsilon \sum_{T \in 2_{fin}^{\mathbb{N}}} \nu(T) \max(T).$$

Consider the skew-product action $G \curvearrowright X \times 2^{\mathbb{N}}$ given by $g(x, T) = (gx, c(g, x)T)$. By Lemma 4.1,

$$\begin{aligned} h_{\mu}(X \times 2^{\mathbb{N}}, \eta \times \omega_{p(\epsilon)}) &= \int_X h_{\mu_x}(2^{\mathbb{N}}, \omega_{p(\epsilon)}) d\eta(x) \\ &\leq \epsilon \int_X \sum_{T \in 2_{fin}^{\mathbb{N}}} \mu_x(T) \max(T) d\eta(x). \end{aligned}$$

Note that, by Proposition 3.1, we know that

$$\begin{aligned} \int_X \sum_{T \in 2_{fin}^{\mathbb{N}}} \mu_x(T) \max(T) d\eta(x) &= \sum_{g \in G} \mu(g) \int_X \max(c(g, x)) d\eta(x) \\ &\leq \sum_{g \in G} \mu(g) \rho(g) \leq 2, \end{aligned}$$

and so

$$\lim_{\epsilon \rightarrow 0} h_{\mu}(X \times 2^{\mathbb{N}}, \eta \times \omega_{p(\epsilon)}) = 0.$$

Since c is not cohomologous to a cocycle with a finite image, Lemma 4.3 implies that each measure $\eta \times \omega_{p(\epsilon)}$ is properly nonsingular, and furthermore almost every measure in its ergodic decomposition is properly nonsingular.

Recall that the entropy of a convex combination of mutually singular measures is the same convex combination of their entropies. Therefore, since the entropy of $\eta \times \omega_{p(\epsilon)}$ is a convex combination of the entropies of its ergodic components, it follows that there exist ergodic, properly nonsingular G -actions with arbitrarily small entropy, and therefore (G, μ) does not have an entropy gap. $\square$

APPENDIX A. GROUPS WITH PROPERTY (T)

Theorem A.1 (Nevo). *Let G be a countable group with property (T), and let μ be a generating measure. Then (G, μ) has an entropy gap.*

The following proof is based on Nevo's [10].

Proof. Let $\bar{\mu}$ be the measure on G given by

$$\bar{\mu} = \sum_{n=0}^{\infty} 2^{-n-1} \mu^n,$$

where μ^n is the convolution of μ with itself n times. Then $h_{\bar{\mu}}(X, \eta) = h_{\mu}(X, \eta)$ (see, e.g., [8], or [6, Section 2.8]). The advantage of $\bar{\mu}$ is that it is supported everywhere on G .

Consider, given a nonsingular action $G \curvearrowright (X, \eta)$, the unitary representation π on $L^2(X, \eta)$ given by

$$\pi(g)f(x) = \sqrt{\frac{dg_*\eta}{d\eta}}(x) f(g^{-1}x).$$

Note that $\pi(g)^* = \pi(g^{-1})$.

It is easy to check that by Jensen's inequality, for any $g \in G$,

$$(1) \quad -2 \log \langle 1, \pi(g)1 \rangle = -2 \log \langle \pi(g^{-1})1, 1 \rangle \leq \int_X -\log \frac{dg_*^{-1}\eta}{d\eta}(x) d\eta(x).$$

Consider the Markov operator $\pi(\bar{\mu}): L^2(X, \eta) \rightarrow L^2(X, \eta)$ given by $\pi(\bar{\mu}) = \sum_{g \in G} \bar{\mu}(g) \pi(g)$. Then (1) yields the bound $-2 \log \|\pi(\bar{\mu})\| \leq h_{\bar{\mu}}(X, \eta)$, by another application of Jensen's inequality.

Denote

$$\|\bar{\mu}\|_T = \sup \{ \|\pi(\bar{\mu})\| : \pi \text{ is a unitary representation of } G \text{ with no invariant vectors} \}.$$

If G has property (T), then $\|\bar{\mu}\|_T < 1$ (see, e.g., Bekka, de La Harpe and Valette [1, Corollary 6.2.3]); here we use the fact that $\bar{\mu}$ is supported everywhere.

When $G \curvearrowright (X, \eta)$ is a properly nonsingular ergodic action, $L^2(X, \eta)$ has no invariant vectors (see [3, Lemma 7.2]). It therefore follows that

$$0 < -2 \log \|\bar{\mu}\|_T \leq -2 \log \|\pi(\bar{\mu})\| \leq h_{\bar{\mu}}(X, \eta) = h_{\mu}(X, \eta),$$

and (G, μ) has an entropy gap, with $\epsilon(\mu) = -2 \log \|\bar{\mu}\|_T$. $\square$

REFERENCES

- [1] B. Bekka, P. de la Harpe, and A. Valette, *Kazhdan's property (T)*, Cambridge University Press, 2008.
- [2] L. Bowen, *Random walks on random coset spaces with applications to Furstenberg entropy*, Invent. Math. **196** (2014), no. 2, 485–510, DOI 10.1007/s00222-013-0473-0. MR3193754
- [3] L. Bowen, Y. Hartman, and O. Tamuz, *Generic stationary measures and actions*, arXiv preprint arXiv:1405.2260, 2014.
- [4] A. Connes and B. Weiss, *Property T and asymptotically invariant sequences*, Israel J. Math. **37** (1980), no. 3, 209–210, DOI 10.1007/BF02760962. MR599455 (82e:28023b)
- [5] H. Furstenberg, *Noncommuting random products*, Trans. Amer. Math. Soc. **108** (1963), 377–428. MR0163345 (29 #648)
- [6] Y. Hartman and O. Tamuz, *Furstenberg entropy realizations for virtually free groups and lamplighter groups*, Journal d'Analyse Mathématique, to appear, 2015.
- [7] V. F. R. Jones and Klaus Schmidt, *Asymptotically invariant sequences and approximate finiteness*, Amer. J. Math. **109** (1987), no. 1, 91–114, DOI 10.2307/2374553. MR878200 (88h:28021)
- [8] V. A. Kaĭmanovich and A. M. Vershik, *Random walks on discrete groups: boundary and entropy*, Ann. Probab. **11** (1983), no. 3, 457–490. MR704539 (85d:60024)
- [9] D. A. Kaĭdan, *On the connection of the dual space of a group with the structure of its closed subgroups* (Russian), Funkcional. Anal. i Priložen. **1** (1967), 71–74. MR0209390 (35 #288)
- [10] A. Nevo, *The spectral theory of amenable actions and invariants of discrete groups*, Geom. Dedicata **100** (2003), 187–218, DOI 10.1023/A:1025839828396. MR2011122 (2004j:22012)
- [11] A. Nevo and R. J. Zimmer, *Rigidity of Furstenberg entropy for semisimple Lie group actions* (English, with English and French summaries), Ann. Sci. École Norm. Sup. (4) **33** (2000), no. 3, 321–343, DOI 10.1016/S0012-9593(00)00113-0. MR1775184 (2001k:22009)
- [12] N. Ozawa, *Noncommutative real algebraic geometry of Kazhdan's property (T)*, J. Inst. Math. Jussieu, FirstView:1–6, 11 2014.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS AT AUSTIN, AUSTIN, TEXAS 78712

DEPARTMENT OF MATHEMATICS, WEIZMANN INSTITUTE OF SCIENCE, 761001 REHOVOT, ISRAEL

MICROSOFT RESEARCH, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MASSACHUSETTS 02139