

A SUBHARMONICITY PROPERTY OF HARMONIC MEASURES

VILMOS TOTIK

(Communicated by Walter Van Assche)

ABSTRACT. Recently it has been established that for compact sets F lying on a circle S , the harmonic measure in the complement of F with respect to any point $a \in S \setminus F$ has convex density on any arc of F . In this note we give an alternative proof of this fact which is based on random walks, and which also yields an analogue in higher dimensions: for compact sets F lying on a sphere S in \mathbf{R}^n , the harmonic measure in the complement of F with respect to any point $a \in S \setminus F$ is subharmonic in the interior of F .

1. THE RESULT IN TWO DIMENSIONS

Let G be a domain $G \subset \mathbf{R}^n$ with compact boundary. In what follows, we denote the n -dimensional harmonic measure for a point $z \in G$ by $\omega(\cdot, z, G)$ (if it exists). So this is a measure on the boundary of G and it is the reproducing measure for harmonic functions in G : if u is harmonic in G (including at infinity if G is unbounded) and continuous on the closure \overline{G} , then

$$u(z) = \int_{\partial G} u(t) d\omega(t, z, G).$$

See [4], [7], [9] or [10] for the notion of harmonic measures.

The following convexity result was proved in [2].

Theorem 1. *Let S be a circle on the plane, and $F \subset S$ a closed subset of S . If $a \in S \setminus F$ and $I \subset F$ is an arc, then the density of the harmonic measure $\omega(\cdot, a, \mathbf{R}^2 \setminus F)$ with respect to arc-measure on S is convex on I .*

Actually, the stronger log-convexity was established (i.e. even the logarithm of the density is convex), and that is what we shall also prove below.

The theorem implies that if S is a circle and $F \subset S$ is a closed set with non-empty (one-dimensional) interior, then the equilibrium measure of F is convex on any subarc of the interior of F ; see [2, Theorem 1.5].

The circle S in these statements can also be a line and the $F \subset S$ a compact subset of that line (just apply the theorem to circles of radius R with $R \rightarrow \infty$). In particular, if $F \subset \mathbf{R}$ is a compact set with non-empty (one-dimensional) interior, then the density of the equilibrium measure of F with respect to Lebesgue-measure on \mathbf{R} is log-convex on any interval I that lies in F .


We also mention that Theorem 1 was extended to Riesz potentials in [3].

In [2] the proof of Theorem 1 was given by an iterated balayage technique. In this paper first we reprove Theorem 1 using the connection between harmonic measures

Received by the editors February 16, 2015 and, in revised form, June 1, 2015.

2010 *Mathematics Subject Classification.* Primary 31C12, 31A15, 60J45.

This work was supported by NSF DMS-1265375.

FIGURE 1. The domain G and its boundary

and random walks. This proof will allow us in the next section to prove an analogue in higher dimensions.

Proof of Theorem 1. Let S^1 be the unit circle and let D_- , resp. D_+ , be the inner, resp. exterior, domains complement to S^1 (i.e. D_- is the open unit disk and D_+ is the exterior of the closed unit disk). For a set $E \subset S^1$ let

$$E^* = \{t \in [0, 2\pi) \mid e^{it} \in E\}.$$

Let $F \subset S^1$ be a closed set and J a closed subarc in the (one-dimensional) interior of F . For some small $\varepsilon > 0$ set

$$G = \{z \mid \text{dist}(z, S^1 \setminus F) < \min\{\frac{1}{2}\text{dist}(z, F), \varepsilon\}\}.$$

Then G is a small neighborhood of $S^1 \setminus F$; see Figure 1. Let Γ be the boundary of G and $\Gamma_{\pm} = \Gamma \cap D_{\pm}$ the part of that boundary that lies in the unit disk and in its exterior, respectively.

Let $a \in S^1 \setminus F$ be fixed. What we want to show is that the harmonic measure $\omega(\cdot, a, \mathbf{R}^2 \setminus F)$ has log-convex density on J . We can formulate the claim as there is a convex function $\sigma_a(t)$ (convexity in the variable t) on J^* such for any Borel set $E \subset J$ we have

$$\omega(E, a, \mathbf{R}^2 \setminus F) = \int_{E^*} \sigma_a(t) dt.$$

By simple approximation from the outside we may assume that F consists of finitely many arcs on S^1 .

Start a two-dimensional Brownian motion $X(t)$, $t \geq 0$, at a : $X(0) = a$. We are going to use Kakutani's theorem that for $E \subset J$ the harmonic measure $\omega(E, a, \mathbf{R}^2 \setminus F)$ is the probability that X leaves the domain $\mathbf{R}^2 \setminus F$ first at a point of E (see [4, Theorem F6, (F.10)], [6], or [9, Section 3.4]). Let

$$T_X = \min\{t \mid X(t) \in F\}.$$

Note that, by the recurrence of the two-dimensional Brownian motion, we have $T_X < \infty$ almost surely (because we assumed that F contains an arc), although we are not going to use that. Let

$$T_X^0 = \sup\{t < T_X \mid X(t) \in S^1 \setminus F\}$$

and

$$T_X^1 = \min\{t \mid t \geq T_X^0, X(t) \in \Gamma\}.$$

Again, $T_X^1 < T_X < \infty$ almost surely, and $X(T_X^1) \in \Gamma$. Suppose that, say, $X(T_X^1) \in \Gamma_-$. Then $\{X(t) \mid t \geq T_X^1\}$ is a Brownian motion starting at $X(T_X^1) \in D_-$ that leaves the domain D_- in a point of F , so the probability that it leaves D_- at a point of a given Borel set $E \subset J$ is (conditional probability)

$$\frac{\omega(E, X(T_X^1), D_-)}{\omega(F, X(T_X^1), D_-)}.$$

Note that here the denominator is positive since F contains an arc of the unit circle. In a similar fashion, if $X(T_X^1) \in \Gamma_+$, then the probability that the Brownian motion $\{X(t) \mid t \geq T_X^1\}$ starting at $X(T_X^1) \in D_+$ leaves the domain D_+ at a point of $E \subset J$ is

$$\frac{\omega(E, X(T_X^1), D_+)}{\omega(F, X(T_X^1), D_+)}.$$

Now

$$\mu_{a,\pm}(A) = \mathbf{P}(X(T_X^1) \in A), \quad A \subset \Gamma_{\pm}, \quad A \text{ Borel},$$

are two positive Borel measures on Γ_{\pm} , respectively, and, according to what we have just explained, we have the formula

$$(1) \quad \omega(E, a, \mathbf{R}^2 \setminus F) = \int_{\Gamma_-} \frac{\omega(E, \zeta, D_-)}{\omega(F, \zeta, D_-)} d\mu_{a,-}(\zeta) + \int_{\Gamma_+} \frac{\omega(E, \zeta, D_+)}{\omega(F, \zeta, D_+)} d\mu_{a,+}(\zeta).$$

But here $\omega(E, \zeta, D_-)$ is given (see [4, Sec. 1.1] or [9, Theorem 3.44]) by the Poisson integral

$$\omega(E, \zeta, D_-) = \int_{E^*} P_{\zeta}(e^{it}) dt$$

with the Poisson kernel

$$P_{\zeta}(e^{it}) = \frac{1}{2\pi} \frac{1 - |\zeta|^2}{|\zeta - e^{it}|^2}, \quad \zeta \in D_-,$$

and similarly

$$\omega(E, \zeta, D_+) = \int_{E^*} P_{\zeta}(e^{it}) dt$$

with the exterior Poisson kernel

$$P_{\zeta}(e^{it}) = \frac{1}{2\pi} \frac{|\zeta|^2 - 1}{|\zeta - e^{it}|^2}, \quad \zeta \in D_+.$$

Thus,

$$\omega(E, a, \mathbf{R}^2 \setminus F) = \int_{E^*} \left(\int_{\Gamma_-} \frac{P_{\zeta}(e^{it})}{\omega(F, \zeta, D_-)} d\mu_{a,-}(\zeta) + \int_{\Gamma_+} \frac{P_{\zeta}(e^{it})}{\omega(F, \zeta, D_+)} d\mu_{a,+}(\zeta) \right) dt,$$

i.e. the density σ_a in question is

$$(2) \quad \sigma_a(t) = \int_{\Gamma_-} \frac{P_{\zeta}(e^{it})}{\omega(F, \zeta, D_-)} d\mu_{a,-}(\zeta) + \int_{\Gamma_+} \frac{P_{\zeta}(e^{it})}{\omega(F, \zeta, D_+)} d\mu_{a,+}(\zeta).$$

Now it is easy to see that the sum, and hence the integral, of log-convex functions is again log-convex (see [2]), so it is sufficient to show that if ε (appearing in the definition of Γ) is sufficiently small, then for all $\zeta \in \Gamma$ the function $P_\zeta(e^{it})$ is log-convex on J^* . But that is simple: the function

$$\frac{1}{1 - 2 \cos v + 1} = \frac{1}{4 \sin^2(v/2)}$$

is strictly log-convex on the open interval $(0, 2\pi)$, hence

$$\frac{1}{1 - 2 \cos(\theta - t) + 1}$$

is strictly log-convex on any interval not containing $\theta \pmod{2\pi}$. This and simple compactness implies that if $\theta \in (S^1 \setminus F)^*$, then for r sufficiently close to 1 the functions (in t)

$$\frac{1}{1 - 2r \cos(\theta - t) + r^2}$$

are log-convex for $t \in J^*$. But if $\zeta = re^{i\theta} \in \Gamma$, then $\theta \in (S^1 \setminus F)^*$ and r is close to 1 ($1 - \varepsilon \leq r \leq 1$), furthermore

$$P_\zeta(e^{it}) = \frac{1}{2\pi} \frac{|1 - r^2|}{1 - 2r \cos(\theta - t) + r^2},$$

so the log-convexity of $P_\zeta(e^{it})$ on J^* for all $\zeta \in \Gamma$ follows. \square

2. HARMONIC MEASURES IN HIGHER DIMENSIONS

In this section, we prove the following higher-dimensional analogue of Theorem 1.

Theorem 2. *Let S be an $(n - 1)$ -dimensional sphere in \mathbf{R}^n , $n \geq 3$, and $F \subset S$ a closed set. If $a \in S \setminus F$, then the density of the harmonic measure $\omega(\cdot, a, \mathbf{R}^n \setminus F)$ is subharmonic on the $((n - 1)$ -dimensional) interior of F .*

An explanation is needed for the statement. We may assume S to be S^{n-1} , the unit sphere in \mathbf{R}^n . Let $\text{Int}(F)$ be the $(n - 1)$ -dimensional interior of F . On S we consider the geodesic distance d and geodesic spheres $S_\rho(P) = \{Q \in S \mid d(Q, P) = \rho\}$ (of dimension $n - 2$) and geodesic balls $B_\rho(P) = \{Q \in S \mid d(Q, P) \leq \rho\}$ (of dimension $n - 1$). Let also Δ be the Laplace-Beltrami operator on S (see e.g. [1, Section 3.1]), which is the restriction to S^{n-1} of the angular part of the Laplacian on \mathbf{R}^n . If dS denotes the surface element on S , then, in the interior of F , the harmonic measure has the form (see the proof below)

$$d\omega(E, a, \mathbf{R}^n \setminus F) = \int_E \sigma_a(\mathbf{x}) dS(\mathbf{x}),$$

where $\sigma_a(\mathbf{x})$ is the density function in the theorem. Now the subharmonicity of σ_a means either of the following:

- (i) σ_a has the submean-value property on every geodesic sphere $S_\rho(P)$ lying in $\text{Int}(F)$ together with its interior (i.e. the value $\sigma_a(P)$ is at most as large as the average of σ_a over $S_\rho(P)$),
- (ii) σ_a has the submean-value property on every geodesic ball $B_\rho(P)$ lying in $\text{Int}(F)$ (i.e. the value $\sigma_a(P)$ is at most as large as the average of σ_a over $B_\rho(P)$),
- (iii) $\Delta \sigma_a(\mathbf{x}) \geq 0$ for all $\mathbf{x} \in \text{Int}(F)$.

Naturally, the averages in question are taken with respect to the corresponding surface or volume elements on $S_\rho(P)$ or $B_\rho(P)$, respectively.

We are going to prove Theorem 2 in the form (iii). From there (ii) and (i) follow from the spherical version of Green's formulae; see e.g. [8, Proposition 2.1, (2.1), (2.2)]. An alternative approach is to apply that Δ -harmonic functions have the mean value property over geodesic balls (see e.g. [5, Corollary X.7.3] as well as the Remark there, or see [11]) and hence also over geodesic spheres, and then showing that (iii) implies that if σ_a agrees with a Δ -harmonic function on a geodesic sphere, then it is below that function inside that sphere (otherwise, if we add to σ_a a small multiple of an appropriate function with strictly positive spherical Laplacian, at a local maximum point (iii) would be violated).

Consider now the equilibrium measure μ_F in the Newtonian case (the kernel is $|z|^{2-n}$) of a non-polar compact set $F \subset \mathbf{R}^n$ in \mathbf{R}^n (see e.g. [9, Sec. 4.3] where it is called harmonic measure from infinity, or [7, Section II.1], where a different normalization is used). Let us record the following

Corollary 3. *If F is a compact subset of a sphere or of \mathbf{R}^{n-1} , then the equilibrium measure (with respect to the Newtonian kernel in \mathbf{R}^n) μ_F of F has subharmonic density on the $((n-1)$ -dimensional) interior of F .*

The density in question is taken with respect to surface measure on the sphere or on \mathbf{R}^{n-1} .

Indeed, the equilibrium measure μ_S of the whole sphere S is the uniform distribution on S , and from it the equilibrium measure μ_F of F is obtained by taking the balayage (see [7, Sect. IV.1]) of $\mu_S|_{S \setminus F}$ onto F (followed by a normalization if one uses mass 1 for the equilibrium measure), i.e. $\mu_F = \nu/\nu(F)$, where

$$\nu(\cdot) = \mu_S|_F(\cdot) + \int_{S \setminus F} \omega(\cdot, a, \mathbf{R}^n \setminus F) d\mu_S(a).$$

In view of this formula, Corollary 3 follows from Theorem 2.

Let us also mention that, just as in [2, Corollary 1.6], Theorem 2 implies the subharmonicity of the harmonic measures $\omega(\cdot, a, \mathbf{R}^n \setminus F)$ for a lying in a relatively large n -dimensional domain that contains $S \setminus F$, but the exact description of that domain is not clear and we do not pursue this direction.

Proof of Theorem 2. As before, we may assume that $S = S^{n-1}$ is the unit sphere in \mathbf{R}^n , and that the $(n-1)$ -dimensional interior of F is not empty (otherwise there is nothing to prove). Also, by approximation from the outside, we may assume (just to avoid irregular sets for the Dirichlet problem) that F is the union of finitely many closed balls.

Again, it is sufficient to show that if the point $(0, \dots, 0, 1)$ belongs to the interior of F , then the density σ_a of $\omega(\cdot, a, \mathbf{R}^n \setminus F)$ satisfies

$$\Delta\sigma_a(0, \dots, 0, 1) > 0.$$

If, in a small neighborhood of $(0, \dots, 0, 1)$, we use the Cartesian coordinates (x_1, \dots, x_{n-1}) as local coordinates for S^{n-1} , then (see e.g. [1, (3.18)] or use [1, (3.4)])

$$\Delta\sigma_a(0, \dots, 0, 1) = \sum_{i=1}^{n-1} \frac{\partial^2 \sigma_a}{\partial x_i^2}(0, \dots, 0, 1),$$

so we need to prove that the right-hand side is positive.

We follow the argument in the preceding proof. Let X be an n -dimensional Brownian motion in \mathbf{R}^n starting at $a \in S \setminus F$. As before, $\omega(E, a, \mathbf{R}^n \setminus F)$ is the probability that X leaves the domain $\mathbf{R}^n \setminus F$ in a point of E (see [9, Section 3.4]).

Let now D_{\pm} denote the interior/exterior domains to S^{n-1} , so D_- is the open unit ball in \mathbf{R}^n and D_+ is the (open) exterior of that ball. Define G and its boundary Γ appropriately as in the preceding proof, and introduce the stopping times T_X, T_X^0, T_X^1 as above. It is no longer true that $T_X < \infty$ almost surely, but we may restrict our attention only to Brownian motions hitting F , in which case T_X^0, T_X^1 are finite (under the just set condition $T_X < \infty$). From our point of view this restriction is irrelevant, since that means that instead of the probability that X exits $\mathbf{R}^n \setminus F$ in a point of $E \subset F$ we consider the same probability under the condition $T_X < \infty$ (which has probability $\omega(F, a, \mathbf{R}^n \setminus F)$), so the two densities differ only in the multiplicative constant $\omega(F, a, \mathbf{R}^n \setminus F)$. We have again formula (1) in the form

$$\frac{\omega(E, a, \mathbf{R}^2 \setminus F)}{\omega(F, a, \mathbf{R}^2 \setminus F)} = \int_{\Gamma_-} \frac{\omega(E, \zeta, D_-)}{\omega(F, \zeta, D_-)} d\mu_{a,-}(\zeta) + \int_{\Gamma_+} \frac{\omega(E, \zeta, D_+)}{\omega(F, \zeta, D_+)} d\mu_{a,+}(\zeta),$$

and here the harmonic measures $\omega(E, \zeta, D_{\pm})$ are given by integrals on E (against $(n-1)$ -dimensional surface elements of S^{n-1}) of the n -dimensional Poisson kernels (see [9, Theorem 3.44])

$$P_{\zeta}(\mathbf{x}) = \tau_n \frac{1 - |\zeta|^2}{|\zeta - \mathbf{x}|^n}, \quad |\zeta| < 1,$$

and

$$P_{\zeta}(\mathbf{x}) = \tau_n \frac{|\zeta|^2 - 1}{|\zeta - \mathbf{x}|^n}, \quad |\zeta| > 1,$$

where τ_n is a normalizing constant, and where $|\cdot|$ denotes the Euclidean norm in \mathbf{R}^n . Therefore, we get again (2) in the form

$$\frac{\sigma_a(\mathbf{x})}{\omega(F, a, \mathbf{R}^2 \setminus F)} = \int_{\Gamma_-} \frac{P_{\zeta}(\mathbf{x})}{\omega(F, \zeta, D_-)} d\mu_{a,-}(\zeta) + \int_{\Gamma_+} \frac{P_{\zeta}(\mathbf{x})}{\omega(F, \zeta, D_+)} d\mu_{a,+}(\zeta)$$

for $\mathbf{x} \in \text{Int}(F)$. With the same argument as in the preceding proof we can see that the claim in the theorem reduces to the fact that for all $\zeta \in \Gamma$ (if the ε in the definition of Γ is sufficiently small) the function $P_{\zeta}(\mathbf{x})$ has positive spherical Laplacian at $(0, \dots, 0, 1)$: $\Delta P_{\zeta}(0, \dots, 0, 1) > 0$. Again, by compactness, it is enough to prove that if $\zeta = (\zeta_1, \dots, \zeta_n) \neq (0, \dots, 0, 1)$ belongs to S^{n-1} and $h(x_1, \dots, x_{n-1})$ is defined as

$$\frac{1}{\left((\zeta_1 - x_1)^2 + (\zeta_2 - x_2)^2 + \dots + (\zeta_{n-1} - x_{n-1})^2 + (\zeta_n - \sqrt{1 - x_1^2 - \dots - x_{n-1}^2})^2 \right)^{n/2}},$$

then

$$(3) \quad \sum_{i=1}^{n-1} \frac{\partial^2 h}{\partial x_i^2}(\mathbf{0}) > 0.$$

Simple calculation shows that

$$\frac{\partial^2 h}{\partial x_i^2}(\mathbf{0}) = \frac{n}{2^{\frac{n}{2}+1}} \cdot \frac{-\zeta_n + \zeta_n^2 + (\frac{n}{2} + 1)\zeta_i^2}{(1 - \zeta_n)^{\frac{n}{2}+2}},$$

so in view of

$$\sum_{i=1}^{n-1} \zeta_i^2 = 1 - \zeta_n^2,$$

(3) reduces to showing that

$$\frac{(\frac{n}{2} + 1) - (n - 1)\zeta_n + (\frac{n}{2} - 2)\zeta_n^2}{(1 - \zeta_n)^{\frac{n}{2} + 2}}$$

is positive for $\zeta_n \in [-1, 1]$, $\zeta_n \neq 1$, which can be readily seen. \square

It is clear from the just given proof that whatever properties one establishes for the “limit” Poisson kernel $1/|\zeta - \mathbf{x}|^n$ at $\mathbf{x} = (0, \dots, 1) \neq \zeta$ on S^{n-1} , the same will be true for the density of harmonic measures $\omega(\cdot, a, \mathbf{R}^n \setminus F)$, $a \in S \setminus F$, in the interior of F , provided the property is preserved under summation and taking a limit.

The author is grateful to B. Simon for the reference [8]. In connection with Theorem 2 the author would like to mention that in 2010 D. Benko announced a subharmonicity property of balayage measures for Riesz kernels, which may be related to that theorem.

REFERENCES

- [1] Kendall Atkinson and Weimin Han, *Spherical harmonics and approximations on the unit sphere: an introduction*, Lecture Notes in Mathematics, vol. 2044, Springer, Heidelberg, 2012. MR2934227
- [2] David Benko, Peter Dragnev, and Vilmos Totik, *Convexity of harmonic densities*, Rev. Mat. Iberoam. **28** (2012), no. 4, 947–960, DOI 10.4171/RMI/698. MR2990129
- [3] David Benko and Peter D. Dragnev, *Balayage ping-pong: a convexity of equilibrium measures*, Constr. Approx. **36** (2012), no. 2, 191–214, DOI 10.1007/s00365-011-9143-x. MR2957308
- [4] John B. Garnett and Donald E. Marshall, *Harmonic measure*, New Mathematical Monographs, vol. 2, Cambridge University Press, Cambridge, 2008. Reprint of the 2005 original. MR2450237 (2009k:31001)
- [5] Sigurdur Helgason, *Differential geometry and symmetric spaces*, Pure and Applied Mathematics, Vol. XII, Academic Press, New York-London, 1962. MR0145455 (26 #2986)
- [6] Shizuo Kakutani, *Two-dimensional Brownian motion and harmonic functions*, Proc. Imp. Acad. Tokyo **20** (1944), 706–714. MR0014647 (7,315b)
- [7] N. S. Landkof, *Foundations of modern potential theory*, Springer-Verlag, New York-Heidelberg, 1972. Translated from the Russian by A. P. Doohovskoy; Die Grundlehren der mathematischen Wissenschaften, Band 180. MR0350027 (50 #2520)
- [8] Lei Ni, *Mean value theorems on manifolds*, Asian J. Math. **11** (2007), no. 2, 277–304, DOI 10.4310/AJM.2007.v11.n2.a6. MR2328895 (2008h:58046)
- [9] Peter Mörters and Yuval Peres, *Brownian motion*, Cambridge Series in Statistical and Probabilistic Mathematics, Cambridge University Press, Cambridge, 2010. With an appendix by Oded Schramm and Wendelin Werner. MR2604525 (2011i:60152)
- [10] Thomas Ransford, *Potential theory in the complex plane*, London Mathematical Society Student Texts, vol. 28, Cambridge University Press, Cambridge, 1995. MR1334766 (96e:31001)
- [11] T. J. Willmore, *Mean value theorems in harmonic Riemannian spaces*, J. London Math. Soc. **25** (1950), 54–57. MR0033408 (11,436f)

MTA-SZTE ANALYSIS AND STOCHASTICS RESEARCH GROUP, BOLYAI INSTITUTE, UNIVERSITY OF SZEGED, SZEGED, ARADI V. TERE 1, 6720, HUNGARY – AND – DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SOUTH FLORIDA, 4202 E. FOWLER AVE, CMC342, TAMPA, FLORIDA 33620-5700

E-mail address: totik@mail.usf.edu