

FORCING WITH MATRICES OF COUNTABLE ELEMENTARY SUBMODELS

BORISA KUZELJEVIC AND STEVO TODORCEVIC

(Communicated by Mirna Džamonja)

ABSTRACT. We analyze the forcing notion \mathcal{P} of finite matrices whose rows consist of isomorphic countable elementary submodels of a given structure of the form H_θ . We show that forcing with this poset adds a Kurepa tree T . Moreover, if \mathcal{P}_c is a suborder of \mathcal{P} containing only continuous matrices, then the Kurepa tree T is almost Souslin, i.e., the level set of any antichain in T is not stationary in ω_1 .

1. INTRODUCTION

In this paper we analyze the forcing notion mentioned in the remark on page 217 of [10]. This is the forcing notion of finite matrices whose rows consist of isomorphic countable elementary submodels of a given structure of the form H_θ . In [10] they were merely meant as side conditions to various proper forcing constructions when one is interested in getting the \aleph_2 -chain condition that can be iterated. Soon afterwards the second author realized that this variation of the original side conditions is as much an interesting forcing notion as the poset of finite chains of countable elementary submodels analyzed briefly in Theorem 6 of [10]. For example, he showed that the poset of finite matrices of row-isomorphic countable elementary submodels always forces CH (so, in particular preserves CH if it is true in the ground model). The second author observed at the same time that this forcing notion gives a natural example of a Kurepa tree. Here we shall explore this further and produce a natural variation of this forcing notion that gives us a Kurepa tree with no stationary antichains. This gives us a quite different forcing construction of such a tree from the previous ones which use countable rather than finite conditions (see [3] and [11]). We believe that there will be other natural variations of this forcing notion with interesting applications. For example, we note that in the recent paper [1] Asperó and Mota have used the poset of finite matrices of elementary submodels to control their iteration scheme which shows that the forcing axiom for the class of all finitely proper posets of size ω_1 is compatible with $2^{\aleph_0} > \aleph_2$. In view of the recent efforts to generalize the side condition method to higher cardinals (see [6, 7]) it would be interesting to also explore the possible higher-cardinal analogues of the

Received by the editors March 28, 2015 and, in revised form, June 19, 2015 and September 2, 2015.

2010 *Mathematics Subject Classification*. Primary 03E57.

Key words and phrases. Proper forcing, side condition, countable elementary submodel.

This paper was conceived during the Summer of 2014 when the first author was visiting Institut de Mathématiques de Jussieu. The first author would like to thank the Equipe de Logique of that Institute for support which made this visit possible. The first author was partially supported by the MPNTR grant ON174006.

posets that we analyze here. This could also be asked for the original side-condition poset of finite elementary chains of countable elementary submodels of $[10]$ which, as shown in Theorem 6 of $[10]$, gives us a natural forcing notion that collapses a given cardinal θ to ω_1 preserving all other cardinals.¹ As far as we know, no higher-cardinal analogue of this poset has been produced. Some research related to this have been recently produced by Aspero $[2]$.

2. PRELIMINARIES

Definition 2.1. Let $\theta \geq \omega_2$ be a regular cardinal. By H_θ we denote the collection of all sets whose transitive closure has cardinality $< \theta$. We consider it as a model of the form $\langle H_\theta, \in, <_\theta \rangle$ where $<_\theta$ is some fixed well-ordering of H_θ that will not be explicitly mentioned. The partial order \mathcal{P} is the set of all functions $p : \omega_1 \rightarrow H_\theta$ satisfying:

- (1) $\text{supp}(p) = \{\alpha < \omega_1 : p(\alpha) \neq \emptyset\}$ is a finite set;
- (2) $p(\alpha)$ is a finite collection of isomorphic countable elementary submodels of H_θ for every $\alpha \in \text{supp}(p)$;
- (3) for each $\alpha, \beta \in \text{supp}(p)$ if $\alpha < \beta$, then $\forall M \in p(\alpha) \exists N \in p(\beta) M \in N$.

The ordering on \mathcal{P} is given by:

$$(2.1) \quad p \leq q \Leftrightarrow \forall \alpha < \omega_1 \quad q(\alpha) \subseteq p(\alpha).$$

The fact that M is a countable elementary submodel of $\langle H_\theta, \in \rangle$ will be denoted by $M \prec H_\theta$. Also, if $M \prec H_\theta$, then $\overline{M} \in H_{\omega_1}$ denotes the transitive collapse of M with π_M being the corresponding isomorphism. For $p \in \mathcal{P}$ and $\alpha \in \text{supp}(p)$ we denote $\delta_\alpha^p = M \cap \omega_1$ where M is some (any) model in $p(\alpha)$. Also, if $M \prec H_\theta$, then δ_M will denote the ordinal $M \cap \omega_1$. We list some standard lemmas concerning countable elementary submodels of H_θ that will be useful throughout the paper.

Lemma 2.2. *Let F be a countable subset of H_θ . Then the set of all ordinals of the form $M \cap \omega_1$ such that M is a countable elementary submodel of H_θ with $F \subseteq M$ contains a club.*

Lemma 2.3. *If $M \prec H_\theta$ contains some element X , then X is countable if and only if $X \subseteq M$.*

Lemma 2.4. *If $M \prec H_\theta$ contains as an element some subset A of ω_1 , then A is uncountable if and only if $A \cap M \cap \omega_1$ is unbounded in $M \cap \omega_1$.*

Lemma 2.5. *If $M \prec H_\theta$, X is in H_θ and X is definable from parameters in M , then $X \in M$.*

Lemma 2.6. *Let $\langle N_\xi : \xi < \omega_1 \rangle$ be a continuous \in -chain of countable elementary submodels of H_θ . Then $\{\xi < \omega_1 : N_\xi \cap \omega_1 = \xi\}$ is a club in ω_1 .*

Lemma 2.7. *Let M_0 and M_1 be isomorphic countable elementary submodels of some H_θ . Let $L_0 = M_0 \cap \omega_2$ and $L_1 = M_1 \cap \omega_2$. Then $L_0 \cap L_1$ is an initial segment of both L_0 and L_1 .*

¹It should be noted that the hypothesis of Theorem 6 of $[10]$ assumes that there is a stationary subset S of $[\theta]^{\aleph_0}$ or cardinality θ , a condition that is satisfied by many cardinals and in particular by all cardinals of uncountable cofinality if 0^\sharp does not exist.

Proof. Without loss of generality, we can assume that both M_0 and M_1 contain the same family of mappings $\langle e_\gamma : \gamma < \omega_2 \rangle$ where each map $e_\gamma : \gamma \rightarrow \omega_1$ is 1-1. Now we will prove that if β is in $L_0 \cap L_1$ and $\alpha < \beta$ is in L_0 , then $\alpha \in L_1$. Consider the map $e_\beta : \beta \rightarrow \omega_1$. Then, in M_0 there is some $\xi < \omega_1$ such that $e_\beta(\alpha) = \xi$, i.e., $\alpha = e_\beta^{-1}(\xi)$. But M_1 knows both e_β and ξ (this is because $M_0 \cong M_1 \Rightarrow M_0 \cap \omega_1 = M_1 \cap \omega_1$). Hence, $\alpha \in M_1 \cap \omega_2 = L_1$. \square

If $\mathcal{G} \subseteq \mathcal{P}$ is a filter in \mathcal{P} generic over V , then we define $G : \omega_1 \rightarrow H_\theta$ as the function satisfying

$$G(\alpha) = \{M \prec H_\theta : \exists p \in \mathcal{G} \text{ such that } M \in p(\alpha)\}.$$

Note that G is a well defined function because \mathcal{G} is a filter. For α in the domain of G we denote $\delta_\alpha = M \cap \omega_1$ for some (any) M in $G(\alpha)$. Further, we denote $A_\gamma = \bigcup_{M \in G(\gamma)} M \cap \omega_2$ for $\gamma < \omega_1$, and note that if $\gamma < \delta$, then $A_\gamma \subseteq A_\delta$. Also, we define the function $g : \omega_1 \rightarrow H_{\omega_1}$ with $g(\alpha) = \overline{M}$ for some (any) model M from $G(\alpha)$ and for $p \in \mathcal{P}$, by \bar{p} we define $\bar{p} : \omega_1 \rightarrow [H_{\omega_1}]^\omega$ as a function with the same support as p which maps $\alpha \in \text{supp}(\bar{p})$ to the transitive collapse of some model from $p(\alpha)$, while for $\alpha \in \omega_1 \setminus \text{supp}(\bar{p})$ take $\bar{p}(\alpha) = \emptyset$.

Lemma 2.8. *Let $p, q \in \mathcal{P}$. If $\bar{p} = \bar{q}$, then p and q are compatible conditions.*

Proof. First note that if $\bar{p} = \bar{q}$, then $\text{supp}(p) = \text{supp}(q)$. Also, if two countable elementary submodels of H_θ , say M_1 and M_2 have the same transitive collapse, then they are isomorphic (the isomorphism is simply $\pi_{M_2}^{-1} \circ \pi_{M_1}$). Now it is clear that the function $r : \omega_1 \rightarrow H_\theta$ defined by $r(\alpha) = p(\alpha) \cup q(\alpha)$ is in \mathcal{P} and that it satisfies $r \leq p, q$. \square

For $p, q \in \mathcal{P}$ we will define their ‘join’ $p \vee q$ as the function from ω_1 to H_θ satisfying $(p \vee q)(\alpha) = p(\alpha) \cup q(\alpha)$ for $\alpha < \omega_1$. Further, if a condition $q \in \mathcal{P}$ and $M \prec H_\kappa$ ($\delta_M = M \cap \omega_1$) for $\kappa \geq \theta$ are given, it is clear what intersection $q \cap M$ represents, and we define the restriction of q to M as a function with finite support $q \upharpoonright M : \omega_1 \rightarrow H_\theta$ satisfying $\text{supp}(q \upharpoonright M) = \text{supp}(q) \cap \delta_M$ and for $\alpha < \delta_M$:

$$(q \upharpoonright M)(\alpha) = \left\{ \varphi_{M'}(N) : M' \in q(\delta_M), \varphi_{M'} : M' \xrightarrow{\cong} M \cap H_\theta, N \in q(\alpha) \cap M' \right\}.$$

Note that the function $q \upharpoonright M$ is in \mathcal{P} . We will also need the following notion which we call ‘the closure of p below δ ’.

Definition 2.9. Let $p \in \mathcal{P}$ and $\delta \in \text{supp}(p)$. Then $\text{cl}_\delta(p) : \omega_1 \rightarrow H_\theta$ is a function such that $\text{supp}(\text{cl}_\delta(p)) = \text{supp}(p)$ and $\text{cl}_\delta(p)(\gamma) = p(\gamma)$ for $\gamma \geq \delta$, while for $\gamma < \delta$ we have

$$\text{cl}_\delta(p)(\gamma) = \left\{ \psi_{N_1, N_2}(M) : M \in p(\gamma) \cap N_1 \ \& \ N_1, N_2 \in p(\delta) \ \& \ \psi_{N_1, N_2} : N_1 \xrightarrow{\cong} N_2 \right\}.$$

We will also need the following standard lemmas later in the paper.

Lemma 2.10. *Suppose that $\theta \geq \omega_2$ is a regular cardinal. If $\mathcal{P} \in H_\theta$, then in $V[\mathcal{G}]$ we have $H_\theta[\mathcal{G}] = \{\text{int}_{\mathcal{G}}(\tau) : \tau \in H_\theta \text{ is a } \mathcal{P}\text{-name}\} = H_\theta^{V[\mathcal{G}]}$.*

Proof. $H_\theta^{V[\mathcal{G}]} \subseteq H_\theta[\mathcal{G}]$ follows from the fact that any $x \in H_\theta^{V[\mathcal{G}]}$ is of the form $\text{int}_{\mathcal{G}}(\tau)$ for some $\tau \in H_\theta$ (see [9, Claim I 5.17]). If $\tau \in H_\theta$ is a \mathcal{P} -name, then from $|\text{trcl}(\tau)| < \theta$ and $\text{rank}(\text{int}_{\mathcal{G}}(\tau)) \leq \text{rank}(\tau)$ we have that $|\text{trcl}(\text{int}_{\mathcal{G}}(\tau))| < \theta$. \square

Lemma 2.11. *If $M \prec H_\theta$ and $\mathcal{P} \in M$, then in $V[\mathcal{G}]$ we have that*

$$M[\mathcal{G}] = \{\text{int}_{\mathcal{G}}(\tau) : \tau \in M \text{ is a } \mathcal{P}\text{-name}\} \prec H_\theta[\mathcal{G}] = H_\theta.$$

Proof. First note that $M[\mathcal{G}] \subseteq H_\theta[\mathcal{G}]$. Now take any $\tau_1, \dots, \tau_n \in M$ and any formula $\varphi(x, x_1, \dots, x_n)$. Assume that there is some $y \in H_\theta[\mathcal{G}]$ such that $H_\theta[\mathcal{G}] \models \varphi(y, \text{int}_{\mathcal{G}}(\tau_1), \dots, \text{int}_{\mathcal{G}}(\tau_n))$. Then there is $p \in \mathcal{G}$ which forces $\varphi(\tau_y, \tau_1, \dots, \tau_n)$ for a \mathcal{P} -name $\tau_y \in H_\theta$ and $M \prec H_\theta$ implies $p \Vdash \varphi(\tau, \tau_1, \dots, \tau_n)$ for a \mathcal{P} -name $\tau \in M$. Hence $H_\theta[\mathcal{G}] \models \varphi(\text{int}_{\mathcal{G}}(\tau), \text{int}_{\mathcal{G}}(\tau_1), \dots, \text{int}_{\mathcal{G}}(\tau_n))$ for a \mathcal{P} -name $\tau \in M$, so $M[\mathcal{G}] \prec H_\theta[\mathcal{G}]$. \square

3. PROPERNESS

In this section we show that \mathcal{P} satisfies the condition stronger than being proper, namely \mathcal{P} is strongly proper forcing. We are not sure who was the first to define this notion, but Mitchell’s paper [5] is a good reference.

Definition 3.1. If P is a forcing notion and X is a set, then we say that p is strongly (X, P) -generic if for any set D which is dense and open in the poset $P \cap X$, the set D is predense in P below p . The poset P is strongly proper if for every large enough regular cardinal κ , there are club many countable elementary submodels M of H_κ such that whenever $p \in M \cap P$, there exists a strongly (M, P) -generic condition below p .

Lemma 3.2. *\mathcal{P} is strongly proper.*

Proof. Let p be a condition in \mathcal{P} and $M \prec H_\kappa$ (for some $\kappa \geq \theta$ which is large enough) such that $p, \mathcal{P} \in M$. Denote $\delta = M \cap \omega_1$. We will show that the condition $p_M = p \cup \{\langle \delta, M \cap H_\theta \rangle\}$ is strongly generic (i.e. if $q \leq p_M$ and $\mathcal{D} \subseteq \mathcal{P} \cap M$ is dense open, then there is some $q' \in \mathcal{D}$ such that $q \not\leq q'$). Let $q \leq p_M$ be arbitrary and $\mathcal{D} \subseteq M \cap \mathcal{P}$ dense open. Consider the condition $q \upharpoonright M \in \mathcal{P} \cap M$. Because $\mathcal{D} \subseteq M$ is dense open, there is some $q' \leq q \upharpoonright M$ which is in \mathcal{D} (hence in M). To finish the proof, we still have to show that q and q' are compatible. First note that $\text{supp}(q \upharpoonright M) = \text{supp}(q) \cap \text{supp}(q')$ and consider the following function $r : \omega_1 \rightarrow H_\theta$ defined on the support $\text{supp}(r) = \text{supp}(q) \cup \text{supp}(q')$:
For $\alpha \geq \delta$ define $r(\alpha) = q(\alpha)$ and for $\alpha < \delta$ define

$$r(\alpha) = (q \vee q')(\alpha) \cup \left\{ \varphi_{N'}^{-1}(N) : N \in q'(\alpha), N' \in q(\delta), \varphi_{N'} : N' \xrightarrow{\cong} M \cap H_\theta \right\}.$$

We will prove that $r \leq q, q'$ is in \mathcal{P} which will finish the proof. Properties (1) and (2) from Definition 2.1 are clear. To see (3) take any $\alpha, \beta < \omega_1$ with $\alpha < \beta$ and any $N \in r(\alpha)$.

If $\alpha \geq \delta$, then $r(\alpha) = q(\alpha)$ and $r(\beta) = q(\beta)$, hence there is clearly some $N' \in r(\beta) = q(\beta)$ such that $N \in N'$.

If $\alpha < \delta$ and $\beta \geq \delta$, then N belongs to some $N' \in q(\delta)$ which belongs to some $N_1 \in q(\beta) = r(\beta)$, hence $N \in N_1$ and the statement is true in this case as well.

Otherwise, note that $\text{supp}(q) \cap \delta \subseteq \text{supp}(q')$ and consider the following two possibilities. The first, that $N \in q'(\alpha)$ and $\beta < \delta$. Then there is a model $N' \in q'(\beta) \subseteq r(\beta)$ such that $N \in N'$. The second case is that $N \in r(\alpha) \setminus q'(\alpha)$ and $\beta < \delta$. If $N = \varphi_{N'}^{-1}(N_1)$ for some $N_1 \in q'(\alpha)$, take $N_2 \in q'(\beta)$ such that $N_1 \in N_2$ and note that $N = \varphi_{N'}^{-1}(N_1) \in \varphi_{N'}^{-1}(N_2) \in r(\beta)$. If not, then $N \in q(\alpha)$, so there is some $N_3 \in q'(\beta)$ such that $\varphi_{N'}(N) \in N_3$, hence $N \in \varphi_{N'}^{-1}(N_3) \in r(\beta)$ and the proof is finished. \square

Lemma 3.3. *Let \mathcal{G} be a filter generic in \mathcal{P} over V , let $M, M' \prec H_{(2^\theta)^+}$ and $p, \mathcal{P} \in M \cap M'$. If $\varphi : M \xrightarrow{\cong} M'$, then for $\delta = M \cap \omega_1 = M' \cap \omega_1$ the condition $p_{MM'} = p \cup \{\langle \delta, \{M \cap H_\theta, M' \cap H_\theta\} \rangle\}$ satisfies:*

$$p_{MM'} \Vdash \check{\varphi}[\check{\mathcal{G}} \cap \check{M}] = \check{\mathcal{G}} \cap \check{M}'.$$

Proof. Assume the contrary, that there is a condition $q \leq p_{MM'}$ and a set x such that $q \Vdash \check{x} \in \check{\mathcal{G}} \cap \check{M} \wedge \check{\varphi}(\check{x}) \notin \check{\mathcal{G}} \cap \check{M}'$. Then we have $q \Vdash \check{x} \in \check{\mathcal{G}} \cap \check{M}$ and $q \Vdash \check{\varphi}(\check{x}) \notin \check{\mathcal{G}} \cap \check{M}'$. From the fact that $q \Vdash \check{x} \in \check{\mathcal{G}} \cap \check{M}$, we have that $q \not\perp x$ (this is true because if $q \perp x$, then it is not possible that $q \Vdash \check{x} \in \check{\mathcal{G}}$). Now take some $q' \leq q, x$ and assume that for all $r \leq q'$ there is some $t \leq r$ such that $t \leq \varphi(x)$ (which implies that $t \Vdash \check{\varphi}(\check{x}) \in \check{\mathcal{G}}$). It follows that the set $\{t \in \mathcal{P} : t \Vdash \check{\varphi}(\check{x}) \in \check{\mathcal{G}}\}$ is dense below q' , which is impossible because it would imply that $q' \Vdash \check{\varphi}(\check{x}) \in \check{\mathcal{G}} \cap \check{M}'$ which is in contradiction with the assumption that $q' \leq q$ and that $q \Vdash \check{\varphi}(\check{x}) \notin \check{\mathcal{G}} \cap \check{M}'$. Hence, we can pick some $r \leq q' \leq q, x$ which is incompatible with $\varphi(x)$. Now, consider the condition $r \upharpoonright M$. We have the following claim.

Claim 3.4. Conditions $\varphi(r \upharpoonright M)$ and r are compatible.

Proof. First note that $\text{supp}(\varphi(r \upharpoonright M)) \subseteq \text{supp}(r)$. We will prove that the condition $s = \varphi(r \upharpoonright M) \vee r$ is in \mathcal{P} which will prove the claim (then s will be below both $\varphi(r \upharpoonright M)$ and r). It is clear that the conditions (1) and (2) from Definition 2.1 are fulfilled so pick arbitrary $\alpha, \beta \in \text{supp}(s)$ with $\alpha < \beta$. If $\alpha \geq \delta$, then every $N \in s(\alpha)$ is in $r(\alpha)$ hence there is some $N' \in r(\beta) = s(\beta)$ such that $N \in N'$. If $\beta \leq \delta$, then for $N \in s(\alpha) \cap r(\alpha)$ there is some $N' \in r(\beta) \subseteq r(s)$ such that $N \in N'$. On the other hand, if $\beta \leq \delta$ and $N \in s(\alpha) \cap \varphi(r \upharpoonright M)(\alpha)$, then $N = \varphi(N_1)$ for some $N_1 \in (r \upharpoonright M)(\alpha)$, and $N_1 = \psi_{N_2, M}(N_3)$ for $N_2 \in r(\delta)$, $N_3 \in r(\alpha) \cap N_2$ and an isomorphism $\psi_{N_2, M} : N_2 \rightarrow M \cap H_\theta$. Now, there is some $N_4 \in r(\beta)$ such that $N_3 \in N_4$. So, clearly we have that $N \in \varphi(\psi_{N_2, M}(N_4)) \in \varphi(r \upharpoonright M)(\beta) \subseteq s(\beta)$. If $\alpha < \delta$ and $\delta \leq \beta$, then for each $N \in s(\alpha) \cap r(\alpha)$ there is some $N' \in r(\beta) = s(\beta)$ such that $N \in N'$. Finally, let $\alpha < \delta$, $\delta \leq \beta$ and $N \in s(\alpha) \cap \varphi(r \upharpoonright M)(\alpha)$. Then $N \in M \cap H_\theta \in s(\delta) \cap r(\delta)$ and clearly there is some $N' \in r(\beta) = s(\beta)$ such that $N \in M \cap H_\theta \in N'$, hence $N \in N'$. \square

Now because $x \in M$ and $r \leq x$ we have that $r \upharpoonright M \leq x$ which implies $\varphi(r \upharpoonright M) \leq \varphi(x)$ (the last implication is true because φ is an automorphism between M and M'). Now from $\varphi(r \upharpoonright M) \leq \varphi(x)$ and $r \perp \varphi(x)$ follows that $\varphi(r \upharpoonright M)$ and r are incompatible which is not possible according to the Claim 3.4. \square

The following lemma will be useful in Section 5 of the paper.

Lemma 3.5. *If $M = M' \cap H_\theta$ for some $M' \prec H_{(2^\theta)^+}$ such that $\mathcal{P} \in M'$ and if $M \in G(\delta)$ for $\delta = M \cap \omega_1$, then in $V[\mathcal{G}]$ we have $M[\mathcal{G}] \cap \omega_1 = \delta$.*

Proof. First note that any $p \in \mathcal{G}$ such that $M \in p(\delta)$ is an (M', \mathcal{P}) -generic condition which forces that $M' \cap \text{Ord} = M'[\mathcal{G}] \cap \text{Ord}$ (see [9, Lemma III 2.6]). Because $\omega_1 \subseteq H_\theta$ this implies that $M[\mathcal{G}] \cap \omega_1 = M \cap \omega_1 = \delta$. \square

4. PRESERVING CH

Lemma 4.1 (CH). \mathcal{P} satisfies ω_2 -c.c.

Proof. Assume that CH holds and that there is a sequence $\{p_\alpha : \alpha < \omega_2\}$ of pairwise incompatible elements of \mathcal{P} . For each $\alpha < \omega_2$ the function \bar{p}_α (the transitive collapse of p_α) is a finite subset of $\omega_1 \times H_{\omega_1}$. Hence, there are some distinct $\alpha, \beta < \omega_2$ such that $\bar{p}_\alpha = \bar{p}_\beta$ (here we are using CH which implies that $|H_{\omega_1}| = \omega_1$). But then conditions p_α and p_β are compatible by Lemma 2.8 which is a contradiction with the choice of the sequence $\{p_\alpha : \alpha < \omega_2\}$. \square

Lemma 4.2. \mathcal{P} preserves CH.

Proof. Assume that CH holds in V and that there is a sequence $\{\tau_\alpha : \alpha < \omega_2\}$ of \mathcal{P} -names and a condition $p \in \mathcal{G}$ such that

$$p \Vdash \langle \tau_\alpha : \alpha < \omega_2 \rangle \text{ is a sequence of pairwise distinct reals}.$$

For each $\alpha < \omega_2$ let M_α be a countable elementary submodel of $H_{(2^\theta)^+}$ containing $\mathcal{P}, \tau_\alpha, p$.

Now, using CH, we conclude that there are $\alpha, \beta < \omega_2$ ($\alpha \neq \beta$) such that there is an automorphism $\varphi : M_\alpha \rightarrow M_\beta$ which satisfies $\varphi(\tau_\alpha) = \tau_\beta$. To see this consider the transitive collapses of M_α 's, they are countable submodels of H_{ω_1} but since we assumed the CH (which implies $|H_{\omega_1}| = \omega_1$) there must be two collapses $\overline{M_\alpha}$ and $\overline{M_\beta}$ which are isomorphic (via isomorphism ϕ). Then the isomorphism φ is given by $\varphi = \pi_{\overline{M_\beta}}^{-1} \circ \phi \circ \pi_{M_\alpha}$. Also, it clearly holds that $\varphi(M_\alpha \cap H_\theta) = M_\beta \cap H_\theta$, and because M_α and M_β are isomorphic we can denote $\delta = M_\alpha \cap \omega_1 = M_\beta \cap \omega_1$.

Now we prove that there is a condition $p_{\alpha\beta} \leq p$ such that $p_{\alpha\beta} \Vdash \tau_\alpha = \tau_\beta$, hence τ_α and τ_β cannot be names for distinct reals in $V[\mathcal{G}]$. First note that φ being an isomorphism, we have

$$(4.1) \quad \forall n < \omega \forall p' \in M_\alpha \cap \mathcal{P} \forall \epsilon < 2 \ (p' \Vdash \tau_\alpha(\check{n}) = \check{\epsilon} \Leftrightarrow \varphi(p') \Vdash \tau_\beta(\check{n}) = \check{\epsilon}).$$

Claim 4.3. If $p_{\alpha\beta} = p \cup \{\delta, \{M_\alpha \cap H_\theta, M_\beta \cap H_\theta\}\}$, then $p_{\alpha\beta} \Vdash \tau_\alpha = \tau_\beta$.

Proof. Assume the contrary, that there is some $q \leq p_{\alpha\beta}$ and $n < \omega$ such that $q \Vdash \tau_\alpha(\check{n}) \neq \tau_\beta(\check{n})$ (suppose that $q \Vdash \tau_\alpha(\check{n}) = \check{0}$ and $q \Vdash \tau_\beta(\check{n}) = \check{1}$). Then by elementarity of M_α there is some $r \in \mathcal{P} \cap M_\alpha$ such that $r \leq q \upharpoonright M$ and $\text{supp}(q \upharpoonright M_\alpha) \sqsubseteq \text{supp}(r)$ (where $a \sqsubseteq b$ denotes that a is an initial segment of b) and which satisfies $r \Vdash \tau_\alpha(\check{n}) = \check{0}$. Again, because φ is an isomorphism we have $\varphi(q \upharpoonright M_\alpha) = q \upharpoonright M_\beta$, so $q \upharpoonright M_\beta$ is compatible with $\varphi(r)$. From $q \upharpoonright M_\beta \not\leq \varphi(r)$ we conclude that $q \not\leq \varphi(r)$. But, then $\varphi(r) \Vdash \tau_\beta(\check{n}) = \check{0}$ (from equation (4.1)) which is in contradiction with the fact that $q \Vdash \tau_\beta(\check{n}) = \check{1}$ (simply because $\varphi(r) \not\leq q$). \square

Hence, according to the Claim 4.3, p cannot force that $\langle \tau_\alpha : \alpha < \omega_2 \rangle$ is a sequence of names for distinct reals in $V[\mathcal{G}]$, which proves the theorem. \square

5. KUREPA TREE

Recall that a Kurepa tree is a tree of height ω_1 , with all levels countable but at least ω_2 branches. In this section we will show that forcing with \mathcal{P} adds a Kurepa tree.

Theorem 5.1. *There is a Kurepa tree in $V[\mathcal{G}]$.*

Proof. For each $\alpha < \omega_2$ define the function $f_\alpha : \omega_1 \rightarrow \omega_1$:

$$(5.1) \quad f_\alpha(\delta) = \begin{cases} \xi, & \text{if there is } M \in G(\delta) \text{ such that } \alpha \in M, \pi_M(\alpha) = \xi; \\ 0, & \text{otherwise.} \end{cases}$$

Note that if there are two isomorphic models $M_1, M_2 \in G(\delta)$ containing $\alpha < \omega_2$, then Lemma 2.7 implies that $\pi_{M_1}(\alpha) = \pi_{M_2}(\alpha)$, so each f_α is well defined. Also, if $\alpha \neq \beta$, then there are some $\delta < \omega_1$ and $M \in G(\delta)$ such that $\{\alpha, \beta\} \in M$, but then $\pi_M(\alpha) \neq \pi_M(\beta)$ (i.e. $f_\alpha(\delta) \neq f_\beta(\delta)$). So for $\alpha \neq \beta$ we have $f_\alpha \neq f_\beta$. If we denote the set of functions coding branches in T by $\mathcal{F} = \{f_\alpha : \alpha < \omega_2\}$, then $\mathcal{F}_\alpha = \{f_\alpha \upharpoonright \delta : \delta < \omega_1\}$ will be the α -th branch and the Kurepa tree will be given by $T = \bigcup_{\delta < \omega_1} T_\delta$, where $T_\delta = \mathcal{F} \upharpoonright \delta = \{f_\alpha \upharpoonright \delta : \alpha < \omega_2\}$. We will show that for each δ , the level T_δ is countable. This will finish the proof. So assume that there are some $p \in \mathcal{G}$ and $\delta' < \omega_1$ such that $p \Vdash \text{“}\dot{T}_{\delta'} \text{ is uncountable”}$. Take a countable elementary submodel M of $H_{(2^\theta)^+}$ such that $p, \mathcal{P}, \delta' \in M$ and denote $\delta = M \cap \omega_1$. Because we have chosen M so that $\delta' \in M$ we have $\delta' < \delta$. Consider the (M, \mathcal{P}) -generic condition $p_M = p \cup \{\delta, \{M \cap H_\theta\}\} \leq p$ (note that $p_M \leq p$ because $p \in M$). The following claim shows that the (M, \mathcal{P}) -generic condition forces that the branches passing through the δ' -th level are indexed only by ordinals less than ω_2 which are already in M .

Claim 5.2. Suppose that $p' \in \mathcal{P}$ is such that $M' \in G(\delta) \cap p'(\delta)$ and $\delta \geq \delta'$. Then $p' \Vdash \dot{T}_{\delta'} = \{ \dot{f}_\alpha \upharpoonright \delta' : \alpha \in \check{M}' \cap \check{\omega}_2 \}$.

Proof. The inclusion “ \supseteq ” is trivial. To prove the reverse inclusion take some $q \leq p'$ and $\alpha' < \omega_2$. Because $q \leq p'$ we have that $q \Vdash \dot{f}_{\alpha'} \upharpoonright \delta' \in \dot{T}_{\delta'}$. In order to finish the proof of the claim, we will find $r \leq q$ and $\alpha \in M' \cap \omega_2$ such that $r \Vdash \dot{f}_{\alpha'} \upharpoonright \delta' = \dot{f}_\alpha \upharpoonright \delta'$. We will consider two cases.

Case I ($\dot{f}_{\alpha'} \upharpoonright \delta' = 0$). Then for $0 \in M' \cap \omega_2$ we have that $f_0 \upharpoonright \delta' = \dot{f}_{\alpha'} \upharpoonright \delta'$. To see this notice that either for every $\gamma < \delta'$ there is some $N \in G(\gamma)$ and then clearly $0 \in N$ and $\pi_N(0) = 0$ or $G(\gamma) = \emptyset$ and again $f_0(\gamma) = 0$. So we clearly have $q \Vdash \dot{f}_{\alpha'} \upharpoonright \delta' = \dot{f}_0 \upharpoonright \delta'$.

Case II. There is $\gamma < \delta'$ such that $f_{\alpha'}(\gamma) \neq 0$. Let γ_0 be minimal such γ . This means that for some $N_0 \in G(\gamma_0)$ we have $\alpha' \in N_0$. Now, there is some $r \leq q \leq p'$ such that $N_0 \in r(\gamma_0)$, hence there is some $N_1 \in r(\delta)$ such that $N_0 \in N_1$ (note that $r(\delta)$ is not empty because $M' \in p'(\delta) \subseteq r(\delta)$). Now we have that $\alpha' \in N_1$ ($N_0 \in N_1$ implies $N_0 \subseteq N_1$). Because $M' \in r(\delta)$ there is an isomorphism $\varphi : N_1 \xrightarrow{\cong} M'$. Denote $\alpha = \varphi(\alpha')$. We show that the condition $r_1 = \text{cl}_\delta(r)$ (note $r_1 \leq r \leq q$) forces “ $\dot{f}_{\alpha'} \upharpoonright \delta' = \dot{f}_\alpha \upharpoonright \delta'$ ”. First, if $\gamma' < \gamma_0$, then $r_1 \Vdash \dot{f}_\alpha(\check{\gamma}') = 0 = \dot{f}_{\alpha'}(\check{\gamma}')$. This is true because if there is some $N' \in r_1(\gamma')$ such that $\alpha = \varphi(\alpha') \in N'$, then there would be some $\varphi^{-1}(N') \in r_1(\gamma')$ such that $\alpha' \in \varphi^{-1}(N')$ which is impossible by the choice of γ_0 (it is a minimal ordinal such that α' belongs to some model on its level in generic filter). If $\gamma \geq \gamma_0$ and $\gamma < \delta'$, let $r_1 \Vdash \dot{f}_{\alpha'}(\check{\gamma}) = \check{\xi} \neq 0$ (it cannot be 0 because $\alpha' \in N_0 \in r_1(\gamma_0)$ so for every $\gamma \geq \gamma_0 \exists N \in r_1(\gamma) \alpha' \in N_0 \in N$ and since $\alpha' \neq 0$ we clearly have $\pi_N(\alpha') \neq 0$) and take $N' \in r_1(\gamma)$ such that $\alpha' \in N'$ and $\pi_{N'}(\alpha') = \xi$. Then $\alpha \in \varphi(N') \in r_1(\gamma)$ and clearly $\pi_{\varphi(N')}(\alpha) = \xi$ which implies $r_1 \Vdash \dot{f}_\alpha(\check{\gamma}) = \check{\xi}$. So $r_1 \leq q$ forces “ $\dot{f}_{\alpha'} \upharpoonright \delta' = \dot{f}_\alpha \upharpoonright \delta'$ ” and the claim is proved. \square

Now according to the Claim 5.2 $p_M \Vdash \dot{T}_{\delta'} = \{ \dot{f}_\alpha \upharpoonright \dot{\delta}' : \check{\alpha} \in \check{M} \cap \check{\omega}_2 \}$ so p cannot force that $T_{\delta'}$ is uncountable (because M is countable and $p_M \leq p$), hence $T_{\delta'}$ is countable in $V[\mathcal{G}]$. \square

Corollary 5.3. *Every uncountable downward closed set $S \subseteq T$ contains a branch of T .*

Proof. First recall that we denoted a branch of T by $\mathcal{F}_\alpha = \{ f_\alpha \upharpoonright \gamma : \gamma < \omega_1 \}$. Take a \mathcal{P} -name \dot{S} for S and $p \in \mathcal{G}$ such that $p \Vdash \text{“}\dot{S} \text{ is downward closed”}$. Now pick $M \prec H_{(2^\theta)^+}$ such that $\dot{S}, \mathcal{P}, p \in M$ and denote $\delta = M \cap \omega_1$. So we have that $S \in M[\mathcal{G}] \prec H_{(2^\theta)^+}[\mathcal{G}] = H_{(2^\theta)^+}^{V[\mathcal{G}]}$ (according to Lemma 2.11). We have already shown that the condition $p_M = p \cup \{ \langle M \cap \omega_1, \{ M \cap H_\theta \} \rangle \}$ is (M, \mathcal{P}) -generic and by Claim 5.2 we have $p_M \Vdash \dot{T}_\delta = \{ \dot{f}_\alpha \upharpoonright \dot{\delta} : \check{\alpha} \in \check{M} \cap \check{\omega}_2 \}$. Note also that according to Lemma 3.5 $p_M \Vdash \check{\delta} = \check{M}[\dot{\mathcal{G}}] \cap \check{\omega}_1$. Now, because S is uncountable and each level in T is countable by Theorem 5.1, there is some $\beta > \delta$ such that $S \cap T_\beta \neq \emptyset$ and because it is downward closed there is some $\alpha \in M \cap \omega_2$ such that $f_\alpha \upharpoonright \delta \in S$. Now the fact that S and \mathcal{F}_α are in $M[\mathcal{G}]$ (for S is clear and for \mathcal{F}_α it follows from Lemma 2.5 and the fact that f_α is defined only from \mathcal{G} and $\alpha \in M$) implies that $S \cap \mathcal{F}_\alpha \in M[\mathcal{G}]$. If the set $S \cap \mathcal{F}_\alpha$ is countable, then by elementarity of $M[\mathcal{G}]$ in $H_{(2^\theta)^+}^{V[\mathcal{G}]}$ and Lemma 2.3 we have $S \cap \mathcal{F}_\alpha \subseteq M[\mathcal{G}]$. Now $\delta = M[\mathcal{G}] \cap \omega_1 = M \cap \omega_1$ and Lemma 2.4 imply that there is some $\gamma' < \delta$ such that for every $\gamma \geq \gamma'$ we have $f_\alpha \upharpoonright \gamma \notin S$ which contradicts the assumption that $f_\alpha \upharpoonright \delta \in S$. So $S \cap \mathcal{F}_\alpha$ is uncountable and if a downward closed set in a tree of height ω_1 intersects a branch at uncountably (hence cofinally) many levels it clearly contains that branch. Hence, $\mathcal{F}_\alpha \subseteq S$. \square

6. ALMOST SOUSLIN TREE

In this section we consider the slightly modified version of the poset \mathcal{P} . Namely, let \mathcal{P}_c be the partial order satisfying all the conditions (1)-(3) from Definition 2.1 together with

- (4) for every $p \in \mathcal{P}_c$ there is a continuous \in -chain $\langle M_\xi : \xi < \omega_1 \rangle$ (i.e. if β is a limit ordinal, then $M_\beta = \bigcup_{\xi < \beta} M_\xi$) of countable elementary submodels of H_θ such that $\forall \xi \in \text{supp}(p) \ M_\xi \in p(\xi)$.

We point out that in this case the generic filter in \mathcal{P}_c is denoted by \mathcal{G}_c and that the function G_c analogous to G is a total function from ω_1 to H_θ . Moreover, the poset \mathcal{P}_c is strongly proper. The proof of this fact needs slight modification of the proof of the Lemma 3.2. Also, the Kurepa tree from the previous paragraph would be obtained in the same way with the poset \mathcal{P}_c . Now we prove that the tree T which we already constructed is an almost Souslin tree in $V[\mathcal{G}_c]$, i.e., if $X \subseteq T$ is an antichain, then $L(X) = \{ \gamma < \omega_1 : X \cap T_\gamma \neq \emptyset \}$ (the level set of X) is not stationary in ω_1 . Hence, to show that T is almost Souslin we have to find a club Γ in ω_1 such that $\Gamma \cap L(X) = \emptyset$. So let $\tau \in H_\theta$ be a \mathcal{P}_c -name and define the set

$$\Gamma_\tau = \{ \gamma < \omega_1 : \exists M \in G_c(\gamma) \ \tau \in M \ \& \ M[\mathcal{G}_c] \cap \omega_1 = M \cap \omega_1 = \gamma \}.$$

Lemma 6.1 (CH). *The set Γ_τ is a club in ω_1 .*

Proof. First we prove that Γ_τ is unbounded in ω_1 . Take $\gamma_1 < \omega_1$ and assume that there is some $p \in \mathcal{G}_c$ such that $p \Vdash \forall \check{\gamma} \in \check{\Gamma}_\tau \ \check{\gamma} < \check{\gamma}_1$. Take elementary

submodel $M \prec H_{(2^\theta)^+}$ such that $p, \mathcal{P}_c, \gamma_1, \tau \in M$. The condition $p_M \leq p$ given by $p_M = p \cup \{\langle M \cap \omega_1, \{M \cap H_\theta\} \rangle\}$ is in \mathcal{P}_c . To see this note that because $p \in \mathcal{P}_c$ there is a continuous chain $\langle M_\xi : \xi < \omega_1 \rangle$ witnessing that condition (4) is satisfied, so by elementarity of M there is a continuous chain $\langle N_\xi : \xi < \delta_M \rangle$ such that $\forall \alpha \in \text{supp}(p) N_\alpha \in p(\alpha)$ and $\bigcup_{\xi < \delta_M} N_\xi = M$. The chain $\langle N'_\xi : \xi < \omega_1 \rangle$ witnessing that $p_M \in \mathcal{P}_c$ is now recursively given by $\forall \xi < \delta N'_\xi = N_\xi, N'_\delta = M \cap H_\theta$, for successor $\alpha + 1 > \delta$ define $N'_{\alpha+1} \prec H_\theta$ as an arbitrary submodel containing N_α while for limit $\alpha > \delta$ define $N'_\alpha = \bigcup_{\xi < \alpha} N'_\xi$. Also, p_M is an (M, \mathcal{P}_c) -generic condition, so we have that $p_M \Vdash \check{M}[\mathcal{G}_c] \cap \check{\omega}_1 = \check{M} \cap \check{\omega}_1$ (see [9, Lemma III 2.6]). This implies that $p_M \Vdash \check{M} \cap \check{\omega}_1 \in \Gamma_\tau$, but from $\gamma_1 \in M$ it follows that $p_M \Vdash \check{\gamma}_1 < \check{M} \cap \check{\omega}_1$, which is in contradiction with the choice of p . So Γ_τ is unbounded in ω_1 .

In order to prove that Γ_τ is a club, it is enough to show that for every ordinal δ such that $\delta = \text{sup}(\Gamma_\tau \cap \delta)$ there is some $M \in G_c(\delta)$ which satisfies $M[\mathcal{G}_c] \cap \omega_1 = M \cap \omega_1 = \delta$. Because $\delta = \text{sup}(\Gamma_\tau \cap \delta)$ there is an increasing sequence $\gamma_n \in \Gamma_\tau \cap \delta$ such that $\text{sup}_{n < \omega} \gamma_n = \delta$. Pick $N_{\gamma_0} \in G_c(\gamma_0)$ such that $\tau \in N_{\gamma_0}$, hence there is some $M \in G_c(\delta)$ such that $\tau \in M$. First we will show that $M \cap \omega_1 = \delta$. Inductively pick models $N_{\gamma_n} \in G_c(\gamma_n)$ such that $N_{\gamma_{n-1}} \in N_{\gamma_n}$. Note that because $\gamma_n \in \Gamma_\tau$, we have $N_{\gamma_n} \cap \omega_1 = \gamma_n$. Now, because $\text{sup}_{n < \omega} \gamma_n = \delta$ and $\forall n < \omega \gamma_n < M \cap \omega_1$ (from $\forall n < \omega \exists N \in G_c(\delta) N_{\gamma_n} \in N$) we have that $M \cap \omega_1 \geq \delta$. So assume that $M \cap \omega_1 = \beta > \delta$. Let $p \in \mathcal{G}_c$ be any condition such that $M \in p(\delta)$.

Claim 6.2. The set $D_\delta = \{q \leq p : \exists \gamma' < \delta \exists N \in q(\gamma') N \cap \omega_1 > \delta\}$ is dense below p in \mathcal{P}_c .

Proof. Take arbitrary $p' \leq p$ and pick a continuous \in -sequence $\langle M_\xi : \xi < \omega_1 \rangle$ such that $\forall \xi \in \text{supp}(p') M_\xi \in p'(\xi)$. Because this chain is continuous, δ is a limit ordinal and $M \in p'(\delta)$ is such that $M \cap \omega_1 = \beta > \delta$, there is some $\xi_0 < \delta$ such that $M_{\xi_0} \cap \omega_1 > \delta$. Now pick any $\xi_1 \in \delta \setminus \max(\delta \cap (\text{supp}(p') \cup \{\xi_0\}))$. Clearly $M_{\xi_1} \cap \omega_1 > \delta$. To extend p' to some $q \in D_\delta$ first denote the ordinal $\alpha = \max(\text{supp}(p') \cap \delta)$. Further, note that for each $N \in p'(\alpha)$ there is some $M_N \in p'(\delta)$ such that $N \in M_N$. Let $\psi_N : M_N \xrightarrow{\cong} M_\delta$ be an isomorphism for each $N \in p'(\alpha)$. Because $\langle M_\xi : \xi < \omega_1 \rangle$ is a continuous chain and δ is a limit ordinal, there is some $\xi_2 < \delta$ such that $M_{\xi_1} \in M_{\xi_2}$ and that moreover $\forall N \in p'(\alpha) \psi_N(N) \in M_{\xi_2}$. Now define

$$q = p' \cup \{ \langle \xi_2, \{M_{\xi_2}\} \cup \{ \psi_N^{-1}(M_{\xi_2}) : N \in p'(\alpha) \cap M_\delta \} \rangle \}.$$

It is clear that $q \in \mathcal{P}$ and the sequence $\langle M_\xi : \xi < \omega_1 \rangle$ witnesses that q satisfies the property (4), hence q is also in \mathcal{P}_c . □

Now, according to Claim 6.2, there is some $q \in \mathcal{G}_c \cap D_\delta$ below p . But this implies that there is some $N \in G_c(\gamma')$ for $\gamma' < \delta$, satisfying $N \cap \omega_1 > \delta$ which is impossible by the choice of γ_n (note that γ_n is cofinal in δ and we would have that there is some $\gamma_{n'} > \gamma'$ such that $N_{\gamma_{n'}} \cap \omega_1 < N \cap \omega_1$ and $N_{\gamma_{n'}} \in G_c(\gamma_{n'})$ and $N \in G_c(\gamma')$). So $M \cap \omega_1 = \delta$.

We still have to prove that also $M[\mathcal{G}_c] \cap \omega_1 = \delta$. Let $\sigma \in M$ be a \mathcal{P}_c -name for a countable ordinal. Because CH holds, the poset \mathcal{P}_c is ω_2 -c.c., Lemma 4.1, so we can assume that σ is a nice name of cardinality at most ω_1 , i.e., $\sigma = \{ \langle \check{\xi}, p_\xi \rangle : \xi < \omega_1 \}$ where $\{p_\xi : \xi < \omega_1\}$ is a maximal antichain in \mathcal{P}_c . Now let $p \in \mathcal{G}_c$ be any condition containing M . Because $p \in \mathcal{P}_c$, there is a continuous chain $\langle M_\xi : \xi < \omega_1 \rangle$ such that $\forall \xi \in \text{supp}(p) M_\xi \in p(\xi)$. Now there is an isomorphism $\varphi : M \xrightarrow{\cong} M_\delta$ and in the

same way as in the proof of Claim 6.2 we show that there is some $q \in \mathcal{G}_c$ and $\xi_1 < \delta$ such that $\varphi(\sigma) \in M_{\xi_1} \in q(\xi_1) \subseteq G_c(\xi_1)$. Because $\delta = \sup(\Gamma_\tau \cap \delta)$ we can assume that $\xi_1 \in \Gamma_\tau$. Now according to Lemma 3.3 and the form of σ and $\varphi(\sigma)$ we have that $\text{int}_{\mathcal{G}_c}(\sigma) = \text{int}_{\mathcal{G}_c}(\varphi(\sigma)) < M_{\xi_1}[\mathcal{G}_c] \cap \omega_1 = M_{\xi_1} \cap \omega_1 < \delta$. Hence $M[\mathcal{G}_c] \cap \omega_1 = \delta$ and the proof is finished. \square

Theorem 6.3 (CH). *The tree T is an almost Souslin tree.*

Proof. Let τ' be a \mathcal{P}_c -name for an antichain X in T . Because CH holds in V , according to Lemma 4.1 \mathcal{P}_c is ω_2 -c.c. so there is a \mathcal{P}_c -name τ for X which is in H_θ . To prove the theorem, we will show that $L(X) \cap \Gamma_\tau = \emptyset$.

So assume that $X \cap T_\delta \neq \emptyset$ for some $\delta \in \Gamma_\tau$. Because $\delta \in \Gamma_\tau$ there is some $M \in G_c(\delta)$ such that $\tau \in M$ and that $M[\mathcal{G}_c] \cap \omega_1 = M \cap \omega_1 = \delta$, so take any $p \in \mathcal{G}_c$ such that $M \in p(\delta)$. Now, in the same way as in the proof of Claim 5.2 we know that p forces that $T_\delta = \{f_\alpha \upharpoonright \delta : \alpha \in M \cap \omega_2\}$, hence there is some $\alpha \in M \cap \omega_2$ such that $f_\alpha \upharpoonright \delta \in X$. Consider the branch \mathcal{F}_α . It is defined solely from $\alpha \in M$ and \mathcal{G}_c , so $\mathcal{F}_\alpha \in M[\mathcal{G}_c]$. Also, because $\tau \in M$ we have that $X \in M[\mathcal{G}_c]$. Consequently $\mathcal{F}_\alpha \cap X \in M[\mathcal{G}_c]$, and from the fact that X is an antichain it follows that this intersection is singleton (i.e. $f_\alpha \upharpoonright \delta$). But, because $M[\mathcal{G}_c] \prec H_\theta^{V[\mathcal{G}_c]}$ (see Lemma 2.11) and the height of $f_\alpha \upharpoonright \delta$ is less than ω_1 , there must exist some element $t \in \mathcal{F}_\alpha \cap X \cap M[\mathcal{G}_c]$ which is of height less than $\delta = M[\mathcal{G}_c] \cap \omega_1$. But then $t < f_\alpha \upharpoonright \delta$ and both $t, f_\alpha \upharpoonright \delta \in X$ which is in contradiction with the fact that X is an antichain. \square

7. CONCLUDING REMARKS

We have seen that both versions of the matrix posets force the Continuum Hypothesis. It turns out that a bit more is true, both versions of the matrix posets force the combinatorial principle \diamond independently of the status of the Continuum Hypothesis in the ground model. In fact, if CH fails in V , that \diamond is forced follows from a slight adaptation of a result of Roslanowski and Shelah [8] to our matrix posets that do not have cardinality 2^{\aleph_0} but are nevertheless 2^{\aleph_0} -centered in the canonical way. So we may concentrate on the case that the ground model V satisfies CH. If CH holds in V , then the following theorem from [12] proves a bit more for the continuous version of the matrix forcing (poset \mathcal{P}_c of Section 6).

Theorem 7.1 (CH). \diamond^+ holds in $V[\mathcal{G}_c]$.

For the convenience of the reader we include the sketch of the proof.

Proof. First denote the generic club by Δ , i.e., Δ is the club contained in the set $\text{Tr}(\mathcal{G}_c) = \left\{M \cap \omega_1 : M \in \bigcup_{p \in \mathcal{G}_c} \text{ran}(p)\right\}$. The key part of the proof is the following claim which shows that the generic club is almost contained in every club from the ground model.

Claim 7.2. Let $C \subseteq \omega_1$ be a club in V . Then there is a countable ordinal δ such that for every $\beta \geq \delta$ if $\beta \in \Delta$, then $\beta \in C$.

Proof. Because $C \in V$ there is some $\alpha < \omega_1$ such that $\exists M \in G_c(\alpha) C \in M$. Denote $\delta = M \cap \omega_1$. By elementarity of M we know that C is a club in δ , so $\delta \in C$. Now take arbitrary $\beta > \delta$ such that $\beta \in \Delta$. This means that for some $\gamma < \omega_1$ ($\alpha < \gamma$) there is some $N' \in G_c(\gamma)$ such that $N' \cap \omega_1 = \beta$. Also, by the definition of \mathcal{P}_c there

is some $N \in G_c(\gamma)$ such that $C \in N$. Now by elementarity of N we conclude that C is a club in $\beta = N \cap \omega_1$ so $\beta \in C$. \square

Claim 7.3. There is, in $V[\mathcal{G}_c]$, a sequence $\langle S_\alpha : \alpha < \omega_1 \rangle$ such that for every $\alpha < \omega_1$ we have $S_\alpha \in [P(\alpha)]^{\leq \omega}$ and that

$$\forall X \in P(\omega_1) \cap V \exists \gamma < \omega_1 \forall \alpha \geq \gamma X \cap \alpha \in S_\alpha.$$

Proof. By CH in V we can find an increasing continuous sequence of countable sets $\langle D_\alpha : \alpha < \omega_1 \rangle$ such that $D_\alpha \subseteq P(\alpha)$ and $\bigcup_{\alpha < \omega_1} D_\alpha = [\omega_1]^{\leq \omega}$. Let $f : \omega_1 \rightarrow \omega_1$ be defined by $f(\alpha) = \min(\Delta \setminus (\alpha + 1))$. Finally, for $\alpha < \omega_1$ we let $S_\alpha = \{X \cap \alpha : X \in D_{f(\alpha)}\}$. To see that the sets S_α ($\alpha < \omega_1$) satisfy the statement of the claim pick any $X \subseteq \omega_1$ which is in V . As in the proof of Lemma 2.1 in [4] there is a club $E \subseteq \omega_1$ which is in V such that $\forall \alpha \in E \forall \beta < \alpha X \cap \beta \in D_\alpha$. Now according to Claim 7.2 there is some $\gamma < \omega_1$ such that $\Delta \setminus \gamma \subseteq E$. For $\gamma \leq \alpha < \omega_1$ it holds $f(\alpha) \in \Delta \setminus \alpha \subseteq \Delta \setminus \gamma \subseteq E$ so as $f(\alpha) > \alpha$, the choice of E ensures that $X \cap \alpha \in D_{f(\alpha)}$. So the claim is proved. \square

Let $\langle S_\alpha : \alpha < \omega_1 \rangle$ be the sequence from the previous claim. For $\alpha < \omega_1$ let $W_\alpha = P(\alpha) \cap (\bigcup_{X \in S_\alpha} L_{\alpha+2}[X, \Delta \cap \alpha])$. Then $\langle W_\alpha : \alpha < \omega_1 \rangle$ is a \diamond^+ sequence. To show this pick arbitrary $A \subseteq \omega_1$. Because \mathcal{P}_c is ω_2 -c.c. there is a name for A which is coded by some $X \subseteq \omega_1$. Hence, $A \in L[X, \Delta]$. By Claim 7.3 there is a $\gamma < \omega_1$ such that $\forall \alpha \geq \gamma X \cap \alpha \in S_\alpha$. By induction we define a normal sequence $\langle \alpha_\xi : \xi < \omega_1 \rangle$ in ω_1 . Let $\alpha_0 > \gamma$ be the least α such that $L_\alpha[X \cap \alpha, \Delta \cap \alpha] \prec L_{\omega_1}[X, \Delta]$. If α_ξ is defined let $\alpha_{\xi+1} > \alpha_\xi$ be the least ordinal such that $L_{\alpha_{\xi+1}}[X \cap \alpha_{\xi+1}, \Delta \cap \alpha_{\xi+1}] \prec L_{\omega_1}[X, \Delta]$. Let $B = \langle \alpha_\xi : \xi < \omega_1 \rangle$. Then it is easily checked by the construction that B is a club in ω_1 . So pick arbitrary $\alpha \in B$ (we will prove that $A \cap \alpha, B \cap \alpha \in W_\alpha$). Because $\alpha > \gamma$ we have $X \cap \alpha \in S_\alpha$, so $P(\alpha) \cap L_{\alpha+2}[X \cap \alpha, \Delta \cap \alpha] \subseteq W_\alpha$. Because $L_\alpha[X \cap \alpha, \Delta \cap \alpha] \prec L_{\omega_1}[X, \Delta]$ we have that $A \cap \alpha$ is first-order definable over $L_\alpha[X \cap \alpha, \Delta \cap \alpha]$ so $A \cap \alpha \in L_{\alpha+1}[X \cap \alpha, \Delta \cap \alpha] \subseteq W_\alpha$. Similarly we would show that $B \cap \alpha \in W_\alpha$ and the theorem is proved. \square

It is clear that this proof adapts to showing that the original matrix poset (poset \mathcal{P} of Section 2) also forces \diamond .

ACKNOWLEDGEMENTS

The authors would like to thank the referee for carefully reading the paper and for many useful suggestions.

NOTE ADDED IN PROOF

After this paper was completed we received some unpublished notes of Professor T. Miyamoto which give a similar analysis of the forcing notion of matrices of elementary submodels.

REFERENCES

[1] David Asperó and Miguel Angel Mota, *Forcing consequences of PFA together with the continuum large*, Trans. Amer. Math. Soc. **367** (2015), no. 9, 6103–6129, DOI 10.1090/S0002-9947-2015-06205-9. MR3356931
 [2] D. Aspero, *A forcing notion collapsing \aleph_3 and preserving all other cardinals*, preprint (2014).
 [3] Mohammad Golshani, *Almost Souslin Kurepa trees*, Proc. Amer. Math. Soc. **141** (2013), no. 5, 1821–1826, DOI 10.1090/S0002-9939-2012-11461-3. MR3020868

- [4] Keith J. Devlin, *Concerning the consistency of the Souslin hypothesis with the continuum hypothesis*, Ann. Math. Logic **19** (1980), no. 1-2, 115–125, DOI 10.1016/0003-4843(80)90022-4. MR593030
- [5] William J. Mitchell, *$I[\omega_2]$ can be the nonstationary ideal on $\text{Cof}(\omega_1)$* , Trans. Amer. Math. Soc. **361** (2009), no. 2, 561–601, DOI 10.1090/S0002-9947-08-04664-3. MR2452816
- [6] Itay Neeman, *Forcing with sequences of models of two types*, Notre Dame J. Form. Log. **55** (2014), no. 2, 265–298, DOI 10.1215/00294527-2420666. MR3201836
- [7] I. Neeman, *Higher analog of the proper forcing axiom*, Talk at the Fields Institute, Toronto, October 2012.
- [8] Andrzej Roslanowski and Saharon Shelah, *More forcing notions imply diamond*, Arch. Math. Logic **35** (1996), no. 5-6, 299–313, DOI 10.1007/s001530050047. MR1420260
- [9] Saharon Shelah, *Proper and improper forcing*, 2nd ed., Perspectives in Mathematical Logic, Springer-Verlag, Berlin, 1998. MR1623206
- [10] Stevo Todorčević, *A note on the proper forcing axiom*, Axiomatic set theory (Boulder, Colo., 1983), Contemp. Math., vol. 31, Amer. Math. Soc., Providence, RI, 1984, pp. 209–218, DOI 10.1090/conm/031/763902. MR763902
- [11] S. Todorcevic, *Kurepa tree with no stationary antichain*, Note of September, 1987.
- [12] S. Todorcevic, *Forcing club with finite conditions*, Note of December, 1982.

MATHEMATICAL INSTITUTE SANU, KNEZA MIHAILA 36, 11001 BELGRADE, SERBIA
E-mail address: `borisa@mi.sanu.ac.rs`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, CANADA, M5S 2E4 –
AND – INSTITUT DE MATHÉMATIQUES DE JUSSIEU, UMR 7586, 2 PL. JUSSIEU, CASE 7012, 75251
PARIS CEDEX 05, FRANCE – AND – MATHEMATICAL INSTITUTE SANU, KNEZA MIHAILA 36, 11001
BELGRADE, SERBIA

E-mail address: `stevo@math.toronto.edu`

E-mail address: `stevo.todorcevic@imj-prg.fr`

E-mail address: `stevo@mi.sanu.ac.rs`