

A SIMPLIFIED KRONECKER RULE FOR ONE HOOK SHAPE

RICKY INI LIU

(Communicated by Harm Derksen)

ABSTRACT. Recently Blasiak has given a combinatorial rule for the Kronecker coefficient $g_{\lambda\mu\nu}$ when μ is a hook shape by defining a set of colored Yamanouchi tableaux with cardinality $g_{\lambda\mu\nu}$ in terms of a process called conversion. We give a characterization of colored Yamanouchi tableaux that does not rely on conversion, which leads to a simpler formulation and proof of the Kronecker rule for one hook shape.

1. INTRODUCTION

Let V_λ be the irreducible representation of the symmetric group S_n corresponding to the partition λ . The *Kronecker coefficient* $g_{\lambda\mu\nu}$ is defined to be the multiplicity of V_ν in the tensor product $V_\lambda \otimes V_\mu$. Despite their simple definition and fundamental importance, the Kronecker coefficients are not well understood: indeed, a longstanding open problem is to give a positive combinatorial formula for $g_{\lambda\mu\nu}$. For some known special cases, see [1, 3–5, 7–10].

In [3], Blasiak gives a combinatorial rule for $g_{\lambda\mu\nu}$ when only one of the partitions is a hook shape, which represents the first nontrivial Kronecker rule when two of the partitions are completely unrestricted. This rule describes the Kronecker coefficients as the cardinalities of certain sets of *colored Yamanouchi tableaux*. The characterization of these tableaux as well as the proof that they are correctly enumerated relies on two processes introduced by Haiman [6] called *mixed insertion* and *conversion*. As a result, it is unfortunately somewhat complicated to describe and apply this Kronecker rule in practice.

In this paper, we give an alternative characterization of the colored Yamanouchi tableaux from [3] that does not make reference to mixed insertion or conversion. The result is a simplified formulation and proof of the Kronecker rule for one hook shape. While we use much of the same notation as [3] for reasons of consistency, the proof contained herein is essentially self-contained.

We begin in Section 2 with some preliminaries about colored tableaux and conversion. In Section 3 we discuss reading words of colored tableaux and prove our main lemma (Lemma 3.1). Finally, in Section 4 we show how Lemma 3.1 implies the Kronecker rule for one hook shape, given in Theorem 4.2.

2. PRELIMINARIES

In this section, we review the necessary background on colored tableaux and conversion.

Received by the editors August 3, 2015.

2010 *Mathematics Subject Classification.* Primary 05E10; Secondary 20C30.

2.1. Young diagrams. Given a positive integer n , a *partition* $\lambda = (\lambda_1, \lambda_2, \dots) \vdash n$ is a weakly decreasing sequence of nonnegative integers summing to n . (We often ignore trailing zeros.) The *Young diagram* of a partition λ is an array of boxes, aligned to the north and west, with λ_i boxes in row i . We typically refer to a partition and its Young diagram interchangeably. Transposing the Young diagram of λ gives the Young diagram of its *conjugate* λ' . Given two partitions λ and μ with $\mu_i \leq \lambda_i$ for all i , the *skew Young diagram* λ/μ is the set of boxes in λ with the boxes of μ removed.

A *ribbon* is a skew Young diagram that does not contain a 2×2 square. Any ribbon can be divided into *connected components*, where boxes x and y lie in the same component if there is a sequence of boxes $x = x_1, x_2, \dots, x_k = y$ in the ribbon such that x_i and x_{i+1} share an edge.

2.2. Colored tableaux. Let \mathcal{A} be the alphabet $\{1, 2, \dots, n, \bar{1}, \bar{2}, \dots, \bar{n}\}$ endowed with the partial order $1 < 2 < \dots < n$ and $\bar{1} < \bar{2} < \dots < \bar{n}$. Note that there are $\binom{2n}{n}$ total orders of \mathcal{A} compatible with this partial order.

For any total order $<$ of \mathcal{A} , a (*semistandard*) *colored tableau* $T = T_<$ of shape λ is a filling of the boxes of λ with letters of \mathcal{A} such that:

- each row and column is weakly increasing (with respect to $<$);
- no two identical unbarred letters appear in the same column; and
- no two identical barred letters appear in the same row.

See Figure 2 for some examples of colored tableaux with respect to different orders.

Observe that if a colored tableau contains only barred letters, then transposing the tableau and removing all bars gives a semistandard tableau containing only unbarred letters.

The *unbarred content* of a colored tableau $T_<$ is the sequence $\alpha(T_<) = (\alpha_1, \alpha_2, \dots)$, where α_i is the number of occurrences of i in $T_<$, while the (*total*) *content* is (c_1, c_2, \dots) , where c_i is the total number of occurrences of i and \bar{i} in $T_<$. The *total color* is the number of barred letters in $T_<$.

If a and \bar{b} are consecutive letters in $<$ (in either order), then for any tableau $T_<$, the letters a and \bar{b} must occupy boxes forming a ribbon. Any connected component of a ribbon can be filled by a and \bar{b} in exactly two ways: these differ only in the northeast corner if $a < \bar{b}$ or in the southwest corner if $a > \bar{b}$.

2.3. Conversion. Suppose $<$ and \prec are total orders of \mathcal{A} that are identical except for the order of a and \bar{b} , say $a < \bar{b}$ but $\bar{b} \prec a$. Then there is a natural bijection between colored tableaux with respect to $<$ and colored tableaux with respect to \prec via a process called *conversion*, introduced by Haiman [6] (see also [2]).

Let $T_<$ be a colored tableau, so that the boxes containing a or \bar{b} form a ribbon. For each component of the ribbon, there is a unique way to refill it with the same number of a 's and \bar{b} 's in a way that is compatible with $\bar{b} \prec a$. (If the northeast corner contains a , then move each a to the bottom of its column; if the northeast box contains \bar{b} , move each a to the right within its row. See Figure 1.) Replacing each component in this manner switches $T_<$ to a colored tableau T_\prec .

Given any two total orders $<$ and \prec of \mathcal{A} , one can be obtained from the other by repeatedly switching the order of a consecutive barred letter and unbarred letter. Therefore, one can iterate the switching process above to convert any tableau $T_<$ to a tableau T_\prec . See Figure 2 for an example. Importantly, the resulting tableau




FIGURE 1. Conversion from $a < \bar{b}$ to $\bar{b} \prec a$. The ribbon above has two connected components with six boxes each.




FIGURE 2. Conversion from the natural order $\bar{1} < 1 < \bar{2} < 2 < \bar{3} < 3$ to the small bar order $\bar{1} \prec \bar{2} \prec \bar{3} \prec 1 \prec 2 \prec 3$ in three steps.

is well defined, that is, it does not depend on the sequence of switches used to transform $<$ to \prec (as follows from the Diamond Lemma, or see [2]).

3. READING WORDS

In this section, we define a particular reading word of a colored tableau and show how it is affected by conversion.

3.1. Ballot sequences. Given a word $w = w_1 \cdots w_m$ in the alphabet $\{1, \dots, n\}$, we say that w is a *ballot sequence* if any initial segment of w contains at least as many occurrences of i as of $i + 1$ for each letter i . (Ballot sequences are also called *lattice words* or *reverse Yamanouchi words*.) For instance, 11232132 is a ballot

sequence, but 11232213 is not because the initial segment 112322 contains more 2's than 1's.

Definition. For a sequence $\alpha = (\alpha_1, \dots, \alpha_n)$, we say that w is an α -ballot sequence if, in any initial segment of w , the number of occurrences of $i+1$ minus the number of occurrences of i is at most $\alpha_i - \alpha_{i+1}$.

When α is a partition, w is an α -ballot sequence if and only if the concatenation

$$\underbrace{11 \cdots 1}_{\alpha_1} \underbrace{22 \cdots 2}_{\alpha_2} \cdots \underbrace{nn \cdots n}_{\alpha_n} w$$

is a ballot sequence.

3.2. Reading words. We specify a particular reading order for colored tableaux as follows.

Definition. Let $T_{<}$ be a colored tableau (with respect to any total order $<$ of \mathcal{A}).

- The *unbarred reverse (row) reading word* $u = u(T_{<})$ of $T_{<}$ is the word obtained by reading its unbarred entries along rows from right to left, top to bottom, skipping over any barred entries.
- The *barred (column) reading word* $\bar{v} = \bar{v}(T_{<})$ of $T_{<}$ is the word obtained by reading the barred entries along columns from bottom to top, left to right, skipping over any unbarred entries. Let $v = v(T_{<})$ be the word obtained from \bar{v} by removing the bar from each letter.
- The *total reverse (row-column) reading word* $w = w(T_{<})$ is the concatenation $w = uv$.

For example, if $T_{<}$ is the first tableau in Figure 2, then $u(T_{<}) = 11221323$, $v(T_{<}) = 113232$, and $w(T_{<}) = 11221323113232$ (which is a ballot sequence).

Two types of total orders will be especially important for determining when the property of w being a ballot sequence is preserved under conversion.

Definition. Let $<$ be a total order on \mathcal{A} .

- The order $<$ is *unbarred-tight* if at most one barred letter occurs between any two consecutive unbarred letters.
- The order $<$ is *barred-tight* if at most one unbarred letter occurs between any two consecutive barred letters.

For instance, the order $1 < \bar{1} < \bar{2} < 2 < \bar{3} < 3$ is barred-tight but not unbarred-tight. Most total orders are neither barred-tight nor unbarred-tight, and few are both. However, among the orders that are both barred-tight and unbarred-tight are the *small bar order*

$$\bar{1} \prec \bar{2} \prec \cdots \prec \bar{n} \prec 1 \prec 2 \prec \cdots \prec n$$

and the *natural order*

$$\bar{1} < 1 < \bar{2} < 2 < \cdots < \bar{n} < n.$$

We are now ready to prove the main lemma needed to derive the hook Kronecker rule.

Lemma 3.1. *Let $<$ and \prec be total orders of \mathcal{A} , and let $T_{<}$ and T_{\prec} be colored tableaux that correspond under conversion with unbarred content $\alpha = \alpha(T_{<}) = \alpha(T_{\prec})$.*

- (a) If $<$ and \prec are both unbarred-tight, then $u(T_{<})$ is a ballot sequence if and only if $u(T_{\prec})$ is.
- (b) If $<$ and \prec are both barred-tight, then $v(T_{<})$ is an α -ballot sequence if and only if $v(T_{\prec})$ is.
- (c) If $<$ and \prec are both unbarred-tight and barred-tight, then $w(T_{<})$ is a ballot sequence if and only if $w(T_{\prec})$ is.

Proof. Part (c) follows immediately from parts (a) and (b): by definition, w is a ballot sequence if and only if u is a ballot sequence and v is an α -ballot sequence. We prove part (a) below; part (b) is similar.

Let $<$ be any unbarred-tight order, and find the smallest pair of consecutive letters of the form $a < \bar{b}$. If $a \neq 1$, then the letter preceding a must be $a - 1$, and if $a \neq n$, then the letter succeeding \bar{b} must be $a + 1$. Then define \prec to be the unbarred-tight order given by the switch

$$\cdots < a - 1 < a < \bar{b} < a + 1 < \cdots \rightarrow \cdots \prec a - 1 \prec \bar{b} \prec a \prec a + 1 \prec \cdots.$$

Switching from $T_{<}$ to T_{\prec} only shifts the occurrences of a within $u(T_{<})$ and $u(T_{\prec})$.

Denote the i th occurrence of a in $u(T_{<})$ or $u(T_{\prec})$ by a_i (and similarly for other letters). Note that for either $T_{<}$ or T_{\prec} , if a_i is the rightmost occurrence of a or \bar{b} in its row, then the occurrences of $a + 1$ that appear before a_i in u are exactly those that lie in a column to the right of a_i . (Since $a + 1$ immediately follows a and \bar{b} in $<$ and \prec , any label directly east of a_i must be at least $a + 1$, so no occurrence of $a + 1$ can be strictly southeast of a_i .) Similarly, if a_i is the leftmost occurrence of a or \bar{b} in its row, then the occurrences of $a - 1$ that appear before a_i in u are exactly those that lie in a column weakly to the right of a_i .

Suppose $u(T_{<})$ is a ballot sequence. When switching from $T_{<}$ to T_{\prec} , each a either moves to the right within its row past some \bar{b} 's, which leaves it in the same position in $u(T_{\prec})$ as in $u(T_{<})$, or it moves down within its column, which shifts it later in $u(T_{\prec})$ than in $u(T_{<})$. Hence we need only verify that a_i occurs before $(a + 1)_i$ in $u(T_{\prec})$ when a_i gets shifted down within its column c_i . In this case, a_i is the rightmost a or \bar{b} in its row in T_{\prec} , so we need at most $i - 1$ occurrences of $a + 1$ to the right of column c_i in T_{\prec} or, equivalently, in $T_{<}$.

In the ribbon containing a and \bar{b} in $T_{<}$, consider the connected component containing a_i . Since a_i gets shifted down, the northeast corner of this component must contain some a_j , $j \leq i$, lying in column $c_j = c_i + i - j$. Since a_j is the rightmost a or \bar{b} in its row and $u(T_{<})$ is a ballot sequence, there are at most $j - 1$ occurrences of $a + 1$ to the right of column c_j . But since each column can contain at most one occurrence of $a + 1$, there can be at most $j - 1 + (c_j - c_i) = i - 1$ occurrences of $a + 1$ to the right of column c_i , as desired.

Conversely, if $u(T_{\prec})$ is a ballot sequence, then we need only verify that a_i occurs after $(a - 1)_i$ in $T_{<}$ when a_i is shifted within its column. This follows from a similar argument as above.

Therefore $u(T_{<})$ is a ballot sequence if and only if $u(T_{\prec})$ is whenever $<$ and \prec are related by the single switch above. But repeatedly applying such switches will transform any unbarred-tight order to the small bar order. It follows that if $u(T_{<})$ is a ballot sequence for any unbarred-tight order, then $u(T_{\prec})$ must be for any unbarred-tight order \prec , as desired. \square

Remark 3.2. The proof of Lemma 3.1 in fact shows slightly more: if $u(T_{<})$ is a ballot sequence for some unbarred-tight order $<$, then $u(T_{\prec})$ is a ballot sequence

for any total order \prec . A similar result holds for v if $<$ is barred-tight and for w if $<$ is both barred-tight and unbarred-tight. However, it is not true that the ballot sequence condition is always preserved by conversion: for instance, the two tableaux $T_{<}$ and T_{\prec} below are related by conversion, but $w(T_{\prec}) = 1212$ is a ballot sequence while $w(T_{<}) = 2112$ is not.

$$\begin{array}{|c|c|} \hline 1 & \bar{2} \\ \hline \bar{1} & 2 \\ \hline \end{array} \longleftrightarrow \begin{array}{|c|c|} \hline 1 & 2 \\ \hline \bar{1} & \bar{2} \\ \hline \end{array}$$

$$1 \prec \bar{1} \prec \bar{2} \prec 2 \qquad 1 < \bar{1} < 2 < \bar{2}$$

Remark 3.3. Although we present Lemma 3.1 using a specific reading order for simplicity, the same result holds for many other reading orders. In particular, let $u'(T_{<})$ be any reading word in which a box containing a is read after any box containing a , $a-1$, or $a+1$ that lies weakly to the northeast or immediately to the northwest. It is easy to show that if $<$ is an unbarred-tight order, then $u(T_{<})$ is a ballot sequence if and only if $u'(T_{<})$ is.

Similarly, for any barred-tight order, v can be replaced with any reading order in which \bar{b} is read after any box containing \bar{b} , $\bar{b}-1$, or $\bar{b}+1$ that lies weakly to the southwest or immediately to the northwest.

Remark 3.4. Conversion does not in general preserve the Knuth equivalence class of the reverse of w even when $<$ and \prec are the natural order and the small bar order: take

$$T_{<} = \begin{array}{|c|c|} \hline \bar{1} & 1 \\ \hline 2 & \bar{3} \\ \hline 3 & \\ \hline \end{array}, \quad T_{\prec} = \begin{array}{|c|c|} \hline \bar{1} & 1 \\ \hline \bar{3} & 2 \\ \hline 3 & \\ \hline \end{array}.$$

The reverse of $w(T_{<})$ and $w(T_{\prec})$ are 31321 and 13321, which are not Knuth equivalent.

4. KRONECKER COEFFICIENTS

We are now ready to prove a combinatorial rule for Kronecker coefficients with one hook shape. For $0 \leq d \leq n-1$, we will denote by $\mu(d)$ the partition $(n-d, \underbrace{1, \dots, 1}_d)$.

First we recall the following well-known result relating colored tableaux to the sum of two Kronecker coefficients. This follows from the classical theory of Littlewood-Richardson coefficients; we sketch the proof below (see also Proposition 3.1 of [3]).

Proposition 4.1. *Let λ and ν be partitions of n , and let \prec be the small bar order on \mathcal{A} given by*

$$\bar{1} \prec \bar{2} \prec \dots \prec \bar{n} \prec 1 \prec 2 \prec \dots \prec n.$$

Then the number of colored tableaux T_{\prec} of content λ , total color d , and shape ν such that the total reverse reading word $w(T_{\prec})$ is a ballot sequence is $g_{\lambda\mu(d)\nu} + g_{\lambda\mu(d-1)\nu}$.

Proof. Let $\langle \cdot, \cdot \rangle$ be the Hall inner product on the ring of symmetric functions, so that, for instance, $\langle s_{\lambda}, s_{\mu} s_{\nu} \rangle$ is the Littlewood-Richardson coefficient $c_{\mu\nu}^{\lambda}$. Let $*$ denote the internal product on symmetric functions defined by $\langle s_{\lambda}, s_{\mu} * s_{\nu} \rangle = g_{\lambda\mu\nu}$.

Then

$$\begin{aligned}
 g_{\lambda\mu(d)\nu} + g_{\lambda\mu(d-1)\nu} &= \langle s_\lambda, s_{\mu(d)} * s_\nu \rangle + \langle s_\lambda, s_{\mu(d-1)} * s_\nu \rangle \\
 &= \langle s_\lambda, (s_{\mu(d)} + s_{\mu(d-1)}) * s_\nu \rangle \\
 &= \langle s_\lambda, (s_{(n-d)} \cdot s_{(1^d)}) * s_\nu \rangle \\
 &= \sum_{\beta \vdash d} \langle s_\lambda, (s_{(n-d)} * s_{\nu/\beta}) (s_{(1^d)} * s_\beta) \rangle \\
 &= \sum_{\beta \vdash d} \langle s_\lambda, s_{\nu/\beta} s_{\beta'} \rangle.
 \end{aligned}$$

Let $\nu/\beta \oplus \beta'$ be the skew Young diagram consisting of the disjoint union of ν/β and β' , translated to lie in distinct rows and columns (with β' to the southwest). By the Littlewood-Richardson rule, $\langle s_\lambda, s_{\nu/\beta} s_{\beta'} \rangle$ is the number of semistandard tableaux T of shape $\nu/\beta \oplus \beta'$ and content λ whose reverse (row) reading word is a ballot sequence. By transposing the part of T that lies in β' , barring each letter within, and combining with the part of T lying in ν/β , we arrive at a colored tableau $T_<$ of content λ , total color d (whose barred letters form the shape β), and shape ν such that $w(T_<)$ is a ballot sequence. Summing over all β gives the desired result. \square

We are now ready to prove a reformulated version of Hook Kronecker Rule I from [3].

Theorem 4.2. *Let λ and ν be partitions of n , and let $<$ be the natural order on \mathcal{A} given by*

$$\bar{1} < 1 < \bar{2} < 2 < \cdots < \bar{n} < n.$$

Then the Kronecker coefficient $g_{\lambda\mu(d)\nu}$ is the number of colored tableaux $T_<$ of content λ , total color d , and shape ν such that the total reverse (row-column) reading word $w(T_<)$ is a ballot sequence and the southwest corner of $T_<$ is unbarred.

Proof. By Lemma 3.1(c) and Proposition 4.1, the number of colored tableaux $T_<$ of content λ , total color d , and shape ν such that $w(T_<)$ is a ballot sequence is $g_{\lambda\mu(d)\nu} + g_{\lambda\mu(d-1)\nu}$. Denote this set by $\text{CYT}_{\lambda,d}^+(\nu)$, and write $\text{CYT}_{\lambda,d}^+(\nu)$ (resp. $\text{CYT}_{\lambda,d}^-(\nu)$) for the subset of these tableaux with barred (resp. unbarred) southwest corner.

Since i immediately follows \bar{i} in $<$ for all unbarred letters $i \in \mathcal{A}$, the southwest corner of $T_<$ can always be toggled from barred to unbarred or vice versa and remain a colored tableau. Moreover, this toggle does not change $w(T_<)$: it simply moves the last letter of $u(T_<)$ to be the first letter of $v(T_<)$ or vice versa. Hence $|\text{CYT}_{\lambda,d}^+(\nu)| = |\text{CYT}_{\lambda,d-1}^-(\nu)|$.

The result now follows easily, say, by induction on d : if $|\text{CYT}_{\lambda,d-1}^-(\nu)| = g_{\lambda\mu(d-1)\nu}$, then

$$\begin{aligned}
 g_{\lambda\mu(d)\nu} &= |\text{CYT}_{\lambda,d}(\nu)| - |\text{CYT}_{\lambda,d-1}^-(\nu)| \\
 &= |\text{CYT}_{\lambda,d}(\nu)| - |\text{CYT}_{\lambda,d}^+(\nu)| = |\text{CYT}_{\lambda,d}^-(\nu)|.
 \end{aligned}
 \quad \square$$

ACKNOWLEDGMENT

The author would like to thank Jonah Blasiak for useful discussions.

REFERENCES

- [1] C. M. Ballantine and R. C. Orellana, *On the Kronecker product $s_{(n-p,p)} * s_\lambda$* , Electron. J. Combin. **12** (2005), Research Paper 28, 26. MR2156682
- [2] Georgia Benkart, Frank Sottile, and Jeffrey Stroomer, *Tableau switching: algorithms and applications*, J. Combin. Theory Ser. A **76** (1996), no. 1, 11–43, DOI 10.1006/jcta.1996.0086. MR1405988
- [3] Jonah Blasiak, *Kronecker coefficients for one hook shape*, to appear in Sem. Lothar. Combin.
- [4] Jonah Blasiak, Ketan D. Mulmuley, and Milind Sohoni, *Geometric complexity theory IV: nonstandard quantum group for the Kronecker problem*, Mem. Amer. Math. Soc. **235** (2015), no. 1109, x+160, DOI 10.1090/memo/1109. MR3338303
- [5] Emmanuel Briand, Rosa Orellana, and Mercedes Rosas, *Quasipolynomial formulas for the Kronecker coefficients indexed by two two-row shapes (extended abstract)*, 21st International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2009), Discrete Math. Theor. Comput. Sci. Proc., AK, Assoc. Discrete Math. Theor. Comput. Sci., Nancy, 2009, pp. 241–252. MR2721516
- [6] Mark D. Haiman, *On mixed insertion, symmetry, and shifted Young tableaux*, J. Combin. Theory Ser. A **50** (1989), no. 2, 196–225, DOI 10.1016/0097-3165(89)90015-0. MR989194
- [7] A. Lascoux, *Produit de Kronecker des représentations du groupe symétrique*, Séminaire d'Algèbre Paul Dubreil et Marie-Paule Malliavin, 32ème année (Paris, 1979), Lecture Notes in Math., vol. 795, Springer, Berlin, 1980, pp. 319–329. MR582085 (82b:20016)
- [8] Jeffrey B. Remmel, *A formula for the Kronecker products of Schur functions of hook shapes*, J. Algebra **120** (1989), no. 1, 100–118, DOI 10.1016/0021-8693(89)90191-9. MR977863
- [9] Jeffrey B. Remmel and Tamsen Whitehead, *On the Kronecker product of Schur functions of two row shapes*, Bull. Belg. Math. Soc. Simon Stevin **1** (1994), no. 5, 649–683. MR1315363
- [10] Mercedes H. Rosas, *The Kronecker product of Schur functions indexed by two-row shapes or hook shapes*, J. Algebraic Combin. **14** (2001), no. 2, 153–173, DOI 10.1023/A:1011942029902. MR1867232

DEPARTMENT OF MATHEMATICS, NORTH CAROLINA STATE UNIVERSITY, RALEIGH, NORTH CAROLINA 27695

E-mail address: riliu@ncsu.edu