



Mariam Beriashvili, Ralf Schindler, Liuzhen Wu, Liang Yu

Hamel bases and well-ordering the continuum

Proceedings of the American Mathematical Society

DOI: 10.1090/proc/14010

Accepted Manuscript

This is a preliminary PDF of the author-produced manuscript that has been peer-reviewed and accepted for publication. It has not been copyedited, proofread, or finalized by AMS Production staff. Once the accepted manuscript has been copyedited, proofread, and finalized by AMS Production staff, the article will be published in electronic form as a “Recently Published Article” before being placed in an issue. That electronically published article will become the Version of Record.

This preliminary version is available to AMS members prior to publication of the Version of Record, and in limited cases it is also made accessible to everyone one year after the publication date of the Version of Record.

The Version of Record is accessible to everyone five years after publication in an issue.

Hamel bases and well-ordering the continuum

Mariam Beriashvili¹

Ivane Javakhishvili Tbilisi State University and Ilia Vekua Institute of Applied
Mathematics
Tbilisi, Georgia

Ralf Schindler²

Institut für Mathematische Logik und Grundlagenforschung, Universität Münster
Einsteinstr. 62, 48149 Münster, Germany

Liuzhen Wu³

Institute of Mathematics, Chinese Academy of Sciences
East Zhong Guan Cun Road No. 55, Beijing 100190, P. R. of China

Liang Yu⁴

Institute of Mathematical Sciences, Nanjing University
Nanjing, Jiangsu Province 210093, P. R. of China

Abstract

In ZF, the existence of a Hamel basis does not yield a well-ordering of \mathbb{R} .

Throughout this paper, by a *Hamel basis* we always mean a basis for \mathbb{R} , construed as a vector space over \mathbb{Q} . We denote by E the *Vitali equivalence relation*, xEy iff $x - y \in \mathbb{Q}$ for $x, y \in \mathbb{R}$. We also write $[x]_E = \{y : yEx\}$ for the E -equivalence class of x . A transversal for the set of all E -equivalence classes picks exactly one member from each $[x]_E$. The range of any such transversal is also called a *Vitali set*.

A set $\Lambda \subset \mathbb{R}$ is a *Luzin set* iff Λ is uncountable but $\Lambda \cap M$ is at most countable for every meager set $M \subset \mathbb{R}$. A set $S \subset \mathbb{R}$ is a *Sierpiński set* iff S is uncountable but $S \cap N$ is at most countable for every null set $N \subset \mathbb{R}$ (“null” in the sense of Lebesgue measure). A set $B \subset \mathbb{R}$ is a *Bernstein set* iff $B \cap P \neq \emptyset \neq P \setminus B$ for every perfect set $P \subset \mathbb{R}$.

It has been well-known for more than a century that the existence of a well-ordering of the reals implies the existence of all these “pathological” sets of reals:

¹The first author gratefully acknowledges support from the Marianne und Dr. Horst Kiesow-Stiftung, Frankfurt a.M.

²The second author was partially supported by the SFB 878 “Groups, geometry, and actions” from the DFG (Deutsche Forschungsgemeinschaft). Most of the results in this paper were obtained while the second author was an invited visiting researcher at the Centre de Recerca Matemàtica (CRM) in Bellaterra (Catalonia), and he would like to thank the CRM for their support and hospitality. This paper was typed while the second author visited Grigor Sargsyan in Warsaw (Poland) in Nov 2016, whom he would like to thank for his hospitality and support. He would also like to thank William Chan for his comments on an earlier version of this paper.

³The third author would like to acknowledge the support through the funding Projects NSFC 11321101 and 11401567.

⁴The fourth author gratefully acknowledges support from the National Natural Science Fund of China, No. 11322112 and 11671196 and from a Humboldt Research Fellowship for Experienced Researchers.

Hamel bases, Vitali sets, Luzin sets, Sierpiński sets,⁵ and Bernstein sets; see e.g. the thorough discussion in [4].⁶

D. Pincus and K. Prikry study the Cohen-Halpern-Lévy model H in [8] and show that there is a Luzin set in H , thereby establishing that in ZF, the existence of a Luzin set does not imply the existence of a well-ordering of the reals. We will recall their proof below, cf. Theorem 1.5.

In ZF, the existence of a Hamel basis implies the existence of a Vitali set of reals, cf. Lemma 1.1 below. Feferman had observed that H has a Vitali set, cf. [8, p. 433]. Pincus and Prikry ask:

“We would be interested in knowing whether a Hamel basis for \mathbb{R} over \mathbb{Q} (the rationals) exists in H or in any other model in which \mathbb{R} cannot be well ordered.” ([8, p. 433])

In [1], A. Blass shows that in ZF, if every vector space has a basis, then the axiom of choice holds true.

In the current paper we answer the question by Pincus and Prikry and show that H does have a Hamel basis. This will also give Feferman’s result as a corollary, cf. Corollary 2.4 below.

We shall also show that H has a Bernstein sets, cf. Theorem 1.7. There is no Sierpiński set in H , though, cf. Lemma 1.6. Therefore, in ZF not even the conjunction of the following statements (1), (3), (4), and (5) implies the existence of a well-ordering of the reals.

- (1) There is a Luzin set.
- (2) There is a Sierpiński set.
- (3) There is a Bernstein set.
- (4) There is a Vitali set.
- (5) There is a Hamel basis.

In a sequel to the current paper, in [10], it is shown that in ZF plus DC, (5) does not yield a well-ordering of the reals.

We would like to thank the referee for her/his comments on the first draft of this paper.

1 Warm ups.

In what follows, we shall sometimes think of reals as elements of the Baire space ${}^\omega\omega$, sometimes as elements of the Cantor space ${}^\omega 2$, and at other times think of them as actual reals. The attentive reader will have no problem sorting this out.

⁵To get Luzin and Sierpiński sets, one needs to make the additional hypothesis that CH holds true, unless e.g. one works with the concept of *generalized* Luzin and Sierpiński sets which arises from the concept of Luzin and Sierpiński sets by replacing “at most countable” with “smaller than the continuum” and works under Martin’s Axiom.

⁶A discussion of “paradoxical” decompositions of the unit ball à la Hausdorff and Banach–Tarski is beyond the scope of this paper, cf. also [4].

Let us first show that (4) implies (5). If X is a set of reals, then we write $\text{span}(X)$ for the set of all $\sum_{n=1}^m q_n \cdot x_n$, where $m \in \mathbb{N}$, $m \geq 1$, $q_n \in \mathbb{Q}$, and $x_n \in X$ for all n , $1 \leq n \leq m$. By convention, we also declare $\text{span}(\emptyset) = \{0\}$.

Lemma 1.1 (*Folklore*) *In ZF, if there is a Hamel basis, then there is a Vitali set.*

Proof. Let B be a Hamel basis. Let $1 = \sum_{k=1}^n q_k \cdot z_k$, where $q_k \in \mathbb{Q} \setminus \{0\}$ and $z_k \in B$ for $1 \leq k \leq n$. It is straightforward to verify that $\text{span}(B \setminus \{z_1\})$ is a Vitali set. \square (Lemma 1.1)

Let us now recall the Cohen–Halpern–Lévy model. We let \mathbb{C} denote Cohen forcing, i.e., the collection of all finite sequences of natural numbers, ordered by end–extension. If I is any index set, then $\mathbb{C}(I)$ denotes the finite support product of I many copies of \mathbb{C} , i.e., $p \in \mathbb{C}(I)$ iff $p(\ell) \in \mathbb{C}$ for $\ell \in I$ and

$$\text{supp}(p) = \{\ell \in I : p(\ell) \neq \emptyset\}$$

is finite. In what follows, $I \subseteq \omega$. If $I \cap J = \emptyset$, then $\mathbb{C}(I \cup J) \cong \mathbb{C}(I) \times \mathbb{C}(J)$.

Let us force with $\mathbb{C}(\omega)$ over L ,⁷ and let g be a generic filter. Let c_n , $n < \omega$, denote the Cohen reals which g adds. Let us write $A = \{c_n : n < \omega\}$ for the set of those Cohen reals. The model

$$H = H(L) = \text{HOD}_{A \cup \{A\}}^{L[g]}$$

of all sets which inside $L[g]$ are hereditarily definable from parameters in $\text{OR} \cup A \cup \{A\}$ is the Cohen–Halpern–Lévy model (over L), cf. [2, pp. 136–141], [3], and [8, p. 429]. As $L \subset H \subset L[g]$ and $\mathbb{C}(\omega)$ is countable, and hence trivially has the c.c.c., L , H , and $L[g]$ all have the same cardinals, and in particular $\omega_1^H = \omega_1^L$. It is well–known that in H , the reals cannot be well–ordered and in fact A has no countable subset, cf. e.g. [2, pp. 136–141] and Lemma 1.2 below. Here and in what follows, a set X is called *countable* iff there is some bijection $f : \omega \rightarrow X$, and X is called *at most countable* iff X is finite or countable.

In particular, the Continuum Hypothesis fails in H : the set $A \subset \mathbb{R} \cap H$ is not countable, but H can see no surjection from A onto $\mathbb{R} \cap H$.

For any finite $a \subset A$, we write $L[a]$ for the model constructed from the finitely many reals in a . Fixing some Gödelization of formulae (or some enumeration of all the rud functions, resp.) at the outset, each $L[a]$ comes with a unique canonical global well–ordering $<_a$ of $L[a]$ by which we mean the one which is induced by the *natural* order of the elements of a and the fixed Gödelization device in the usual fashion. The assignment $a \mapsto <_a$, $a \in [A]^{<\omega}$, is hence in H .⁸ This is a crucial fact.

Let us fix a bijection

$$(1) \quad e : \omega \rightarrow \omega \times \omega,$$

and let us write $((n)_0, (n)_1) = e(n)$.

We shall also make use the following.

⁷We might as well force over V rather than L , but forcing over L will simplify the notation a bit.

⁸More precisely, the ternary relation consisting of all (a, x, y) such that $x <_a y$ is definable over H .

Lemma 1.2 (1) Let $a \in [A]^{<\omega}$ and $X \subset L[a]$, $X \in H$, say $X \in \text{HOD}_{b \cup \{A\}}^{L[g]}$, where $b \supseteq a$, $b \in [A]^{<\omega}$. Then $X \in L[b]$.

(2) There is no well-ordering of the reals in H .

(3) A has no countable subset in H .

(4) $[A]^{<\omega}$ has no countable subset in H .

Proof sketch. (1) Every permutation $\pi: \omega \rightarrow \omega$ induces an automorphism e_π of $\mathbb{C}(\omega)$ by sending p to q , where $q(\pi(n)) = p(n)$ for all $n < \omega$. It is clear that no e_π moves the canonical name for A , call it \dot{A} . Let us also write \dot{c}_n for the canonical name for c_n , $n < \omega$. Now if a , and b are as in the statement of (1), say $b = \{c_{n_1}, \dots, c_{n_k}\}$, if $p, q \in \mathbb{C}(\omega)$, if $\pi \upharpoonright \{n_1, \dots, n_k\} = \text{id}$, $p \upharpoonright \{n_1, \dots, n_k\}$ is compatible with $q \upharpoonright \{n_1, \dots, n_k\}$, and $\text{supp}(\pi(p)) \cap \text{supp}(q) \subseteq \{n_1, \dots, n_k\}$, if $x \in L$, if $\alpha_1, \dots, \alpha_m$ are ordinals, and if φ is a formula, then

$$p \Vdash_L^{\mathbb{C}(\omega)} \varphi(\check{x}, \check{\alpha}_1, \dots, \check{\alpha}_m, \dot{c}_{n_1}, \dots, \dot{c}_{n_k}, \dot{A}) \iff \pi(p) \Vdash_L^{\mathbb{C}(\omega)} \varphi(\check{x}, \check{\alpha}_1, \dots, \check{\alpha}_m, \dot{c}_{n_1}, \dots, \dot{c}_{n_k}, \dot{A})$$

and $\pi(p)$ is compatible with q , so that the statement $\varphi(\check{x}, \check{\alpha}_1, \dots, \check{\alpha}_m, \dot{c}_{n_1}, \dots, \dot{c}_{n_k}, \dot{A})$ will be decided by conditions $p \in \mathbb{C}(\omega)$ with $\text{supp}(p) \subseteq \{n_1, \dots, n_k\}$. But every set in $L[b]$ is coded by a set of ordinals, so if X is as in (1), this shows that $X \in L[b]$.

(2) Every real is a subset of L . Hence by (1), if $L[g]$ had a well-ordering of the reals in $\text{HOD}_{a \cup \{A\}}^{L[g]}$, some $a \in [A]^{<\omega}$, then every real of H would be in $L[a]$, which is nonsense.

(3) Assume that $f: \omega \rightarrow A$ is injective, $f \in H$. Let $x \in {}^\omega\omega$ be defined by $x(n) = f((n)_0)((n)_1)$, so that $x \in H$. By (1), $x \in L[a]$ for some $a \in [A]^{<\omega}$. But then $\text{ran}(f) \subset L[a]$, which is nonsense, as there is some $n < \omega$ such that $c_n \in \text{ran}(f) \setminus a$.

(4) This readily follows from (3). □ (Lemma 1.2)

Let us recall another standard fact.

(2) If $a, b \in [A]^{<\omega}$, then $L[a] \cap L[b] = L[a \cap b]$.

To see this, let us assume without loss of generality that $a \setminus b \neq \emptyset \neq b \setminus a$, and say $a \setminus b = \{c_n : n \in I\}$ and $b \setminus a = \{c_n : n \in J\}$, where I and J are non-empty disjoint finite subsets of ω . Then $\mathbb{C}(I) \cong \mathbb{C} \cong \mathbb{C}(J)$, and $a \setminus b$ and $b \setminus a$ are mutually \mathbb{C} -generic over $L[a \cap b]$. But then $L[a] \cap L[b] = L[a \cap b][a \setminus b] \cap L[a \cap b][b \setminus a] = L[a \cap b]$, cf. [9, Problem 6.12].

For any $a \in [A]^{<\omega}$, we write $\mathbb{R}_a = \mathbb{R} \cap L[a]$ and $\mathbb{R}_a^+ = \mathbb{R}_a \setminus \bigcup \{\mathbb{R}_b : b \subsetneq a\}$. By [2, pp. 136–141], $(\mathbb{R}_a^+ : a \in [A]^{<\omega})$ is a partition of \mathbb{R} : By Lemma 1.2 (1),

(3)
$$\mathbb{R} \cap H = \bigcup \{\mathbb{R}_a^+ : a \in [A]^{<\omega}\},$$

and $\mathbb{R}_a \cap \mathbb{R}_b = \mathbb{R}_{a \cap b}$ by (2), so that

(4)
$$\mathbb{R}_a^+ \cap \mathbb{R}_b^+ = \emptyset \text{ for } a, b \in [A]^{<\omega}, a \neq b.$$

For $x \in \mathbb{R}$, we shall also write $a(x)$ for the unique $a \in [A]^{<\omega}$ such that $x \in \mathbb{R}_a^+$, and we shall write $\#(x) = \text{Card}(a(x))$.

Adrian Mathias showed that there is an H -definable function which assigns to each $x \in H$ an ordering $<_x$ such that $<_x$ is a well-ordering iff x can be well-ordered in H , cf. [6, p. 182]. This gives the following as a special simple case.

Lemma 1.3 (*A. Mathias*) *In H , the union of countably many countable sets of reals is countable.*

Proof. Let us work inside H . Let $(A_n : n < \omega)$ be such that for each $n < \omega$, $A_n \subset \mathbb{R}$ and there exists some surjection $f : \omega \rightarrow A_n$. For each such pair n , f let $y_{n,f} \in {}^\omega \omega$ be such that $y_{n,f}(m) = f((m)_0)((m)_1)$. If $a \in [A]^{<\omega}$ and $y_{n,f} \in \mathbb{R}_a$, then $A_n \in L[a]$. By (2), for each n there is a unique $a_n \in [A]^{<\omega}$ such that $A_n \in L[a_n]$ and $b \supset a_n$ for each $b \in [A]^{<\omega}$ such that $A_n \in L[b]$. Notice that A_n is also countable in $L[a_n]$.

Using the function $n \mapsto a_n$, an easy recursion yields a surjection $g : \omega \rightarrow \bigcup \{a_n : n < \omega\}$: first enumerate the finitely many elements of a_0 according to their natural order, then enumerate the finitely many elements of a_1 according to their natural order, etc. As A has no countable subset, $\bigcup \{a_n : n < \omega\}$ must be finite, say $a = \bigcup \{a_n : n < \omega\} \in [A]^{<\omega}$. But then $\{A_n : n < \omega\} \subset L[a]$. (We don't claim $(A_n : n < \omega) \in L[a]$.)

For each $n < \omega$, we may now let f_n the $<_a$ -least surjection $f : \omega \rightarrow A_n$. Then $f(n) = f_{(n)_0}((n)_1)$ for $n < \omega$ defines a surjection from ω onto $\bigcup \{A_n : n < \omega\}$, as desired. \square (Lemma 1.3)

Lemma 1.4 (1) ([5, Theorem 3.20]) *Let $a \in [A]^{<\omega}$. Then \mathbb{R}_a is a null set in H .*

(2) *If $B \subset \mathbb{R} \cap H$, $B \in H$, and B is countable in $L[g]$, then B is a null set in H .*

Proof sketch. (1) Let $\mathbb{R} = {}^\omega 2$ in this argument, with the addition $+$ being the componentwise addition in $\mathbb{Z}/2\mathbb{Z}$. Let $n < \omega$ be such that $c_n \notin a$. It suffices to prove that \mathbb{R}_a is null in $L[a \cup \{c_n\}]$.

In $L[a]$, let $\mathbb{R}_a = N \cup M$, where N is G_δ and null set, and M is F_σ and meager, cf. e.g. [7]. Inside $L[a \cup \{c_n\}]$, let us consider $N^* + c_n = \{x + c_n : x \in N^*\}$, where N^* is $L[a \cup \{c_n\}]$'s version of N .

Let $x \in \mathbb{R}_a$. As N is comeager in $L[a]$, $N + x$ is also comeager in $L[a]$, so that $c_n \in (N + x)^* = N^* + x$, see [9, Lemma 8.9 (2)], and hence $x \in N^* + c_n$. So $\mathbb{R}_a \subseteq N^* + c_n$. But N is null in $L[a]$, and hence N^* and $N^* + c_n$ are null in $L[a \cup \{c_n\}]$. \mathbb{R}_a is therefore contained in a null set of $L[a \cup \{c_n\}]$ and is hence itself null.

(2) Say $f : \omega \rightarrow B$, $f \in L[g]$, is an enumeration of B , and let $\tau \in L^{\mathbb{C}(\omega)}$ be such that $\tau^g = f$. Let us write $\tau(n)$ for the canonical name for $f(n)$ induced by τ . We aim to find $N \in H$, a G_δ null set in H with a code in L such that $B \subset N$. Let $h : \mathbb{C}(\omega) \times \omega \rightarrow \omega$ be bijective.

Let $m < \omega$. Set $\epsilon^m = \frac{1}{m+1}$ and $\epsilon_n^m = \frac{1}{2^{n+1}} \cdot \epsilon^m$ for $n < \omega$, so that $\sum_{n=0}^{\infty} \epsilon_n^m = \epsilon^m$.

Working in L , for each pair $(p, k) \in \mathbb{C}(\omega) \times \omega$, write $n = h((p, k))$, and let us pick some $q \in \mathbb{C}(\omega)$, $q \leq p$, and some $s \in {}^{<\omega} \omega$ such that $q \Vdash_L^{\mathbb{C}(\omega)} \check{s} \subset \tau(k)$, and $\mu(U_s) \leq \epsilon_n^m$, and write $\mathcal{O}_n^m = U_s$. (Here, U_s is the basis clopen set $\{x : x \supset s\}$.)

Set $\mathcal{O}^m = \bigcup \{\mathcal{O}_n^m : n < \omega\}$. For a given $k < \omega$, the set $\{q \in \mathbb{C}(\omega) : \exists n q \Vdash_L^{\mathbb{C}(\omega)} \tau(k) \in \mathcal{O}_n^m\}$ is dense, so that $f(k) = (\tau(k))^g \in \mathcal{O}_n^m$ for some n . In other words, $B \subset \mathcal{O}^m$.

Set $N = \bigcap_{m < \omega} \mathcal{O}^m$, to be interpreted in H . We have that N is a G_δ null set inside H with a code in L , and $B \subset N$. \square (Lemma 1.4)

Theorem 1.5 (*D. Pincus, K. Prikry*) *In H , there is a Luzin set.*

Proof. Let $\Lambda \in L$ be such that $L \models$ “ Λ is a Luzin set.” We aim to verify that Λ is Luzin in H . Λ is uncountable in L , so that also H can see a bijection of Λ with its own ω_1 , as $\omega_1^H = \omega_1^L$. In particular, Λ is uncountable in H .

By Lemma 1.3, it suffices to verify that inside H ,

$$(5) \quad \Lambda \setminus \mathcal{O} \text{ is at most countable,}$$

whenever \mathcal{O} is a dense union of countably many open intervals with rational end-points.

Let $((p_n, q_n) : n < \omega)$ be an enumeration of all open intervals with rational end-points, and let $X \subset \omega$, $X \in H$, be such that

$$H \models \text{“} \mathcal{O} = \bigcup \{(p_n, q_n) : n \in X\} \text{ is dense.”}$$

Let us suppose that (5) were not true in H for this fixed \mathcal{O} . As $\Lambda \in L$, inside H there must then be a bijection from ω_1 onto $\Lambda \setminus \mathcal{O}$, so that by $\omega_1^{L[g]} = \omega_1^H$ also

$$(6) \quad \Lambda \setminus \mathcal{O} \text{ is uncountable in } L[g].$$

Let $\tau \in L^{\mathbb{C}(\omega)}$ be a name for X , and let $p \in g$ be such that

$$p \Vdash_L^{\mathbb{C}(\omega)} \text{“} \Lambda \setminus \bigcup \{(p_n, q_n) : n \in \tau\} \text{ is uncountable.”}$$

As $\mathbb{C}(\omega)$ is countable, we may work in $L[g]$ and find some $q \in g$, $q \leq p$, such that for uncountably many $x \in \mathbb{R} \cap L$,

$$(7) \quad q \Vdash_L^{\mathbb{C}(\omega)} \text{“} \check{x} \in \Lambda \setminus \bigcup \{(p_n, q_n) : n \in \tau\} \text{.”}$$

Let us write U for the set of all $x \in \mathbb{R} \cap L$ with (7), so that U is an uncountable set of reals in L , and let

$$\mathcal{O}^* = \bigcup \{(p_n, q_n) : \exists r \leq q r \Vdash_L^{\mathbb{C}(\omega)} n \in \tau\},$$

as being defined in L .

Of course, $\mathcal{O}^* \supseteq \mathcal{O} \cap L$, so that \mathcal{O}^* is open and dense in L . As Λ is a Luzin set in L , $\Lambda \setminus \mathcal{O}^*$ must be countable in L .

We have a contradiction with (6).

\square (Theorem 1.5)

Lemma 1.6 *In H , there is no Sierpiński set.*

Proof. We shall prove that there is no set $S \in H$ of reals such that S is not at most countable in H and for each null set N of H , $S \cap N$ is at most countable.

Let us suppose that $S \in H$ is such a set. By Lemma 1.4, we cannot have that $S \subseteq \mathbb{R}_a$ for some $a \in [A]^{<\omega}$, because if this were true, then $S \cap \mathbb{R}_a = S$ and S itself would have to be at most countable.

Therefore, the set

$$F = \{a \in [A]^{<\omega} : S \cap \mathbb{R}_a^+ \neq \emptyset\}$$

is not finite. We may then inside H define the function $f: F \rightarrow \mathbb{R} \cap H$ by setting $f(a)$ to be the $<_a$ -least element of $S \cap \mathbb{R}_a^+$.

Write $B = \text{ran}(f)$. Then $B \in H$, and B is countable inside $L[g]$. By Lemma 1.4 (2), B is then a null set in H . Therefore, $B = S \cap B$ must be countable in H , i.e., there is some bijective $h \in H$, $h: \omega \rightarrow B$.

However, $((a, \mathbb{R}_a^+) : a \in [A]^{<\omega}) \in H$, so that $x \mapsto a(x)$ is in H , and hence $a \circ h \in H$, where $(a \circ h)(n) = a(h(n))$, $n < \omega$. Then $a \circ h: \omega \rightarrow [A]^{<\omega}$ is injective, which contradicts Lemma 1.2 (4). \square (Lemma 1.6)

Theorem 1.7 *In H , there is a Bernstein set.*

Proof. In this proof, let us think of reals as elements of the Cantor space ${}^\omega 2$. Let us work in H .

We let

$$B = \{x \in \mathbb{R} : \exists \text{ even } n (2^n < \#(x) \leq 2^{n+1})\} \quad \text{and} \\ B' = \{x \in \mathbb{R} : \exists \text{ odd } n (2^n < \#(x) \leq 2^{n+1})\}.$$

Obviously, $B \cap B' = \emptyset$.

Let $P \subseteq \mathbb{R}$ be perfect. We aim to see that $P \cap B \neq \emptyset \neq P \cap B'$.

Say $P = [T] = \{x \in {}^\omega 2 : \forall n x \upharpoonright n \in T\}$, where $T \subseteq {}^{<\omega} 2$ is a perfect tree. Modulo some fixed natural bijection ${}^{<\omega} 2 \leftrightarrow \omega$, we may identify T with a real. By (3), we may pick some $a \in [A]^{<\omega}$ such that $T \in L[a]$. Say $\text{Card}(a) < 2^n$, where n is even.

Let $b \in [A]^{2^{n+1}}$, $b \supset a$, and let $x \in \mathbb{R}_b^+$. In particular, $\#(x) = 2^{n+1}$. It is easy to work in $L[b]$ and construct some $z \in [T]$ such that $x \leq_T z \oplus T$,⁹ e.g., arrange that if $z \upharpoonright m$ is the k^{th} splitting node of T along z , where $k \leq m < \omega$, then $z(m) = 0$ if $x(k) = 0$ and $z(m) = 1$ if $x(k) = 1$.

If we had $\#(z) \leq 2^n$, then $\#(z \oplus T) \leq \#(z) + \#(T) < 2^n + 2^n = 2^{n+1}$, so that $\#(x) < 2^{n+1}$ by $x \leq_T z \oplus T$. Contradiction! Hence $\#(z) > 2^n$. By $z \in L[b]$, $\#(z) \leq 2^{n+1}$. Therefore, $z \in P \cap B$.

The same argument shows that $P \cap B' \neq \emptyset$. B (and also B') is thus a Bernstein set. \square (Theorem 1.7)

2 A Hamel basis.

The following is the main theorem of the current paper. Recall that for any $a \in [A]^{<\omega}$, we write $\mathbb{R}_a = \mathbb{R} \cap L[a]$. Let us now also write $\mathbb{R}_{<a} = \text{span}(\bigcup\{\mathbb{R}_b : b \subsetneq a\})$,

⁹Here, $(x \oplus y)(2n) = x(n)$ and $(x \oplus y)(2n+1) = y(n)$, $n < \omega$.

and $\mathbb{R}_a^* = \mathbb{R}_a \setminus \mathbb{R}_{<a}$. In particular, $\mathbb{R}_{<\emptyset} = \{0\}$ by our above convention that $\text{span}(\emptyset) = \{0\}$, and $\mathbb{R}_\emptyset^* = (\mathbb{R} \cap L) \setminus \{0\}$.

The proof of Claim 2.2 below will show that

$$(8) \quad \mathbb{R} \cap H = \text{span}\left(\bigcup\{\mathbb{R}_a^* : a \in [A]^{<\omega}\}\right).$$

Also, we have that $\mathbb{R}_a^* \subset \mathbb{R}_a^+$, so that by (3),

$$(9) \quad \mathbb{R}_a^* \cap \mathbb{R}_b^* = \emptyset \text{ for } a, b \in [A]^{<\omega}, a \neq b.$$

Theorem 2.1 *In H , there is a Hamel basis.*

Proof. We call $X \subset \mathbb{R}_a^*$ *linearly independent over $\mathbb{R}_{<a}$* iff whenever

$$\sum_{n=1}^m q_n \cdot x_n \in \mathbb{R}_{<a},$$

where $m \in \mathbb{N}$, $m \geq 1$, and $q_n \in \mathbb{Q}$ and $x_n \in X$ for all n , $1 \leq n \leq m$, then $q_1 = \dots = q_m = 0$. In other words, $X \subset \mathbb{R}_a^*$ is linearly independent over $\mathbb{R}_{<a}$ iff

$$\text{span}(X) \cap \mathbb{R}_{<a} = \{0\}.$$

We call $X \subset \mathbb{R}_a^*$ *maximal linearly independent over $\mathbb{R}_{<a}$* iff X is linearly independent over $\mathbb{R}_{<a}$ and no $Y \supsetneq X$, $Y \subset \mathbb{R}_a^*$ is still linearly independent over $\mathbb{R}_{<a}$. In particular, $X \subset \mathbb{R}_\emptyset^* = (\mathbb{R} \cap L) \setminus \{0\}$ is linearly independent over $\mathbb{R}_{<\emptyset} = \{0\}$ iff X is a Hamel basis for $\mathbb{R} \cap L$.

For any $a \in [A]^{<\omega}$, we let b_a denote the $<a$ -least set $X \subset \mathbb{R}_a^*$, $X \in L[a]$, which is maximal linearly independent over $\mathbb{R}_{<a}$. By the above crucial fact, the function $a \mapsto b_a$ is well-defined and *exists inside H* . In particular,

$$B = \bigcup\{b_a : a \in [A]^{<\omega}\}$$

is an element of H .

We claim that B is a Hamel basis for the reals of H , which will be established by Claims 2.2 and 2.3.

Claim 2.2 $\mathbb{R} \cap H \subset \text{span}(B)$.

Proof of Claim 2.2. Assume not, and let $n < \omega$ be the least size of some $a \in [A]^{<\omega}$ such that $\mathbb{R}_a^* \setminus \text{span}(B) \neq \emptyset$. Pick $x \in \mathbb{R}_a^* \setminus \text{span}(B) \neq \emptyset$, where $\text{Card}(a) = n$.

We must have $n > 0$, as b_\emptyset is a Hamel basis for the reals of L . Then, by the maximality of b_a , while b_a is linearly independent over $\mathbb{R}_{<a}$, $b_a \cup \{x\}$ cannot be linearly independent over $\mathbb{R}_{<a}$. This means that there are $q \in \mathbb{Q}$, $q \neq 0$, $m \in \mathbb{N}$, $m \geq 1$, and $q_n \in \mathbb{Q} \setminus \{0\}$ and $x_n \in b_a$ for all n , $1 \leq n \leq m$, such that

$$z = q \cdot x + \sum_{n=1}^m q_n \cdot x_n \in \mathbb{R}_{<a}.$$

By the definition of $\mathbb{R}_{<a}$ and the minimality of n , $z \in \text{span}(\bigcup\{b_c : c \subsetneq a\})$, which then clearly implies that $x \in \text{span}(\bigcup\{b_c : c \subseteq a\}) \subset \text{span}(B)$.

This is a contradiction!

□ (Claim 2.2)

Claim 2.3 B is linearly independent.

Proof of Claim 2.3. Assume not. This means that there are $1 \leq k < \omega$, $a_i \in [A]^{<\omega}$ pairwise different, $m_i \in \mathbb{N}$, $m_i \geq 1$ for $1 \leq i \leq k$, and $q_n^i \in \mathbb{Q} \setminus \{0\}$ and $x_n^i \in b_{a_i}$ for all n , $1 \leq n \leq m_i$, and i , $1 \leq i \leq k$, such that

$$(10) \quad \sum_{n=1}^{m_1} q_n^1 \cdot x_n^1 + \dots + \sum_{n=1}^{m_k} q_n^k \cdot x_n^k = 0.$$

By the properties of b_{a_i} , $\sum_{n=1}^{m_i} q_n^i \cdot x_n^i \in \mathbb{R}_{a_i}^*$, so that (10) buys us that there are $z_i \in \mathbb{R}_{a_i}^*$, $z_i \neq 0$, $1 \leq i \leq k$, such that

$$(11) \quad z_1 + \dots + z_k = 0.$$

There must be some i such that there is no j with $a_j \supseteq a_i$, which implies that $a_j \cap a_i \subsetneq a_i$ for all $j \neq i$. Let us assume without loss of generality that $a_j \cap a_1 \subsetneq a_1$ for all j , $1 < j \leq k$.

Let $a_1 = \{c_\ell : \ell \in I\}$, where $I \in [\omega]^{<\omega}$, and let $a_j \cap a_1 = \{c_\ell : \ell \in I_j\}$, where $I_j \subsetneq I$, for $1 < j \leq k$.

In what follows, a nice name τ for a real is a name of the form

$$(12) \quad \tau = \bigcup_{n,m < \omega} \{(n, m)^\vee\} \times A_{n,m},$$

where each $A_{n,m}$ is a maximal antichain of conditions of the forcing in question deciding that $\tau(\check{n}) = \check{m}$.

We have that z_1 is $\mathbb{C}(I)$ -generic over L , so that we may pick a nice name $\tau_1 \in L^{\mathbb{C}(I)}$ for z_1 with $(\tau_1)^{g \upharpoonright I} = z_1$. Similarly, for $1 < j \leq k$, z_j is $\mathbb{C}(I_j)$ -generic over $L[g \upharpoonright (\omega \setminus I)]$, so that we may pick a nice name $\tau_j \in L[g \upharpoonright (\omega \setminus I)]^{\mathbb{C}(I_j)}$ for z_j with $(\tau_j)^{g \upharpoonright I_j} = z_j$. We may construe each τ_j , $1 < j \leq k$, as a name in $L[g \upharpoonright (\omega \setminus I)]^{\mathbb{C}(I)}$ by replacing each $p: I_j \rightarrow \mathbb{C}$ in an antichain as in (12) by $p': I \rightarrow \mathbb{C}$, where $p'(\ell) = p(\ell)$ for $\ell \in I_j$ and $p'(\ell) = \emptyset$ otherwise. Let $p \in g \upharpoonright I$ be such that

$$p \Vdash_{L[g \upharpoonright (\omega \setminus I)]}^{\mathbb{C}(I)} \tau_1 + \tau_2 + \dots + \tau_k = 0.$$

We now have that inside $L[g \upharpoonright (\omega \setminus I)]$, there are nice $\mathbb{C}(I)$ -names τ'_j , $1 < j \leq k$ (namey, τ_j , $1 < j \leq k$), such that still inside $L[g \upharpoonright (\omega \setminus I)]$

$$(1) \quad p \Vdash^{\mathbb{C}(I)} \tau_1 + \tau'_2 + \dots + \tau'_k = 0, \text{ and}$$

$$(2) \quad \text{for all } j, 1 < j \leq k \text{ and for all } p \text{ in one of the antichains of the nice name } \tau'_j, \text{ supp}(p) \subseteq I_j.$$

Both (1) and (2) are arithmetic in real codes for $\tau_1, \tau'_2, \dots, \tau'_k$, so that by $\tau_1 \in L^{\mathbb{C}(I)}$ and Σ_1^1 -absoluteness between L and $L[g \upharpoonright (\omega \setminus I)]$ there are inside L nice $\mathbb{C}(I)$ -names τ'_j , $1 < j \leq k$, such that in L , (1) and (2) hold true. But then, writing $z'_j = (\tau'_j)^{g \upharpoonright I}$, we have by (2) that $z'_j \in \mathbb{R}_{I_j}$ for $1 < j \leq k$, and $z_1 + z'_2 + \dots + z'_k = 0$ by (1). But then $z_1 \in \mathbb{R}_I^* \cap \mathbb{R}_{<I}$, which is absurd. \square (Claim 2.3)

This finishes the proof of Theorem 2.1. \square (Theorem 2.1)

In the light of Lemma 1.1, Theorem 2.1 reproves Feferman's result.

Corollary 2.4 (*S. Feferman*) *In H , there is a Vitali set.*

References

- [1] Blass, A., *Existence of bases implies the axiom of choice*, Contemporary Mathematics **31** (1984), pp. 31–33.
- [2] Cohen, P., *Set theory and the continuum hypothesis*, Benjamin, New York 1966.
- [3] Halpern, J.D., and Lévy, A., *The Boolean prime ideal theorem does not imply the axiom of choice*, Proc. Sympos. Pure Math. **13** part I, Amer. Math. Soc., Providence, R.I., 1971, pp. 83–134.
- [4] Herrlich, H., *Axiom of choice*, Lecture Notes in Mathematics **1876**, Springer–Verlag 2006.
- [5] Kunen, K., *Random and Cohen reals*, Handbook of set–theoretic topology (K. Kunen and J. Vaughan, eds.), North Holland, Amsterdam, 1984, pp. 887–911.
- [6] Mathias, A., *The order extension principle*, in: Proceedings of Symposia in Pure Mathematics vol. 13 part II: Axiomatic Set Theory, T. Jech (ed.), American Mathematical Society, 1974.
- [7] Oxtoby, J.C., *Measure and Category*, Springer–Verlag 1971.
- [8] Pincus, D., and Prikry, K., *Luzin sets and well ordering the continuum*, Proc. Americ. Math. Soc. **49** (2), 1975, pp. 429–435.
- [9] Schindler, R., *Set theory. Exploring independence and truth*, Springer–Verlag 2012.
- [10] Schindler, R., Wu, L., and Yu, L., *Hamel bases and the principle of dependent choice*, preprint, available at https://ivv5hpp.uni-muenster.de/u/rds/hamel_basis_2.pdf