RELAXATION OSCILLATIONS¹

BY

J. LASALLE

Princeton University

1. The equation

$$\frac{d^2x}{d\tau^2} + \mu f(x) \frac{dx}{d\tau} + x = 0 \quad (1)$$

has been studied by E. and H. Cartan [1], van der Pol [2]–[4], Liénard [5], Levinson and Smith [6] and Haag [7], among others. A large number of intriguing and interesting applications of (1) may be found in the work of van der Pol and in [8].

In this paper we study a class of relaxation oscillations related to (1) which is simpler though not included in the general equation of Levinson and Smith [6]. As has been done for van der Pol's special case, $f(x) = x^2 - 1$, by Flanders and Stoker [9], we give the explicit construction of regions $E(\mu)$ of the phase plane each of which contains a single periodic solution. We have not strived for generality nor included possible refinements of the regions but hope to illustrate the relationship between solving differential inequalities and the construction of such regions. We do abandon the assumption that $f(x)$ be even, as the asymmetric case is of practical interest.

In Sec. 2 we consider a simple physical example of the equation to be studied and anticipate some of the results by heuristic arguments. The regions are constructed in Sec. 3 and the uniqueness of the periodic solutions within each region is proved in Sec. 4. In Sec. 5 we show how bounds on periods and amplitudes and their asymptotic values can be computed.

2. Consider the differential equation

$$L \frac{d^2q}{dt'^2} + F\left(\frac{dq}{dt'}\right) + q/C = 0 \quad (2)$$

for an LRC circuit with R non-linear and non-passive. $F(i)$ is the voltage-current characteristic of R , and q is the charge. A suitable choice of units for q and t' (time)— $q = (LC)^{1/2} z$ and $t' = (LC)^{1/2} \tau$ —gives Rayleigh's equation

$$\frac{d^2z}{d\tau^2} + \mu F\left(\frac{dz}{d\tau}\right) + z = 0, \quad \text{where} \quad \mu = (C/L)^{1/2}. \quad (3)$$

Liénard's equation (1) is equivalent to (3). It can be obtained from (3) by differentiating

¹Received Aug. 19, 1948. This work was carried out under an Office of Naval Research Contract. The author wishes in particular to thank Professor Lefschetz for his assistance and constant encouragement.

and placing $x = dz/d\tau$ and $f(x) = F'(x)$ ($f(x)$ is the dynamic resistance of R). Conversely, (2) can be obtained from (3) by integration. Also (1) and (3) are equivalent to the systems

$$\frac{dw}{dt} = -f(x)w - \epsilon^2 x \quad (4)$$

$$\frac{dx}{dt} = w,$$

$$\frac{dx}{dt} = -F(x) - y = G(x) - y \quad (5)$$

$$\frac{dy}{dt} = \epsilon^2 x,$$

FIG. 1.

where $\epsilon = 1/\mu = (L/C)^{1/2}$, $t = \mu\tau = t'/C$, $y = \mu z = q/L$, $x = dz/d\tau = dq/dt'$, $G(x) = -F(x)$, and $f(x) = F'(x)$. In the physical problem x represents current. Its units have remained unchanged.

Although it will be more convenient to work in the (x, y) phase plane, let us first examine the behavior of the paths (solutions) in the (x, w) plane with $f(x) = x^2 - 1$, van der Pol's case. The dotted curve in Fig. 1 is the isocline Γ_0 , corresponding to $dw/dt = 0$; i.e. $w = -\epsilon^2 x/f(x)$. Along the x -axis, $dx/dt = 0$, and the slopes of the paths have the general aspect indicated by the arrows. For ϵ quite small the curve Γ_0 approaches the lines $x = \pm 1$ and $w = 0$. When $(1 - x^2) \gg \epsilon^2 x$, i.e., not too close to Γ_0 , then $dw/dx = (1 - x^2)$ and $w = x - x^3/3 + c$. Starting with a path through a point such as A , the path is approximately $w = x - x^3/3 + c$ until reaching the point B . From B the path is confined to move between $w = 0$ and Γ_0 until reaching C at $(1, 0)$. Here the path leaves Γ_0 and is now approximately $w = x - x^3/3 - 2/3$ from C through D until reaching $w = 0$ at E . As before, the path is practically horizontal from E to C' where it follows $w = x - x^3/3 + 2/3$ until reaching E' . Here it begins to repeat the cycle Γ through $DED'E'$. Similarly, starting inside Γ the path would approach Γ , and the limit cycle is stable. Thus we expect the amplitude of the oscillation (the maximum value of $|x|$) to approach 2 as ϵ becomes small. From E' to C , dx/dt is small and x decreases slowly from 2 to 1 and then drops rapidly from 1 to -2 , giving rise to the relaxation oscillation pictured in Fig. 2.

FIG. 2.

The (x, y) phase plane has the advantage that the isoclines $dx/dt = 0$ and $dy/dt = 0$ do not depend upon ϵ , and the curve $w = -\int f(x) dx = G(x) + c$ becomes a horizontal line. From Eqs. (5) we see that the curve

$$\Delta : y = G(x)$$

corresponds to the isocline $dx/dt = 0$, and $x = 0$ to $dy/dt = 0$. Consider a $G(x)$ such as that in Fig. 3.

In the region I (above Δ and $x > 0$) and in the region III (below Δ and $x < 0$) the paths have positive slopes. In regions II and IV the slopes are negative, and the paths encircle the origin in a counter-clockwise direction. When ϵ is very small we see from

(5) that the path through a point such as A will be almost horizontal until reaching Δ . If the slope of Δ is positive, the path will cross Δ and continue moving horizontally until approaching Δ again. At B the slope of Δ is negative and the path tends to follow Δ ;

FIG. 3.

since if it moved away from Δ it would tend to move horizontally to the left and hence back towards Δ . Upon reaching a maximum point P_2 , the path leaves Δ , is horizontal until returning to Δ and so on. Thus the path follows Γ_2 and we expect there to be periodic solutions (limit cycles) which lie near Γ_2 when ϵ is small. The limit cycles which approach Γ_2 would correspond to stable² periodic solutions. In a similar manner by letting $t \rightarrow -\infty$, i.e., moving in the direction of decreasing time, a curve such as Γ_1 is obtained and we expect it to be approached by unstable periodic solutions as $\epsilon \rightarrow 0$. The function $G(x)$ of Fig. 3 represents a case of so called "hard oscillating" conditions (see [8]). In order that the system approach a steady oscillation, it must initially be driven outside the region enclosed by Γ_1 .

3. Let us make two simplifying observations. We see that Eqs. (5) are unchanged if y is replaced by $-y$, x by $-x$, and $G(-x)$ by $-G(x)$. Geometrically, this means turning our picture of the phase plane upside down (through 180°). Hence we can limit our discussion to the right half-plane $x \geq 0$, and then by this transformation apply the result to the left half-plane $x \leq 0$. Also if in (5) t is replaced by $-t$ and y by $-y$, we obtain a system of the same form with $-G(x)$ in place of $G(x)$. Thus, upon replacing $G(x)$ by $-G(x)$, in a theorem which concerns stable periodic solutions, we obtain a corresponding theorem for unstable periodic solutions.

²Throughout we use the term *stability* to mean "absolute orbital stability" as defined in [11]. By unstable we mean that stability is obtained if t is replaced by $-t$.

The term *enclosure* will refer to a region of the (x, y) phase plane which contains a periodic solution. To be more specific, our enclosures will be annular closed regions E (see Fig. 4) whose boundaries are a pair of simple, closed, sectionally-smooth curves C_0 and C_i , with C_i inside C_0 .

FIG. 4.

These boundaries are constructed to have the property that a path γ through a point P of the boundary enters E ; i.e., if at $t = t_0$, γ is at P , then for all $t > t_0$ γ is in E . If there are no critical points in E , it follows from the Poincaré-Bendixson theorem (see, e.g., [11]), that E is an enclosure.

We assume the behavior of Fig. 5, i.e. that:

- I. $G'(x)$ is continuous for all x , and $G(0) = 0$;
- II. there exist points $a_1 < a_2 < 0 < a_3 < a_4$ such that
 - (1) $G(a_1) = G(a_3)$ and $G(a_2) = G(a_4)$,
 - (2) $G(a_2) \leq G(x) \leq G(a_3)$ for $a_1 \leq x \leq a_4$,
 - (3) $G'(a_1) < 0$ and $G'(a_4) < 0$.

These properties of $G(x)$ enable us to construct the following simple, closed curve $\Gamma(a)$. Define

$$\begin{aligned}\Phi_a^+(x) &= \text{Max}_{\sigma \leq x \leq a} [G(\sigma)], \\ \Phi_a^-(x) &= \text{Min}_{a \leq \sigma \leq x} [G(\sigma)].\end{aligned}\tag{6}$$

$\Gamma(a)$ is defined to be the simple, closed curve formed by $\Phi_a^+(x)$ and $\Phi_a^-(x)$ (see Fig. 5). Γ_2 of Fig. 3 was an example of such a curve. We use the letter "a" to refer to the

dependence of $\Gamma(a)$ upon the particular set $a = (a_1, a_2, a_3, a_4)$ satisfying II (1) and (2). The manner in which II (3) enters the discussion will be seen later.

FIG. 5.

Let S and T be two regions of the phase plane. Define

$$d(S, T) = \text{l.u.b.}_{s \in S} \left\{ \text{g.l.b.}_{t \in T} d(s, t) \right\}^3$$

and

$$\delta(S, T) = d(S, T) + d(T, S).$$

By definition $S(\epsilon) \rightarrow T$ as $\epsilon \rightarrow 0$ if and only if $\delta(S(\epsilon), T) \rightarrow 0$ as $\epsilon \rightarrow 0$.

Theorem 1. If $G(x)$ satisfies I and II and $0 < \epsilon < \epsilon_0$, an enclosure $E(a, \epsilon)$ containing a stable periodic solution of (5) can be constructed. As $\epsilon \rightarrow 0$, $E(a, \epsilon) \rightarrow \Gamma(a)$.

Proof. Let us represent a simple closed curve C in the form

$$\begin{aligned} x &= f(\theta), \\ C : \quad y &= \psi(\theta) + g(x). \end{aligned} \tag{7}$$

Then

$$\bar{n} = \left(\frac{dy}{d\theta}, -\frac{dx}{d\theta} \right)$$

is a normal vector to C , and

$$\bar{v} = \left(\frac{dx}{dt}, \frac{dy}{dt} \right) = (G(x) - y, \epsilon^2 x)$$

is the phase velocity vector for a path γ through (x, y) . Define $\lambda(\theta)$ to be the scalar product of $\bar{n}$ and $\bar{v}$, i.e.,

³ $d(s, t)$ is the ordinary (Euclidean) distance between two points.

$$\begin{aligned}\lambda = [G(x) - \psi(\theta) - g(x)][\psi'(\theta) + g'(x)f'(\theta)] \\ - \epsilon^2 f(\theta)f'(\theta).\end{aligned}\quad (8)$$

The sign of λ is the sign of the projection of $\bar{v}$ on $\bar{n}$. The curves of interest to us are those for which $\lambda \geq 0$ or $\lambda \leq 0$. In the construction of the enclosure for this theorem we confine ourselves to the simpler solutions of the differential inequality.

Solution 1. Take $f(\theta) = \theta$ and $g(x) = 0$. Then

$$\lambda = [G(x) - \psi(x)]\psi'(x) - \epsilon^2 x.$$

(a) For $\psi(\theta) = A$, we have $D = -\epsilon^2 x$, and

$$C : y = A. \quad (9)$$

Hence $\lambda \leq 0$ for $x \geq 0$, and since the normal to C is vertical and downward, the paths for $x > 0$ cross horizontal lines from below. This, of course, we knew already. (See Fig. 3).

(b) Taking $\psi(x)\psi'(x) + \epsilon^2 x = A\psi'(x)$, we have

$$C : y = A \pm \epsilon(c^2 - x^2)^{1/2} \quad (10)$$

and

$$\lambda = \psi'(x)[G(x) - A] = \frac{\mp x}{(c^2 - x^2)^{1/2}} [G(x) - A].$$

The curve C is an ellipse with center at $(0, A)$ with c and ϵc as horizontal and vertical semi-axes. Along the lower-half of the ellipse the normal vector $\bar{n}$ is outward and on the upper-half, inward. Thus, for a particular x , we see from the expression for λ that the paths enter the ellipse if the horizontal axis of the ellipse is above Δ and leave the ellipse if the horizontal axis is below Δ .

Solution 2. Let $f(\theta) = \theta$ and $g(x) = \Phi_\alpha^+(x)$ (see (6)).

(a) $\psi(x) = 0$. Then

$$C : y = \Phi_\alpha^+(x), \quad 0 \leq x \leq \alpha,$$

$$\lambda = -\epsilon^2 x$$

and $\lambda \leq 0$ for $x \geq 0$.

(b) $\psi(x) = \epsilon(\alpha^2 - x^2)^{1/2}$, i.e., $\psi(x)\psi'(x) + \epsilon^2 x = 0$. Here

$$\lambda = -\epsilon(\alpha^2 - x^2)^{1/2}\Phi_\alpha^{+'}(x) + \frac{\epsilon x}{(\alpha^2 - x^2)^{1/2}} [\Phi_\alpha^+(x) - G(x)]$$

and from the definition of $\Phi_\alpha^+(x)$ it is clear that $\lambda \geq 0$ when $x \geq 0$. Hence the paths cross

$$C : y = \Phi_\alpha^-(x) + \epsilon(\alpha^2 - x^2)^{1/2}, \quad 0 \leq x \leq \alpha, \quad (11)$$

from above.

The inner boundary (see Fig. 6). The inner boundary will be a simple, closed (sectionally smooth) curve $C_i(a, \epsilon)$ with the paths crossing $C_i(a, \epsilon)$ from its interior to exterior, the paths being tangent ($\lambda = 0$) at only a finite number of points.

(a) *Below Δ and $x \geq 0$. (From A to B).* Draw the line L_1 through $A : (0, G(a_2))$ with slope ϵ . The first intersection of L_1 with Δ is called $B : (b, G(b))$. It follows from II (2) that for ϵ sufficiently small the horizontal line from B to M lies below Δ and $b > a_3$. Take

FIG. 6.

$$C_i(a, \epsilon) : y = G(b) - \epsilon(b^2 - x^2)^{1/2}, \quad 0 \leq x \leq b.$$

This is an ellipse with center at M and semi-axes MA and MB .

(b) *Above or along Δ and $x \geq 0$. (From B to C).*

$$C_i(a, \epsilon) : y = \Phi_b^+(x), \quad 0 \leq x \leq b.$$

(c) *Above Δ and $x \leq 0$. (From C to D).* Draw the line L_2 of slope ϵ through $C : (0, G(a_3))$. Its first intersection with Δ is $D : (d, G(d))$. For ϵ sufficiently small DN is above Δ and we take

$$C_i(a, \epsilon) : y = G(d) + \epsilon(d^2 - x^2)^{1/2}, \quad d \leq x \leq 0.$$

(d) *Below or along Δ and $x \leq 0$. (From D to A).*

$$C_i(a, \epsilon) : y = \Phi_d^-(x), \quad d \leq x \leq 0.$$

From solutions 1(b) and 2(a), and the remark made at the beginning of this section concerning the relation between the regions $x \geq 0$ and $x \leq 0$, the above is seen to be a suitable inner boundary.

The outer boundary: The outer boundary $C_0(a, \epsilon)$ will also be a simple closed (sectionally smooth) curve with the paths crossing $C_0(a, \epsilon)$ from its exterior to interior.

Again $\lambda = 0$ at only a finite number of points. Referring to Fig. 6 and II (3), we note first that within neighborhoods of $(a_1, G(a_1))$ and $(a_4, G(a_4))$ the inverse functions $G^{-1}(y)$ exist and have continuous derivatives. Consider the curves

$$C_1 : \xi = -G^{-1}(\eta + G(a_1))$$

and

$$C_4 : \xi = G^{-1}(-\eta + G(a_4)).$$

Figure 7 is the case $-a_1 = a_4$. Draw the line L of slope $d\xi/d\eta = 1/\epsilon$. Since the slopes of C_1 and C_4 are bounded within a neighborhood of $\eta = 0$, it will be true for ϵ sufficiently small that, starting from 0 and going out along L , L will intersect both C_1 and C_4 and these intersections will be consecutive (or coincide). We assume that L intersects C_1 first and then C_4 or that they coincide, and begin the construction at A (Fig. 6). Otherwise we would start at C .

(a) Below Δ and $x \geq 0$. (From P to Q).

FIG. 7.*

*In Fig. 7, the relation between a_1 and a_4 should be $-a_1 = a_4$.

The line L of slope ϵ through A intersects Δ for ϵ sufficiently small. Call this first intersection $Q : (q, G(q))$.

$$C_0(a, \epsilon) : y = G(q), \quad 0 \leq x \leq q$$

($q = \overline{PQ}$ of Fig. 6 or 7).

(b) *Above Δ and $x \geq 0$. (From Q to R).*

$$C_0(a, \epsilon) : y = G(q) + \epsilon(q^2 - x^2)^{1/2}, \quad 0 \leq x \leq q.$$

(c) *Above Δ and $x \leq 0$. (From R to S). The horizontal line from R to the left (Fig. 6) will intersect Δ for ϵ sufficiently small. Let this intersection be $S : (s, G(s))$. As noted above we have for ϵ sufficiently small that $-s = \overline{RS} < q = \overline{PQ}$. Take*

$$C_0(a, \epsilon) : y = G(s) = G(a_3) + \epsilon q, \quad s \leq x \leq 0.$$

(d) *Below Δ and $x \leq 0$. (From S to T).*

$$C_0(a, \epsilon) : y = \Phi_s^-(x) - \epsilon(s^2 - x^2)^{1/2}, \quad s \leq x \leq 0.$$

T is the point $(0, G(a_2) + \epsilon s)$.

(e) *Along $y = 0$. The curve is closed by the vertical line segment from T to P . Since $-s < q$, T is above P .*

From solutions 1(a) and 2(b) and the manner in which the paths cross the y -axis, the above is a suitable outer boundary.

The only critical point of the system (5) is the origin $(0, 0)$ and this point is not in $E(a, \epsilon)$. Therefore, as discussed previously, $E(a, \epsilon)$ contains a stable periodic solution and is an enclosure. Also it is clear that $E(a, \epsilon) \rightarrow \Gamma(a)$ as $\epsilon \rightarrow 0$. As a matter of fact $\delta(\Gamma(a), E(a, \epsilon)) = O(\epsilon)$.

For any given case the ϵ_0 of Th. 1 is the least upper bound of the ϵ meeting the conditions stated in the above construction. For van der Pol's equation ($G(x) = x - x^3/3$),

$$\epsilon_0 = \frac{2(3)^{1/2}}{9} < \frac{1}{2.6}.$$

Fig. 8 shows the enclosure for $\epsilon = 1/3$.

4. In van der Pol's special case it is possible to construct an enclosure for any $\epsilon > 0$ and thus to prove the existence of a stable periodic solution for each $\epsilon > 0$. Liénard [5] has, however, proved more; namely, that there exists for each $\epsilon > 0$ a unique periodic solution, and that this solution is stable. It is certainly not true in general that the periodic solution is unique. We might, however, expect it to be unique within $E(a, \epsilon)$. We now prove that this is the case.

Along a path γ of the system of equations

$$\frac{dx}{dt} = G(x) - y \tag{5}$$

$$\frac{dy}{dt} = \epsilon^2 x,$$

$$\frac{1}{2} \frac{d}{dt} (\epsilon^2 x^2 + y^2) = \epsilon^2 x G(x). \tag{12}$$

FIG. 9*

Theorem 2. If $G(x)$ satisfies I of Sec. 3 and if, in addition,

$$(a) \quad G(x) \geq G(c_0) \quad \text{for} \quad 0 \leq x \leq c_0,$$

$$G(x) \leq G(e_0) \quad \text{for} \quad e_0 \leq x \leq 0,$$

$$(b) \quad G(x) < G(c_0) \quad \text{for} \quad c_0 < x < c_1,$$

$$G(x) > G(e_0) \quad \text{for} \quad e_1 < x < e_0;$$

$$\text{then} \quad U(b) < U(b_0) \quad \text{for} \quad b_1 \leq b < b_0 < 0.$$

Proof. Define (see Fig. 9)

$$U^+(b) = \int_B^D G(x) dy$$

and

$$U^-(b) = \int_D^F G(x) dy,$$

so that

$$U(b) = U^+(b) + U^-(b).$$

*The point F in Fig. 9 should be marked $F:(0, f)$.

Note also that

$$U^+(b) = \int_B^D [G(x) - G(c_0)] dy + \delta(b)G(c_0), \quad (15)$$

where $\delta(b) = d - b$ and the integral is along $\gamma(b)$. The paths $\gamma(b)$ do not intersect, and therefore $\delta(b)$ increases as b decreases. Let $\gamma_1(b)$ be that part of $\gamma(b)$ corresponding to $0 < x < c_0$ and $\gamma_2(b)$ that part of $\gamma(b)$ corresponding to $x > c_0$. On $\gamma_1(b)$, $G(x) - G(c_0) \geq 0$ and for the same value of x , we see from (5), since $c > c_0$, that

$$\frac{dy}{dx} (\text{on } \gamma_1(b)) < \frac{dy}{dx} (\text{on } \gamma(b_0)), \quad \text{above } \Delta$$

and

$$\frac{-dy}{dx} (\text{on } \gamma_1(b)) < \frac{-dy}{dx} (\text{on } \gamma(b_0)), \quad \text{below } \Delta.$$

Hence

$$\int_{\gamma_1(b)} [G(x) - G(c_0)] dy \leq \int_{\gamma_1(b_0)} [G(x) - G(c_0)] dy,$$

where the positive directions along the curves are those of increasing time (counterclockwise). Thus from (14) and (15)

$$\begin{aligned} U^+(b) &\leq U^+(b_0) + \int_{\gamma_2(b)} [G(x) - G(c_0)] dy \\ &\quad + G(c_0)[\delta(b) - \delta(b_0)]. \end{aligned}$$

Now by (b), $G(x) - G(c_0) < 0$ on $\gamma_2(b)$; $G(c_0) \leq G(0) = 0$ by (a), and $\delta(b) - \delta(b_0) \geq 0$ since $b \leq b_0$. Hence

$$U^+(b) < U^+(b_0), \quad b_1 \leq b < b_0 < 0.$$

In an exactly analogous fashion

$$U^-(b) < U^-(b_0), \quad b_1 \leq b < b_0 < 0,$$

from which the conclusion of the theorem follows.

We can infer from this theorem that if $\gamma(b_0)$ were a periodic solution, then it would be the only periodic solution crossing the lower half of the y -axis between B_0 and B_1 . A more important corollary of Th. 2 is

Theorem 3. If (a) $G(x)$ satisfies I of Section 3,

(b) $G(x) \geq G(c_2)$, $0 \leq x \leq c_2$,

$G(x) \leq G(e_2)$, $e_2 \leq x \leq 0$,

(c) $G(x)$ is monotone decreasing for $c_2 \leq x \leq c_3$ and $e_3 \leq x \leq e_2$;

then there is at most a single periodic solution $\gamma(b)$ for $b_3 \leq b \leq b_2 < 0$. If the solution exists, it is stable.⁴

Proof. If $b_3 \leq b_1 < b_0 \leq b_2$, then b_1 and b_0 satisfy the conditions of Th. 2 and $U(b_0) > U(b_1)$. Hence $U(b)$ is monotone increasing and has at most one zero for $b_3 \leq$

⁴A similar theorem holds for the unstable case and can be obtained by replacing $G(x)$ by $-G(x)$.

FIG. 11.

places bounds on its amplitude.⁵ It is clear that these bounds are determined by the intersections (at Q , B , S , and D , Fig. 6) of $E(a, \epsilon)$ with Δ . With $G(x)$ given they could be readily computed. In van der Pol's case, $G(x) = x - x^3/3$ and

$$2 - \frac{2}{3}\epsilon - \frac{4}{27}\epsilon^2 \leq \text{amp} \leq 2 + \frac{2}{3}\epsilon - \frac{4}{27}\epsilon^2 \quad (16)$$

when $\epsilon \leq \epsilon_0 = 2(3)^{1/2}/9$. Hence $\text{amp} \rightarrow 2$ as $\epsilon \rightarrow 0$.

Of even greater importance is to investigate the asymptotic behavior of the period T of $\gamma(a, \epsilon)$. Van der Pol in [2] obtains graphically the result that $\epsilon^2 T$ is⁶ approximately 2 for $\epsilon = 1/10$. Liénard in [5] gives an inconclusive proof that $\epsilon^2 T \rightarrow 3 - 2 \log 2 = 1.6 +$ and Kazakevich [12] gives the result as $\epsilon^2 T \rightarrow 2$. As we shall see, Liénard's result is correct.⁷

Let $\gamma(b, \epsilon)$ be, as in Fig. 12, the path through the point $B : (0, b)$, $b < 0$.⁸ In order to assure that for all $\epsilon > 0$, $\gamma(b, \epsilon)$ intersects Δ we assume that for some $x = c_3 > 0$, $G(c_3) = b$. Let T_1 be the time for the representative point to move from B to the point C on Δ . L_1 is the line through B with slope ϵ . C_1 is the first intersection ($x > 0$) of L_1

⁵The amplitude is the maximum of $|x|$.

⁶Our units differ from van der Pol's by the factor $1/\epsilon$.

⁷A refinement of this result is given by Haag [7].

⁸The case $b > 0$ can be treated in a similar manner, as was pointed out at the beginning of Sec. 3.

FIG. 12.

with Δ , and $D: (d, G(d))$ is an absolute maximum of $G(x)$ for $0 \leq x \leq c$. Then by solution (1) of Th. 1 $\gamma(a, \epsilon)$ lies between the ellipses⁹

$$\begin{aligned} x &= c_1 \cos \theta \\ E_1 : \\ y &= G(c_1) + \epsilon c_1 \sin \theta, \quad -\frac{\pi}{2} \leq \theta \leq 0 \end{aligned}$$

and

$$\begin{aligned} x &= \frac{d-b}{\epsilon} \cos \varphi \\ E_2 : \\ y &= d + (d-b) \sin \varphi, \quad -\frac{\pi}{2} \leq \varphi \leq -\frac{\pi}{2} + \varphi_2 \end{aligned}$$

where $\varphi = -\pi/2 + \varphi_2$ corresponds to the intersection C_2 of E_2 with Δ . Thus $\varphi_2 = \tan^{-1}(\epsilon \tan \alpha)$, where α is the angle BMC_2 . From Eq. (5)

$$dt = \frac{dy}{\epsilon^2 x}$$

⁹We are assuming ϵ to be so small that the horizontal axis of E_1 lies below Δ .

and hence

$$\int_{E_2} \frac{dy}{x} \leq \epsilon^2 T_1 = \int_{\gamma(b)} \frac{dy}{x} \leq \int_{E_1} \frac{dy}{x}.$$

It then follows that

$$\epsilon \varphi_2 \leq \epsilon^2 T_1 \leq \epsilon \frac{\pi}{2}. \quad (17)$$

In order to treat the case where the path is above Δ we need the following solution to $\lambda \geq 0$.

Solution 3. Taking $f(\theta) = \theta$, $g(x) = \Phi_c^+(x)$ and $\psi = k_c(c^2 - x^2)^{1/2}$ in Eq. (7), we obtain

$$\Gamma : y = \Phi_c^+(x) + \epsilon k_c(c^2 - x^2)^{1/2}, \quad 0 \leq x \leq c \quad (18)$$

and

$$\lambda = x \left\{ \epsilon^2 k_c^2 + \left[\frac{\Phi_c^+(x) - G(x)}{(c^2 - x^2)^{1/2}} - \frac{\Phi_c^{+'}(x)(c^2 - x^2)^{1/2}}{x} \right] \epsilon k_c - \epsilon^2 \right\}.$$

Noting that

$$\lim_{x \rightarrow c} - \frac{\Phi_c^+(x) - G(x)}{(c^2 - x^2)^{1/2}} = 0,$$

we define

$$2M_c = \max_{0 \leq x \leq c} \left[\frac{\Phi_c^+(x) - G(x)}{(c^2 - x^2)^{1/2}} - \frac{\Phi_c^{+'}(x)(c^2 - x^2)^{1/2}}{x} \right] < \infty$$

and obtain

$$\lambda \leq x(\epsilon^2 k_c^2 + 2M_c \epsilon k_c - \epsilon^2).$$

Picking

$$k_c = -\frac{M_c}{\epsilon} + \left[\left(\frac{M_c}{\epsilon} \right)^2 + 1 \right]^{1/2} = \frac{\epsilon}{2M_c} - \left(\frac{\epsilon}{2M_c} \right)^3 + O(\epsilon^5),$$

we find that $\lambda \leq 0$ along Γ .

Therefore from the above and the solution (2) of Th. 1, the curve $\gamma(c, \epsilon)$ (Fig. 13) from $C : (c, G(c))$ to the y -axis at H lies between Γ of (18) and

$$\Gamma_1 : y = \Phi_c^+(x) + \epsilon(c^2 - x^2)^{1/2}. \quad (19)$$

Let D be the absolute maximum point of Δ , $0 \leq x \leq c$, which is closest to the y -axis; and E the intersection with $\gamma(c, \epsilon)$ of the horizontal line through D .

Now by (5)

$$\epsilon^2 T_2 = \int_{C_{\gamma(c, \epsilon)}}^E \frac{dy}{x} + \int_{E_{\gamma(c, \epsilon)}}^F \frac{dy}{x} + \epsilon^2 \int_{F_{\gamma(c, \epsilon)}}^H \frac{dx}{G(x) - y},$$

FIG. 13.

where F corresponds to the point $x = d/2$ on $\gamma(c, \epsilon)$. But

$$\int_{C_{\gamma(c, \epsilon)}}^E \frac{dy}{x} \leq \int_{C\Gamma}^D \frac{dy}{x} = \int_c^d \frac{\Phi_c^{+'}(x)}{x} dx + k_c \epsilon \int_c^d \frac{dx}{(c^2 - x^2)^{1/2}}$$

$$\int_{E_{\gamma(c, \epsilon)}}^F \frac{dy}{x} \leq \int_{G(d)}^{G(d) + \epsilon(c^2 - d^2/4)^{1/2}} 2 \frac{dy}{d} = \frac{2\epsilon}{d} (c^2 - d^2/4)^{1/2}$$

$$\begin{aligned} \int_{F_{\gamma(c, \epsilon)}}^H \frac{dx}{G(x) - y} &\leq \int_0^{d/2} \frac{dx}{G(d) - G(x) + k_c \epsilon (c^2 - x^2)^{1/2}} \\ &\leq \int_0^{d/2} \frac{dx}{G(d) - G(x)}, \end{aligned}$$

and hence

$$\epsilon^2 T_2 \leq \int_c^d \frac{\Phi_c^{+'}(x)}{x} dx + O(\epsilon). \quad (20)$$

Since $\gamma(c, \epsilon)$ lies below Γ_1 of (19),

$$\epsilon^2 T_2 = \int_{\gamma(c, \epsilon)}^E \frac{dy}{x} \geq \int_{C\Gamma_1}^{E_1} \frac{dy}{x} = \int_c^{e_1} \frac{\Phi_c^{+'}(x)}{x} dx + \epsilon \cos^{-1} \frac{e_1}{c} \quad (21)$$

where E_1 is the point on Γ_1 for which $y = G(d)$ and e_1 is the x -coordinate of E_1 . Hence

$$- \int_{e_1}^c \frac{\Phi_e^{+'}(x)}{x} dx + O(\epsilon) \leq \epsilon^2 T_2 \leq - \int_d^c \frac{\Phi_e^{+'}(x)}{x} dx + O(\epsilon). \quad (22)$$

Note that $e_1 \rightarrow d$ as $\epsilon \rightarrow 0$. Bounds on the times when $x < 0$ can be obtained in a similar manner. Thus we see from (17) and (22) that the time $\epsilon^2 T$ which the path spends in the region above where $G'(x) < 0$ for $x > 0$ and below where $G'(x) < 0^{10}$ for $x < 0$ is $O(1)$. Otherwise the time is $O(\epsilon)$. Thus for small ϵ we obtain the characteristic of a relaxation, with x changing relatively slowly in some regions and rapidly elsewhere.

Let $T(a, \epsilon)$ be the period of the periodic solution within $E(a, \epsilon)$. Referring to (22) and Fig. 6, we see that

$$\epsilon^2 T(a, \epsilon) \rightarrow \int_{a_1}^{a_2} \frac{\Phi_{a_1}^{+'}(x) - \Phi_{a_2}^{+'}(x)}{x} dx \quad \text{as} \quad \epsilon \rightarrow 0. \quad (23)$$

In van der Pol's case ($G(x) = x - x^3/3$) we obtain

$$\epsilon^2 T \rightarrow 2 \int_1^2 \left(x - \frac{1}{x} \right) dx = 3 - 2 \log 2 = 1.614+,$$

which confirms the result given by Liénard.

BIBLIOGRAPHY

1. E. and H. Cartan, *Note sur la génération des oscillations entretenues*, Ann. des Post Télégr. Téléph. **12**, 1196-1207 (1925).
2. B. van der Pol, *On "relaxation oscillations"*, Phil. Mag. **27**, 978-992 (1926).
3. B. van der Pol, *Oscillations sinusoidales et de relaxation*, l'Onde Electrique **9**, 293-312 (1930).
4. B. van der Pol, *The non-linear theory of electrical oscillations*, Proc. Instit. Radio Eng. **22**, 1051-1086 (1934).
5. A. Liénard, *Etude des oscillations entretenues*, Rev. Gén. de l'Elec. **23**, 901-912 and 946-954 (1928).
6. N. Levinson and O. Smith, *A general equation for relaxation oscillations*, Duke Math. Jour. **9**, 382-403 (1942).
7. J. Haag, *Etude asymptotique des oscillations de relaxation*, Ann. Sci. École Norm. Sup. (3) **60**, 35-64, 65-111 and **61**, pp. 73-117 (1944).
8. A. Andronow and C. Chaikin, *Theory of oscillations*, English Language Edition, Princeton Univ., 1948.
9. D. Flanders and J. Stoker, *The limit case of relaxation oscillations*, *Studies in Non-Linear Vibration Theory*, New York Univ., 1946.
10. N. Levinson, *Perturbations of discontinuous solutions of non-linear systems of differential equations*, Proc. Nat. Acad. Sci. **33**, 214-218 (1947).
11. S. Lefschetz, *Lectures on differential equations*, Princeton Univ., 1946.
12. U. Kazakevich, *On approximate integration of van der Pol's equation*, Doklady Akad. Nauk SSSR (N.S.) **49**, 414-417 (1945).

¹⁰ $G'(x) < 0$ states that the dynamic resistance is positive.