QUARTERLY OF APPLIED MATHEMATICS

Vol. XIII

OCTOBER, 1955

No. 3

LINEAR PROGRAMMING*

BY

KURT EISEMANN

International Business Machines Corporation, New York

1. Economic foundation. The growing application of mathematical techniques to problems of economics has brought about, in the last decade, development of a practical and promising method for providing answers to certain, hitherto unanswerable questions. Where applicable, "Linear Programming" is able to divulge the most profitable or least costly mode of operations (within given ranges of conditions) for a variety of industrial operations. At times, unexpected results obtained thereby represented an improvement over past operations and permitted substantial savings thereafter. The mathematical formulation and solution of problems expressible in Linear Programming form promises to grow into a useful servant to the economic planning of many an industry.

Among the questions to which Linear Programming can help provide answers, are these:

1. Within specified ranges of conditions, what quantities of what items should be manufactured so as to yield maximum profit?
2. Within stipulated ranges of specifications, what mixture of what compositions should be marketed so as to result in maximum profit? Or in maximum sales volume?
3. Within definite ranges of market conditions, from what suppliers shall what items be distributed to what consumers so as to minimize costs of distribution?
4. Within specified limitations of labor requirements, what allocation of what labor categories to what jobs will require a minimum of operating personnel? Or minimum expenditure? Or assure maximum efficiency?
5. Given the nutritional values of various animal feeds, what quantities of what feeds should be selected so as to satisfy minimum nutritional requirements and minimize overall cost of raising livestock?

With complicated interdependent industrial operations, a change in one operating characteristic may propagate as a network of readjustments in other associated factors, the tracing of which may become excessively intricate. Trial and error methods then prove impractical and a systematic procedure is called for.

The restrictions or constraints imposed may be of a most diverse nature and may include the following: raw material availability; plant capacity; transportation facilities; freight rates for ships, railroads, trucks, and pipelines; lengths of shipping routes; storage capacities and costs; product disposability; time limitations on supply, processing, storage, and shipments; balance between input and output; physical, chemical, or tech-

*Received March 11, 1954; revised manuscript received Sept. 24, 1954.

variables, can always be reduced to the present form by the substitution $x_i = y_i - z_i$ with $y_i, z_i \geq 0$ so that restriction (2) now applies to the new variables.

Any inequality of the form $x_i \geq b_i$ may be converted, by the substitution $x_i = b_i + y_i$ into $y_i \geq 0$, which is of the form (2). It will thus be respected automatically by the subsequent procedure and need no longer be carried explicitly under (1).

Illustrative example. We shall illustrate our discussion by carrying along and solving one numerical example, which is tiny in magnitude but which illustrates the procedure to be followed for large-scale problems.

Suppose for extreme simplicity that a manufacturer produces four items, x_1 to x_4 . Let the economic limitations take the form,

$$\begin{aligned} 2x_3 + x_4 &= 10, \\ -7x_1 + x_2 + 2x_3 - 5x_4 &= 14, \\ 6x_1 - 4x_3 + x_4 &\leq 4. \end{aligned} \quad (2.6)$$

Non-negative production requirements give the conditions,

$$\text{all } x_i \geq 0. \quad (2.7)$$

Let the profit for each manufactured item be as follows:

Item	x_1	x_2	x_3	x_4
Profit (in thousands of dollars)	4	-5 (= loss)	8	38

Of all possible production schedules meeting these conditions it is desired to find that one which will maximize profit, i.e. for which

$$f = 4x_1 - 5x_2 + 8x_3 + 38x_4 = \text{a maximum}. \quad (2.8)$$

Note that the values c_i are here interpreted as profit coefficients.

3. Conversion to standard equalities. The given inequalities (2.1) are first converted into equalities with non-negative right-hand sides as follows.

Rule 1. Multiply through by -1 where necessary to make all $b_i \geq 0$.

Rule 2. Where $b_i = 0$, multiply through by -1 where necessary to convert inequalities $\sum a_{ij} x_j \geq 0$ into the form $\sum a'_{ij} x_j \leq 0$.

$$\begin{array}{ll} \text{Rule 3. Replace inequalities} & a_{i1}x_1 + \cdots + a_{in}x_n \leq b_i \\ \text{by the equalities} & a_{i1}x_1 + \cdots + a_{in}x_n + x_{n+i} = b_i \end{array} \quad (3.1)$$

$$\begin{array}{ll} \text{and replace inequalities} & a_{i1}x_1 + \cdots + a_{in}x_n \geq b_i \\ \text{by the equalities} & a_{i1}x_1 + \cdots + a_{in}x_n - x_{n+i} = b_i \end{array} \quad (3.2)$$

The additional variables x_{n+i} take up the "slack" by which the inequalities are allowed to depart from equalities and are therefore called "slack variables". From (1) and (2) they are subject, like all other variables x_i , to condition (2.2) and may constitute an integral part of the final solution. By (2.3) they will be able to make no positive contribution to f and must therefore be adjoined to the right hand side of (2.3) with associated coefficients $c_{n+i} = 0$. No slack variable is introduced for any equality of (2.1).

With the inclusion of original as well as slack variables, our problem now reads:

$$A_1x_1 + \cdots + A_nx_n = B, \quad \text{where all } b_i \geq 0, \quad (3.3)$$

$$x_i \geq 0 \quad \text{for all } i, \quad (3.4)$$

$$f = c_1x_1 + \cdots + c_nx_n = \max. \quad (3.5)$$

4. Solvability theorems. We are confronted with the simple problem of maximizing a linear function of several variables subject to side conditions, apparently amenable to the standard analytical treatment with Lagrange multipliers. However, an analytical maximum may well have some or all of the variables negative, violating restrictions (3.4). If we should try to satisfy (3.4) by the substitution $x_i = y_i^2$ and then apply Lagrange multipliers, the derivatives of (3.5) will turn out to be linear, but the constraint relations (3.3), which will specifically be needed for solution, will remain non-linear, furnishing a system of simultaneous non-linear equations. A different approach will therefore be taken.

We define:

A "feasible" solution is a set of x_i which satisfies (3.3) and (3.4) but not necessarily (3.5).

A "maximal" solution satisfies (3.3), (3.4) and (3.5).

A "basic feasible" solution is a feasible solution with *not more than* m non-zero (and therefore positive) values x_i .

A "basic maximal" solution is a maximal solution with not more than m non-zero (and therefore positive) values x_i .

The following existence theorems apply to our problem:

Theorem 1. Whenever there exists a feasible solution, there also exists a basic feasible solution.

Theorem 2. If f remains finite for all possible feasible solutions, then there exists a basic maximal solution.

Solution of the linear programming problem is accomplished by the so-called "simplex" procedure. First a basic feasible solution is constructed. This is then replaced by another basic feasible solution for which f is larger than before. The process is continued until either

(i) f can no longer increase, indicating attainment of a maximal solution; or

(ii) certain characteristics in the procedure indicate that $f_{\max} = \infty$.

The number of iterations required will be shown to be finite.

For the sake of illustration, we shall for the moment assume that our procedure admits of a solution and that $f_{\max}$ = finite. Other situations will be discussed further below.

5. Unit basis. If the first m column vectors A_i are linearly independent, they provide a basis in terms of which all columns of the coefficient matrix can be expressed as

$$A_j = d_{1j}A_1 + d_{2j}A_2 + \cdots + d_{mj}A_m. \quad (5.1)$$

If we can furthermore arrange it that $A_1, A_2, \cdots, A_m$ are the m columns I_i of the m th order identity matrix (we shall call these vectors "unit columns", collectively forming a "unit basis"), we shall gain two vital advantages.

First, to obtain the d_{ij} , which will be required in (7.1), (8.1) and (8.2) below, it would normally be necessary to solve Eq. (1), which, written in terms of the components

$a_{1i} \cdots a_{mi}$ of the A_i , represents m equations in the m unknowns d_{ij} . Solutions will be required for $j = m + 1$ to n and it would thus be necessary to solve $n - m$ systems of m equations each. With $A_i = I_i$, this labor will be completely eliminated, for now

$$A_j = \begin{bmatrix} a_{1j} \\ a_{2j} \\ \cdot \\ \cdot \\ \cdot \\ a_{mj} \end{bmatrix} = a_{1j} \begin{bmatrix} 1 \\ 0 \\ 0 \\ \cdot \\ \cdot \\ 0 \end{bmatrix} + a_{2j} \begin{bmatrix} 0 \\ 1 \\ 0 \\ \cdot \\ \cdot \\ 0 \end{bmatrix} + \cdots + a_{mj} \begin{bmatrix} 0 \\ 0 \\ \cdot \\ \cdot \\ 0 \\ 1 \end{bmatrix} \quad [\text{any } j], \quad (5.2)$$

so that $d_{ij} = a_{ij}$, which are given.

$$\left. \begin{array}{l} \text{Second, with} \quad x_i = 0 \quad \text{for} \quad i > m, \\ (3.3) \text{ then gives} \quad x_i = b_i \geq 0 \quad (i = 1 \text{ to } m), \end{array} \right\} \quad (5.3)$$

immediately furnishing a basic feasible solution.

6. Initial set-up. We first wish to set up a starting unit basis. In (3.3) several of the originally given A_j may happen to be unit columns. Furthermore, each slack variable x_{n+i} supplied by (3.1) [but not by (3.2)] introduces a corresponding unit column A_{n+i} .

We now apply

Rule 4. Augment the coefficient matrix by further unit columns I_j until A contains all the m possible unit columns of an $m \times m$ identity submatrix.

Equation (3.3) is thus replaced by

$$A_1 x_1 + \cdots + A_n x_n + A_{n'+1} x_{n'+1} + \cdots + A_{n'+n'} x_{n'+n'} = B, \quad (6.1)$$

where the new unit columns and quantities $x_{n'+i}$ are called "artificial" vectors and variables.

Our example, by the introduction of slack and artificial variables, now takes the matrix form

$$\begin{bmatrix} 0 & 0 & 2 & 1 & 0 & 1 \\ -7 & 1 & 2 & -5 & 0 & 0 \\ 6 & 0 & -4 & 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{bmatrix} = \begin{bmatrix} 10 \\ 14 \\ 4 \end{bmatrix}. \quad (6.2)$$

Here

A_5 = slack vector

A_6 = artificial vector

x_5 = slack variable

x_6 = artificial variable

All 3 columns of the 3rd-order identity matrix now appear in A . Any inequality of the type (3.2) introduces a slack vector $-I_i$ into the matrix of coefficients and therefore will in addition require the introduction of an artificial vector $+I_i$.

Our original equations (2.1) permit certain amounts of slack, which may well appear in the final solution. Since they do not make any contribution to the function f , the c_i coefficients corresponding to *slack* variables x_i were taken as zero.

The *artificial* variables, however, are introduced only for convenience in getting started and have no place in a final solution. To assure their subsequent elimination we attach an unduly large penalty to their presence. By setting the c_i coefficients corresponding to *artificial* variables x_i equal to $-M$ (a very large negative number), we shall insure that f cannot possibly reach its maximum as long as any feasible solution still contains any of the artificial variables. Their elimination by the routine procedure below is thus insured automatically.

It is convenient to apply

Rule 5. Re-arrange the A -columns and variables x_i so as to display the identity sub-matrix on the left.

Redefining accordingly A_i , x_i , c_i , and n , our problem takes on the final formulation,

$$\left. \begin{aligned} A_1x_1 + A_2x_2 + \cdots + A_nx_n &= B \\ (A_j = I_j \quad \text{for } j \leq m; \quad \text{all } b_i &\geq 0) \end{aligned} \right\}, \quad (6.3)$$

$$\text{all } x_i \geq 0, \quad (6.4)$$

$$f = c_1x_1 + c_2x_2 + \cdots + c_nx_n = \max. \quad (6.5)$$

By (5.3), an initial basic feasible solution is immediately obtained from (6.3) as

$$\left. \begin{aligned} x_i &= b_i \quad \text{for } i \leq m \\ &= 0 \quad \text{for } i > m, \end{aligned} \right\} \quad (6.6)$$

which satisfies

$$A_1x_1 + \cdots + A_mx_m = B \quad [A_j = I_j \text{ for all } j], \quad (6.7)$$

and

$$f = c_1x_1 + \cdots + c_mx_m \quad (\text{not yet necessarily a maximum}). \quad (6.8)$$

Our example, on defining

$$\begin{aligned} x'_1 &= x_6 & x'_2 &= x_2 & x'_3 &= x_5 \\ x'_4 &= x_4 & x'_5 &= x_1 & x'_6 &= x_3 \end{aligned} \quad (6.9)$$

now takes on the final form

$$\begin{bmatrix} 1 & 0 & 0 & 1 & 0 & 2 \\ 0 & 1 & 0 & -5 & -7 & 2 \\ 0 & 0 & 1 & 1 & 6 & -4 \end{bmatrix} \begin{bmatrix} x'_1 \\ x'_2 \\ \cdot \\ \cdot \\ \cdot \\ x'_6 \end{bmatrix} = \begin{bmatrix} 10 \\ 14 \\ 4 \end{bmatrix} \quad (6.10)$$

$$\text{all } x'_i \geq 0 \quad (6.11)$$

$$f = -Mx'_1 - 5x'_2 + 0x'_3 + 38x'_4 + 4x'_5 + 8x'_6 = \max, \quad (6.12)$$

with the initial basic feasible solution

$$x'_1 = 10, \quad x'_2 = 14, \quad x'_3 = 4, \quad x'_4 = x'_5 = x'_6 = 0. \quad (6.13)$$

7. Change of basis. In conformity with *Existence Theorem 1*, solution (6.6) involves non-zero variables x_i at most for $i = 1$ to m . We shall now try to determine x_r ($r \leq m$) and some value $x_k \geq 0$ ($k > m$) such that, in (6.3), replacement of the term $A_r x_r$ by a term $A_k x_k$ ($A_k =$ given) will leave all previous conditions satisfied and also *increase* f . This new solution, again involving only m variables x_i , will thus also be "basic feasible", but the increase in f will represent an improvement.

Instead of concentrating on the variables x_i , we can express our intentions in terms of the given vectors A_i : desired to replace A_r ($r \leq m$), one of the unit base vectors, by A_k ($k > m$), one of the matrix columns not in the basis, and to determine corresponding r and k , such that (6.3) and (6.4) will remain satisfied and (6.5) will increase. This process is called "introducing vector A_k into the basis".

From (5.2), for $j = k$,

$$A_k - (A_1 a_{1k} \cdots + A_m a_{mk}) = 0. \quad (7.1)$$

In order to introduce A_k into the basis, we add θ times identity (1) to (6.7):

$$A_k \theta + A_1(x_1 - \theta a_{1k}) + \cdots + A_r(x_r - \theta a_{rk}) + \cdots + A_m(x_m - \theta a_{mk}) = B \quad (7.2)$$

Lemma 1. For $i = k$, restriction (6.4) applied to (2) requires $\theta \geq 0$.

How will this modification affect f ? From (6.8),

$$f = c_1 x_1 + \cdots + c_m x_m. \quad (7.3)$$

In order to construct the new f corresponding to (2), we analogously define

$$f_j = c_1 a_{1j} + \cdots + c_m a_{mj} \quad [j = 1 \text{ to } n]. \quad (7.4)$$

Putting herein $j = k$, subtracting θ times (4) from (3), and adding the term $c_k \theta$ to each side, gives

$$f' \equiv f - \theta(f_k - c_k) = c_k \theta + c_1(x_1 - \theta a_{1k}) + \cdots + c_m(x_m - \theta a_{mk}). \quad (7.5)$$

The right-hand side is precisely the new f for (2); and we derive

Lemma 2. For $\theta > 0$ we can make $f' > f$ only if k is so chosen that $f_k - c_k < 0$.

Lemma 3. When eventually no negative $f_k - c_k$ remains, no further increase in f is possible. It will indicate termination of our procedure owing to attainment of a *maximal* solution.

Choice of k . In view of Lemma 2, it is convenient to replace (4) by the definition

$$g_j \equiv f_j - c_j = c_1 a_{1j} + \cdots + c_m a_{mj} - c_j \quad [j = 1 \text{ to } n]. \quad (7.6)$$

In theory, the largest increase in f , for one step, is obtained in (5) when $-\theta(f_k - c_k)$ is as large as possible. In practice we do the following, which is almost equally satisfactory and requires far less labor.

Rule 6. Examine only the negative $g_j \equiv f_j - c_j$, choose the most negative one, and take that j as k .

Choice of r . For all $i \leq m$, requirement (6.4) applied to the coefficients of A_i in (2) becomes $x_i - \theta a_{ik} \geq 0$.

Lemma 4. For $a_{ik} \leq 0$, this requirement is automatically satisfied for any $\theta \geq 0$, since $x_i \geq 0$.

Lemma 5. For $a_{ik} > 0$, we must make $\theta \leq x_i/a_{ik}$ for all $i \leq m$.

Lemma 6. Elimination of A_r requires $x_r - \theta a_{rk} = 0$, i.e. for some r , $\theta = x_r/a_{rk}$. Lemmas (1), (5), (6) will be simultaneously satisfied by

Rule 7. For the chosen k , select only those a_{ik} which are > 0 to form the quotients $q_i = x_i/a_{ik} (\geq 0)$. Take $r =$ the i corresponding to the smallest q_i .

Hence

$$\theta = \frac{x_r}{a_{rk}} = \min_{a_{ik} > 0} \frac{x_i}{a_{ik}}. \quad (7.7)$$

Example. It proves efficient to arrange numerical problems in the following table form, which here represents (6.10) to (6.12).

TABLE 1.

c_i of basis	Basis	$c_i :$	0	$-M$	-5	0	38	4	8
		Vector: $B = A_0$	A_1	A_2	A_3	A_4	A_5	A_6	
$-M$	A_1	10	1	0	0	1	0	2	
-5	A_2	14	0	1	0	-5	-7	2	
0	A_3	4	0	0	1	1	6	-4	
$g_j = (\sum_i c_i a_{ij}) - c_j :$		$-10M$ -70	0	0	0	$-M$ -13	31	$-2M$ -18	

Since $b_i = x_i = a_{i0}$, the insertion $c_0 = 0$ gives $g_0 = \sum c_i a_{i0} - 0 = \sum c_i x_i = f$ so that our table also displays f . Rule 6 gives $k = 6$; rule 7 gives $r = 1$; (7) gives $\theta = 5$.

In rule 6, $j = 0$ is excluded from consideration; in rule 7, $i = m + 1$ is excluded.

8. Transformation to new basis. Our change of the basis $A_1 \cdots A_r \cdots A_m$ consisted in "introducing A_k into the basis" and removing A_r . Each A_i , expressed in terms of the old basis by (5.1), must now be converted into a new form A'_i which similarly refers to the new basis.

By (5.2), for any j ,

$$A_j = a_{1j}A_1 + \cdots + a_{rj}A_r + \cdots + a_{mj}A_m. \quad (8.1)$$

Putting $j = k$ and solving for A_r , we obtain $A_r = (1/a_{rk}) (A_k - \sum_{i \neq r} a_{ik}A_i)$. Substitution into (1) will express any A_i in terms of the new basis, viz,

$$A'_i = \frac{a_{ri}}{a_{rk}} A_k + \sum_{i \neq r} \left(a_{ij} - \frac{a_{ri}}{a_{rk}} a_{ik} \right) A_i \quad [\text{any } j]. \quad (8.2)$$

This transformation likewise applies to the vector $B \equiv A_0$. For $j = k$, (2) gives

$$A'_k = 1 \cdot A_k + \sum 0 \cdot A_i \quad (8.3)$$

as should be.

Restoration to unit form. The new basis now contains $A_k = [a_{1k}, \cdots, a_{mk}]$ and is therefore no longer a unit basis. However, a notational trick restores unit form as follows:

In (5.2), the r th component of any vector indicated the coefficient of A_r , which

henceforth will be zero. On the other hand we must adjoin to (5.2) an additional term $a'_{ki}A'_k$. We now utilize the available vacancy of the r th component a'_{ri} of each vector to indicate $a'_{ki} = a_{ri}/a_{rk}$, the coefficient of A_k [from (2)]. In this new notation, (3) becomes $A'_i = [0, \dots, 0, 1, 0, \dots, 0] = I_r$, where the component 1 occurs in row r and which is now *in unit form*.

For the other unit vectors A_j remaining in the basis ($j \leq m, j \neq r$), we have

$$a_{ii} \begin{cases} = 0 & \text{when } i \neq j \\ = 1 & \text{when } i = j \end{cases}$$

and their new form is given by (2) as $A'_i = 0 + \sum_i (a_{ii} - 0) A_i = A_i = I_i$ [$j \leq m, j \neq r$]. The transformation thus leaves them unchanged in their original unit form.

However, the former unit base vector $A_r (j = r)$ is turned by (2) into $A'_r = (1/a_{rk}) A_k + \sum_{i \neq r} (-a_{ik}/a_{rk}) A_i$, and is thus no longer in unit form.

With our notation, (2) becomes, analogously to (5.2),

$$A'_i = \begin{bmatrix} a'_{1i} \\ \vdots \\ \cdot \\ \vdots \\ a'_{mi} \end{bmatrix} = \frac{a_{ri}}{a_{rk}} I_r + \sum_{i \neq r} \left(a_{ii} - \frac{a_{ri}}{a_{rk}} a_{ik} \right) I_i$$

and therefore, as in (5.2),

$$a'_{ij} \begin{cases} = a_{ri}/a_{rk} & \text{for } i = r \\ = a_{ij} - (a_{ri}/a_{rk})a_{ik} & \text{for } i \neq r \end{cases} \quad [\text{any } j], \quad (8.4)$$

which gives the law of transformation of individual components.

Transformation of g_i and f . Equation (5.2) gives rise to the definition (7.6),

$$g_i = \sum_{i=1}^m c_i a_{ii} - c_i.$$

Similarly from (2),

$$g'_i = c_k a'_{ri} + \sum_{i \neq r} c_i a'_{ij} - c_i \quad (8.5)$$

which, by (4), becomes

$$g'_i = c_k \frac{a_{ri}}{a_{rk}} + \sum_{i \neq r} c_i \left(a_{ii} - \frac{a_{ri}}{a_{rk}} a_{ik} \right) - c_i.$$

Adding the identity

$$0 \equiv c_r a_{ri} - (a_{ri}/a_{rk}) \cdot c_r a_{rk},$$

$$g'_i = \left(\sum_{i=1}^m c_i a_{ii} - c_i \right) - \frac{a_{ri}}{a_{rk}} \left(\sum_{i=1}^m c_i a_{ik} - c_k \right),$$

i.e.,

$$g'_i = g_i - \frac{a_{ri}}{a_{rk}} g_k. \quad (8.6)$$

For $j = 0$, $b_i = a_{i0}$ and $c_0 = 0$, (6) also gives f' .

From (7.6), $g_i = 0$ for $j = 1$ to m .

From (6), for the new basis vector k , likewise $g'_k = 0$ but for the vector r removed from the basis, the new value $g'_r = 0 - (1/a_{rk}) g_k$ which (since $g_k < 0$, $a_{rk} > 0$) is > 0 .

The identity of this transformation relation (6) with (4) (case $i \neq r$) allows us to establish

Rule 8. Treating the table of $B \equiv A_0$, the A_i , and the row of g_i (including $j = 0$) all as one single matrix, apply transformation (8.4) as follows. Divide through row r by the pivotal element a_{rk} , giving the new row r . Replace each remaining element a_{ij} of the matrix, including the g_i row and including the column $j = 0$, by $a'_{ij} = a_{ij} - a'_{ri} a_{rk}$, which involves precisely the four corner points of a rectangle built on the diagonal line from a_{ij} to the pivotal element. A_k then becomes I_r , g'_k becomes 0, basis vectors other than A_r remain unaltered. In the "basis identification" columns to the left of the matrix, replace c_r by c_k and A_r by A_k .

After c_r in the left-hand margin has been replaced by c_k , an arithmetic check can be obtained by comparing the g'_i with those computed independently from (5), which now becomes

$$g'_i = \sum_{j=1}^m c'_j a'_{ij} - c_i. \quad (8.7)$$

Example. One iteration of Table 1 by means of Rule 8 and check (7) results in

TABLE 2.

c_i of basis	Basis	c_i :	0	$-M$	-5	0	38	4	8
		Vector:	$B \equiv A_0$	A_1	A_2	A_3	A_4	A_5	A_6
8	A_6		5	$1/2$	0	0	$1/2$	0	1
-5	A_2		4	-1	1	0	-6	-7	0
0	A_3		24	2	0	1	3	6	0
$g_i = (\sum_j c_i a_{ij}) - c_i :$			20	$M + 9$	0	0	-4	31	0

$f = g_0$ is shown to be 20, A_0 and the identifying "basis" column display the improved basic feasible solution

$$x'_6 = 5, \quad x'_2 = 4, \quad x'_3 = 24, \quad x'_1 = x'_4 = x'_5 = 0$$

and the presence of a negative g_i indicates, by Lemma 2, that further improvement is possible.

Subsequent iterations. After completing any iteration, there is no need to rearrange columns, as the above theory still holds for the columns as now standing, by identification of the I_i with the appropriate A_i . Further iterations are carried out by application of rules (6), (7), (8) to the successively obtained tables, without any alteration of the rules, till no more negative g_i remain, indicating attainment of a desired maximal solution.

One further iteration of our example gives

TABLE 3.

c_i of basis	Basis	$c_j :$	0	$-M$	-5	0	38	4	8
		Vector:	$B = A_0$	A_1	A_2	A_3	A_4	A_5	A_6
8	A_6		1	1/6	0	$-1/6$	0	-1	1
-5	A_2		52	3	1	2	0	5	0
38	A_4		8	2/3	0	1/3	1	2	0
$g_i = (\sum_j c_j a_{ij}) - c_i :$			52	$M + 11\frac{2}{3}$	0	4/3	0	39	0

All g_i are now ≥ 0 , indicating termination. Column A_0 gives

$$x'_6 = 1, \quad x'_2 = 52, \quad x'_4 = 8, \quad x'_1 = x'_3 = x'_5 = 0,$$

so that (6.9) finally gives the desired maximal solution of problem (2.6) as

$$x_1 = 0, \quad x_2 = 52, \quad x_3 = 1, \quad x_4 = 8, \quad (x_5 = x_6 = 0). \quad (8.8)$$

The maximum profit attainable under the stipulated restrictions is indicated in the final table as $f = g_0 = 52$, which figure can be checked by substituting (8) into (2.8).

It is noteworthy in this example that realization of maximum overall profit necessitates inclusion in the production program of a large quantity of item x_2 , although this item, taken by itself, causes the producer to sustain a loss of $1000c_2 = \$5000$ per unit quantity.

9. Interpretation of the g_i . A significant question is: If we were to depart from the maximal solution by replacing $x_k = 0$ by some non-zero (i.e. positive) value for x_k , and adjustment of all other x_i so as still to satisfy inequalities (2.1) and (2.2), what would be the attending decrease in the value of f ?

At any stage of iteration, departure from a feasible solution by adjoining the value $x_k = \theta$, and leaving all other conditions satisfied, corresponds to introduction of A_k into the basis with coefficient θ as per (7.1) and (7.2). By (7.5), the decrease in f will then be $f - f' = \theta g_k$. Hence,

Lemma 7. g_i represents the decrease in f per unit increase of x_i (x_i not in the basis).

At every stage, including the final one, the table thus reveals at a glance the effect of departing from the listed basic solution. The g_i of a slack variable not occurring in the final solution will indicate the cost of tightening or relaxing that particular restriction by unity.

Furthermore, solutions "almost as good" as the maximal solution may sometimes be preferred by virtue of extraneous considerations. While the utilization of any one of the otherwise excluded variables will necessitate readjustment of all other variables, the composite effect on f can be determined instantly by a mere inspection of the g_i .

10. Check row. A very convenient check on arithmetic may be obtained by

Rule 9. Carry the values $a_{m+2,j} = 1 - \sum_{i=1}^{m+1} a_{i,j}$ as a special row of A . As a check, we must have after every matrix transformation $\sum_{i=1}^{m+2} a'_{i,j} - 1 = 0$ for each j .

This amounts to the interpretation of the g_i -row as an additional restriction and taking all $c_i = 1$ for checking purposes only.

11. Degeneracy. Whenever, in Rule 7, the smallest positive q_i occurs for several i , then by (7.7) more than one term in (7.2) will become zero, indicating degeneracy. Furthermore, by (7.5), f would show no increase at that stage, which opens the door to the disastrous possibility of "cycling": introducing and eliminating vectors until eventually arriving at a previous basis, whence the cycle is repeated indefinitely. The following "perturbation" treatment resolves the dilemma.

Suppose that at any stage the sequence of matrix columns is $P_1 \cdots P_n$. The unit basis $A_1 \cdots A_m$ no longer coincides with $P_1 \cdots P_m$ but is irregularly distributed over the matrix. Letting ϵ be a small positive number, consider B replaced by

$$B' = B + \sum_{i=1}^n \epsilon^i P_i, \quad (11.1)$$

where ϵ^j denotes the j th power of ϵ . This modified version will include the original problem by the specialization $\epsilon \rightarrow 0$. Using (5.2) for the P_i , (6.7) will be replaced by

$$\left. \begin{aligned} &A_1(x_1 + \epsilon p_{11} + \epsilon^2 p_{12} + \cdots + \epsilon^n p_{1n}) \\ &\quad + A_2(x_2 + \epsilon p_{21} + \epsilon^2 p_{22} + \cdots + \epsilon^n p_{2n}) \\ &\quad \dots\dots\dots \\ &\quad + A_m(x_m + \epsilon p_{m1} + \epsilon^2 p_{m2} + \cdots + \epsilon^n p_{mn}) \\ &\quad = B + \epsilon P_1 + \epsilon^2 P_2 + \cdots + \epsilon^n P_n \end{aligned} \right\} \quad (11.2)$$

Abbreviating

$$\xi_i = x_i + \epsilon p_{i1} + \epsilon^2 p_{i2} + \cdots + \epsilon^n p_{in} \quad (11.3)$$

(6.8) will be replaced, in view of (7.4), by

$$f = c_1 \xi_1 + \cdots + c_m \xi_m = (c_1 x_1 + \cdots + c_m x_m) + \epsilon f_1 + \epsilon^2 f_2 + \cdots + \epsilon^n f_n. \quad (11.4)$$

As in the derivation of (7.2), the addition of θ times the identity (7.1) to (2) gives

$$A_k \cdot \theta + A_1(\xi_1 - \theta a_{1k}) + \cdots + A_r(\xi_r - \theta a_{rk}) \cdots + A_m(\xi_m - \theta a_{mk}) = B' \quad (11.5)$$

and the subtraction of θ times (7.4) [for $j = k$] from (11.4) replaces (7.5) by

$$f' = f - \theta(f_k - c_k) = c_k \theta + c_1(\xi_1 - \theta a_{1k}) + \cdots + c_m(\xi_m - \theta a_{mk}). \quad (11.6)$$

Suppose rule 7 had produced "ties" for the i -values $i_1 < i_2 < \cdots < i_r < \cdots$ and is now applied to (5). For those i which gave $x_{i_1}/a_{i_1k} = x_{i_2}/a_{i_2k} = \cdots$ we now have to compare

$$\begin{aligned} q_{i_1} &= \frac{\xi_{i_1}}{a_{i_1k}} = \frac{x_{i_1}}{a_{i_1k}} + \epsilon \frac{p_{i_11}}{a_{i_1k}} + \epsilon^2 \frac{p_{i_12}}{a_{i_1k}} + \cdots + \epsilon^n \frac{p_{i_1n}}{a_{i_1k}}, \\ q_{i_2} &= \frac{\xi_{i_2}}{a_{i_2k}} = \frac{x_{i_2}}{a_{i_2k}} + \epsilon \frac{p_{i_21}}{a_{i_2k}} + \epsilon^2 \frac{p_{i_22}}{a_{i_2k}} + \cdots + \epsilon^n \frac{p_{i_2n}}{a_{i_2k}} \text{ etc.} \end{aligned} \quad (11.7)$$

Now, for sufficiently small ϵ , P_1 will in general make some of the factors of ϵ in (7) larger than others and thus eliminate the corresponding rows from further competition involving higher powers of ϵ . P_2 will then, in general, by a similar effect of ϵ^2 , eliminate further rows from subsequent competition; etc. At no stage can *all* of several remaining tied rows be thus eliminated. On reaching comparison for factors of ϵ^j , whenever P_i is one of the unit columns with its non-zero element in one of the tied rows, say $P_i = I_{i_r}$ ($p_{i_r} = 1$ for $i = i_r$, $p_{i_r} = 0$ for $i \neq i_r$), if i_r has not already been eliminated by this process, it will now certainly be eliminated. And since all I_{i_r} (for all tied rows i_r) are present among the P_i , the process must eventually resolve all ties.

There will thus result a uniquely determined smallest q_i , giving

$$\theta = \min_{r=1,2,\dots} \frac{1}{a_{i_r,k}} (x_{i_r} + \epsilon p_{i_r,1} + \epsilon^2 p_{i_r,2} + \dots + \epsilon^n p_{i_r,n}) \quad (11.8)$$

and among the coefficients $\xi_{i_r} - \theta a_{i_r,k}$, which had previously all been equal to zero, there will now be precisely one smallest = 0, all others being > 0 . In (5), only one of the basis vectors A_{i_r} will now be eliminated. *Degeneracy is thus avoided.*

In case ties are again encountered during a later iteration, it is merely necessary, for all such ties, to take the sequence of P_i , for (3), in the same order $P_1 \dots P_n$ as before, i.e. again simply from left to right.

It can be shown that no difficulties will be encountered by the use of this procedure if the original unit basis $A_1 \dots A_m$ is written at the left-hand side of matrix A , through which the sequence of P_i will begin its progression in the case of degeneracy.

In practice, there is no need whatever to carry through the preceding analysis, as the emerging procedure can be simply summarized by

Rule 10. In the case of ties for r , consider P_i and A_k , starting with $t = 1$. Only for those i which are tied, form the quotients $q_i = p_{i,t}/a_{i,k}$ and take $r =$ the i corresponding to the algebraically smallest q_i (negative $p_{i,t}$ may now make q_i negative). If ties remain, consider only the i for the remaining ties, increase t by 1, and repeat. After r is uniquely determined, continue with the transformation as usual.

As stated, the presence of m unit vectors guarantees that any ties must be broken before t reaches the value n . Usually $t = 1$ or 2 already does the trick.

12. Special situations. (i) All $g_i \geq 0$. We have seen in Lemma 3 that this represents attainment of a maximal solution.

(ii) For some negative g_i , all corresponding $a_{i,j}$ are ≤ 0 . Eq. (7.5) and Lemma 4 show that we may take $\theta \rightarrow \infty$, leading to a feasible solution that includes both x_r and x_k and for which we obtain

Lemma 8. If, for any k , $g_k < 0$ and all $a_{i,k} \leq 0$, then $f_{\max} = \infty$ with $x_k = \infty$.

If desired, this situation can be excluded *a priori*, and finiteness of all variables insured, by adjoining to the original set of inequalities the additional condition $\sum_{i=1}^n x_i \leq N$, where N is a sufficiently large number. If the slack variable x_{n+1} corresponding to this additional restriction appears in the final basis, we shall know that all variables would have remained finite anyhow. If x_{n+1} does not appear in the final basis, and some of the other variables x_i are large, we shall know that the growth of one or more variables towards infinity was arrested solely by the present extra restriction.

(iii) $\theta = 0$. This will occur whenever the corresponding x_i in (7.7) is = 0. By (6.7), B is then expressed in terms of less than m basis vectors A_i and we again have degeneracy. In this case, from (8.6), $g'_0 = g_0$ and $g'_k = 0$ so that f does not increase. Here again,

application of the foregoing perturbation method is equivalent, by (11.8), to making θ non-zero, resulting in a corresponding infinitesimal increase in f . Rules 7, 8, and 10 are applied to this case without modification.

(iv) *Multiple solutions.* On termination, when all $g_i \geq 0$, we know that no feasible solution can be found for which f would be larger yet. However, it is quite possible that other maximal solutions exist which possess an identical value for f . It may be desirable to determine these alternative maximal solutions.

At first we determine alternative maximal solutions which are *basic*. This corresponds to a matrix transformation as before, with no change in f . By (7.5) and (7.7) this will be possible whenever either

(1) there exist $g_k = 0$ associated with vectors not in the basis (i.e. A_k not a unit column) with $a_{ik} > 0$ for some i ; or

(2) one or more values $b_r (= x_r$ in the basis) are $= 0$ with $a_{rj} > 0$ for some j .

With such values of k or r , all possible basic maximal solutions can be obtained by transformation as before. Non-basic maximal solutions can then be obtained as linear combinations of basic maximal solutions. Further solutions that are "almost maximal" with slightly lower f can then be found by Sec. 9.

(v) *The final basis includes an artificial vector with associated variable different from zero.* This result indicates incompatibility of the original set of inequalities.

13. Convergence. At each iteration, by Lemma 2, f increases progressively; hence no solution can reappear at a later iteration. Furthermore, the n column vectors admit at most $\binom{n}{m}$ different bases and, from (6.7), any solution for any one basis must be unique. Hence there can exist at most $\binom{n}{m}$ different basic feasible solutions; and since none can reappear, *our procedure will terminate in a finite number of steps*, either because some $g_i < 0$ with all corresponding $a_{ij} \geq 0$, indicating by Lemma 8 that $f_{\max} = \infty$, or because all $g_i \geq 0$, indicating that $f_{\max}$ has been reached. Experience shows that the maximal solution is reached, on the average, in m to $2m$ iterations. It must be emphasized that although no previous solution can reappear, it may well happen that individual vectors are removed from the basis and later reintroduced; or are introduced into the basis and later on removed again. It is not the presence of any particular basis vector, but the totality of basis vectors, that is non-repetitive.

14. Alternative methods. Two approaches other than the simplex method are available for the solution of linear programming problems. First, it can be shown that an equivalent problem is the determination of the best strategies for playing a "zero-sum two-person game", enabling the application of game theory. Second, successive approximations may be obtained by a relaxation process, for which, however, a very large number of iterations may be required.

For efficient solution, these two methods are of subordinate importance, as experience on the various methods of solution has demonstrated the decisive superiority of the simplex procedure over other methods.

15. The transportation problem. Three typical problems in this category are the following:

(i) Given m sources of a certain commodity to be shipped to n destinations. Source i supplies quantity s_i , destination j requires quantity d_j . Shipment from source i to destination j costs c_{ij} per item and is made for quantity x_{ij} . How many items should be shipped from any source to any destination so as to minimize overall transportation costs?

it will be seen that the transformation (8.4) for the t th iteration may be written

$$A' = {}_tE A. \quad (16.3)$$

For iterations 1, 2, $\dots$, t , successive application of (3) shows that

$$A' = ({}_1E {}_{t-1}E \cdots {}_2E {}_1E)A. \quad (16.4)$$

At this stage, the columns of the original identity submatrix (basis) I will have been converted into the new $m \times m$ submatrix V , say. The new identity submatrix (current basis) J now consists of columns which were, in the original system, submatrix U , say. Other columns, which originally formed submatrix P , will have been converted into Q . We can therefore write, original $A = (I, U, P)$, current $A' = (V, J, Q)$, where the transformation has resulted in

$$I \rightarrow V, \quad U \rightarrow J, \quad P \rightarrow Q. \quad (16.5)$$

In this notation, several columns of A may be recorded twice, once under I and once under U ; correspondingly for V and J . Similarly, P (or Q) need not be restricted to columns other than I, U (or V, J); any statement for P and Q will include columns I, U and V, J as special cases.

By (4), every column of A is transformed in one and the same manner by matrix premultiplication. Equation (5) therefore gives

$$V = \left(\prod_{i=1}^t {}_iE \right) I, \quad (16.6)$$

$$J = VU, \quad (16.7)$$

$$Q = VP. \quad (16.8)$$

J is the current identity submatrix with scrambled column order. Denoting the transpose of J by J^T , we have $JJ^T = J^TJ = I$. Combination of these relations with (7) shows that

$$\begin{aligned} U &= V^{-1}J, & U^{-1} &= J^T V, \\ V &= JU^{-1}, & V^{-1} &= UJ^T, \end{aligned} \quad (16.9)$$

i.e. after appropriate rearrangement of columns or rows, U and V give the inverses of each other.

Let

$$\begin{aligned} V^i &= i\text{th row of } V \\ P_j &= j\text{th column of } P \\ q_{ij} &= \text{typical element of } Q \\ g_i^Q &= \text{the row of } g_i \text{ underneath matrix } Q \\ g_i^Q &= \text{the element of row } g_i^Q \text{ in column } j. \end{aligned}$$

If V is kept up to date, there is no need to compute Q for each iteration, as the final Q on termination of the procedure will follow from (8) as $q_{ij} = V^i P_j$. It is on this crucial fact that the modified simplex method is based.

In the sequel we shall denote a zero row by $0'$, a zero column by 0 .

We have seen from (8.4) and (8.6) that g_i may be treated as the $(m+1)$ 'th component of A_i . On defining ${}_tE'$ as the $(m+1) \times (m+1)$ matrix corresponding to (2), the

overall transformation from A to any A' (now of $m + 1$ rows each) may be taken as equivalent to

$$A' = \left(\prod_{r=1}^i E' \right) A = \begin{pmatrix} V & 0 \\ g^V & 1 \end{pmatrix} A, \quad (16.10)$$

where, on the right, V has been bordered by row g^V at the bottom and by the $(m + 1)$ unit column on the right.

The cumulative effect of any number of transformations on the entire simplex table may therefore be represented by the following product of partitioned matrices,

$$\begin{pmatrix} V & J & Q \\ g^V & 0' & g^Q \end{pmatrix} = \begin{pmatrix} V & 0 \\ g^V & 1 \end{pmatrix} \cdot \begin{pmatrix} I & U & P \\ 0' & g^U & g^P \end{pmatrix}. \quad (16.11)$$

It is seen from this representation that $g^Q = g^V P + g^P$. Hence, for column j , the formula

$$g_i^Q = g^V \cdot P_i + g_i^P \quad (16.12)$$

furnishes the values of all g_i .

For any one iteration, only the following quantities need be evaluated:

Rule 11. (i) Given V , Q_0 , and all g_i , determine k as usual.

(ii) Obtain only column k from $q_{ik} = V^i P_k$ ($i = 1$ to m).

(iii) Determine r as usual.

(iv) Transform V and g^V as usual by (8.4) or the $(m + 1)$ dimensional form of (16.3).

(v) Obtain only the new column Q'_0 from $q'_{i0} = V^i P_0$ ($i = 1$ to m).

(vi) Obtain all new g_i from $g_i^Q = g^V P_i + g_i^P$.

(vii) Repeat.

At the end of the procedure, the complete final matrix Q may be obtained from (11).

The modified method has the following advantages over the regular simplex procedure:

(i) Each iteration step requires transformation only of P_k , V , g^V , Q_0 , and the row of g_i , but *not* of Q (and U). This generally amounts to a considerable saving of labor.

(ii) No reliance need be placed on data that bear accumulated round-off errors. g^Q , Q_0 , P_k and Q are obtained by an application of V and g^V to the original data. And any accumulated round-off errors in V and g^V may be eliminated at any desired stage by recalculation by the $(m + 1)$ dimensional analog of (9), viz.

$$\begin{pmatrix} V & 0 \\ g^V & 1 \end{pmatrix} = \begin{pmatrix} J & 0 \\ 0' & 1 \end{pmatrix} \text{ times the inverse of } \begin{pmatrix} U & 0 \\ g^U & 1 \end{pmatrix}. \quad (16.13)$$

(iii) For each transformation, continued use is made of the large blocks of zeros usually present in the original matrix, which would otherwise rapidly fill up.

17. Determinants. From (16.2) and (16.1),

$$|E'| = \frac{1}{a_{rk}}$$

so that (16.6) gives

$$|V| = \frac{1}{\prod_r a_{rk}}$$

and by (16.7),

$$|U| = \frac{|J|}{|V|} = |J| \cdot \prod_{r=1}^l a_{rk}, \quad \text{where} \quad |J| = \pm 1. \quad (17.1)$$

Hence the determinant of U is given in magnitude by the product of the pivotal elements, always non-zero and positive, arising in the simplex procedure. The non-singularity of successive $|U|$ is to be expected, since the final basis arises from a non-singular overall transformation of linearly independent vectors.

18. Matrix inversion and simultaneous linear algebraic equations. Given A for inversion, or $Ay = b$ for solution, let $s_i = \sum_j a_{ij}$ and $e =$ a column of m components, each equal to 1.

We first set up the problem

$$Ax = s; \quad x \geq 0 \quad (18.1)$$

which we know has the unique, finite, non-negative solution $x = e$. To render all $s_i \geq 0$, multiply appropriate equations through by -1 . This corresponds to premultiplication by a matrix I' , which is an identity matrix with plus or minus "ones" on the diagonal. Denoting the changed matrices by primes,

$$A' = I'A \quad (18.2)$$

and we have the problem

$$A'x = s' \quad (s' \geq 0) \quad x \geq 0. \quad (18.3)$$

From (16.5) and (16.9), we can accomplish the inversion of $A' (= U)$ if matters can be so arranged that the final basis J will involve precisely the m columns of A' and no others. This end result may be forced on the problem at the outset as follows,

- (i) Augment A' by a complete identity submatrix I of artificial vectors.
- (ii) Require $f = \sum c_i x_i = \text{maximum}$.
- (iii) Specify $c_i \begin{cases} = 1 \text{ (say) for columns of } A' \\ = -M \text{ (a large negative number) for all other columns.} \end{cases}$

As explained earlier (Sec. 6), f then cannot possibly reach its maximum as long as any variable x_i with $c_i = -M$ still remains in the basis. Since $x = e$, we know that the maximum of f exists and is finite ($= m$). The simplex procedure applied to the present linear programming problem must therefore necessarily lead to a final basis involving precisely the m columns of A' , so that the entire process can be represented as $(s', A', b', I) \rightarrow (e, J, p, V)$ where, in particular,

$$s' \rightarrow e, \quad A' \rightarrow J, \quad b' \rightarrow p, \quad I \rightarrow V \quad (18.4)$$

and where, by (16.9) and (2), $V = JA'^{-1} = JA^{-1}I'$, whence

$$A^{-1} = J^T V I'. \quad (18.5)$$

Similarly $p = Vb' = JA^{-1}I'(I'b)$, whence

$$y \equiv A^{-1}b = J^T p. \quad (18.6)$$

For pure matrix inversion, column b' is omitted.

These products do not require actual matrix multiplication. Premultiplication by J^T merely amounts to a rearrangement of rows, while post-multiplication by I' merely changes the signs of certain columns.

If each iteration step introduces into the basis a column of A' and removes from

the basis a column not in A' , then the simplex process requires approximately the same number of operations as ordinary elimination. Selection of k and r corresponds, in elimination, to proper choices for avoiding very small divisors.

The simplex procedure of linear programming may thus be said to represent a generalization of matrix inversion, which it includes as a special case.

19. Duality. In matrix form, we have so far treated the "direct" problem

$$Ax \leq b, \quad x \geq 0, \quad f \equiv c \cdot x = \text{maximum}. \quad (19.1)$$

A related problem, called the "dual" of (1), reads

$$A^T y \geq c, \quad y \geq 0, \quad \phi \equiv b \cdot y = \text{minimum}, \quad (19.2)$$

where A^T is the transpose of A ; A is of order $m \times n$; x, c are $n \times 1$; and y, b are $m \times 1$.

It can be shown that finite solutions exist either for both or for none of these two problems and when they exist,

$$\max f = \min \phi. \quad (19.3)$$

Solution of (1) simultaneously provides the solution of (2). Using the notation of the preceding section, the values y_i are contained in the final optimal table for (1) as

$$y = g^v. \quad (19.4)$$

If the direct problem is solved with the aid of artificial vectors and the solution y for the dual is desired, the row of g_i should be split into two distinct rows: the first, to list the coefficients of M ; and the second, to record values free of M . In the final optimal table for (1), Eq. (4), giving the values of y , then refers to the second g_i row only.

If the system (1) contains several inequalities of the form (3.2), the corresponding dual (2) may be obtained as follows. Multiply such reversed inequalities through by -1 . All inequalities are now in the form (1), which may then be translated into its corresponding dual problem.

Any equality $\sum a_{r,i} x_i = b_r$ may be converted into inequality form by replacing it by the pair of inequalities $\sum a_{r,i} x_i \leq b_r$ together with $\sum a_{r,i} x_i \geq b_r$, i.e. $\sum (-a_{r,i}) x_i \leq -b_r$. Each equation j of the corresponding dual will be of the form

$$\sum_{i \neq r} y_i a_{i,j} + (y_{r_1} - y_{r_2}) a_{r,j} \geq c_j.$$

This may be considered to represent the dual of the original matrix (a_{ij}) , where y_r has been replaced by an "unrestricted" variable $y'_r = y_{r_1} - y_{r_2}$, that may be positive or negative. Thus an equation of the original "direct" system corresponds to an unrestricted variable in the "dual".

A direct problem that permits some of the x_i to be unrestricted (i.e. to take on positive as well as negative values) may be converted to the standard form (1) by the substitution $x_k = x_{k_1} - x_{k_2}$ ($x_{k_1} \geq 0, x_{k_2} \geq 0$). Matrix A will then contain two identical columns with opposite signs throughout the transformations. Since they are linearly dependent, only one of them can be in the basis at any one time and only one need be recorded explicitly. Corresponding to the columns k_1, k_2 , the dual will contain the two inequalities $\sum y_i a_{i,k_1} \geq c_k$ and $\sum y_i (-a_{i,k_2}) \geq c_k$, whence $\sum y_i a_{i,k_1} = c_k$. Hence an unrestricted variable x_k of the direct problem corresponds, in the dual, to an equation instead of an inequality.

Whenever most inequalities of a problem are of the form (3.2), or whenever the number of restrictions greatly exceeds the number of variables, the matrices involved

in the calculation may be materially reduced by replacing such problem by its dual, possibly making use of (2.5) or its converse.

20. Partitioned problems. Many problems arising in applications result in matrices primarily consisting of several compact blocks of coefficients with blanks in the remainder of the corresponding rows. As a tiny example, consider the problem

TABLE 4.

$c_j :$	1	1	3	1	1	1	$-M$		
	A_1	A_2	A_3	A_4	A_5	A_6	A_7	A_0	
	1	0	1				(0)	x_1	≤ 2
	-1	1	3				(0)	x_2	≤ 3
				0	1	-2	(0)	x_3	≤ 4
				1	-1	7	(0)	x_4	≤ 5
	1	1	1	1	1	1	(-1)	x_5	≤ 16
								x_6	
								(x_7)	

where all $x_i \geq 0$ and $f = \sum c_j x_j = \text{maximum}$. $A_7 x_7$ is not part of the problem but is included merely for the discussion below.

We may first of all solve the smaller, mutually independent problems given by the individual blocks of submatrices:

$$\begin{array}{l} \text{Problem 1.} \\ \begin{array}{ccc} A_1 & A_2 & A_3 \\ 1 & 0 & 1 \leq 2 \\ -1 & 1 & 3 \leq 3 \end{array} \end{array} \quad (20.2)$$

$$\begin{array}{l} \text{Problem 2.} \\ \begin{array}{ccc} A_4 & A_5 & A_6 \\ 0 & 1 & -2 \leq 4 \\ 1 & -1 & 7 \leq 5 \end{array} \end{array} \quad (20.3)$$

On introduction of slack variables, the simplex procedure applied to each of these two independent problems furnishes the partial maximal solutions

$$x_1 = 2; \quad x_2 = 5; \quad x_3 = 0; \quad x_4 = 9; \quad x_5 = 4; \quad x_6 = 0. \quad (20.4)$$

For partial Problem 1, the basis vectors are thus given as A_1 and A_2 . The remaining vector (or vectors) follow from application of (5.1) to (2)

$$\begin{pmatrix} 1 \\ 3 \end{pmatrix} = d_{13} \begin{pmatrix} 1 \\ -1 \end{pmatrix} + d_{23} \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

whence

$$A_3 = 1 \cdot A_1 + 4 \cdot A_2 \quad (20.5)$$

and for the slack vectors,

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix} = s_1 = A_1 + A_2, \quad (20.6)$$

$$\begin{pmatrix} 0 \\ 1 \end{pmatrix} = s_2 = 0 \cdot A_1 + A_2. \quad (20.7)$$

In practice, these relations appear explicitly in the final optimal table of the partial problem by virtue of (5.2).

Similarly for (3),

$$A_6 = 5 \cdot A_4 - 2 \cdot A_5, \quad (20.8)$$

with slacks

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix} = s_3 = A_4 + A_5, \quad (20.9)$$

$$\begin{pmatrix} 0 \\ 1 \end{pmatrix} = s_4 = A_4 + 0 \cdot A_5. \quad (20.10)$$

The values (4) are now taken as a starting solution of (1). Since $\sum x_i = 20 > 16$, the present x_i can be made to satisfy the last inequality of (1) only on introduction of an artificial vector A_7 , as shown in the table, with an associated artificial variable $x_7 \geq 0$.

A_3 considered as a 2-component vector requires satisfaction of (5). As a 5-component vector its last non-zero component may be expressed in terms of the additional basis vector A_7 . Equation (5) then becomes

$$\begin{pmatrix} 1 \\ 3 \\ 0 \\ 0 \\ 1 \end{pmatrix} = 1 \begin{pmatrix} 1 \\ -1 \\ 0 \\ 0 \\ 1 \end{pmatrix} + 4 \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \\ 1 \end{pmatrix} + 4 \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ -1 \end{pmatrix},$$

i.e.

$$A_3 = 1 \cdot A_1 + 4 \cdot A_2 + 4 \cdot A_7 \quad (20.11)$$

and similarly

$$s_1 = 1 \cdot A_1 + 1 \cdot A_2 + 2 \cdot A_7, \quad (20.12)$$

$$s_2 = 1 \cdot A_2 + 1 \cdot A_7. \quad (20.13)$$

In the same way (8), (9), (10) are augmented to

$$A_6 = 5A_4 - 2A_5 + 2A_7, \quad (20.14)$$

$$s_3 = A_4 + A_5 + 2A_7, \quad (20.15)$$

$$s_4 = A_4 + A_7, \quad (20.16)$$

(4) gives

$$A_0 = 2A_1 + 5A_2 + 9A_4 + 4A_5 + 4A_7, \quad (20.17)$$

while from (1)

$$s_5 = -A_7. \quad (20.18)$$

These relations express the vectors of (1) in terms of A_1, A_2, A_4, A_5, A_7 taken as basis.

For the overall problem (1), we now wish to use the latter basis vectors as *unit* basis for the initial solution (4) and therefore write them in the form

$$A_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}; \quad A_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}; \quad A_4 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}; \quad A_5 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}; \quad A_7 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}. \quad (20.19)$$

This change of notation requires corresponding changes in the notation of all other column vectors such that the fundamental "basis relations" (11) to (18) remain unaltered. With notation (19), they become

$$\begin{aligned} A_3 &= \begin{bmatrix} 1 \\ 4 \\ 0 \\ 0 \\ 4 \end{bmatrix}; & s_1 &= \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \\ 2 \end{bmatrix}; & s_2 &= \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \\ 1 \end{bmatrix}; & A_6 &= \begin{bmatrix} 0 \\ 0 \\ 5 \\ -2 \\ 2 \end{bmatrix}; \\ s_3 &= \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \\ 2 \end{bmatrix}; & s_4 &= \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \\ 1 \end{bmatrix}; & A_0 &= \begin{bmatrix} 2 \\ 5 \\ 9 \\ 4 \\ 4 \end{bmatrix}; & s_5 &= \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ -1 \end{bmatrix}. \end{aligned} \quad (20.20)$$

Equation (1), converted to equalities and transformed to the new starting basis, now takes on the form

TABLE 5

c_j of basis	Basis	$c_j : 1$	1	3	1	1	1	$-M$	0	0	0	0	0	0
		A_1	A_2	A_3	A_4	A_5	A_6	A_7	s_1	s_2	s_3	s_4	s_5	A_0
1	A_1	1	0	1	0	0	0	0	1	0	0	0	0	= 2
1	A_2	0	1	4	0	0	0	0	1	1	0	0	0	= 5
1	A_4	0	0	0	1	0	5	0	0	0	1	1	0	= 9
1	A_5	0	0	0	0	1	-2	0	0	0	1	0	0	= 4
$-M$	A_7	0	0	4	0	0	2	1	2	1	2	1	-1	= 4
$g_j :$		0	0	$-4M$ +2	0	0	$-2M$ +2	0	$-2M$ +2	$-M$ +1	$-2M$ +2	$-M$ +1	$+M$	$-4M$ +20

From here on, a single iteration step yields the final solution $x_1 = 1, x_2 = 1, x_3 = 1, x_4 = 9, x_5 = 4, x_6 = 0, x_7 = 0, f_{\max} = 18$. Large problems may thus often be broken down into several smaller ones, for which the maximal solutions can be found comparatively easily. With the aid of the resulting partial optimal solutions, satisfaction of the remaining conditions, that tie the individual sections together, will then usually require only a few iterations of the re-formulated large original problem.

21. High-speed electronic computing. The advent of high-speed electronic computers renders the solution of large-scale linear programming problems economically feasible. A standard program was designed by the writer to carry-through the unmodified simplex procedure on the IBM 701, and is now used successfully. Data are handled in floating-decimal form, eliminating scaling analysis for each new problem as well as the common incidence of machine difficulties resulting from unforeseeable fluctuations in the size of numbers. With the present program, any size matrix (after augmentation by slack and artificial vectors) up to 50×100 can be accommodated. In cases of degeneracy, ties are resolved automatically by the procedure of Rule 10. Numerous checks are built into the program, which render undetected machine errors virtually impossible. The time required for an $m \times n$ matrix (all operations in floating-decimal form) is about $(0.5/1000) mn$ minutes per iteration.

The time-saving features in case of error detection may be noteworthy. Whenever one of the checks detects a calculation error, the program automatically restores certain data, depending on the subprogram concerned, skips back and tries again. If corrected on the second try, automatic continuation is assured without stopping. On persistence of an error, or on indication of infinite solutions, or on reaching the optimal solution, or if the number of completed iterations has reached a pre-assigned bound, the program automatically prints out the matrix for the last successfully completed iteration together with an identification of the nature and location of the condition that caused print-out.

The value of f for each iteration is likewise printed out to display an indication of the progress of convergence.

Equation (8.4) stipulates special treatment for the r th element of each column. When the transformation is carried out columnwise, this would require a special test for each i to determine whether or not it equals r . The elimination of such a test for each element at the cost of a few additional operations per column would contribute to greater speed. A little artifice was found to accomplish this purpose: at the start of the iteration, column k and the iteration divisor $a_{rk} = d$ are set aside and *the element a_{rk} of the transferred column k is replaced by zero*. Transformation of any column j then proceeds thus: (i) replace a_{ri} by $a'_{ri} = a_{ri}/d$; (ii) replace a_{ij} by $a'_{ij} = a_{ij} - a'_{ri}a_{rk}$ for all i . It is readily seen that this will automatically give the correct result for all i including $i = r$ and extend uniformity so far—should this be desired—as to include the case $j = k$ (the non-transferred unaltered column k) both for $i \neq r$ and for $i = r$.

22. Application to optimum gasoline blending. Application of Linear Programming to an industrial problem of wide scope is found in the determination of optimal blends of gasoline components to maximize profit within the framework of meeting specifications on technical properties and availability. The imposed constraints primarily stipulate ranges for vapor pressures, volatilities and octane numbers at various temperatures for the various grades of finished gasoline in terms of data given for the components, as well as supply limitations on each of the components.

As a typical instance, a problem expressed by an 18×34 matrix was solved by

machine in 29 iterations within 20 minutes. As is common with problems solved by means of Linear Programming, the maximum realizable profit was shown to be greater than the yield of past operations.

Conclusion. The impressive power of Linear Programming holds the promise of a significant contribution to the economic machinery of human society.

Acknowledgment. Grateful acknowledgment is made to Dr. A. Charnes of Carnegie Institute of Technology, to whom the writer is indebted for much valuable information and instruction, especially on the contents of Secs. 10, 19 and 20.

BIBLIOGRAPHY

1. A. Charnes, W. W. Cooper and A. Henderson: *An introduction to Linear Programming*, J. Wiley & Sons, 1953
2. *Symposium on linear inequalities and programming—Project SCOOP (Scientific computation of optimum programs)*, 1952, Manual No. 10, published by Directorate of Management Analysis Service, DCS/Comptroller, Hq. U. S. Air Force, Washington, D. C. (in particular the paper *Application of the simplex method to a variety of matrix problems* by Alex Orden)
3. *Activity analysis of production and allocation*, edited by T. C. Koopmans, J. Wiley & Sons, 1951
4. G. B. Dantzig, *Computational algorithm of the revised simplex method*, RAND Corp. publ. P-394-1 (1953)
5. Wm. Orchard—Hays: *Revisions and extensions to the simplex method*, RAND Corp. publ. P-562 (1954)
6. *Set-theoretic methods in economic analysis*, Notes of lectures by Prof. A. Charnes, Carnegie Institute of Technology, 1953
7. Hoffman, Mannos, Sokolowsky & Wiegmann, *Computational experience in solving linear programs*, J. Soc. Indust. and Appl. Math., 1, No. 1 (Sept. 1953)
8. Proceedings of the Association for Computing Machinery, May 1952, pp. 91-98
9. A. Charnes and W. W. Cooper, *The stepping stone method of explaining linear programming calculations in transportation problems*, Management Science 1, 49-69 (1954).