—NOTES—

ON THE MATERIAL TIME DERIVATIVE OF ARBITRARY ORDER*

BY L. N. TAO (*Illinois Institute of Technology*)

Abstract. The material time derivative of arbitrary order of a function of space and time is established. In the case of a function of a single space variable, it is reduced to Faa de Bruno's formula for derivatives of a function of a function.

This note is concerned with the material time derivative of arbitrary order of a function of space and time. These derivatives, in addition to their own merits, are needed in problems with non-stationary boundaries [1, 2]. The result is quite different from the Rivlin-Ericksen tensors for material time derivatives of the square of an infinitesimal arc length.

Let $f = f(x_1, x_2, x_3, t)$ and $x_i = x_i(t)$. Using a multinomial expansion, we have the Taylor's series

$$f = \sum_{m_1=0}^{\infty} \sum_{m_2=0}^{\infty} \sum_{m_3=0}^{\infty} \partial_t^{m_0} \partial_{x_1}^{m_1} \partial_{x_2}^{m_2} \partial_{x_3}^{m_3} f|_0 h^{m_0} k_1^{m_1} k_2^{m_2} k_3^{m_3} / (m_0! m_1! m_2! m_3!) \quad (1)$$

where $\partial_t \equiv \partial/\partial t$, $\partial_{x_i} \equiv \partial/\partial x_i$, $h = t - t_0$ and $k_i = x_i - x_i(t_0)$. The Taylor's series of $k_1^{m_1}$ is

$$k_1^{m_1} = \left(\sum_1^{\infty} D^p x_1(t_0) h^p / p! \right)^{m_1} = \sum_{r_1=m_1}^{\infty} m_1! A_{m_1}^{r_1}(x_1)|_0 h^{r_1} \quad (2)$$

where $D \equiv d/dt$. The coefficient of h^{r_1} in this series is

$$m_1! A_{m_1}^{r_1}(x_1)|_0 = \sum_{\alpha_k} \frac{m_1!}{\alpha_1! \alpha_2! \cdots \alpha_{r_1}!} (Dx_1|_0)^{\alpha_1} (D^2x_1|_0/2!)^{\alpha_2} \cdots (D^{r_1}x_1|_0/r_1!)^{\alpha_{r_1}}. \quad (3)$$

The sum is extended to all values of $\alpha_k = 0, 1, 2, \cdots$ which satisfy

$$\alpha_1 + \alpha_2 + \cdots + \alpha_{r_1} = m_1, \quad (4)$$

$$\alpha_1 + 2\alpha_2 + \cdots + r_1\alpha_{r_1} = r_1.$$

Similarly, we obtain

$$k_2^{m_2} = \sum_{r_2=m_2}^{\infty} m_2! A_{m_2}^{r_2}(x_2)|_0 h^{r_2}, \quad k_3^{m_3} = \sum_{r_3=m_3}^{\infty} m_3! A_{m_3}^{r_3}(x_3)|_0 h^{r_3} \quad (5)$$

where

$$A_{m_2}^{r_2}(x_2)|_0 = \sum_{\beta_k} \prod_1^{r_2} [(D^k x_2|_0/k!)^{\beta_k} / \beta_k!], \quad (6)$$

$$A_{m_3}^{r_3}(x_3)|_0 = \sum_{\gamma_k} \prod_1^{r_3} [(D^k x_3|_0/k!)^{\gamma_k} / \gamma_k!]$$

* Received September 30, 1977.

subject to

$$\begin{aligned}\sum \beta_k &= m_2, & \sum k\beta_k &= r_2, \\ \sum \gamma_k &= m_3, & \sum k\gamma_k &= r_3.\end{aligned}\quad (7)$$

Substitutions of (2) and (5) into (1) yield

$$f = \sum_{m_0=0}^{\infty} \sum_{m_1=0}^{\infty} \sum_{m_2=0}^{\infty} \sum_{r_1=m_1}^{\infty} \sum_{r_2=m_2}^{\infty} A_{m_1}^{r_1}(x_1)|_0 A_{m_2}^{r_2}(x_2)|_0 A_{m_3}^{r_3}(x_3)|_0 \partial_t^{m_0} \partial_{x_1}^{m_1} \partial_{x_2}^{m_2} \partial_{x_3}^{m_3} f|_0 h^N / m_0! \quad (8)$$

where $N = m_0 + r_1 + r_2 + r_3$. A change of the order of summations yields

$$f = \sum_{m_0=0}^{\infty} \sum_{r_1=0}^{\infty} \sum_{r_2=0}^{\infty} \sum_{m_1=0}^{r_1} \sum \cdots \quad (9)$$

The function f can also be written as

$$f = \sum_{N=0}^{\infty} D^N f|_0 h^N / N!. \quad (10)$$

Matching the coefficients of equal powers of h and using the fact that they are valid for arbitrary t_0 , we obtain

$$D^N f = \sum_{m_0=0}^N \sum_{r_1=0}^{N-m_0} \sum_{r_2=0}^{N-m_0-r_1} \sum_{m_1=0}^{r_1} \sum_{m_2=0}^{r_2} \sum_{m_3=0}^{r_3} \frac{N!}{m_0!} A_{m_1}^{r_1}(x_1) A_{m_2}^{r_2}(x_2) A_{m_3}^{r_3}(x_3) \partial_t^{m_0} \partial_{x_1}^{m_1} \partial_{x_2}^{m_2} \partial_{x_3}^{m_3} f \quad (11)$$

where $r_3 = N - m_0 - r_1 - r_2$. This is the required result.

In the one-dimensional case as needed in [1, 2], this result is reduced to

$$D^N f = \sum_{n=0}^N \sum_{m=0}^{N-n} (N!/n!) A_m^{N-n}(x) \partial_t^n \partial_x^m f. \quad (12)$$

Also, if f does not depend on t explicitly, then $n = 0$. It is further reduced to Faa de Bruno's formula of derivatives of a function of a function [3],

$$D^N f = \sum_0^N N! A_m^N(x) d^m f / dx^m. \quad (13)$$

REFERENCES

- [1] D. D. Joseph and R. L. Fosdick, *The free surface on a liquid between cylinders rotating at different speeds—I*, Arch. Rat. Mech. Anal. **49**, 321–380 (1973)
- [2] L. N. Tao, *The Stefan problem with arbitrary initial and boundary conditions*, Quart. Appl. Math., this issue
- [3] C. Jordan, *Calculus of finite differences*, Chelsea, 1965