UNIFORM STABILIZATION OF THE HIGHER-DIMENSIONAL SYSTEM OF THERMOELASTICITY WITH A NONLINEAR BOUNDARY FEEDBACK

By

WEI-JIU LIU (*Department of Mathematics and Statistics, Dalhousie University, Halifax, Nova Scotia, Canada*)

AND

ENRIQUE ZUAZUA (*Departamento de Matemática Aplicada, Universidad Complutense de Madrid, Spain*)

Abstract. Using multiplier techniques and Lyapunov methods, we derive explicit decay rates for the energy in the higher-dimensional system of thermoelasticity with a nonlinear velocity feedback on part of the boundary of a thermoelastic body, which is clamped along the rest of its boundary.

1. Introduction. Let Ω be a bounded domain in $\mathbf{R}^n$ with smooth boundary $\Gamma = \partial\Omega$ of class C^2 , and consider an n -dimensional linear, homogeneous, isotropic, and thermoelastic body occupying Ω in its non-deformed state. For a material point with configuration $x = (x_1, \dots, x_n)$ at time t , let $u(x, t) = (u_1(x, t), \dots, u_n(x, t))$ and $\theta(x, t)$ denote the displacement and temperature deviation, respectively, from the natural state of the reference configuration. Then, in the absence of external forces and heat sources, u and θ satisfy the system of equations of thermoelasticity

$$\begin{cases} u'' - \mu \Delta u - (\lambda + \mu) \nabla \operatorname{div} u + \alpha \nabla \theta = 0 & \text{in } \Omega \times (0, \infty), \\ \theta' - \Delta \theta + \beta \operatorname{div} u' = 0 & \text{in } \Omega \times (0, \infty), \\ u(0) = u^0, \quad u'(0) = u^1, \quad \theta(0) = \theta^0 & \text{in } \Omega, \end{cases} \quad (1.1)$$

where $\alpha, \beta > 0$ are the coupling parameters and λ, μ are Lamé's constants satisfying

$$\mu > 0, \quad n\lambda + (n+1)\mu > 0. \quad (1.2)$$

Received November 24, 1998.

2000 *Mathematics Subject Classification.* Primary 35B35, 74B05.

Key words and phrases. Thermoelasticity, uniform boundary stabilization, Lyapunov methods.

The work of the first author was supported in part by the Killam Postdoctoral Fellowship and done in part while the author was with UC San Diego.

The work of the second author was supported by grant PB96-0663 of the DGES (Spain).

E-mail address: weiliu@mathstat.dal.ca

E-mail address: zuazua@sunma4.mat.ucm.es

By ' we denote the derivative with respect to the time variable. Δ , ∇ , div denote the Laplace, gradient, and divergence operators in the space variables respectively. $u(0)$, $u'(0)$ and $\theta(0)$ denote the functions $x \rightarrow u(x, 0)$, $x \rightarrow u'(x, 0)$ and $x \rightarrow \theta(x, 0)$, respectively. For the derivation of (1.1), we refer to [24] and [44].

Note that condition (1.2) is weaker than the following usual condition on the Lamé coefficients (see [44] and [13, p. 414])

$$n\lambda + 2\mu > 0. \quad (1.3)$$

Extensive work has been done on the problem of stabilization for system (1.1) (see [5, 9, 7, 12, 17, 20, 27, 33, 37, 40, 42, 43, 45, 46, 47, 48]). We give here a brief description of the existing literature. For a detailed survey, we refer to [37, 38].

The thermoelastic energy of (1.1) can be defined as

$$\begin{aligned} E(u, \theta, t) = & \frac{1}{2} \int_{\Omega} \left[|u'(x, t)|^2 + \mu |\nabla u(x, t)|^2 \right. \\ & \left. + (\lambda + \mu) |\operatorname{div} u(x, t)|^2 + \frac{\alpha}{\beta} |\theta(x, t)|^2 \right] dx. \end{aligned} \quad (1.4)$$

Here we have used the notation

$$|\nabla u(x, t)|^2 = \sum_{i,j=1}^n \left| \frac{\partial u_i}{\partial x_j} \right|^2. \quad (1.5)$$

Under the Dirichlet-Dirichlet boundary conditions

$$u = 0, \quad \theta = 0 \quad \text{on } \Gamma \times (0, \theta), \quad (1.6)$$

it is easy to verify that

$$E'(u, \theta, t) = -\frac{\alpha}{\beta} \int_{\Omega} |\nabla \theta(x, t)|^2 dx. \quad (1.7)$$

Therefore, the energy $E(u, \theta, t)$ decreases on $(0, \infty)$. Furthermore, in the case of one space dimension, it has been shown (see [9, 17, 20, 40]) that the energy $E(u, \theta, t)$ of system (1.1) associated with various boundary conditions decays to zero exponentially. In the case of multi-space dimension, Dafermos in his pioneering work [12] showed that, generically with respect to the domain, the energy of every solution of (1.1) and (1.6) tends to zero as $t \rightarrow \infty$. However, he also pointed out that, when Ω is a ball, non-decaying solutions do exist. More recently, Lebeau and Zuazua [33] proved that the decay rate is never uniform when Ω is convex. Thus, in order to ensure the uniform decay rate for such convex domains, additional damping mechanisms are necessary. In this aspect, the first author (see [37]) introduced a linear boundary velocity feedback acting on the elastic component of the system and established the uniform decay rate.

The purpose of this paper is to introduce a nonlinear boundary feedback which allows us to test the robustness of the damping mechanisms. Under the classical polynomial growth assumption on the nonlinear boundary feedback near the origin, by using multiplier techniques and Lyapunov methods, we show that the energy in the multi-dimensional system of thermoelasticity decays to zero at an exponential or polynomial rate.

Further, even if the nonlinearity does degenerate at the origin faster than any polynomial, we show that the decay rate is governed by a dissipative ordinary differential equation. This allows us to show, in particular, that if the nonlinearity degenerates at the origin exponentially, then we obtain a logarithmic decay rate. To do that, we proceed as in [41] where the simpler case of the wave equation with internal damping is addressed. This type of result was obtained earlier by Lasiecka *et al* [32] for the wave equation with nonlinear boundary feedback. However, our approach, even if it uses some ingredients as in [32] (for instance, Jensen's inequality), relies essentially on the generalized Young's inequality and it is simpler. This allows us to get more explicit expressions for the nonlinearity entering in the differential inequality governing the decay of the energy.

The rest of this paper is organized as follows. In Section 2, we present our main results. In Section 3, we prove that the system of thermoelasticity with a nonlinear boundary feedback is well-posed by using the theory of nonlinear semigroups. Then, borrowing Lyapunov methods and multiplier techniques, we prove our main results in Section 4. Finally, in Section 5, we briefly discuss some special case and pose an open problem.

2. Main results. Let Ω be a bounded domain in $\mathbf{R}^n$ with smooth boundary $\Gamma = \partial\Omega$ of class C^2 . Set

$$\Gamma_1 = \{x \in \Gamma : m(x) \cdot \nu(x) \leq 0\}, \quad (2.1)$$

$$\Gamma_2 = \{x \in \Gamma : m(x) \cdot \nu(x) > 0\}, \quad (2.2)$$

where

$$m(x) = x - x^0 = (x_1 - x_1^0, \dots, x_n - x_n^0) \quad (2.3)$$

for some $x^0 \in \mathbf{R}^n$, $\nu = (\nu_1, \dots, \nu_n)$ denotes the unit normal on Γ directed towards the exterior of Ω and

$$m \cdot \nu = m(x) \cdot \nu(x) = \sum_{i=1}^n (x_i - x_i^0) \nu_i. \quad (2.4)$$

Γ_1 is assumed either to be empty or to have a nonempty interior relative to Γ .

In what follows, $H^s(\Omega)$ denotes the usual Sobolev space (see [1]) for any $s \in \mathbf{R}$. For $s \geq 0$, $H_0^s(\Omega)$ denotes the completion of $C_0^\infty(\Omega)$ in $H^s(\Omega)$, where $C_0^\infty(\Omega)$ denotes the space of all infinitely differentiable functions on Ω with compact support in Ω . Let X be a Banach space. We denote by $C^k([0, T]; X)$ the space of all k times continuously differentiable functions defined on $[0, T]$ with values in X , and write $C([0, T]; X)$ for $C^0([0, T]; X)$.

We further introduce other function spaces as follows:

$$H_{\Gamma_1}^1(\Omega) = \{u \in H^1(\Omega) : u = 0 \text{ on } \Gamma_1\}, \quad (2.5)$$

$$\mathcal{H} = (H_{\Gamma_1}^1(\Omega))^n \times (L^2(\Omega))^n \times L^2(\Omega). \quad (2.6)$$

We consider the thermoelastic system with a nonlinear boundary feedback

$$\begin{cases} u'' - \mu \Delta u - (\lambda + \mu) \nabla \operatorname{div} u + \alpha \nabla \theta = 0 & \text{in } \Omega \times (0, \infty), \\ \theta' - \Delta \theta + \beta \operatorname{div} u' = 0 & \text{in } \Omega \times (0, \infty), \\ \theta = 0 & \text{on } \Gamma \times (0, \infty), \\ u = 0 & \text{on } \Gamma_1 \times (0, \infty), \\ \mu \frac{\partial u}{\partial \nu} + (\lambda + \mu) \operatorname{div} (u) \nu \\ \quad + am \cdot \nu u + m \cdot \nu g(u') = 0 & \text{on } \Gamma_2 \times (0, \infty), \\ u(0) = u^0, \quad u'(0) = u^1, \quad \theta(0) = \theta^0 & \text{in } \Omega, \end{cases} \quad (2.7)$$

where $a = a(x)$ is a given nonnegative function on Γ_2 with

$$a(x) \in C^1(\Gamma_2), \quad (2.8)$$

and $g(u) = (g_1(u_1, \dots, u_n), \dots, g_n(u_1, \dots, u_n)) \in (C(\mathbf{R}^n))^n$ is a given vector function. Similar nonlinear boundary feedbacks were introduced for the wave equation (see [11, 31, 32, 50, 51]) and the equations of elasticity (see [28, 29]).

The elastic Lamé operator $\mu \Delta u - (\lambda + \mu) \nabla \operatorname{div} u$ may be written in divergence form $\operatorname{div} [\sigma_{ij}(u)]$, where the stress tensor $\sigma_{ij}(u)$ is given by

$$\sigma_{ij}(u) = 2\mu \varepsilon_{ij}(u) + \lambda \delta_{ij} \sum_{k=1}^n \varepsilon_{kk}(u), \quad (2.9)$$

and the linearized strain tensor $\varepsilon_{ij}(u)$ is given by

$$\varepsilon_{ij}(u) = \frac{1}{2} \left(\frac{\partial u_i}{\partial x_j} + \frac{\partial u_j}{\partial x_i} \right). \quad (2.10)$$

The boundary conditions describing the surface forces are given by

$$\sigma_{ij}(u) \nu_j + am \cdot \nu u_i = f_i. \quad (2.11)$$

Since

$$\mu \frac{\partial u_i}{\partial \nu} + (\lambda + \mu) \operatorname{div} (u) \nu_i = \sigma_{ij}(u) \nu_j + \mu \left(\frac{\partial u_j}{\partial x_j} \nu_i - \frac{\partial u_j}{\partial x_i} \nu_j \right), \quad (2.12)$$

the boundary condition in (2.7)

$$\mu \frac{\partial u}{\partial \nu} + (\lambda + \mu) \operatorname{div} (u) \nu + am \cdot \nu u + m \cdot \nu g(u') = 0 \quad (2.13)$$

can be easily transferred into (2.11) with

$$f_i(u) = -m \cdot \nu g_i(u') - \mu \left(\frac{\partial u_j}{\partial x_j} \nu_i - \frac{\partial u_j}{\partial x_i} \nu_j \right). \quad (2.14)$$

Therefore, system (2.7) may be viewed as the system of thermoelasticity subject to a boundary feedback force of the form (2.14). This feedback mechanism is however non-optimal since, due to the presence of the first-order space derivatives, its regularity is not sharp (one cannot guarantee that it belongs to $L^2(\partial\Omega \times (0, T))$ for finite energy solutions). Therefore, the question of analyzing the stabilization under the weaker feedback forces of the form

$$f_i(u) = -m \cdot \nu g_i(u'), \quad (2.15)$$

in which the last term in (2.14) has been dropped, is an interesting open problem. This analysis has been performed in [3, 19] in the context of the system of elasticity but, to our knowledge, this issue has not been addressed for the system of thermoelasticity.

Throughout this paper, we assume that

$$\Gamma_1 \neq \emptyset \quad \text{or} \quad a(x) \not\equiv 0. \quad (2.16)$$

We refer to Section 5 for a brief discussion of the case where $\Gamma_1 = \emptyset$ and $a(x) \equiv 0$. Under assumptions (1.2) and (2.16), one can easily show that the following norm on $(H_{\Gamma_1}^1(\Omega))^n$

$$\begin{aligned} \|u\|_{(H_{\Gamma_1}^1(\Omega))^n} &= \left(\int_{\Omega} [\mu |\nabla u|^2 + (\lambda + \mu) |\operatorname{div}(u)|^2] dx \right. \\ &\quad \left. + \int_{\Gamma_2} am \cdot \nu |u|^2 d\Gamma \right)^{1/2} \end{aligned} \quad (2.17)$$

is equivalent to the usual one induced by $(H^1(\Omega))^n$ (see [37]). Indeed, it suffices to note the following fact:

$$\mu |\nabla u|^2 + (\lambda + \mu) |\operatorname{div} u|^2 \geq \mu |\nabla u|^2, \quad \text{for } \lambda + \mu \geq 0, \quad (2.18)$$

and for $\lambda + \mu < 0$,

$$\begin{aligned} \mu |\nabla u|^2 + (\lambda + \mu) |\operatorname{div} u|^2 &= \mu \sum_{i,j=1}^n \left| \frac{\partial u_i}{\partial x_j} \right|^2 + (\lambda + \mu) \sum_{i,j=1}^n \frac{\partial u_i}{\partial x_i} \frac{\partial u_j}{\partial x_j} \\ &\geq \mu \sum_{i,j=1}^n \left| \frac{\partial u_i}{\partial x_j} \right|^2 + (\lambda + \mu) n \sum_i \left| \frac{\partial u_i}{\partial x_i} \right|^2 \\ &= \mu \sum_{i \neq j} \left| \frac{\partial u_i}{\partial x_j} \right|^2 + [n\lambda + (n+1)\mu] \sum_i \left| \frac{\partial u_i}{\partial x_i} \right|^2 \\ &\geq \min\{\mu, n\lambda + (n+1)\mu\} |\nabla u|^2. \end{aligned} \quad (2.19)$$

In the sequel, we use the following energy norm on $\mathcal{H}$:

$$\begin{aligned} \|(u, v, \theta)\|_{\mathcal{H}} &= \left(\int_{\Omega} [\mu |\nabla u|^2 + (\lambda + \mu) |\operatorname{div}(u)|^2 + |v|^2 + \frac{\alpha}{\beta} |\theta|^2] dx \right. \\ &\quad \left. + \int_{\Gamma_2} am \cdot \nu |u|^2 d\Gamma \right)^{1/2} \end{aligned} \quad (2.20)$$

for $(u, v, \theta) \in \mathcal{H}$, which is equivalent to the usual one induced by $(H^1(\Omega))^n \times (L^2(\Omega))^n \times L^2(\Omega)$.

In Section 3, we will prove that system (2.7) generates a nonlinear continuous semi-group $S(t)$ on $\mathcal{H}$. Thus, system (2.7) is well-posed.

In order to state our main results, we first introduce some constants. In what follows, we denote by $\|\cdot\|$ the norm of $L^2(\Omega)$ or $(L^2(\Omega))^n$. Set

$$R_0 = \max_{x \in \bar{\Omega}} |m(x)| = \max_{x \in \bar{\Omega}} \left| \sum_{k=1}^n (x_k - x_k^0)^2 \right|^{1/2}, \quad (2.21)$$

$$a_0 = \max_{x \in \Gamma_2} a(x), \quad (2.22)$$

$$K(a) = \frac{2a_0^2 R_0^2}{\mu} + (2-n)a_0. \quad (2.23)$$

Let γ be the smallest positive constant such that

$$\int_{\Gamma_2} |u|^2 d\Gamma \leq \gamma^2 \|u\|_{(H_{\Gamma_1}^1(\Omega))^n}^2, \quad \forall u \in (H_{\Gamma_1}^1(\Omega))^n. \quad (2.24)$$

Let λ_0 and λ_1 be the smallest positive constants, respectively, such that

$$\|u\| \leq \lambda_0 \|u\|_{(H_{\Gamma_1}^1(\Omega))^n}, \quad \forall u \in (H_{\Gamma_1}^1(\Omega))^n, \quad (2.25)$$

and

$$\|u\| \leq \lambda_1 \|\nabla u\|, \quad \forall u \in H_0^1(\Omega). \quad (2.26)$$

Since the decay rates depend also on the constant appearing in the following technical lemma, we present it before stating our main results. This lemma is helpful for dealing with the case where the potential a is large.

LEMMA 2.1. There exists a constant $k_1 > 0$, independent of u , such that the solution φ of

$$\begin{cases} \mu \Delta \varphi + (\lambda + \mu) \nabla \operatorname{div} \varphi = 0 & \text{in } \Omega, \\ \varphi = u & \text{on } \Gamma, \end{cases} \quad (2.27)$$

satisfies

$$\|\varphi\|^2 \leq k_1^2 \int_{\Gamma} |u|^2 d\Gamma, \quad (2.28)$$

and

$$\int_{\Omega} [\mu \nabla u \cdot \nabla \varphi + (\lambda + \mu) \operatorname{div}(u) \operatorname{div}(\varphi)] dx \geq 0, \quad (2.29)$$

for all $u \in (H^1(\Omega))^n$, where

$$\nabla u \cdot \nabla \varphi = \sum_{i=1}^n \nabla u_i \cdot \nabla \varphi_i. \quad (2.30)$$

Proof. For any $f \in (L^2(\Omega))^n$, we consider

$$\begin{cases} \mu \Delta \psi + (\lambda + \mu) \nabla \operatorname{div} \psi = f & \text{in } \Omega, \\ \psi = 0 & \text{on } \Gamma. \end{cases} \quad (2.31)$$

Multiplying (2.31) by φ and integrating over Ω by parts, we obtain

$$\int_{\Omega} \varphi \cdot f dx = \int_{\Gamma} \left[\mu \frac{\partial \psi}{\partial \nu} + (\lambda + \mu) \operatorname{div}(\psi) \nu \right] \cdot u d\Gamma. \quad (2.32)$$

On the other hand, there exists a constant $c > 0$ such that

$$\left\| \mu \frac{\partial \psi}{\partial \nu} + (\lambda + \mu) \operatorname{div}(\psi) \nu \right\|_{(L^2(\Gamma))^n} \leq c \|f\|, \quad \forall f \in (L^2(\Omega))^n. \quad (2.33)$$

To prove this inequality, we let $h = (h_1, \dots, h_n)$ be a vector field in $(C^1(\bar{\Omega}))^n$ satisfying

$$h = \nu \quad \text{on } \Gamma. \quad (2.34)$$

Multiplying (2.31) by $h \cdot \nabla \psi$ and integrating over Ω by parts, we obtain

$$\begin{aligned} \int_{\Omega} \Delta \psi_i h_k \frac{\partial \psi_i}{\partial x_k} &= \frac{1}{2} \int_{\Gamma} h \cdot \nu \left| \frac{\partial \psi_i}{\partial \nu} \right|^2 d\Gamma + \frac{1}{2} \int_{\Omega} \operatorname{div}(h) |\nabla \psi_i|^2 dx \\ &\quad - \int_{\Omega} \frac{\partial \psi_i}{\partial x_j} \frac{\partial \psi_i}{\partial x_k} \frac{\partial h_k}{\partial x_j} \end{aligned} \quad (2.35)$$

and

$$\begin{aligned} \int_{\Omega} \frac{\partial \operatorname{div}(\psi)}{\partial x_i} h_k \frac{\partial \psi_i}{\partial x_k} &= \frac{1}{2} \int_{\Gamma} h \cdot \nu |\operatorname{div}(\psi)|^2 d\Gamma + \frac{1}{2} \int_{\Omega} \operatorname{div}(h) |\operatorname{div}(\psi)|^2 dx \\ &\quad - \int_{\Omega} \operatorname{div}(\psi) \frac{\partial \psi_i}{\partial x_k} \frac{\partial h_k}{\partial x_i}. \end{aligned} \quad (2.36)$$

Here we have used the summation convention for repeated indices. It therefore follows that there exists a constant $c > 0$ such that

$$\begin{aligned} \int_{\Gamma} \left[\mu \left| \frac{\partial \psi}{\partial \nu} \right|^2 + (\lambda + \mu) |\operatorname{div}(\psi)|^2 \right] d\Gamma \\ \leq c \left[\int_{\Omega} (\mu |\nabla \psi_i|^2 + (\lambda + \mu) |\operatorname{div}(\psi)|^2) dx + \|f\|^2 \right]. \end{aligned} \quad (2.37)$$

In addition, multiplying (2.31) by ψ and integrating over Ω by parts, we deduce that there exists a constant $c > 0$ such that

$$\int_{\Omega} (\mu |\nabla \psi_i|^2 + (\lambda + \mu) |\operatorname{div}(\psi)|^2) dx \leq c \|f\|^2. \quad (2.38)$$

Hence, we have

$$\int_{\Gamma} \left[\mu \left| \frac{\partial \psi}{\partial \nu} \right|^2 + (\lambda + \mu) |\operatorname{div}(\psi)|^2 \right] d\Gamma \leq c \|f\|^2, \quad (2.39)$$

which implies (2.33). Consequently, (2.28) follows from (2.32) and (2.33).

To prove inequality (2.29), we multiply (2.27) by u and φ , respectively, and integrate over Ω by parts. This gives

$$\begin{aligned} \int_{\Omega} [\mu \nabla u \cdot \nabla \varphi + (\lambda + \mu) \operatorname{div}(u) \operatorname{div}(\varphi)] dx &= \int_{\Gamma} \left[\mu \frac{\partial \varphi}{\partial \nu} \cdot u + (\lambda + \mu) \nu \cdot u \operatorname{div}(\varphi) \right] d\Gamma \\ &= \int_{\Omega} [\mu |\nabla \varphi|^2 + (\lambda + \mu) |\operatorname{div}(\varphi)|^2] dx \\ &\geq 0. \end{aligned} \quad (2.40)$$

□

The thermoelastic energy of (2.7) is defined by

$$E(t) = E(u, \theta, t) = \frac{1}{2} \|(u(t), u'(t), \theta(t))\|_{\mathcal{H}}^2. \quad (2.41)$$

By a straightforward calculation, we obtain

$$E'(t) = - \int_{\Gamma_2} m \cdot \nu g(u'(t)) \cdot u'(t) d\Gamma - \frac{\alpha}{\beta} \|\nabla \theta(t)\|^2. \quad (2.42)$$

If g satisfies that $g(u) \cdot u \geq 0$ for all $u \in \mathbf{R}^n$, then the energy $E(t)$ decreases in $(0, \infty)$.

What is more, we have the following decay rates. These are our main results of this paper.

THEOREM 2.2. Let Γ_1 and Γ_2 be given by (2.1) and (2.2), respectively, satisfying

$$\Gamma_1 \cap \bar{\Gamma}_2 = \emptyset. \quad (2.43)$$

Suppose that (2.16) holds. Let the function $g \in (C(\mathbf{R}^n))^n$ satisfy the following conditions:

$$g(0) = 0, \quad (2.44)$$

$$|g(u^1) - g(u^2)| \leq k_2 |u^1 - u^2|^q, \quad \forall u^1, u^2 \in \mathbf{R}^n \text{ with } |u^1 - u^2| \leq 1, \quad (2.45)$$

$$|g(u^1) - g(u^2)| \leq k_2 |u^1 - u^2|, \quad \forall u^1, u^2 \in \mathbf{R}^n \text{ with } |u^1 - u^2| \geq 1, \quad (2.46)$$

$$g(u) \cdot u \geq k_3 |u|^{p+1}, \quad \forall u \in \mathbf{R}^n \text{ with } |u| \leq 1, \quad (2.47)$$

$$g(u) \cdot u \geq k_3 |u|^2, \quad \forall u \in \mathbf{R}^n \text{ with } |u| \geq 1, \quad (2.48)$$

$$0 \leq [g(u^1) - g(u^2)] \cdot (u^1 - u^2), \quad \forall u^1, u^2 \in \mathbf{R}^n, \quad (2.49)$$

for some constants $k_2, k_3 > 0$ and p, q with $0 < q \leq 1$. Suppose the function $a(x)$ satisfies

$$2K(a)R_0\gamma^2 < 1, \quad \text{for } n \leq 2, \quad (2.50)$$

or

$$a_0 \leq \frac{(n-2)\mu}{2R_0^2}, \quad \text{for } n \geq 3. \quad (2.51)$$

Then we have

(1) If $p = q = 1$, there exist some constants $M \geq 1$ and $\omega > 0$, independent of (u^0, u^1, θ^0) , such that

$$E(t) \leq ME(0)e^{-\omega t}, \quad \forall t \geq 0, \quad (2.52)$$

for all solutions of (2.7) with $(u^0, u^1, \theta^0) \in \mathcal{H}$.

(2) If $p+1 > 2q$, there exists $\tau > 0$, depending on (u^0, u^1, θ^0) , such that

$$E(t) \leq 4E(0) \left(1 + \frac{p+1-2q}{2q} \tau t \right)^{-\frac{2q}{p+1-2q}}, \quad \forall t \geq 0, \quad (2.53)$$

for all solutions of (2.7) with $(u^0, u^1, \theta^0) \in \mathcal{H}$. Further, the constants M , ω and τ can be explicitly given by

$$\sigma_0 = \frac{p+1-2q}{2q}, \quad (2.54)$$

$$\varepsilon = \begin{cases} (1-2K(a)R_0\gamma^2)/8, & n \leq 2, \\ 1/8, & n \geq 3, \end{cases} \quad (2.55)$$

$$C_1 = \frac{2R_0}{\sqrt{\mu}} + (n-1)\lambda_0, \quad (2.56)$$

$$C_2 = \frac{\alpha^2 R_0^2}{\mu\varepsilon} + \frac{\alpha^2(n-1)^2\lambda_0^2}{4\varepsilon} + \frac{\alpha\lambda_1^2}{\beta}, \quad (2.57)$$

$$C_3 = \frac{2R_0^2}{\mu} + \frac{(n-1)^2 R_0\gamma^2}{4\varepsilon}, \quad (2.58)$$

$$C_4 = 1 + C_3 k_2^2, \quad (2.59)$$

$$b_1 = \frac{1}{2}C_1^{-1}E^{-\sigma_0}(0), \quad (2.60)$$

$$b_2 = \alpha/[E^{\sigma_0}(0)(\sigma_0\alpha C_1 + \beta C_2)], \quad (2.61)$$

$$b_3 = k_3/[E^{\sigma_0}(0)(C_4 + \sigma_0 C_1 k_3)], \quad (2.62)$$

$$b_4 = k_3(p+1)^{(p+1)/(2q)}/[C_1\sigma_0 E^{\sigma_0}(0)k_3(p+1)^{(p+1)/(2q)} + q(2C_4)^{(p+1)/(2q)}[R_0\text{mes}(\Gamma_2)(p+1-2q)]^{\sigma_0}], \quad (2.63)$$

$$\delta_1 = \min\{1/(2C_1), \alpha/(\beta C_2), k_3/C_4\}, \quad (2.64)$$

$$\delta_2 = \min\{b_1, b_2, b_3, b_4\}, \quad (2.65)$$

$$M = (1 + \delta_1 C_1)/(1 - \delta_1 C_1) \leq 3, \quad (2.66)$$

$$\omega = \delta_1/(1 + \delta_1 C_1), \quad (2.67)$$

$$\tau = \delta_2/[2(2E(0))^{-\sigma_0}[1 + \delta_2 C_1 E^{\sigma_0}(0)]^{\sigma_0+1}]. \quad (2.68)$$

If the potential a does not satisfy the smallness conditions (2.50) and (2.51), then we do not know whether or not Theorem 2.2 still holds. However, we can deal with large potentials a under an additional condition on Γ_2 . Namely, we have

THEOREM 2.3. Assume that Γ_1, Γ_2 and g satisfy the conditions of Theorem 2.2. We further assume that

$$m \cdot \nu \geq \eta > 0 \quad \text{on } \Gamma_2. \quad (2.69)$$

Suppose (2.16) holds. Then the decay properties (2.52) and (2.53) hold.

Furthermore, the explicit values of the constants M , ω and τ are as follows:

$$\sigma_0 = \frac{p+1-2q}{2q} \quad (2.70)$$

$$\varepsilon = 1/8, \quad (2.71)$$

$$C_0 = \max \left\{ 0, \frac{2a_0 R_0^2}{\mu} + 2 - n \right\}, \quad (2.72)$$

$$C'_1 = \frac{2R_0}{\sqrt{\mu}} + (n-1)\lambda_0 + C_0 k_1 \gamma, \quad (2.73)$$

$$C'_2 = \frac{\alpha^2 R_0^2}{\mu \varepsilon} + \frac{\alpha^2 (n-1)^2 \lambda_0^2}{4\varepsilon} + \frac{C_0^2 \alpha^2 k_1^2 \gamma^2}{4\varepsilon} + \frac{\alpha \lambda_1^2}{\beta}, \quad (2.74)$$

$$C'_3 = \frac{2R_0^2}{\mu} + \frac{(n-1+C_0)^2 R_0 \gamma^2}{4\varepsilon} \quad (2.75)$$

$$C'_4 = R_0 + \frac{C_0^2 k_1^2}{4\varepsilon} + C'_3 k_2^2 R_0, \quad (2.76)$$

$$b'_1 = \frac{1}{2} C_1'^{-1} E^{-\sigma_0}(0), \quad (2.77)$$

$$b'_2 = \alpha / [E^{\sigma_0}(0)(\sigma_0 \alpha C'_1 + \beta C'_2)], \quad (2.78)$$

$$b'_3 = k_3 \eta / [E^{\sigma_0}(0)(C'_4 + \sigma_0 C'_1 k_3 R_0)], \quad (2.79)$$

$$b'_4 = \eta k_3 (p+1)^{(p+1)/(2q)} / [C'_1 \sigma_0 E^{\sigma_0}(0) R_0 k_3 (p+1)^{(p+1)/(2q)} + q[2C'_4]^{(p+1)/(2q)} [\text{mes}(\Gamma_2)(p+1-2q)]^{\sigma_0}], \quad (2.80)$$

$$\delta'_1 = \min\{1/(2C'_1), \alpha/(\beta C'_2), \eta k_3/C'_4\}, \quad (2.81)$$

$$\delta'_2 = \min\{b'_1, b'_2, b'_3, b'_4\}, \quad (2.82)$$

$$M = (1 + \delta'_1 C'_1) / (1 - \delta'_1 C'_1) \leq 3, \quad (2.83)$$

$$\omega = \delta'_1 / (1 + \delta'_1 C'_1), \quad (2.84)$$

$$\tau = \delta'_2 / [2(2E(0))^{-\sigma_0} [1 + \delta'_2 C'_1 E^{\sigma_0}(0)]^{\sigma_0+1}]. \quad (2.85)$$

EXAMPLE 2.4. It is easy to see that the following function

$$g(u) = g(u_1, u_2, \dots, u_n) = \begin{cases} (u_1^2 + u_2^2 + \dots + u_n^2)u, & \text{if } |u| \leq 1, \\ u, & \text{if } |u| \geq 1, \end{cases} \quad (2.86)$$

satisfies the conditions of Theorem 2.2 with $p = 3$ and any $0 < q \leq 1$. In fact, (2.44), (2.46), (2.47) and (2.48) are obvious. In addition, it is easy to verify that g is globally Lipschitz. Thus, for any $u^1, u^2 \in \mathbf{R}^n$ with $|u^1 - u^2| \leq 1$, we have

$$|g(u^1) - g(u^2)| \leq c|u^1 - u^2| \leq c|u^1 - u^2|^q. \quad (2.87)$$

To show (2.49), it suffices to show that the Jacobian matrix

$$\left(\frac{\partial g_i}{\partial u_j} \right) = \begin{pmatrix} \frac{\partial g_1}{\partial u_1} & \dots & \frac{\partial g_1}{\partial u_n} \\ \vdots & & \vdots \\ \frac{\partial g_n}{\partial u_1} & \dots & \frac{\partial g_n}{\partial u_n} \end{pmatrix} \quad (2.88)$$

is positive semi-definite. If $|u| \geq 1$, this is obvious. If $|u| \leq 1$, then

$$\left(\frac{\partial g_i}{\partial u_j} \right) = \begin{pmatrix} 3u_1^2 & 2u_1u_2 & \cdots & 2u_1u_n \\ 2u_1u_2 & 3u_2^2 & \cdots & 2u_2u_n \\ \cdots & \cdots & \cdots & \cdots \\ 2u_1u_n & 2u_2u_n & \cdots & 3u_n^2 \end{pmatrix} \quad (2.89)$$

is also positive semi-definite since we have

$$\begin{aligned} \begin{vmatrix} 3u_1^2 & 2u_1u_2 & \cdots & 2u_1u_k \\ 2u_1u_2 & 3u_2^2 & \cdots & 2u_2u_k \\ \cdots & \cdots & \cdots & \cdots \\ 2u_1u_k & 2u_2u_k & \cdots & 3u_k^2 \end{vmatrix} &= u_1^2u_2^2 \cdots u_k^2 \begin{vmatrix} 3 & 2 & \cdots & 2 \\ 2 & 3 & \cdots & 2 \\ \cdots & \cdots & \cdots & \cdots \\ 2 & 2 & \cdots & 3 \end{vmatrix} \\ &= u_1^2u_2^2 \cdots u_k^2 [3 + 2(k-1)] \\ &\geq 0. \end{aligned} \quad (2.90)$$

On the other hand, if we take $g(u) = (h(u_1), \dots, h(u_n))$, where

$$h(s) = \begin{cases} |s|^{p-1}s & \text{if } |s| \leq 1, \\ s, & \text{if } |s| \geq 1, \end{cases} \quad (2.91)$$

is a function on $\mathbf{R}$, then g does not satisfy (2.47) and (2.48). But one can expect that this kind of functions should produce good decay rates. Therefore, we amend the conditions on g in Theorems 2.2 and 2.3 and obtain the following theorem.

THEOREM 2.5. Let $h(s)$ be a continuous function on $\mathbf{R}$ satisfying the following conditions:

$$h(0) = 0, \quad (2.92)$$

$$|h(s_1) - h(s_2)| \leq k_2|s_1 - s_2|^q, \quad \forall s_1, s_2 \in \mathbf{R} \text{ with } |s_1 - s_2| \leq 1, \quad (2.93)$$

$$|h(s_1) - h(s_2)| \leq k_2|s_1 - s_2|, \quad \forall s_1, s_2 \in \mathbf{R} \text{ with } |s_1 - s_2| \geq 1, \quad (2.94)$$

$$h(s)s \geq k_3|s|^{p+1}, \quad \forall s \in \mathbf{R} \text{ with } |s| \leq 1, \quad (2.95)$$

$$h(s)s \geq k_3|s|^2, \quad \forall s \in \mathbf{R} \text{ with } |s| \geq 1, \quad (2.96)$$

$$0 \leq [h(s^1) - h(s^2)](s^1 - s^2), \quad \forall s^1, s^2 \in \mathbf{R}, \quad (2.97)$$

for some constants $k_2, k_3 > 0$ and p, q with $0 < q \leq 1$. Suppose that

$$g(u) = (h(u_1), \dots, h(u_n)). \quad (2.98)$$

Assume that the conditions on g in Theorem 2.2 are replaced by the above conditions and that the other conditions in Theorems 2.2 and 2.3 are kept unchanged. Then Theorems 2.2 and 2.3 still hold with the same constants except that b_4 is replaced by

$$\begin{aligned} &k_3(p+1)^{(p+1)/(2q)} / [C_1\sigma_0 E^{\sigma_0}(0)k_3(p+1)^{(p+1)/(2q)} \\ &+ q(2C_4)^{(p+1)/(2q)} [nR_0 \text{mes}(\Gamma_2)(p+1-2q)]^{\sigma_0}], \end{aligned} \quad (2.99)$$

and b'_4 is replaced by

$$\begin{aligned} &\eta k_3(p+1)^{(p+1)/(2q)} / [C'_1\sigma_0 E^{\sigma_0}(0)R_0k_3(p+1)^{(p+1)/(2q)} \\ &+ q[2C'_4]^{(p+1)/(2q)} [n \text{mes}(\Gamma_2)(p+1-2q)]^{\sigma_0}]. \end{aligned} \quad (2.100)$$

Furthermore, if $p > 1$ and $q = 1/p$, then the decay rate (2.53) can be refined to

$$E(t) \leq 4E(0) \left(1 + \frac{p-1}{2}\tau t\right)^{-\frac{2}{p-1}}, \quad \forall t \geq 0, \quad (2.101)$$

with the same constants except the following changes:

$$\sigma_0 = \frac{p-1}{2}, \quad (2.102)$$

$$b_4 = (p+1)^{(p+1)/2} / [C_1 \sigma_0 E^{\sigma_0}(0) (p+1)^{(p+1)/2} + (2C_5)^{(p+1)/2} [R_0 \text{mes}(\Gamma_2)(p-1)]^{\sigma_0}], \quad (2.103)$$

$$b'_4 = \eta(p+1)^{(p+1)/2} / [C'_1 \sigma_0 E^{\sigma_0}(0) R_0 (p+1)^{(p+1)/2} + [2C'_5]^{(p+1)/2} [\text{mes}(\Gamma_2)(p-1)]^{\sigma_0}], \quad (2.104)$$

$$C_5 = \frac{1 + C_3 k_2 k_3}{k_3}, \quad (2.105)$$

$$C'_5 = \frac{4\varepsilon R_0 + C_0 k_1^2 + 4\varepsilon C'_3 k_2 k_3 R_0}{k_3}. \quad (2.106)$$

REMARK 2.6. Note that condition (2.95) on h implies that

$$|h(s)| \geq k_3 |s|^p, \quad \forall s \in \mathbf{R} \text{ with } |s| \leq 1. \quad (2.107)$$

This means that $h(s)$ cannot degenerate at the origin faster than $|s|^p$. The following theorem provides a decay rate when the nonlinearity h degenerates faster than any polynomial.

THEOREM 2.7. Let $h(s)$ be a continuous function on $\mathbf{R}$ satisfying the following conditions:

$$h(0) = 0, \quad (2.108)$$

$$|h(s_1) - h(s_2)| \leq k_2 |s_1 - s_2|^q, \quad \forall s_1, s_2 \in \mathbf{R} \text{ with } |s_1 - s_2| \leq 1, \quad (2.109)$$

$$|h(s_1) - h(s_2)| \leq k_2 |s_1 - s_2|, \quad \forall s_1, s_2 \in \mathbf{R} \text{ with } |s_1 - s_2| \geq 1, \quad (2.110)$$

$$h(s)s \geq k_3 |s|^2, \quad \forall s \in \mathbf{R} \text{ with } |s| \geq 1, \quad (2.111)$$

$$0 \leq [h(s^1) - h(s^2)](s^1 - s^2), \quad \forall s^1, s^2 \in \mathbf{R}, \quad (2.112)$$

for some constants $k_2, k_3 > 0$ and $0 < q \leq 1$. Suppose that

$$g(u) = (h(u_1), \dots, h(u_n)). \quad (2.113)$$

Assume that (2.16), (2.43) and (2.69) hold. Let $\varphi(s)$ denote an increasing and convex function defined on $[0, \infty]$ and twice differentiable outside $s = 0$ such that $\varphi(s^{2q}) \leq h(s)s$ on $[-1, 1]$. Then the energy $E(t)$ of solutions of (2.7) with $(u^0, u^1, \theta^0) \in \mathcal{H}$ satisfies the following decay rate:

$$E(t) \leq 2V(t), \quad \text{for } t \geq 0, \quad (2.114)$$

where $V(t)$ is the solution of the following differential equation:

$$V'(t) = -\frac{\delta V(t)}{2b} \varphi' \left(\frac{aV(t)}{b} \right) - n\delta(R_0 + 2C_0^2 k_1^2 + C_3' k_2^2 R_0) \text{mes}(\Gamma_2) \varphi \left(\frac{aV(t)}{b} \right), \quad (2.115)$$

where δ is a sufficiently small positive constant and

$$a = \frac{1}{2n(R_0 + 2C_0^2 k_1^2 + C_3' k_2^2 R_0) \text{mes}(\Gamma_2)}, \quad (2.116)$$

$$b = 1 + \delta C_1' \varphi'(aE(0)). \quad (2.117)$$

Furthermore, we have

$$\lim_{t \rightarrow \infty} E(t) = \lim_{t \rightarrow \infty} V(t) = 0. \quad (2.118)$$

REMARK 2.8. The function φ that satisfies the conditions of Theorem 2.7 always exists. For example, we set

$$\bar{\varphi}(s) = \text{conv}[\min\{s^{1/(2q)}h(s^{1/(2q)}), -s^{1/(2q)}h(-s^{1/(2q)})\}], \quad \text{for } 0 \leq s \leq 1, \quad (2.119)$$

and extend it to $[0, \infty)$. Here conv denotes the convex envelope of a function. Then we can take an increasing, convex and twice differentiable function $\varphi(s)$ such that $\varphi(s) \leq \bar{\varphi}(s)$.

In the special case where $s^{1/(2q)}h(s^{1/(2q)})$ is convex and $h(s)$ is odd, we have:

COROLLARY 2.9. Assume that all the conditions of Theorem 2.7 hold. If, further, h is odd in $[-1, 1]$ and $s^{1/(2q)}h(s^{1/(2q)})$ is convex on $[0, 1]$, then the energy $E(t)$ of (2.7) satisfies the following decay rate:

$$E(t) \leq 2V(t), \quad \text{for } t \geq 0, \quad (2.120)$$

where, for t large enough, $V(t)$ satisfies the following differential equation:

$$V'(t) = -K_1 V^{1/(2q)} h\left(\left(\frac{aV}{b}\right)^{1/(2q)}\right) - K_2 V^{1/q} h'\left(\left(\frac{aV}{b}\right)^{1/(2q)}\right), \quad (2.121)$$

where K_1, K_2 are positive constants independent of V .

COROLLARY 2.10. Assume that all the conditions of Theorem 2.7 hold. If, further, h satisfies (2.95), then the decay properties (2.52) and (2.53) hold.

Corollary 2.10 shows that Theorem 2.5 is covered by Theorem 2.7 when (2.69) holds. Since Theorem 2.7 does not include the case where (2.69) does not hold, we separate it from Theorem 2.7.

We now give an example of logarithmic decay rate which complements the example of polynomial decay rate existing in the literature.

EXAMPLE 2.11. *Logarithmic Decay Rate.* Let $h(s)$ satisfy

$$h(s) = s^3 e^{-1/s^2}, \quad |s| \leq 1, \quad (2.122)$$

$$c_1 |s| \leq h(s) \leq |s|, \quad |s| \geq 1. \quad (2.123)$$

It is easy to see that h satisfies all the conditions of Corollary 2.9 with $q = 1$. Consequently, for t large enough, by (2.121), V satisfies

$$V'(t) \leq -\omega V^2 e^{-\frac{b}{aV}}, \quad (2.124)$$

which is the same as

$$(e^{\frac{b}{aV}})' \geq \frac{b\omega}{a}, \quad (2.125)$$

where ω is a positive constant independent of V . Solving the inequality, we obtain the logarithmic decay rate

$$V(t) \leq \frac{b}{a} \left[\log \left(\frac{b\omega}{a} t + e^{\frac{b}{aV(0)}} \right) \right]^{-1}. \quad (2.126)$$

REMARK 2.12. The decay rate of the form (2.101) has been established for the wave equation [21, p. 127] and the compactly coupled wave equations [22]. It can be seen from the proof of Theorem 2.5 at the end of section 4 that the key point to obtain this decay rate $t^{-2/(p-1)}$ is to enlarge the following inequality:

$$s^2 + |h(s)|^2 \leq c(h(s)s)^{2/(p+1)}. \quad (2.127)$$

This can be done as follows. We first deduce from (2.95) that

$$s^2 \leq c(h(s)s)^{2/(p+1)}. \quad (2.128)$$

Next, we have

$$|h(s)| \leq c|s|^{1/p}, \quad (2.129)$$

$$|h(s)|^{2p/(p+1)} \leq c|s|^{2/(p+1)}, \quad (2.130)$$

$$|h(s)|^{(2p+2-2)/(p+1)} \leq c|s|^{2/(p+1)}, \quad (2.131)$$

$$|h(s)|^2 \leq c|s|^{2/(p+1)}|^{2/(p+1)}|h(s)|^{2/(p+1)} = c|h(s)s|^{2/(p+1)}. \quad (2.132)$$

In the situation of Theorems 2.2 and 2.3, since, for the vector function $g(u)$ and vector u , we have

$$|u|^{2/(p+1)}|g(u)|^{2/(p+1)} \geq |u \cdot g(u)|^{2/(p+1)}, \quad (2.133)$$

the analogous inequality of (2.127) for g no longer holds. Thus, the decay rate (2.53) of Theorems 2.2 and 2.3 cannot be refined to (2.101) in the case where $p > 1$ and $q = 1/p$.

REMARK 2.13. Observe that (2.44), (2.45) and (2.47) imply that $q \geq p$. Therefore, under the assumption $0 < q \leq 1$, we have $2q \leq p+1$. If $2q = p+1$, then $q-1 = p-q \geq 0$. Hence we have $q = p = 1$. Therefore, we have only two cases: $q = p = 1$ and $2q < p+1$, as stated in Theorem 2.2.

REMARK 2.14. Under condition (2.43), assumptions (2.1) and (2.2) imply that the domain Ω is simply connected and star-shaped with respect to $x^0 \in \Omega$ (in which case $\Gamma_1 = \emptyset$) or $\Omega = \Omega_1 - \Omega_2$, both Ω_1 and Ω_2 being star-shaped with respect to x^0 . If $\Gamma_1 \cap \bar{\Gamma}_2 \neq \emptyset$, it is well known that the solution of (2.7) is not regular enough (see [14]) to perform the integrations by parts that we will do later. Thus, the obtention of decay rates in this case is an open problem. The extension of the results of this paper to the case $\Gamma_1 \cap \bar{\Gamma}_2 \neq \emptyset$ requires a careful analysis of the singularities that the solution may develop on $\Gamma_1 \cap \bar{\Gamma}_2$ as in [16].

REMARK 2.15. The expressions of ω and τ in the decay rates may look complex. But they are useful since they provide explicit estimates of the dependence of the decay rates ω and τ on the various parameters $\alpha, \beta, \Omega, k_2, k_3, \dots$, etc. For example, if we take a linear boundary feedback $g(s) = k_3 s$ and let $k_3 \rightarrow 0$, then it follows from (2.64) and (2.67) that $\omega \rightarrow 0$. In addition, by these expressions, we can analyze the limit of the polynomial

decay rate $(1 + \frac{p+1-2q}{2q}\tau t)^{-\frac{2q}{p+1-2q}}$ as p, q tend to 1 and recover the exponential decay of the case $p = q = 1$. Indeed, it is easy to see that

$$\lim_{p, q \rightarrow 1} \delta_2(p, q) = \min\{\frac{1}{2}C_1, \alpha/(\beta C_2), k_3/C_4\} = \delta_1, \quad (2.134)$$

$$\lim_{p, q \rightarrow 1} \tau(p, q) = \delta_1[2(1 + \delta_1 C_1)] = \omega/2, \quad (2.135)$$

$$\lim_{p, q \rightarrow 1} \left(1 + \frac{p+1-2q}{2q}\tau t\right)^{-\frac{2q}{p+1-2q}} = e^{-\omega t/2}. \quad (2.136)$$

REMARK 2.16. For the linear elastodynamic system, the uniform stabilization with the boundary feedback of the form

$$\sigma_{ij}(u)\nu_j = u'_i \quad (2.137)$$

was established by Horn [18] by developing microlocal estimates (see [19]) for tangential derivatives of the solutions of the elastodynamic system. However, this remains to be done for the system of thermoelasticity.

3. Well-posedness. In this section, we use the theory of nonlinear semigroups to treat the problem of well-posedness of (2.7). Therefore, we formulate (2.7) as an abstract Cauchy problem.

Let $\langle \cdot, \cdot \rangle$ denote the duality pairing between $(H_{\Gamma_1}^1(\Omega))^n$ and $[(H_{\Gamma_1}^1(\Omega))^n]'$ or $H_0^1(\Omega)$ and $H^{-1}(\Omega)$. We define the duality operator $A : (H_{\Gamma_1}^1(\Omega))^n \rightarrow [(H_{\Gamma_1}^1(\Omega))^n]'$ by

$$\langle Au, v \rangle = (u, v)_{(H_{\Gamma_1}^1(\Omega))^n}, \quad \forall u, v \in (H_{\Gamma_1}^1(\Omega))^n, \quad (3.1)$$

and the duality operator $A_0 : H_0^1(\Omega) \rightarrow H^{-1}(\Omega)$ by

$$\langle A_0 u, v \rangle = (u, v)_{H_0^1(\Omega)}, \quad \forall u, v \in H_0^1(\Omega). \quad (3.2)$$

The Riesz representation theorem ensures that A and A_0 are isometric isomorphisms of $(H_{\Gamma_1}^1(\Omega))^n$ onto $[(H_{\Gamma_1}^1(\Omega))^n]'$ and $H_0^1(\Omega)$ onto $H^{-1}(\Omega)$, respectively.

Also, we define a nonlinear operator B by

$$\langle Bu, v \rangle = \int_{\Gamma_2} m \cdot \nu g(u) \cdot v \, d\Gamma, \quad \forall u, v \in (H_{\Gamma_1}^1(\Omega))^n. \quad (3.3)$$

LEMMA 3.1. Suppose that the function $g \in (C(\mathbf{R}^n))^n$. If there exist constants $c > 0$ and $\sigma \geq 0$ such that for $u \in \mathbf{R}^n$

$$|g(u)| \leq \begin{cases} c[1 + |u|^{n/(n-2)}], & n \geq 3, \\ c(1 + |u|^\sigma), & n = 2, \end{cases} \quad (3.4)$$

then B maps $(H_{\Gamma_1}^1(\Omega))^n$ into $[(H_{\Gamma_1}^1(\Omega))^n]'$. Furthermore, B is hemicontinuous, that is, we have

$$\lim_{t \rightarrow 0} \langle B(u^1 + tu^2), v \rangle = \langle Bu^1, v \rangle \quad (3.5)$$

for any $u^1, u^2, v \in (H_{\Gamma_1}^1(\Omega))^n$.

Proof. If $n \geq 3$, then, by the trace theorem (see [36, Chap. 1]) and the imbedding theorem (see [1, p. 217]), we have

$$H_{\Gamma_1}^1(\Omega) \hookrightarrow H^{1/2}(\Gamma) \subset L^{(2n-2)/(n-2)}(\Gamma). \quad (3.6)$$

Thus, we have

$$\begin{aligned} |\langle Bu, v \rangle| &= \left| \int_{\Gamma_2} m \cdot \nu g(u) \cdot v \, d\Gamma \right| \\ &\leq c \left(\int_{\Gamma_2} |g(u)|^{(2n-2)/n} \, d\Gamma \right)^{n/(2n-2)} \left(\int_{\Gamma_2} |v|^{(2n-2)/(n-2)} \, d\Gamma \right)^{(n-2)/(2n-2)} \\ &\leq c \left[1 + \left(\int_{\Gamma_2} |u|^{(2n-2)/(n-2)} \, d\Gamma \right)^{n/(2n-2)} \right] \|v\|_{(H_{\Gamma_1}^1(\Omega))^n} \\ &\leq c[1 + \|u\|_{(H_{\Gamma_1}^1(\Omega))^n}^{n/(n-2)}] \|v\|_{(H_{\Gamma_1}^1(\Omega))^n}. \end{aligned} \quad (3.7)$$

This shows that $Bu \in [(H_{\Gamma_1}^1(\Omega))^n]'$.

If $n = 2$, then, by the trace theorem (see [36, Chap. 1]) and the imbedding theorem (see [1, p. 217]), we have

$$H_{\Gamma_1}^1(\Omega) \hookrightarrow H^{1/2}(\Gamma) \subset L^r(\Gamma) \quad (3.8)$$

for any $2 \leq r < \infty$. In addition, we may as well assume that $\sigma \geq 1$ since, by condition (3.4), we have

$$|g(u)| \leq 2c \leq 2c(1 + |u|^{\sigma+1}), \quad \text{for } |u| \leq 1, \quad (3.9)$$

and

$$|g(u)| \leq c(1 + |u|^{\sigma+1}) \leq 2c(1 + |u|^{\sigma+1}), \quad \text{for } |u| \geq 1. \quad (3.10)$$

It therefore follows that

$$\begin{aligned} |\langle Bu, v \rangle| &= \left| \int_{\Gamma_2} m \cdot \nu g(u) \cdot v \, d\Gamma \right| \\ &\leq c \left(\int_{\Gamma_2} |g(u)|^2 \, d\Gamma \right)^{1/2} \left(\int_{\Gamma_2} |v|^2 \, d\Gamma \right)^{1/2} \\ &\leq c \left[1 + \left(\int_{\Gamma_2} |u|^{2\sigma} \, d\Gamma \right)^{1/2} \right] \|v\|_{(H_{\Gamma_1}^1(\Omega))^n} \\ &\leq c[1 + \|u\|_{(H_{\Gamma_1}^1(\Omega))^n}^\sigma] \|v\|_{(H_{\Gamma_1}^1(\Omega))^n}. \end{aligned} \quad (3.11)$$

This shows that $Bu \in [(H_{\Gamma_1}^1(\Omega))^n]'$.

If $n = 1$, this is just the consequence of the following embedding:

$$H_{\Gamma_1}^1(\Omega) \hookrightarrow C(\bar{\Omega}). \quad (3.12)$$

It remains to prove that B is hemicontinuous. By the continuity of g , we deduce that for any $u^1, u^2, v \in (H_{\Gamma_1}^1(\Omega))^n$

$$g(u^1 + tu^2) \rightarrow g(u^1) \quad \text{as } t \rightarrow 0 \text{ a.e. on } \Gamma. \quad (3.13)$$

It therefore follows from Lebesgue's dominated convergence theorem and (3.4) that

$$\begin{aligned}\lim_{t \rightarrow 0} \langle B(u^1 + tu^2), v \rangle &= \int_{\Gamma_2} m \cdot \nu g(u^1 + tu^2) \cdot v \, d\Gamma \\ &= \int_{\Gamma_2} m \cdot \nu g(u^1) \cdot v \, d\Gamma \\ &= \langle Bu^1, v \rangle.\end{aligned}\tag{3.14}$$

This shows that B is hemicontinuous.

Note that if g satisfies (2.45) and (2.46) of Theorem 2.2 (or (2.93) and (2.94) of Theorem 2.5), then g satisfies (3.4). Thus B maps $(H_{\Gamma_1}^1(\Omega))^n$ into $[(H_{\Gamma_1}^1(\Omega))^n]'$.

Using the operators A , A_0 and B , we can formally transform problem (2.7) into an abstract Cauchy problem. In doing so, we multiply the first equation of (2.7) by $v \in (H_{\Gamma_1}^1(\Omega))^n$ and integrate over Ω by parts. This gives

$$\begin{aligned}0 &= \int_{\Omega} (u'' - \mu \Delta u - (\lambda + \mu) \nabla \operatorname{div} u + \alpha \nabla \theta) \cdot v \, dx \\ &= \int_{\Omega} u'' \cdot v \, dx - \mu \int_{\Gamma} \frac{\partial u}{\partial \nu} \cdot v \, d\Gamma - (\lambda + \mu) \int_{\Gamma} v \cdot \nu \operatorname{div} (u) \, d\Gamma \\ &\quad + \int_{\Omega} (\mu \nabla u \cdot \nabla v + (\lambda + \mu) \operatorname{div} u \operatorname{div} v) \, dx + \int_{\Omega} \alpha \nabla \theta \cdot v \, dx \\ &= \int_{\Omega} u'' \cdot v \, dx + \int_{\Gamma} am \cdot \nu u \cdot v \, d\Gamma + \int_{\Gamma} m \cdot \nu g(u') \cdot v \, d\Gamma \\ &\quad + \int_{\Omega} (\mu \nabla u \cdot \nabla v + (\lambda + \mu) \operatorname{div} u \operatorname{div} v) \, dx + \int_{\Omega} \alpha \nabla \theta \cdot v \, dx \\ &= \langle u'', v \rangle + \langle Au, v \rangle + \langle Bu', v \rangle + \langle \alpha \nabla \theta, v \rangle,\end{aligned}\tag{3.15}$$

and therefore,

$$u'' + Au + Bu' + \alpha \nabla \theta = 0.\tag{3.16}$$

Similarly, multiplying the second equation of (2.7) by $v \in H_0^1(\Omega)$ and integrating over Ω by parts, we obtain

$$\theta' + A_0 \theta + \beta \operatorname{div} u' = 0.\tag{3.17}$$

Setting

$$\Phi = (u, u', \theta),\tag{3.18}$$

$$\Lambda \Phi = (-u', Au + Bu' + \alpha \nabla \theta, A_0 \theta + \beta \operatorname{div} u'),\tag{3.19}$$

we transform (2.7) into

$$\begin{cases} \Phi' + \Lambda \Phi = 0, & t > 0, \\ \Phi(0) = (u^0, u^1, \theta^0). \end{cases}\tag{3.20}$$

This leads us to define the solution of (2.7) as that of (3.20).

Consider the nonlinear operator Λ on $\mathcal{H} = (H_{\Gamma_1}^1(\Omega))^n \times (L^2(\Omega))^n \times L^2(\Omega)$ with the domain

$$D(\Lambda) = \{(u, v, \theta) \in \mathcal{H} : v \in (H_{\Gamma_1}^1(\Omega))^n, Au + Bv \in (L^2(\Omega))^n, \theta \in H^2(\Omega) \cap H_0^1(\Omega)\}. \quad (3.21)$$

We are going to prove that $-\Lambda$ is m -dissipative. For the definition of m -dissipativeness, we refer to [6, p. 71]. $\square$

In what follows, we denote by $(\cdot, \cdot)$ the inner product in $L^2(\Omega)$ or $(L^2(\Omega))^n$.

LEMMA 3.2. Suppose that g satisfies (2.44), (2.49) (or (2.92), (2.97)) and (3.4). Then the operator $-\Lambda$ is m -dissipative on $\mathcal{H}$.

Proof. By (2.49) (or (2.97)), we obtain that for any $(u^1, v^1, \theta^1), (u^2, v^2, \theta^2) \in D(\Lambda)$

$$\begin{aligned} & (\Lambda(u^1, v^1, \theta^1) - \Lambda(u^2, v^2, \theta^2), (u^1, v^1, \theta^1) - (u^2, v^2, \theta^2))_{\mathcal{H}} \\ &= (v^2 - v^1, u^1 - u^2)_{(H_{\Gamma_1}^1(\Omega))^n} + \frac{\alpha}{\beta}(A_0(\theta^1 - \theta^2) + \beta \operatorname{div}(v^1 - v^2), \theta^1 - \theta^2) \\ & \quad + (A(u^1 - u^2) + Bv^1 - Bv^2 + \alpha \nabla(\theta^1 - \theta^2), v^1 - v^2) \\ &= \int_{\Gamma_2} m \cdot \nu [g(v^1) - g(v^2)] \cdot (v^1 - v^2) d\Gamma + \frac{\alpha}{\beta} \|\nabla(\theta^1 - \theta^2)\|^2 \\ &\geq 0. \end{aligned} \quad (3.22)$$

Thus, $-\Lambda$ is dissipative.

It remains to show that

$$(I + \Lambda)(D(\Lambda)) = \mathcal{H}. \quad (3.23)$$

Namely, we want to prove that

$$\begin{cases} u - v = \varphi, \\ v + Au + Bv + \alpha \nabla \theta = \psi, \\ \theta + A_0 \theta + \beta \operatorname{div} v = \xi \end{cases} \quad (3.24)$$

has a solution $(u, v, \theta) \in D(\Lambda)$ for every $(\varphi, \psi, \xi) \in \mathcal{H}$. For this, it suffices to prove that the following problem

$$\begin{cases} v + Av + Bv + \alpha \nabla \theta = \psi - A\varphi, \\ \theta + A_0 \theta + \beta \operatorname{div} v = \xi \end{cases} \quad (3.25)$$

has a solution $v \in (H_{\Gamma_1}^1(\Omega))^n$ and $\theta \in H^2(\Omega) \cap H_0^1(\Omega)$ for every $(\varphi, \psi, \xi) \in \mathcal{H}$. In fact, if this has been done, then, by setting $u = v + \varphi$, it is easy to see that (u, v, θ) satisfies (3.24). Further, we have

$$Au + Bv = \psi - v - \alpha \nabla \theta \in (L^2(\Omega))^n. \quad (3.26)$$

Thus, we have $(u, v, \theta) \in D(\Lambda)$.

To prove that (3.25) has a required solution, it suffices to show that the nonlinear operator $\mathcal{A}$ defined by

$$\mathcal{A}(v, \theta) = (v + Av + Bv + \alpha \nabla \theta, \frac{\alpha}{\beta} \theta + \frac{\alpha}{\beta} A_0 \theta + \alpha \operatorname{div} v) \quad (3.27)$$

maps $(H_{\Gamma_1}^1(\Omega))^n \times H_0^1(\Omega)$ onto $[(H_{\Gamma_1}^1(\Omega))^n]' \times H^{-1}(\Omega)$, that is,

$$\mathcal{A}((H_{\Gamma_1}^1(\Omega))^n \times H_0^1(\Omega)) = [(H_{\Gamma_1}^1(\Omega))^n]' \times H^{-1}(\Omega). \quad (3.28)$$

In fact, if this has been done, then, since $\psi - A\varphi \in [(H_{\Gamma_1}^1(\Omega))^n]'$ and $\frac{\alpha}{\beta}\xi \in L^2(\Omega)$, there exist $(v, \theta) \in (H_{\Gamma_1}^1(\Omega))^n \times H_0^1(\Omega)$ such that (3.25) holds. Further, since $A_0\theta = \xi - \theta - \beta \operatorname{div} v \in L^2(\Omega)$, we have $\theta \in H^2(\Omega) \cap H_0^1(\Omega)$.

To prove (3.28), by Theorem 1.3 of [6, p. 40], it suffices to show $\mathcal{A}$ is monotone, coercive and hemicontinuous. For any $(v^1, \theta^1), (v^2, \theta^2) \in (H_{\Gamma_1}^1(\Omega))^n \times H_0^1(\Omega)$, we have

$$\begin{aligned} & \langle \mathcal{A}(v^1, \theta^1) - \mathcal{A}(v^2, \theta^2), (v^1, \theta^1) - (v^2, \theta^2) \rangle \\ &= \langle v^1 - v^2, v^1 - v^2 \rangle + \langle A(v^1 - v^2), v^1 - v^2 \rangle + \langle Bv^1 - Bv^2, v^1 - v^2 \rangle \\ &+ \langle \alpha \nabla(\theta^1 - \theta^2), v^1 - v^2 \rangle + \frac{\alpha}{\beta} \langle \theta^1 - \theta^2, \theta^1 - \theta^2 \rangle \\ &+ \frac{\alpha}{\beta} \langle A_0(\theta^1 - \theta^2), \theta^1 - \theta^2 \rangle + \alpha \langle \operatorname{div}(v^1 - v^2), \theta^1 - \theta^2 \rangle \\ &\geq 0. \end{aligned} \quad (3.29)$$

So $\mathcal{A}$ is monotone. Similarly, we have

$$\begin{aligned} \langle \mathcal{A}(v, \theta), (v, \theta) \rangle &= \langle v, v \rangle + \langle Av, v \rangle + \langle Bv, v \rangle \\ &+ \frac{\alpha}{\beta} \langle \theta, \theta \rangle + \frac{\alpha}{\beta} \langle A_0\theta, \theta \rangle \\ &\geq \|v\|_{(H_{\Gamma_1}^1(\Omega))^n}^2 + \frac{\alpha}{\beta} \|\theta\|_{H_0^1(\Omega)}^2. \end{aligned} \quad (3.30)$$

Thus $\mathcal{A}$ is coercive. On the other hand, by Lemma 3.1, we can readily deduce that $\mathcal{A}$ is hemicontinuous. That is, we have

$$\lim_{t \rightarrow 0} \langle \mathcal{A}[(v^1, \theta^1) + t(v^2, \theta^2)], (v, \theta) \rangle = \langle \mathcal{A}(v^1, \theta^1), (v, \theta) \rangle \quad (3.31)$$

for any $(v^1, \theta^1), (v^2, \theta^2), (v, \theta) \in (H_{\Gamma_1}^1(\Omega))^n \times H_0^1(\Omega)$. This completes the proof. $\square$

LEMMA 3.3. Suppose that (2.43) holds. If g satisfies (2.44) and (3.4), then $D(\Lambda)$ is dense in $\mathcal{H}$. Further, if g satisfies (2.45) and (2.46) (or (2.93) and (2.94)), we have

$$\begin{aligned} D(\Lambda) \subset D_0 &= \{(u, v, \theta) \in \mathcal{H} : u \in (H^s(\Omega) \cap H_{\Gamma_1}^1(\Omega))^n, v \in (H_{\Gamma_1}^1(\Omega))^n, \\ &\theta \in H^2(\Omega) \cap H_0^1(\Omega), \\ &\mu \frac{\partial u}{\partial \nu} + (\lambda + \mu)\nu \operatorname{div} u + am \cdot \nu u + m \cdot \nu g(v) = 0 \quad \text{on } \Gamma_2\}, \end{aligned} \quad (3.32)$$

for some $s > 3/2$.

Proof. Set

$$\begin{aligned} D &= \{(u, v, \theta) \in \mathcal{H} : u \in (H^2(\Omega) \cap H_{\Gamma_1}^1(\Omega))^n, v \in (H_0^1(\Omega))^n, \\ &\theta \in H^2(\Omega) \cap H_0^1(\Omega), \\ &\mu \frac{\partial u}{\partial \nu} + (\lambda + \mu)\nu \operatorname{div} u + am \cdot \nu u = 0 \quad \text{on } \Gamma_2\}. \end{aligned} \quad (3.33)$$

To prove that $D(\Lambda)$ is dense in $\mathcal{H}$, it suffices to prove that D is dense in $\mathcal{H}$ and $D \subset D(\Lambda)$. Firstly, we prove that D is dense in $\mathcal{H}$. For this, it suffices to prove that

$$W = \{w \in (H^2(\Omega) \cap H_{\Gamma_1}^1(\Omega))^n : \mu \frac{\partial w}{\partial \nu} + (\lambda + \mu)\nu \operatorname{div} w + am \cdot \nu w = 0 \quad \text{on } \Gamma_2\} \quad (3.34)$$

is dense in $(H_{\Gamma_1}^1(\Omega))^n$. Let $v \in (H_{\Gamma_1}^1(\Omega))^n$ be such that

$$(v, w)_{(H_{\Gamma_1}^1(\Omega))^n} = 0, \quad w \in W. \quad (3.35)$$

For any fixed $f \in (L^2(\Omega))^n$, we consider the following elliptic problem:

$$\begin{cases} -\mu \Delta u - (\lambda + \mu) \nabla \operatorname{div} u = f & \text{in } \Omega, \\ u = 0, & \text{on } \Gamma_1, \\ \mu \frac{\partial u}{\partial \nu} + (\lambda + \mu) \nu \operatorname{div} u + am \cdot \nu u = 0 & \text{on } \Gamma_2. \end{cases} \quad (3.36)$$

By the elliptic regularity theory, problem (3.36) has a solution $u \in W$. Thus, by (3.35), we have

$$(v, f) = (v, u)_{(H_{\Gamma_1}^1(\Omega))^n} = 0. \quad (3.37)$$

Hence, we deduce that $v = 0$. It therefore follows from the Hahn-Banach theorem that W is dense in $(H_{\Gamma_1}^1(\Omega))^n$.

Next, we prove that $D \subset D(\Lambda)$. Let $(u, v, \theta) \in D$. To prove that $(u, v, \theta) \in D(\Lambda)$, it suffices to prove that $Au + Bv \in (L^2(\Omega))^n$. For this, let $w \in (H_{\Gamma_1}^1(\Omega))^n$. By the definition of A and B , we have

$$\begin{aligned} \langle Au + Bv, w \rangle &= (u, w)_{(H_{\Gamma_1}^1(\Omega))^n} + \int_{\Gamma_2} m \cdot \nu g(v) \cdot w \, d\Gamma \\ &= \int_{\Omega} (\mu \nabla u \cdot \nabla w + (\lambda + \mu) \operatorname{div} u \operatorname{div} w) \, dx \\ &\quad + \int_{\Gamma_2} am \cdot \nu u \cdot w \, d\Gamma \quad (\text{use } g(0) = 0) \\ &= - \int_{\Omega} (\mu \Delta u + (\lambda + \mu) \nabla \operatorname{div} u) \cdot w \, dx, \end{aligned} \quad (3.38)$$

since $u \in (H^2(\Omega))^n$. It therefore follows that there exists a constant c , depending on u , such that

$$|\langle Au + Bv, w \rangle| \leq c \|w\|, \quad (3.39)$$

which implies that $Au + Bv \in (L^2(\Omega))^n$. Thus $(u, v, \theta) \in D(\Lambda)$.

It remains to prove that $D(\Lambda) \subset D_0$ if g satisfies (2.45) and (2.46) (or (2.93) and (2.94)). Then it is sufficient to prove that $u \in (H^s(\Omega) \cap H_{\Gamma_1}^1(\Omega))^n$ for some $s > 3/2$ if $(u, v, \theta) \in D(\Lambda)$. For this, let $f = Au + Bv$ and $h = -m \cdot \nu g(v)$. Then, for any $w \in (H_{\Gamma_1}^1(\Omega))^n$, we have

$$\begin{aligned} \langle f, w \rangle &= \int_{\Omega} (\mu \nabla u \cdot \nabla w + (\lambda + \mu) \operatorname{div} u \operatorname{div} w) \, dx \\ &\quad + \int_{\Gamma_2} am \cdot \nu u \cdot w \, d\Gamma + \int_{\Gamma_2} m \cdot \nu g(v) \cdot w \, d\Gamma. \end{aligned} \quad (3.40)$$

This shows that u is a weak solution of the following elliptic problem:

$$\begin{cases} -\mu\Delta u - (\lambda + \mu)\nabla\operatorname{div} u = f & \text{in } \Omega, \\ u = 0 & \text{on } \Gamma_1, \\ \mu\frac{\partial u}{\partial\nu} + (\lambda + \mu)\nu\operatorname{div} u + am \cdot \nu u = h & \text{on } \Gamma_2. \end{cases} \quad (3.41)$$

By assumption, we have

$$f = Au + Bv \in (L^2(\Omega))^n. \quad (3.42)$$

We further prove that

$$h \in (H^\sigma(\Gamma))^n \quad (3.43)$$

for some $0 < \sigma < q/2$. For this, we let

$$S_1 = \{(x, y) \in \Gamma \times \Gamma : |x - y| \leq 1\}, \quad (3.44)$$

$$S_{11} = \{(x, y) \in S_1 : |v(x) - v(y)| \leq 1\}, \quad (3.45)$$

$$S_{12} = \{(x, y) \in S_1 : |v(x) - v(y)| \geq 1\}. \quad (3.46)$$

To prove (3.43), by definition (see [15, p. 20]), it suffices to prove

$$\iint_{\Gamma \times \Gamma} \frac{|g(v(x)) - g(v(y))|^2}{|x - y|^{n-1+2\sigma}} d\Gamma(x) d\Gamma(y) < \infty. \quad (3.47)$$

The integral over $\Gamma \times \Gamma - S_1$ is bounded above by

$$c \iint_{\Gamma \times \Gamma} |g(v(x)) - g(v(y))|^2 d\Gamma(x) d\Gamma(y) \leq c \int_{\Gamma} |g(v(x))|^2 d\Gamma(x) < \infty, \quad (3.48)$$

since $g(v(x)) \in L^2(\Gamma)$ which is true because $v \in (H^{1/2}(\Gamma))^n$ and g satisfies (2.45) and (2.46) (or (2.93) and (2.94)). Therefore, it suffices to prove

$$\begin{aligned} I &= \iint_{S_1} \frac{|g(v(x)) - g(v(y))|^2}{|x - y|^{n-1+2\sigma}} d\Gamma(x) d\Gamma(y) \\ &= \iint_{S_{11}} \frac{|g(v(x)) - g(v(y))|^2}{|x - y|^{n-1+2\sigma}} d\Gamma(x) d\Gamma(y) \\ &\quad + \iint_{S_{12}} \frac{|g(v(x)) - g(v(y))|^2}{|x - y|^{n-1+2\sigma}} d\Gamma(x) d\Gamma(y) \\ &= I_1 + I_2 \\ &< \infty. \end{aligned} \quad (3.49)$$

By (2.45) and Hölder's inequality, we have

$$\begin{aligned} I_1 &\leq \iint_{S_{11}} \frac{|v(x) - v(y)|^{2q}}{|x - y|^{n-1+2\sigma}} d\Gamma(x) d\Gamma(y) \\ &\leq \left(\iint_{S_{11}} \frac{|v(x) - v(y)|^2}{|x - y|^2} d\Gamma(x) d\Gamma(y) \right)^q \\ &\quad \times \left(\iint_{S_1} \frac{1}{|x - y|^{(n-1+2\sigma-nq)/(1-q)}} d\Gamma(x) d\Gamma(y) \right)^{1-q}. \end{aligned} \quad (3.50)$$

Since $v \in (H_{\Gamma_1}^1(\Omega))^n$, by the trace theorem, we have $v \in (H^{1/2}(\Gamma))^n$, and then we have

$$\iint_{S_1} \frac{|v(x) - v(y)|^2}{|x - y|^n} d\Gamma(x) d\Gamma(y) < \infty. \quad (3.51)$$

In addition, since for $0 < 2\sigma < q$

$$\frac{n - 1 + 2\sigma - nq}{1 - q} = n + \frac{2\sigma - 1}{1 - q} < n - 1, \quad (3.52)$$

we have

$$\iint_{S_1} \frac{1}{|x - y|^{(n-1+2\sigma-nq)/(1-q)}} d\Gamma(x) d\Gamma(y) < \infty. \quad (3.53)$$

It therefore follows that $I_1 < \infty$. On the other hand, by (2.46) and (3.51), we have

$$\begin{aligned} I_2 &\leq \iint_{S_{12}} \frac{|v(x) - v(y)|^2}{|x - y|^{n-1+2\sigma}} d\Gamma(x) d\Gamma(y) \\ &\leq \iint_{S_{12}} \frac{|v(x) - v(y)|^2}{|x - y|^n} d\Gamma(x) d\Gamma(y) \\ &< \infty. \end{aligned} \quad (3.54)$$

Thus we have proved (3.49). In a similar way, we can show that $I < \infty$ if (2.93) holds.

Finally, we prove that

$$u \in (H^s(\Omega) \cap H_{\Gamma_1}^1(\Omega))^n \quad (3.55)$$

with

$$s = \frac{3}{2} + \sigma > \frac{3}{2}. \quad (3.56)$$

Let $\nu = (\nu_1, \nu_2, \dots, \nu_n)$ denote the unit normal on Γ directed to the exterior of Ω and consider the following system:

$$\sum_{j=1}^n [(\lambda + \mu)\nu_i\nu_j + \delta_{ij}\mu]\xi_j = h_i, \quad i = 1, \dots, n, \quad (3.57)$$

where δ_{ij} denotes the Kronecker symbol, i.e.,

$$\delta_{ij} = \begin{cases} 1, & i = j, \\ 0, & i \neq j. \end{cases} \quad (3.58)$$

It is easy to see that the system has a solution $\xi = (\xi_1, \dots, \xi_n) \in (H^\sigma(\Gamma))^n$ for $h \in (H^\sigma(\Gamma))^n$. By the trace theorem (see, e.g., [36, p. 39, Theorem 8.3]), there exists $\varphi \in (H^s(\Omega) \cap H_{\Gamma_1}^1(\Omega))^n$ such that

$$\frac{\partial \varphi}{\partial \nu} = \xi, \quad \varphi = 0 \quad \text{on } \Gamma. \quad (3.59)$$

Let $\{\tau^k(x)\}_{k=1}^{n-1}$ be a tangential vector field such that $\{\nu(x), \tau^1(x), \dots, \tau^{n-1}(x)\}$ forms an orthonormal basis in $\mathbf{R}^n$ for almost all $x \in \Gamma$. Hence, there exist $\gamma^{k,j}$ ($j = 1, 2, \dots, n$; $k = 1, 2, \dots, n-1$) depending on $\{\nu(x), \tau^1(x), \dots, \tau^{n-1}(x)\}$ such that

$$\begin{aligned} \frac{\partial \varphi_j}{\partial x_j} &= \nu_j \frac{\partial \varphi_j}{\partial \nu} + \sum_{k=1}^{n-1} \gamma^{k,j} \frac{\partial \varphi_j}{\partial \tau^k} \\ &= \nu_j \frac{\partial \varphi_j}{\partial \nu} \\ &= \nu_j \xi_j \quad \text{on } \Gamma, \quad j = 1, 2, \dots, n. \end{aligned} \quad (3.60)$$

It therefore follows from (3.57) that

$$\mu \frac{\partial \varphi}{\partial \nu} + (\lambda + \mu) \nu \operatorname{div} \varphi + am \cdot \nu \varphi = h \quad \text{on } \Gamma_2. \quad (3.61)$$

Setting

$$\psi = u - \varphi, \quad (3.62)$$

then ψ satisfies

$$\begin{cases} -\mu \Delta \psi - (\lambda + \mu) \nabla \operatorname{div} \psi = F & \text{in } \Omega, \\ \psi = 0 & \text{on } \Gamma_1, \\ \mu \frac{\partial \psi}{\partial \nu} + (\lambda + \mu) \nu \operatorname{div} \psi + am \cdot \nu \psi = 0 & \text{on } \Gamma_2, \end{cases} \quad (3.63)$$

where

$$F = f - \mu \Delta \varphi - (\lambda + \mu) \nabla \operatorname{div} \varphi \in (H^{s-2}(\Omega))^n. \quad (3.64)$$

Thus, problem (3.41) is equivalent to (3.63). By the classical variational methods (see, e.g., [13]), for every $F \in ((H_{\Gamma_1}^1(\Omega))')^n$, problem (3.63) has a unique weak solution $\psi \in (H_{\Gamma_1}^1(\Omega))^n$ in the sense of distribution

$$\begin{aligned} &\int_{\Omega} (\mu \nabla \psi \cdot \nabla \phi + (\lambda + \mu) \operatorname{div} \psi \operatorname{div} \phi) dx + \int_{\Gamma_2} am \cdot \nu \psi \cdot \phi d\Gamma \\ &= \int_{\Omega} F \cdot \phi dx \quad \forall \phi \in (H_{\Gamma_1}^1(\Omega))^n. \end{aligned} \quad (3.65)$$

Moreover, by the classical Nirenberg's translation method (see, e.g., [2, p. 107, Lemma 9.2] or [36, p. 124]), we prove that, if $F \in (L^2(\Omega))^n$, then $\psi \in (H^2(\Omega) \cap H_{\Gamma_1}^1(\Omega))^n$. Our proof is essentially the same as that of Lemma 9.2 of [2, p. 107]. Since the regularity is local property, it suffices to prove that, for any $x \in \bar{\Omega}$, there exists a neighborhood $O(x)$

such that $\psi \in x \in \Omega$ is $\psi \in (H^2(O(x) \cap \Omega))^n$. We consider the case $x \in \Gamma$ since the case $x \in \Omega$ is much easier. For simplicity, we may as well assume that $x = 0 \in \Gamma_2$ and that the boundary is flat with the normal oriented in the direction x_n since the general case can be transformed into the special case by a mapping of class C^2 (see Remark 3.4 below). Therefore, there exists a hemisphere $G_R = \{x : |x| < R, x_n > 0\}$ such that $G_R \subset \Omega$ and $\Gamma_{2G} = \{x \in \bar{G}_R : x_n = 0\} \subset \Gamma_2$. Let $0 < R' < R$ and $R'' = \frac{1}{2}(R' + R)$, and let ζ denote a real function that is infinitely differentiable on $\mathbf{R}^n$ and such that $\zeta = 1$ on $G_{R'}$ and $\zeta = 0$ outside $G_{R''}$. Note that ζ need not vanish on the flat part Γ_{2G} of the boundary of G_R . Let $C_0^\infty(G_R \cup \Gamma_{2G})$ denote the function space of all infinitely differentiable functions with support in $G_R \cup \Gamma_{2G}$. By (3.65), we have for any $\phi \in (C_0^\infty(G_R \cup \Gamma_{2G}))^n$ that

$$\int_{G_R} (\mu \nabla \psi \cdot \nabla \phi + (\lambda + \mu) \operatorname{div} \psi \operatorname{div} \phi) dx + \int_{\Gamma_{2G}} am \cdot \nu \psi \cdot \phi d\Gamma = \int_{G_R} F \cdot \phi dx. \quad (3.66)$$

Define the bilinear form $B(\psi, \phi)$ by

$$\begin{aligned} B(\psi, \phi) &= \int_{G_R} (\mu \nabla \psi \cdot \nabla \phi + (\lambda + \mu) \operatorname{div} \psi \operatorname{div} \phi) dx \\ &\quad + \int_{\Gamma_{2G}} am \cdot \nu \psi \cdot \phi d\Gamma. \end{aligned} \quad (3.67)$$

Then we have

$$|B(\psi, \phi)| \leq \|F\| \|\phi\| \quad \forall \phi \in (C_0^\infty(G_R \cup \Gamma_{2G}))^n. \quad (3.68)$$

For a real number, we define the difference operator δ_h^i by

$$\delta_h^i u = h^{-1} [u(x_1, \dots, x_i + h, \dots, x_n) - u(x_1, \dots, x_i, \dots, x_n)], \quad i = 1, 2, \dots, n.$$

We now want to estimate the difference quotients $\delta_h^i(\zeta\psi)$ for $i = 1, 2, \dots, n$. Since

$$\begin{aligned}
B(\delta_h^i(\zeta\psi), \phi) &= \int_{G_R} (\mu \nabla(\delta_h^i(\zeta\psi)) \cdot \nabla \phi + (\lambda + \mu) \operatorname{div}(\delta_h^i(\zeta\psi)) \operatorname{div} \phi) dx \\
&\quad + \int_{\Gamma_{2G}} am \cdot \nu(\delta_h^i(\zeta\psi)) \cdot \phi d\Gamma \\
&= \int_{G_R} (\mu \delta_h^i(\nabla(\zeta\psi)) \cdot \nabla \phi + (\lambda + \mu) \delta_h^i(\operatorname{div}(\zeta\psi)) \operatorname{div} \phi) dx \\
&\quad + \int_{2G} am \cdot \nu(\delta_h^i(\zeta\psi)) \cdot \phi d\Gamma \\
&= \int_{G_R} (\mu \delta_h^i(\psi_j \nabla \zeta + \zeta \nabla \psi_j) \cdot \nabla \phi_j + (\lambda + \mu) \delta_h^i(\nabla \zeta \cdot \psi + \zeta \operatorname{div} \psi) \operatorname{div} \phi) dx \\
&\quad + \int_{\Gamma_{2G}} am \cdot \nu(\delta_h^i(\zeta\psi)) \cdot \phi d\Gamma \\
&= \mu \int_{G_R} (\delta_h^i(\psi_j \nabla \zeta) \cdot \nabla \phi_j + \zeta \nabla \psi_j \cdot \delta_{-h}^i(\nabla \phi_j)) dx \\
&\quad + (\lambda + \mu) \int_{G_R} (\delta_h^i(\nabla \zeta \cdot \psi) \operatorname{div} \phi + \zeta \operatorname{div} \psi \delta_{-h}^i(\operatorname{div} \phi)) dx \\
&\quad + \int_{\Gamma_{2G}} am \cdot \nu \zeta \psi \cdot \delta_{-h}^i(\phi) d\Gamma \\
&= \mu \int_{G_R} (\delta_h^i(\psi_j \nabla \zeta) \cdot \nabla \phi_j + \nabla \psi_j \cdot \nabla(\zeta \delta_{-h}^i \phi_j) - \nabla \psi_j \cdot \nabla \zeta \delta_{-h}^i \phi_j) dx \\
&\quad + (\lambda + \mu) \int_{G_R} (\delta_h^i(\nabla \zeta \cdot \psi) \operatorname{div} \phi + \operatorname{div} \psi \operatorname{div}(\zeta \delta_{-h}^i \phi) - \operatorname{div} \psi \nabla \zeta \cdot \delta_{-h}^i \phi) dx \\
&\quad + \int_{\Gamma_{2G}} am \cdot \nu \psi \cdot (\zeta \delta_{-h}^i \phi) d\Gamma \\
&= B(\psi, \zeta \delta_{-h}^i \phi) + \mu \int_{G_R} (\delta_h^i(\psi_j \nabla \zeta) \cdot \nabla \phi_j - \nabla \psi_j \cdot \nabla \zeta \delta_{-h}^i \phi_j) dx \\
&\quad + (\lambda + \mu) \int_{G_R} (\delta_h^i(\nabla \zeta \cdot \psi) \operatorname{div} \phi - \operatorname{div} \psi \nabla \zeta \cdot \delta_{-h}^i \phi) dx.
\end{aligned} \tag{3.69}$$

It therefore follows from (3.68) that

$$\begin{aligned}
|B(\delta_h^i(\zeta\psi), \phi)| &\leq \|F\| \|\zeta \delta_{-h}^i \phi\| + C \|\phi\|_{(H^1(G_R))^n} \|\psi\|_{(H^1(G_R))^n} \\
&\leq C(\|F\| + \|\psi\|_{(H^1(G_R))^n}) \|\phi\|_{(H^1(G_R))^n}.
\end{aligned} \tag{3.70}$$

Let $(H_{\Gamma_{2G}}^1(G_R))^n$ be the completion of $(C_0^\infty(G_R \cup \Gamma_{2G}))^n$ in $(H^1(G_R))^n$. Then by a density argument, we obtain for any $\phi \in (H_{\Gamma_{2G}}^1(G_R))^n$

$$|B(\delta_h^i(\zeta\psi), \phi)| \leq C(\|F\| + \|\psi\|_{(H^1(G_R))^n}) \|\phi\|_{(H^1(G_R))^n}. \tag{3.71}$$

Since $\delta_h^i(\zeta\psi) \in (H_{\Gamma_{2G}}^1(G_R))^n$ if h is small enough, we deduce

$$|B(\delta_h^i(\zeta\psi), \delta_h^i(\zeta\psi))| \leq C(\|F\| + \|\psi\|_{(H^1(G_R))^n}) \|\delta_h^i(\zeta\psi)\|_{(H^1(G_R))^n}. \tag{3.72}$$

On the other hand, it is clear that

$$|B(\delta_h^i(\zeta\psi), \delta_h^i(\zeta\psi))| \geq C\|\delta_h^i(\zeta\psi)\|_{(H^1(G_R))^n}^2.$$

Hence it follows from (3.72) that

$$\|\delta_h^i(\zeta\psi)\|_{(H^1(G_R))^n} \leq C(\|F\| + \|\psi\|_{(H^1(G_R))^n}). \quad (3.73)$$

Since $\zeta = 1$ on $G_{R'}$, by Theorem 3.16 of [2, p. 45], we deduce that $\frac{\partial\psi_i}{\partial x_j} \in (H^1(G_{R'}))^n$ for all $i = 1, \dots, n, j = 1, \dots, n-1$. It remains to show that $\frac{\partial^2\psi}{\partial x_n^2} \in (L^2(G_{R'}))^n$. To do this we have to distinguish the components ψ_i for $i = 1, \dots, n-1$ and for $i = n$. In what concerns $i = 1, \dots, n-1$, we have

$$-\mu \frac{\partial^2\psi_i}{\partial x_n^2} = \mu\Delta'\psi_i + (\lambda + \mu) \frac{\partial}{\partial x_i} (\operatorname{div} \psi) + F_i \in L^2(G_{R'}),$$

while

$$-(\lambda + 2\mu) \frac{\partial^2\psi_n}{\partial x_n^2} = \mu\Delta'\psi_n + (\lambda + \mu) \frac{\partial}{\partial x_n} \left(\frac{\partial\psi_1}{\partial x_1} + \dots + \frac{\partial\psi_{n-1}}{\partial x_{n-1}} \right) + F_n \in L^2(G_{R'}),$$

where $\Delta' = \frac{\partial^2}{\partial x_1^2} + \dots + \frac{\partial^2}{\partial x_{n-1}^2}$.

By interpolation (see, e.g., [36, p. 29, Theorem 6.2]) for $F \in (H^{s-2}(\Omega))^n$ we have $\psi \in (H^s(\Omega) \cap H_{\Gamma_1}^1(\Omega))^n$, and then $u \in (H^s(\Omega) \cap H_{\Gamma_1}^1(\Omega))^n$. This completes the proof. $\square$

REMARK 3.4. Let us briefly explain how to deal with the case where the boundary is not flat by a flattening procedure. Suppose that $x^0 \in \Gamma_2$ and the boundary of Ω near x^0 can be expressed by

$$\Phi(x_1, x_2, \dots, x_n) = 0.$$

Let

$$y_i = y_i(x_1, x_2, \dots, x_n) = x_i, \quad i = 1, \dots, n-1, \quad (3.74)$$

$$y_n = y_n(x_1, x_2, \dots, x_n) = \Phi(x_1, x_2, \dots, x_n) \quad (3.75)$$

be the invertible mapping of class C^2 that transforms Ω into $\tilde{\Omega}$. We denote by $x = x(y)$ the inverted mapping of the above mapping $y = y(x)$ and set

$$\tilde{\psi}(y) = \psi(x(y)).$$

By calculations, we obtain

$$\frac{\partial\psi_i}{\partial x_j} = \sum_{k=1}^n \frac{\partial\tilde{\psi}_i}{\partial y_k} \frac{\partial y_k}{\partial x_j}, \quad (3.76)$$

$$\frac{\partial^2\psi_i}{\partial x_j^2} = \sum_{k,l=1}^n \frac{\partial^2\tilde{\psi}_i}{\partial y_k \partial y_l} \frac{\partial y_k}{\partial x_j} \frac{\partial y_l}{\partial x_j} + \sum_{k=1}^n \frac{\partial\tilde{\psi}_i}{\partial y_k} \frac{\partial^2 y_k}{\partial x_j^2}. \quad (3.77)$$

Therefore, the first equation of (3.63) is transformed into

$$\begin{aligned} \tilde{F}_i(y) = & -\mu \left(\sum_{j,k,l=1}^n \frac{\partial^2 \tilde{\psi}_i}{\partial y_k \partial y_l} \frac{\partial y_k}{\partial x_j} \frac{\partial y_l}{\partial x_j} + \sum_{j,k=1}^n \frac{\partial \tilde{\psi}_i}{\partial y_k} \frac{\partial^2 y_k}{\partial x_j^2} \right) \\ & - (\lambda + \mu) \left(\sum_{j,k,l=1}^n \frac{\partial^2 \tilde{\psi}_j}{\partial y_k \partial y_l} \frac{\partial y_k}{\partial x_j} \frac{\partial y_l}{\partial x_i} + \sum_{j,k=1}^n \frac{\partial \tilde{\psi}_j}{\partial y_k} \frac{\partial^2 y_k}{\partial x_j \partial x_i} \right), \quad i = 1, \dots, n, \end{aligned} \quad (3.78)$$

where $\tilde{F}_i(y) = F_i(x(y))$. Moreover, since the unit normal $\nu(x)$ on the boundary near x^0 is equal to

$$\nu(x) = \frac{\nabla y_n}{|\nabla y_n|},$$

we have

$$\frac{\partial \psi_i}{\partial \nu} = \sum_{j=1}^n \frac{\partial \psi_i}{\partial x_j} \nu_j = \sum_{j,k=1}^n \frac{\partial \tilde{\psi}_i}{\partial y_k} \frac{\partial y_k}{\partial x_j} \frac{\partial y_n}{\partial x_j} |\nabla y_n|^{-1}, \quad (3.79)$$

$$\nu_i \operatorname{div} \psi = \sum_{j=1}^n \frac{\partial \psi_j}{\partial x_j} \nu_i = \sum_{j,k=1}^n \frac{\partial \tilde{\psi}_j}{\partial y_k} \frac{\partial y_k}{\partial x_j} \frac{\partial y_n}{\partial x_i} |\nabla y_n|^{-1}. \quad (3.80)$$

Hence, the boundary condition of (3.63) is transformed into the following form:

$$\mu \sum_{j,k=1}^n \frac{\partial \tilde{\psi}_i}{\partial y_k} \frac{\partial y_k}{\partial x_j} \frac{\partial y_n}{\partial x_j} + (\lambda + \mu) \sum_{j,k=1}^n \frac{\partial \tilde{\psi}_j}{\partial y_k} \frac{\partial y_k}{\partial x_j} \frac{\partial y_n}{\partial x_i} + \tilde{a} \tilde{\psi}_i = 0, \quad i = 1, \dots, n, \quad (3.81)$$

where $\tilde{a} = \tilde{a}(y) \geq 0$. Since the already known H^1 -regularity guarantees that the first-order derivative terms of the unknown $\tilde{\psi}$ in (3.78) are in L^2 , they may be put on the right-hand side and it is then sufficient to keep the second-order terms on the left-hand side. Therefore, since $\nu(y) = (0, \dots, 0, 1)$ on the boundary of $\tilde{\Omega}$ near $y^0 = y(x^0)$, it follows that the bilinear form corresponding to equation (3.78) and boundary condition (3.81) is

$$\begin{aligned} \tilde{B}(\tilde{\psi}, \tilde{\phi}) = & \mu \int_{\tilde{G}_R} \sum_{i,j,k,l=1}^n \frac{\partial \tilde{\psi}_i}{\partial y_k} \frac{\partial \tilde{\phi}_i}{\partial y_l} \frac{\partial y_k}{\partial x_j} \frac{\partial y_l}{\partial x_j} dy + (\lambda + \mu) \int_{\tilde{G}_R} \sum_{i,j,k,l=1}^n \frac{\partial \tilde{\psi}_j}{\partial y_k} \frac{\partial \tilde{\phi}_i}{\partial y_l} \frac{\partial y_k}{\partial x_j} \frac{\partial y_l}{\partial x_i} dy \\ & + \int_{\tilde{\Gamma}_{2G}} \tilde{\alpha} \tilde{\psi} \cdot \tilde{\phi} d\Gamma, \end{aligned} \quad (3.82)$$

where $\tilde{G}_R$ and $\tilde{\Gamma}_{2G}$ denote the same subsets in the y -space as G_R and Γ_{2G} . It is clear that the bilinear $\tilde{B}(\tilde{\psi}, \tilde{\phi})$ has the same properties as $B(\psi, \phi)$. Therefore, the above proof of regularity is valid for the general case (see, e.g., [8]).

By the classical theory of nonlinear semigroups (see [6, Chap. 3]), we have

THEOREM 3.5. Let Γ_1 and Γ_2 be given by (2.1) and (2.2). Let (2.16) and (2.43) hold. Suppose that the function g satisfies (2.44), (2.49) (or (2.92), (2.97)) and (3.4). Then we have

(i) For every initial condition $(u^0, u^1, \theta^0) \in \mathcal{H}$, problem (2.7) has a unique mild solution satisfying

$$(u, u', \theta) \in C([0, \infty); \mathcal{H}). \quad (3.83)$$

Moreover, if (u, θ) and (v, ξ) are two solutions corresponding to initial states (u^0, u^1, θ^0) and (v^0, v^1, ξ^0) , respectively, then we have for every $t \in [0, \infty)$

$$\|u(t), u'(t), \theta(t) - (v(t), v'(t), \xi(t))\|_{\mathcal{H}} \leq \|(u^0, u^1, \theta^0) - (v^0, v^1, \xi^0)\|_{\mathcal{H}}. \quad (3.84)$$

(ii) Further, if g satisfies (2.45) and (2.46) (or (2.93) and (2.94)), then for every initial condition $(u^0, u^1, \theta^0) \in D(\Lambda)$, problem (2.7) has a unique classical solution satisfying

$$u \in L^\infty([0, \infty); (H^s(\Omega) \cap H_{\Gamma_1}^1(\Omega))^n), \quad (3.85)$$

$$u' \in L^\infty([0, \infty); (H_{\Gamma_1}^1(\Omega))^n), \quad (3.86)$$

$$u'' \in L^\infty([0, \infty); (L^2(\Omega))^n), \quad (3.87)$$

$$\theta \in L^\infty([0, \infty); H^2(\Omega) \cap H_0^1(\Omega)), \quad (3.88)$$

$$\theta' \in L^\infty([0, \infty); L^2(\Omega)), \quad (3.89)$$

for some $s > 3/2$.

4. Proofs of the main results. In this section, we borrow the Lyapunov method to prove our main results.

Let u, θ be the solution of (2.7) and φ the solution of (2.27) corresponding to the present solution u of (2.7) (note that φ depends on t as u does). Let δ be any positive number. We define a functional V on $\mathcal{H}$ by

$$F(t) = F(u, \theta, t) = \int_{\Omega} [2u'_i m \cdot \nabla u_i + (n-1)u'_i u_i + Cu'_i \varphi_i] dx \quad (4.1)$$

$$V(t) = V(u, \theta, t) = E(t) + \delta[E(t)]^\sigma F(t), \quad (4.2)$$

where C and σ are nonnegative constants that will be determined later. Here we have used the summation convention for repeated indices. Thus, equation (4.1) means that

$$F(t) = \sum_{i=1}^n \int_{\Omega} [2u'_i m \cdot \nabla u_i + (n-1)u'_i u_i + C\varphi_i u'_i] dx. \quad (4.3)$$

Evidently, the functional V is actually a generalized energy functional which is closely related to the energy functional $E(t)$. Such similar functionals were constructed for the wave equation (see [10, 25, 51]), thermoelastic plate models (see [42, 43]), viscoelasticity (see [34, 35]) and thermoviscoelasticity (see [39]).

We will see below that the term $Cu'_i \varphi_i$ plays a key role in dealing with the case where the potential a is large.

We are going to show that V satisfies

$$V'(t) \leq -cV^{\sigma+1}(t), \quad (4.4)$$

where c is a positive constant. Then, by solving differential inequality (4.4), Theorems 2.2, 2.3 and 2.5 are proved.

We first show that, by choosing δ sufficiently small, V and E are equivalent.

LEMMA 4.1. Let Γ_1 and Γ_2 be given by (2.1) and (2.2), respectively. Suppose that (2.16) holds. Let the function $g \in (C(\mathbf{R}^n))^n$ satisfy (2.44), (2.49) (or (2.92), (2.97)) and (3.4). Then we have

$$(1 - \delta C_1'' E^\sigma(t))E(t) \leq V(t) \leq (1 + \delta C_1'' E^\sigma(t))E(t), \quad (4.5)$$

for all solutions u, θ of (2.7), where the positive constant C_1'' is given by

$$C_1'' = \frac{2R_0}{\sqrt{\mu}} + (n-1)\lambda_0 + Ck_1\gamma. \quad (4.6)$$

Proof. It is easy to see that

$$\left| 2 \int_{\Omega} u_i' m \cdot \nabla u_i \, dx \right| \leq \frac{R_0}{\sqrt{\mu}} [\|u_i'(t)\|^2 + \mu \|\nabla u_i(t)\|^2] \leq \frac{2R_0}{\sqrt{\mu}} E(t). \quad (4.7)$$

Using (2.25), we obtain

$$\left| \int_{\Omega} u_i' u_i \, dx \right| \leq \frac{\lambda_0}{2} [\|u'(t)\|^2 + \|u(t)\|_{(H_{\Gamma_1}^1(\Omega))^n}^2] \leq \lambda_0 E(t). \quad (4.8)$$

Using (2.28) and (2.24), we have

$$\begin{aligned} \left| \int_{\Omega} u_i' \varphi_i \, dx \right| &= \frac{k_1\gamma}{2} \left| \int_{\Omega} 2u_i' \frac{\varphi_i}{k_1\gamma} \, dx \right| \\ &\leq \frac{k_1\gamma}{2} [\|u_i'(t)\|^2 + \frac{1}{k_1^2\gamma^2} \|\varphi_i(t)\|^2] \\ &\leq \frac{k_1\gamma}{2} [\|u_i'(t)\|^2 + \frac{1}{\gamma^2} \int_{\Gamma_2} |u_i|^2 \, d\Gamma] \\ &\leq \frac{k_1\gamma}{2} [\|u'(t)\|^2 + \|u(t)\|_{(H_{\Gamma_1}^1(\Omega))^n}^2] \\ &\leq k_1\gamma E(t). \end{aligned} \quad (4.9)$$

Noting (4.6), we deduce from (4.7), (4.8) and (4.9) that

$$|F(t)| \leq C_1'' E(t), \quad (4.10)$$

which implies (4.5). $\square$

We then estimate $F'(t)$.

LEMMA 4.2. Let Γ_1 and Γ_2 be given by (2.1) and (2.2), respectively, satisfying (2.43). Suppose that (2.16) holds and the function $g \in (C(\mathbf{R}^n))^n$ satisfies (2.44), (2.45), (2.46) and (2.49) (or (2.92), (2.93), (2.94) and (2.97)). Let u, θ be a classical solution of (2.7) with $(u^0, u^1, \theta^0) \in D(\Lambda)$. Then we have

$$\begin{aligned} F'(t) &\leq -E(t) + C_2'' \|\nabla \theta(t)\|^2 \\ &\quad + \int_{\Gamma_2} \left[\left(m \cdot \nu + \frac{C^2 k_1^2}{4\varepsilon} \right) |u'|^2 + C_3'' m \cdot \nu |g(u')|^2 \right] d\Gamma, \end{aligned} \quad (4.11)$$

where

$$C_2'' = \frac{\alpha^2 R_0^2}{\mu \varepsilon} + \frac{\alpha^2 (n-1)^2 \lambda_0^2}{4\varepsilon} + \frac{C^2 \alpha^2 k_1^2 \gamma^2}{4\varepsilon} + \frac{\alpha \lambda_1^2}{\beta}, \quad (4.12)$$

$$C_3'' = \frac{2R_0^2}{\mu} + \frac{(n-1+C)^2 R_0 \gamma^2}{4\varepsilon}, \quad (4.13)$$

and the constant C is given by

$$C = C_0 = \max \left\{ 0, \frac{2a_0 R_0^2}{\mu} + 2 - n \right\}. \quad (4.14)$$

If, further, the function $a(x)$ satisfies (2.50) or (2.51), then we can take

$$C = 0. \quad (4.15)$$

The constant ε is given by

$$\varepsilon = \begin{cases} 1/8, & \text{if } C = C_0, \\ 1/8, & \text{if } C = 0 \text{ and } a(x) \text{ satisfied (2.51),} \\ (1 - 2K(a)R_0\gamma^2)/8, & \text{if } C = 0 \text{ and } a(x) \text{ satisfies (2.50).} \end{cases} \quad (4.16)$$

Proof. Since $(u^0, u^1, \theta^0) \in D(\Lambda)$, we have $u \in L^\infty([0, \infty); (H^s(\Omega) \cap H_{\Gamma_1}^1(\Omega))^n)$ for some $s > 3/2$ and $\theta \in L^\infty([0, \infty); H^2(\Omega) \cap H_0^1(\Omega))$. Hence, the following integrations by parts are valid.

By (4.1), we have

$$\begin{aligned} F'(t) &= \int_{\Omega} 2u_i'' m \cdot \nabla u_i \, dx + \int_{\Omega} 2u_i' m \cdot \nabla u_i' \, dx \\ &\quad + \int_{\Omega} (n-1)u_i u_i'' \, dx + (n-1)\|u_i'\|^2 \\ &\quad + C \int_{\Omega} \varphi_i u_i'' \, dx + C \int_{\Omega} \varphi_i' u_i' \, dx. \end{aligned} \quad (4.17)$$

We now estimate every integral in (4.17) as follows. Since $u = 0$ on Γ_1 , we have

$$\frac{\partial u_i}{\partial x_k} = \frac{\partial u_i}{\partial \nu} \nu_k \quad \text{on } \Gamma_1. \quad (4.18)$$

Thus, we obtain

$$\begin{aligned}
& 2 \int_{\Omega} u_i'' m \cdot \nabla u_i dx \\
&= 2 \int_{\Omega} \left[\mu \Delta u_i + (\lambda + \mu) \frac{\partial}{\partial x_i} (\operatorname{div} u) - \alpha \frac{\partial \theta}{\partial x_i} \right] m \cdot \nabla u_i dx \\
&= \mu \int_{\Gamma} \left[2 \frac{\partial u_i}{\partial \nu} m \cdot \nabla u_i - m \cdot \nu |\nabla u_i|^2 \right] d\Gamma + (n-2)\mu \|\nabla u_i\|^2 \\
&\quad + (\lambda + \mu) \int_{\Gamma} \left[2 \operatorname{div}(u) m_k \nu_i \frac{\partial u_i}{\partial x_k} - m \cdot \nu |\operatorname{div} u|^2 \right] d\Gamma \\
&\quad + (n-2)(\lambda + \mu) \|\operatorname{div} u\|^2 - 2\alpha \int_{\Omega} (m \cdot \nabla u_i) \frac{\partial \theta}{\partial x_i} dx \\
&= \int_{\Gamma_1} m \cdot \nu \left[\mu \left| \frac{\partial u_i}{\partial \nu} \right|^2 + (\lambda + \mu) |\operatorname{div} u|^2 \right] d\Gamma \\
&\quad + 2 \int_{\Gamma_2} \left[\mu \frac{\partial u_i}{\partial \nu} + (\lambda + \mu) \nu_i \operatorname{div} u \right] m \cdot \nabla u_i d\Gamma \\
&\quad - \int_{\Gamma_2} m \cdot \nu [\mu |\nabla u_i|^2 + (\lambda + \mu) |\operatorname{div} u|^2] d\Gamma \\
&\quad + (n-2)[\mu \|\nabla u_i\|^2 + (\lambda + \mu) \|\operatorname{div} u\|^2] \\
&\quad - 2\alpha \int_{\Omega} (m \cdot \nabla u_i) \frac{\partial \theta}{\partial x_i} dx.
\end{aligned} \tag{4.19}$$

Since $m \cdot \nu \leq 0$ on Γ_1 , we have

$$\int_{\Gamma_1} m \cdot \nu \left[\mu \left| \frac{\partial u_i}{\partial \nu} \right|^2 + (\lambda + \mu) |\operatorname{div} u|^2 \right] d\Gamma \leq 0. \tag{4.20}$$

In addition, we have

$$-2\alpha \int_{\Omega} (m \cdot \nabla u_i) \frac{\partial \theta}{\partial x_i} dx \leq \varepsilon \mu \|\nabla u_i(t)\|^2 + \frac{\alpha^2 R_0^2}{\mu \varepsilon} \|\nabla \theta(t)\|^2. \tag{4.21}$$

It therefore follows from (4.19)–(4.21) that

$$\begin{aligned}
& 2 \int_{\Omega} u_i'' m \cdot \nabla u_i dx \\
&\leq 2 \int_{\Gamma_2} \left[\mu \frac{\partial u_i}{\partial \nu} + (\lambda + \mu) \nu_i \operatorname{div} u \right] m \cdot \nabla u_i d\Gamma \\
&\quad - \int_{\Gamma_2} m \cdot \nu [\mu |\nabla u_i|^2 + (\lambda + \mu) |\operatorname{div} u|^2] d\Gamma \\
&\quad + (n-2)[\mu \|\nabla u_i\|^2 + (\lambda + \mu) \|\operatorname{div} u\|^2] \\
&\quad + \varepsilon \mu \|\nabla u_i(t)\|^2 + \frac{\alpha^2 R_0^2}{\mu \varepsilon} \|\nabla \theta(t)\|^2.
\end{aligned} \tag{4.22}$$

For the second integral in (4.17), we have

$$2 \int_{\Omega} u_i' m \cdot \nabla u_i' dx = -n \|u_i'(t)\|^2 + \int_{\Gamma_2} m \cdot \nu |u_i'|^2 d\Gamma. \tag{4.23}$$

Using (2.25), we deduce

$$\begin{aligned}
& (n-1) \int_{\Omega} u_i u_i'' dx \\
&= (n-1) \int_{\Omega} u_i \left[\mu \Delta u_i + (\lambda + \mu) \frac{\partial}{\partial x_i} (\operatorname{div} u) - \alpha \frac{\partial \theta}{\partial x_i} \right] dx \\
&= (n-1) \int_{\Gamma_2} \left[\mu \frac{\partial u_i}{\partial \nu} + (\lambda + \mu) \nu_i \operatorname{div} u \right] u_i d\Gamma \\
&\quad - (n-1) [\mu \|\nabla u_i(t)\|^2 + (\lambda + \mu) \|\operatorname{div} u(t)\|^2] \\
&\quad - (n-1) \alpha \int_{\Omega} u_i \frac{\partial \theta}{\partial x_i} dx \\
&\leq (n-1) \int_{\Gamma_2} \left[\mu \frac{\partial u_i}{\partial \nu} + (\lambda + \mu) \nu_i \operatorname{div} u \right] u_i d\Gamma \\
&\quad - (n-1) [\mu \|\nabla u_i(t)\|^2 + (\lambda + \mu) \|\operatorname{div} u(t)\|^2] \\
&\quad + (n-1)^2 \frac{\alpha^2 \lambda_0^2}{4\varepsilon} \|\nabla \theta(t)\|^2 + \varepsilon \|u(t)\|_{(H_{\Gamma_1}^1(\Omega))^n}^2.
\end{aligned} \tag{4.24}$$

By (2.28) and (2.29), we have

$$\begin{aligned}
& C \int_{\Omega} \varphi_i u_i'' dx \\
&= C \int_{\Omega} \varphi_i \left[\mu \Delta u_i + (\lambda + \mu) \frac{\partial}{\partial x_i} (\operatorname{div} u) - \alpha \frac{\partial \theta}{\partial x_i} \right] dx \\
&= C \int_{\Omega_2} \left[\mu \frac{\partial u_i}{\partial \nu} + (\lambda + \mu) \nu_i \operatorname{div} u \right] \varphi_i d\Gamma \\
&\quad - C \int_{\Omega} [\mu \nabla u_i(t) \nabla \varphi_i(t) + (\lambda + \mu) \operatorname{div} u(t) \operatorname{div} \varphi(t)] dx \\
&\quad - C \alpha \int_{\Omega} \varphi_i \frac{\partial \theta}{\partial x_i} dx \\
&\leq C \int_{\Gamma_2} \left[\mu \frac{\partial u_i}{\partial \nu} + (\lambda + \mu) \nu_i \operatorname{div} u \right] u_i d\Gamma \\
&\quad + \frac{C^2 \alpha^2 k_1^2 \gamma^2}{4\varepsilon} \|\nabla \theta(t)\|^2 + \frac{\varepsilon}{k_1^2 \gamma^2} \|\varphi_i(t)\|^2 \\
&\leq C \int_{\Gamma_2} \left[\alpha \frac{\partial u_i}{\partial \nu} + (\lambda + \mu) \nu_i \operatorname{div} u \right] u_i d\Gamma \\
&\quad + \frac{C^2 \alpha^2 k_1^2 \gamma^2}{4\varepsilon} \|\nabla \theta(t)\|^2 + \varepsilon \|u(t)\|_{(H_{\Gamma_1}^1(\Omega))^n}^2.
\end{aligned} \tag{4.25}$$

Using (2.28), we have

$$\begin{aligned}
C \int_{\Omega} \varphi_i' u_i' dx &\leq \frac{C^2}{4\varepsilon} \|\varphi_i'(t)\|^2 + \varepsilon \|u_i'(t)\|^2 \\
&\leq \frac{C^2 k_1^2}{4\varepsilon} \int_{\Gamma_2} |u_i'(t)|^2 d\Gamma + \varepsilon \|u_i'(t)\|^2.
\end{aligned} \tag{4.26}$$

It therefore follows from (4.17) and (4.22)–(4.26) that

$$\begin{aligned}
F'(t) &\leq -[\mu\|\nabla u_i\|^2 + (\lambda + \mu)\|\operatorname{div} u\|^2 + \|u'_i(t)\|^2] \\
&\quad + 3\varepsilon\|u(t)\|_{(H^1_{\Gamma_1}(\Omega))^n}^2 + \varepsilon\|u'_i(t)\|^2 \\
&\quad + \left[\frac{\alpha^2 R_0^2}{\mu\varepsilon} + \frac{(n-1)^2\alpha^2\lambda_0^2}{4\varepsilon} + \frac{C^2\alpha^2 k_1^2\gamma^2}{4\varepsilon} \right] \|\nabla\theta(t)\|^2 \\
&\quad + 2 \int_{\Gamma_2} \left[\mu \frac{\partial u_i}{\partial \nu} + (\lambda + \mu)\nu_i \operatorname{div} u \right] m \cdot \nabla u_i \, d\Gamma \quad (= I_1) \\
&\quad - \int_{\Gamma_2} m \cdot \nu [\mu|\nabla u_i|^2 + (\lambda + \mu)|\operatorname{div} u|^2] \, d\Gamma \quad (= I_2) \\
&\quad + \int_{\Gamma_2} \left[m \cdot \nu + \frac{C^2 k_1^2}{4\varepsilon} \right] |u'_i(t)|^2 \, d\Gamma \quad (= I_3) \\
&\quad + (n-1+C) \int_{\Gamma_2} \left[\mu \frac{\partial u_i}{\partial \nu} + (\lambda + \mu)\nu_i \operatorname{div} u \right] u_i \, d\Gamma \quad (= I_4) \\
&\leq -2E(t) + 6\varepsilon E(t) \\
&\quad + \left[\frac{\alpha^2 R_0^2}{\mu\varepsilon} + \frac{(n-1)^2\alpha^2\lambda_0^2}{4\varepsilon} + \frac{C^2\alpha^2 k_1^2\gamma^2}{4\varepsilon} \right] \|\nabla\theta(t)\|^2 + \frac{\alpha}{\beta} \|\theta(t)\|^2 \\
&\quad + I_1 + I_2 + I_3 + I_4 \\
&\quad + \int_{\Gamma_2} am \cdot \nu |u_i|^2 \, d\Gamma \quad (= I_5).
\end{aligned} \tag{4.27}$$

Note that we add here two terms

$$\frac{\alpha}{\beta} \|\theta(t)\|^2, \quad \int_{\Gamma_2} am \cdot \nu |u_i|^2 \, d\Gamma \tag{4.28}$$

to make up $[\mu\|\nabla u_i\|^2 + (\lambda + \mu)\|\operatorname{div} u\|^2 + \|u'_i(t)\|^2]$ into $E(t)$. Since

$$\begin{aligned}
I_1 &= -2 \int_{\Gamma_2} [am \cdot \nu u_i + m \cdot \nu g_i(u')] m \cdot \nabla u_i \, d\Gamma \\
&\leq \int_{\Gamma_2} m \cdot \nu \left[\frac{2\alpha^2 R_0^2}{\mu} |u_i|^2 + \frac{2R_0^2}{\mu} |g(u')|^2 + \mu |\nabla u_i|^2 \right] \, d\Gamma,
\end{aligned} \tag{4.29}$$

we have

$$I_1 + I_2 \leq \int_{\Gamma_2} m \cdot \nu \left[\frac{2\alpha^2 R_0^2}{\mu} |u_i|^2 + \frac{2R_0^2}{\mu} |g(u')|^2 \right] \, d\Gamma. \tag{4.30}$$

In addition,

$$\begin{aligned}
 I_4 + I_5 &= -(n-1+C) \int_{\Gamma_2} [am \cdot \nu u_i + m \cdot \nu g_i(u')] u_i d\Gamma + \int_{\Gamma_2} am \cdot \nu |u_i|^2 d\Gamma \\
 &\leq -(n-2+C) \int_{\Gamma_2} am \cdot \nu |u_i|^2 d\Gamma \\
 &\quad + \int_{\Gamma_2} m \cdot \nu \left[\frac{(n-1+C)^2 R_0 \gamma^2}{4\varepsilon} |g_i(u')|^2 + \frac{\varepsilon}{R_0 \gamma^2} |u_i|^2 \right] d\Gamma \\
 &\leq -(n-2+C) \int_{\Gamma_2} am \cdot \nu |u_i|^2 d\Gamma \\
 &\quad + \frac{(n-1+C)^2 R_0 \gamma^2}{4\varepsilon} \int_{\Gamma_2} m \cdot \nu |g(u')|^2 d\Gamma + \varepsilon \|u(t)\|_{(H_{\Gamma_1}^1(\Omega))^n}^2
 \end{aligned} \tag{4.31}$$

Noting definitions (4.12) and (4.13) of C_2'' and C_3'' , it therefore follows from (4.27)–(4.31) that

$$\begin{aligned}
 F'(t) &\leq -2E(t) + 8\varepsilon E(t) + C_2'' \|\nabla \theta(t)\|^2 \\
 &\quad + \int_{\Gamma_2} \left[m \cdot \nu + \frac{C_1^2 k_1^2}{4\varepsilon} \right] |u'|^2 d\Gamma + C_3'' \int_{\Gamma_2} m \cdot \nu |g(u')|^2 d\Gamma \\
 &\quad + \int_{\Gamma_2} am \cdot \nu \left[\frac{2aR_0^2}{\mu} - n + 2 - C \right] |u_i|^2 d\Gamma \quad (= I).
 \end{aligned} \tag{4.32}$$

Thus, by taking $C = C_0$ and $\varepsilon = 1/8$, we have $I \leq 0$, and then we deduce (4.11).

If, further, the function $a(x)$ satisfies (2.51), we can take $C = 0$ and we still have $I \leq 0$. Then (4.11) follows.

Likewise, if the function $a(x)$ satisfies (2.50), then we can also take $C = 0$. However, in this case, since I is no longer negative, we estimate I as follows. Let $K(a)$ be given by (2.23). Using (2.24), we deduce

$$\begin{aligned}
 I &\leq K(a) \int_{\Gamma_2} m \cdot \nu |u|^2 d\Gamma \\
 &\leq K(a) R_0 \gamma^2 \|u(t)\|_{(H_{\Gamma_1}^1(\Omega))^n}^2 \\
 &\leq 2K(a) R_0 \gamma^2 E(t).
 \end{aligned} \tag{4.33}$$

Thus, by taking $\varepsilon = (1 - 2K(a)R_0\gamma^2)/8$, we also deduce (4.11). This completes the proof of Lemma 4.2. $\square$

We are now ready to prove our main results. The method of the proof is analogous to the one developed by the second author in [51].

Proof of Theorem 2.2. We first assume that $(u^0, u^1, \theta^0) \in D(\Lambda)$. Then Lemma 4.2 is valid. We note that $C = 0$ because the function $a(x)$ satisfies (2.50) or (2.51). Then the constant C_1'' in Lemma 4.1 and the constants C_2'' and C_3'' in Lemma 4.2 become C_1, C_2, C_3

in Theorem 2.2, respectively. By (2.42), (4.10) and Lemmas 4.1 and 4.2, we have

$$\begin{aligned}
 V'(t) &= E'(t) + \delta \frac{d}{dt}(E^\sigma(t)F(t)) \\
 &= E'(t) + \delta \sigma E^{\sigma-1}(t)E'(t)F(t) + \delta E^\sigma(t)F'(t) \\
 &\leq [1 - \delta C_1 \sigma E^\sigma(t)]E'(t) - \delta E^{\sigma+1}(t) \\
 &\quad + \delta C_2 E^\sigma(t) \|\nabla \theta(t)\|^2 + \delta E^\sigma(t) \int_{\Gamma_2} m \cdot \nu [|u'|^2 + C_3 |g(u')|^2] d\Gamma \\
 &= -\delta E^{\sigma+1}(t) + \left[\delta E^\sigma(t) \left(\frac{\sigma \alpha C_1}{\beta} + C_2 \right) - \frac{\alpha}{\beta} \right] \|\nabla \theta(t)\|^2 \\
 &\quad - [1 - \delta C_1 \sigma E^\sigma(t)] \int_{\Gamma_2} m \cdot \nu g(u') \cdot u' d\Gamma \\
 &\quad + \delta E^\sigma(t) \int_{\Gamma_2} m \cdot \nu [|u'|^2 + C_3 |g(u')|^2] d\Gamma.
 \end{aligned} \tag{4.34}$$

We now distinguish the cases $p = q = 1$ and $p + 1 > 2q$.

Case I: $p = q = 1$. In this case, we take $\sigma = 0$ and $\delta = \delta_1$ (see (2.64)) in (4.34). By definitions (2.64) and (2.59) of δ_1 and C_4 , it follows from (4.34) and (2.49) that

$$\begin{aligned}
 V'(t) &\leq -\delta_1 E(t) - \int_{\Gamma_2} m \cdot \nu g(u') \cdot u' d\Gamma \\
 &\quad + \delta_1 C_4 \int_{\Gamma_2} m \cdot \nu |u'|^2 d\Gamma \\
 &\leq -\delta_1 E(t) + [\delta_1 C_4 - k_3] \int_{\Gamma_2} m \cdot \nu |u'|^2 d\Gamma \\
 &\leq -\delta_1 E(t) \quad (\text{use (4.5)}) \\
 &\leq -\frac{\delta_1}{1 + \delta_1 C_1} V(t).
 \end{aligned} \tag{4.35}$$

Solving this differential inequality and using (4.5), we obtain

$$\begin{aligned}
 E(t) &\leq \frac{1}{1 - \delta_1 C_1} V(t) \\
 &\leq \frac{1}{1 - \delta_1 C_1} V(0) e^{-\delta_1 t / (1 + \delta_1 C_1)} \\
 &\leq \frac{1 + \delta_1 C_1}{1 - \delta_1 C_1} E(0) e^{-\delta_1 t / (1 + \delta_1 C_1)}.
 \end{aligned} \tag{4.36}$$

This is (2.52).

Case II: $p + 1 > 2q$. We first estimate

$$E^\sigma(t) \int_{\Gamma_2 \cap \{|u'| \leq 1\}} m \cdot \nu [|u'|^2 + C_3 |g(u')|^2] d\Gamma. \tag{4.37}$$

Applying Young's inequality

$$ab \leq \frac{a^s}{s} + \frac{b^\tau}{\tau}, \quad \forall a, b \geq 0, \quad \text{and } s, \tau > 0 \text{ with } \frac{1}{s} + \frac{1}{\tau} = 1, \tag{4.38}$$

we have for any $b > 0$

$$\begin{aligned}
& E^\sigma \int_{\Gamma_2 \cap [|u'| \leq 1]} m \cdot \nu |u'|^{2q} d\Gamma \\
& \leq \frac{p+1-2q}{p+1} b^{(p+1)/(p+1-2q)} E^{\sigma(p+1)/(p+1-2q)} \\
& \quad + \frac{2q}{(p+1)b^{(p+1)/(2q)}} \left(\int_{\Gamma_2 \cap [|u'| \leq 1]} m \cdot \nu |u'|^{2q} d\Gamma \right)^{(p+1)/(2q)} \\
& \leq \frac{p+1-2q}{p+1} b^{(p+1)/(p+1-2q)} E^{\sigma(p+1)/(p+1-2q)} \\
& \quad + \frac{2q}{(p+1)b^{(p+1)/(2q)}} \left(\int_{\Gamma_2 \cap [|u'| \leq 1]} m \cdot \nu d\Gamma \right)^{(p+1-2q)/(2q)} \\
& \quad \times \int_{\Gamma_2 \cap [|u'| \leq 1]} m \cdot \nu |u'|^{p+1} d\Gamma \\
& \leq \frac{p+1-2q}{p+1} b^{(p+1)/(p+1-2q)} E^{\sigma(p+1)/(p+1-2q)} \\
& \quad + \frac{2q[R_0 \text{mes}(\Gamma_2)]^{(p+1-2q)/(2q)}}{k_3(p+1)b^{(p+1)/(2q)}} \int_{\Gamma_2 \cap [|u'| \leq 1]} m \cdot \nu g(u') \cdot u' d\Gamma.
\end{aligned} \tag{4.39}$$

Thus, by taking

$$b = \left(\frac{p+1}{2C_4(p+1-2q)} \right)^{(p+1-2q)/(p+1)}, \tag{4.40}$$

we obtain

$$\begin{aligned}
& E^\sigma(t) \int_{\Gamma_2 \cap [|u'| \leq 1]} m \cdot \nu [|u'|^2 + C_3 |g(u')|^2] d\Gamma \\
& \text{(note that } |u'|^2 \leq |u'|^{2q} \text{ since } q \leq 1) \\
& \leq C_4 E^\sigma(t) \int_{\Gamma_2 \cap [|u'| \leq 1]} m \cdot \nu |u'|^{2q} d\Gamma \\
& \leq \frac{1}{2} E^{\sigma(p+1)/(p+1-2q)} \\
& \quad + \frac{q(2C_4)^{(p+1)/(2q)} [R_0 \text{mes}(\Gamma_2)(p+1-2q)]^{(p+1-2q)/(2q)}}{k_3(p+1)^{(p+1)/(2q)}} \\
& \quad \times \int_{\Gamma_2 \cap [|u'| \leq 1]} m \cdot \nu g(u') \cdot u' d\Gamma.
\end{aligned} \tag{4.41}$$

It therefore follows from (4.34) and (4.41) that

$$\begin{aligned}
 V'(t) &\leq -\delta E^{\sigma+1}(t) + \left[\delta E^{\sigma}(t) \left(\frac{\sigma \alpha C_1}{\beta} + C_2 \right) - \frac{\alpha}{\beta} \right] \|\nabla \theta(t)\|^2 \\
 &\quad + \left[\delta C_1 \sigma E^{\sigma}(0) + \frac{\delta C_4 E^{\sigma}(0)}{k_3} - 1 \right] \int_{\Gamma_2 \cap [|u'| \geq 1]} m \cdot \nu g(u') \cdot u' d\Gamma \\
 &\quad + [\delta C_1 \sigma E^{\sigma}(0) - 1] \int_{\Gamma_2 \cap [|u'| \leq 1]} m \cdot \nu g(u') \cdot u' d\Gamma \\
 &\quad + \delta E^{\sigma}(t) \int_{\Gamma_2 \cap [|u'| \leq 1]} m \cdot \nu [|u'|^2 + C_3 |g(u')|^2] d\Gamma \\
 &\leq -\delta E^{\sigma+1}(t) + \frac{\delta}{2} E^{\sigma(p+1)/(p+1-2q)} \\
 &\quad + \left[\delta E^{\sigma}(t) \left(\frac{\sigma \alpha C_1}{\beta} + C_2 \right) - \frac{\alpha}{\beta} \right] \|\nabla \theta(t)\|^2 \\
 &\quad + \left[\delta C_1 \sigma E^{\sigma}(0) + \frac{\delta C_4 E^{\sigma}(0)}{k_3} - 1 \right] \int_{\Gamma_2 \cap [|u'| \geq 1]} m \cdot \nu g(u') \cdot u' d\Gamma \\
 &\quad + [\delta C_1 \sigma E^{\sigma}(0) - 1] \int_{\Gamma_2 \cap [|u'| \leq 1]} m \cdot \nu g(u') \cdot u' d\Gamma \\
 &\quad + \delta \frac{q(2C_4)^{(p+1)/(2q)} [R_0 \text{mes}(\Gamma_2)(p+1-2q)]^{(p+1-2q)/(2q)}}{k_3(p+1)^{(p+1)/(2q)}} \\
 &\quad \times \int_{\Gamma_2 \cap [|u'| \leq 1]} m \cdot \nu g(u') \cdot u' d\Gamma.
 \end{aligned} \tag{4.42}$$

We now choose σ so that

$$\sigma + 1 = \frac{\sigma(p+1)}{p+1-2q}. \tag{4.43}$$

Then

$$\sigma = \frac{p+1-2q}{2q} = \sigma_0. \tag{4.44}$$

By taking $\delta = \delta_2$ (see (2.65)) and $\sigma = \sigma_0$ in (4.42), we obtain

$$V'(t) \leq -\frac{\delta_2}{2} E^{\sigma_0+1}(t). \tag{4.45}$$

By (4.5), we have

$$V(t) \leq [1 + \delta_2 C_1 E^{\sigma_0}(0)] E(t). \tag{4.46}$$

It therefore follows from (4.45) that

$$V'(t) \leq -\frac{\delta_2}{2[1 + \delta_2 C_1 E^{\sigma_0}(0)]^{\sigma_0}} V^{\sigma_0+1}(t). \tag{4.47}$$

Solving this differential inequality and noting definition (2.65) of δ_2 , we obtain

$$\begin{aligned}
 V(t) &\leq \left[(V(0))^{-\sigma_0} + \frac{\sigma_0 \delta_2 t}{2[1 + \delta_2 C_1 E^{\sigma_0}(0)]^{\sigma_0+1}} \right]^{-1/\sigma_0} \\
 &\leq \left[(E(0) + \delta_2 C_1 E^{\sigma_0+1}(0))^{-\sigma_0} + \frac{\sigma_0 \delta_2 t}{2[1 + \delta_2 C_1 E^{\sigma_0}(0)]^{\sigma_0+1}} \right]^{-1/\sigma_0} \\
 &\leq 2E(0) \left[1 + \frac{\sigma_0 \delta_2 t}{2(2E(0))^{-\sigma_0}[1 + \delta_2 C_1 E^{\sigma_0}(0)]^{\sigma_0+1}} \right]^{-1/\sigma_0},
 \end{aligned} \tag{4.48}$$

which, combined with (4.5), implies (2.53).

Finally, if $(u^0, u^1, \theta^0) \in \mathcal{H}$, in view of (3.84), we can show that Theorem 2.2 still holds by a density argument. $\square$

Proof of Theorem 2.3. We first note that $C = C_0$ and the constant C_1'' in Lemma 4.1 and the constants C_2'' and C_3'' in Lemma 4.2 become C_1', C_2', C_3' in Theorem 2.3, respectively. In this case, (4.34) becomes

$$\begin{aligned}
 V'(t) &\leq -\delta E^{\sigma+1}(t) + \left[\delta E^\sigma(t) \left(\frac{\sigma \alpha C_1'}{\beta} + C_2' \right) - \frac{\alpha}{\beta} \right] \|\nabla \theta(t)\|^2 \\
 &\quad - [1 - \delta C_1' \sigma E^\sigma(t)] \int_{\Gamma_2} m \cdot \nu g(u') \cdot u' d\Gamma \\
 &\quad + \delta E^\sigma(t) \int_{\Gamma_2} \left[\left(m \cdot \nu + \frac{C_0^2 k_1^2}{4\varepsilon} \right) |u'|^2 + C_3' m \cdot \nu |g(u')|^2 \right] d\Gamma.
 \end{aligned} \tag{4.49}$$

Because the additional term $\frac{C_0^2 k_1^2}{4\varepsilon}$ does not contain the factor $m \cdot \nu$, we have to impose condition (2.69) on Γ_2 so that the sum of the integrals on Γ_2 becomes negative if δ is sufficiently small. The rest of the proof is the same as that of Theorem 2.2 except that C_4 is replaced by C_4' and the integral $\int_{\Gamma_2} m \cdot \nu g(u') \cdot u' d\Gamma$ is enlarged to $R_0 \int_{\Gamma_2} g(u') \cdot u' d\Gamma$ or reduced to $\eta \int_{\Gamma_2} g(u') \cdot u' d\Gamma$. $\square$

Proof of Theorem 2.5. The proof of the first part of Theorem 2.5 is the same as that of Theorems 2.2 and 2.3 except that all u' and $g(u')$ are replaced by u'_i and $h(u'_i)$, respectively. Also, in this case, the constant b in (4.40) is taken as

$$b = \left(\frac{p+1}{2C_4 n(p+1-2q)} \right)^{(p+1-2q)/(p+1)}, \tag{4.50}$$

and (4.41) becomes

$$\begin{aligned}
 E^\sigma(t) &\int_{\Gamma_2 \cap \{|u'_i| \leq 1\}} m \cdot \nu [|u'_i|^2 + C_3 |h(u'_i)|^2] d\Gamma \\
 &\leq \frac{1}{2n} E^{\sigma(p+1)/(p+1-2q)} \\
 &\quad + \frac{q(2C_4)^{(p+1)/(2q)} [n R_0 \text{mes}(\Gamma_2) (p+1-2q)]^{(p+1-2q)/(2q)}}{k_3(p+1)^{(p+1)/(2q)}} \\
 &\quad \times \int_{\Gamma_2 \cap \{|u'_i| \leq 1\}} m \cdot \nu h(u'_i) u'_i d\Gamma.
 \end{aligned} \tag{4.51}$$

Note that the space dimension n appears here.

However, the proof of the decay rate (2.101) is a bit different. Hence we present it as follows.

We first look at the case corresponding to Theorem 2.2. Since $p > 1$ and $q = 1/p$, as in Remark 2.12, we have

$$|s|^2 + C_3|h(s)|^2 \leq C_5(h(s)s)^{2/(p+1)}, \quad |s| \leq 1, \quad (4.52)$$

where

$$C_5 = \frac{1 + C_3 k_2 k_3}{k_3}. \quad (4.53)$$

It therefore follows that for any $b > 0$

$$\begin{aligned} E^\sigma(t) & \int_{\Gamma_2 \cap [|u'_i| \leq 1]} m \cdot \nu [|u'_i|^2 + C_3 |h(u'_i)|^2] d\Gamma \\ & \leq C_5 \frac{p-1}{p+1} b^{(p+1)/(p-1)} E^{\sigma(p+1)/(p-1)} \\ & \quad + \frac{2C_5}{(p+1)b^{(p+1)/2}} \left(\int_{\Gamma_2 \cap [|u'_i| \leq 1]} m \cdot \nu (u'_i h(u'_i))^{2/(p+1)} \Gamma \right)^{(p+1)/2} \\ & \leq C_5 \frac{p-1}{p+1} b^{(p+1)/(p-1)} E^{\sigma(p+1)/(p-1)} \\ & \quad + \frac{2C_5}{(p+1)b^{(p+1)/2}} \left(\int_{\Gamma_2 \cap [|u'_i| \leq 1]} m \cdot \nu d\Gamma \right)^{(p-1)/2} \\ & \quad \times \int_{\Gamma_2 \cap [|u'_i| \leq 1]} m \cdot \nu u'_i h(u'_i) d\Gamma \\ & \leq C_5 \frac{p-1}{p+1} b^{(p+1)/(p-1)} E^{\sigma(p+1)/(p-1)} \\ & \quad + \frac{2C_5 [R_0 \text{mes}(\Gamma_2)]^{(p-1)/2}}{(p+1)b^{(p+1)/2}} \int_{\Gamma_2 \cap [|u'_i| \leq 1]} m \cdot \nu u'_i h(u'_i) d\Gamma. \end{aligned} \quad (4.54)$$

Thus, by taking

$$b = \left(\frac{p+1}{2C_5(p-1)} \right)^{(p-1)/(p+1)}, \quad (4.55)$$

we obtain

$$\begin{aligned} E^\sigma(t) & \int_{\Gamma_2 \cap [|u'_i| \leq 1]} m \cdot \nu [|u'_i|^2 + C_3 |h(u'_i)|^2] d\Gamma \\ & \leq \frac{1}{2} E^{\sigma(p+1)/(p-1)} + \frac{(2C_5)^{(p+1)/2} [R_0 \text{mes}(\Gamma_2)(p-1)]^{(p-1)/2}}{(p+1)^{(p+1)/2}} \\ & \quad \times \int_{\Gamma_2 \cap [|u'_i| \leq 1]} m \cdot \nu u'_i h(u'_i) d\Gamma. \end{aligned} \quad (4.56)$$

It therefore follows from (4.34) that

$$\begin{aligned}
 V'(t) &\leq -\delta E^{\sigma+1}(t) + \left[\delta E^{\sigma}(t) \left(\frac{\sigma \alpha C_1}{\beta} + C_2 \right) - \frac{\alpha}{\beta} \right] \|\nabla \theta(t)\|^2 \\
 &\quad + \left[\delta C_1 \sigma E^{\sigma}(0) + \frac{\delta C_4 E^{\sigma}(0)}{k_3} - 1 \right] \int_{\Gamma_2 \cap \{|u'_i| \geq 1\}} m \cdot \nu u'_i h(u'_i) d\Gamma \\
 &\quad + [\delta C_1 \sigma E^{\sigma}(0) - 1] \int_{\Gamma_2 \cap \{|u'_i| \leq 1\}} m \cdot \nu u'_i h(u'_i) d\Gamma \\
 &\quad + \delta E^{\sigma}(t) \int_{\Gamma_2 \cap \{|u'_i| \leq 1\}} m \cdot \nu [|u'_i|^2 + C_3 |h(u'_i)|^2] d\Gamma \\
 &\leq -\delta E^{\sigma+1}(t) + \frac{\delta}{2} E^{\sigma(p+1)/(p-1)} \\
 &\quad + \left[\delta E^{\sigma}(t) \left(\frac{\sigma \alpha C_1}{\beta} + C_2 \right) - \frac{\alpha}{\beta} \right] \|\nabla \theta(t)\|^2 \\
 &\quad + \left[\delta C_1 \sigma E^{\sigma}(0) + \frac{\delta C_4 E^{\sigma}(0)}{k_3} - 1 \right] \int_{\Gamma_2 \cap \{|u'_i| \geq 1\}} m \cdot \nu u'_i h(u'_i) d\Gamma \\
 &\quad + [\delta C_1 \sigma E^{\sigma}(0) - 1] \int_{\Gamma_2 \cap \{|u'_i| \leq 1\}} m \cdot \nu u'_i h(u'_i) d\Gamma \\
 &\quad + \delta \frac{(2C_5)^{(p+1)/2} [R_0 \text{mes}(\Gamma_2)(p-1)]^{(p-1)/2}}{(p+1)^{(p+1)/2}} \\
 &\quad \times \int_{\Gamma_2 \cap \{|u'_i| \leq 1\}} m \cdot \nu u'_i h(u'_i) d\Gamma.
 \end{aligned} \tag{4.57}$$

By taking $\delta = \delta_2$ (see (2.65)) and $\sigma = (p-1)/2$, we obtain

$$V'(t) \leq -\frac{\delta_2}{2} E^{(p+1)/2}(t). \tag{4.58}$$

The remainder of the proof is the same as that of Theorem 2.2.

The proof of the case corresponding to Theorem 2.3 is similar. $\square$

We now use the method developed in [41] to prove Theorem 2.7.

Proof of Theorem 2.7. If $E(t_0) = 0$ for some $t_0 \geq 0$, then, by (2.42), we have $E(t) \equiv 0$ for $t \geq t_0$ and then the theorem holds. Therefore, we may assume that $E(t) > 0$ for $t \geq 0$. This assumption ensures that, in the following proof, $\varphi''(aE(t))$ makes sense since we have assumed that $\varphi(s)$ is twice differentiable outside $s = 0$.

We first note that $C = C_0$ and the constant C_1'' in Lemma 4.1 and the constants C_2'' and C_3'' in Lemma 4.2 become C_1', C_2', C_3' in Theorem 2.3, respectively. Set

$$V(t) = E(t) + \delta \varphi'(aE(t)) F(t), \tag{4.59}$$

where the constant a is given by (2.116). By (2.42), (4.10) and Lemmas 4.1 and 4.2, we have

$$\begin{aligned}
V'(t) &= E'(t) + \delta a \varphi''(aE(t))E'(t)F(t) + \delta \varphi'(aE(t))F'(t) \\
&\leq [1 - \delta a C_1' \varphi''(aE(0))E(0)]E'(t) - \delta \varphi'(aE(t))E(t) \\
&\quad + \delta C_2' \varphi'(aE(0)) \|\nabla \theta(t)\|^2 \\
&\quad + \delta \varphi'(aE(t)) \int_{\Gamma_2} [(m \cdot \nu + 2C_0^2 k_1^2 |u'|^2 + C_3' m \cdot \nu \sum_{i=1}^n |h(u'_i)|^2) d\Gamma \\
&\leq -\delta \varphi'(aE(t))E(t) + [\delta C_2' \varphi'(aE(0)) + \alpha \delta a C_1' \varphi''(aE(0))E(0)/\beta - \alpha/\beta] \|\nabla \theta(t)\|^2 \\
&\quad + [\eta \delta a C_1' \varphi''(aE(0))E(0) + \delta \varphi'(aE(0))[R_0 + 2C_0^2 k_1^2 + C_3' k_2^2 R_0]/k_3 - \eta] \\
&\quad \times \sum_{i=1}^n \int_{\Gamma_2 \cap [|u'_i| \geq 1]} h(u'_i) u'_i d\Gamma \\
&\quad + [\eta \delta a C_1' \varphi''(aE(0))E(0) - \eta] \sum_{i=1}^n \int_{\Gamma_2 \cap [|u'_i| \leq 1]} h(u'_i) u'_i d\Gamma \\
&\quad (\text{note that } |u'|^2 \leq |u'|^{2q} \text{ since } q \leq 1 \text{ and } |u'_i| \leq 1) \\
&\quad + \delta [R_0 + 2C_0^2 k_1^2 + C_3' k_2^2 R_0] \varphi'(aE(t)) \sum_{i=1}^n \int_{\Gamma_2 \cap [|u'_i| \leq 1]} |u'_i|^{2q} d\Gamma \\
&\leq -\delta \varphi'(aE(t))E(t) \\
&\quad + [\eta \delta a C_1' \varphi''(aE(0))E(0) - \eta] \sum_{i=1}^n \int_{\Gamma_2 \cap [|u'_i| \leq 1]} h(u'_i) u'_i d\Gamma \\
&\quad + \delta [R_0 + 2C_0^2 k_1^2 + C_3' k_2^2 R_0] \varphi'(aE(t)) \sum_{i=1}^n \int_{\Gamma_2 \cap [|u'_i| \leq 1]} |u'_i|^{2q} d\Gamma,
\end{aligned} \tag{4.60}$$

if δ is small enough. Let φ^* denote the dual of φ in the sense of Young (for definition, see [4, p. 64]). Then, by Young's inequality [4, p. 64] and Jensen's inequality [49], we deduce

$$\begin{aligned}
&\varphi'(aE(t)) \int_{\Gamma_2 \cap [|u'_i| \leq 1]} |u'_i|^{2q} d\Gamma \\
&= \text{mes}(\Gamma_2 \cap [|u'_i| \leq 1]) \varphi'(aE) \frac{1}{\text{mes}(\Gamma_2 \cap [|u'_i| \leq 1])} \int_{\Gamma_2 \cap [|u'_i| \leq 1]} |u'_i|^{2q} d\Gamma \\
&\leq \text{mes}(\Gamma_2 \cap [|u'_i| \leq 1]) \left[\varphi^*(\varphi'(aE)) + \varphi \left(\frac{1}{\text{mes}(\Gamma_2 \cap [|u'_i| \leq 1])} \int_{\Gamma_2 \cap [|u'_i| \leq 1]} |u'_i|^{2q} d\Gamma \right) \right] \\
&\leq \text{mes}(\Gamma_2) \varphi^*(\varphi'(aE)) + \int_{\Gamma_2 \cap [|u'_i| \leq 1]} \varphi(|u'_i|^{2q}) d\Gamma \\
&\leq \text{mes}(\Gamma_2) \varphi^*(\varphi'(aE)) + \int_{\Gamma_2 \cap [|u'_i| \leq 1]} u'_i h(u'_i) dx.
\end{aligned} \tag{4.61}$$

Hence, if δ is small enough, then it follows from (4.60) and (4.61) that

$$V'(t) \leq -\delta\varphi'(aE(t))E(t) + n\delta[R_0 + 2C_0^2k_1^2 + C_3'k_2^2R_0]\text{mes}(\Gamma_2)\varphi^*(\varphi'(aE)). \quad (4.62)$$

By the definition of the dual $\varphi^*(t)$ of $\varphi(s)$, $\varphi^*(t)$ is the Legendre transform of $\varphi(s)$, which is given by (see [4, pp. 61–62])

$$\varphi^*(t) = t\varphi'^{-1}(t) - \varphi[\varphi'^{-1}(t)]. \quad (4.63)$$

It therefore follows from (4.62) that

$$\begin{aligned} V'(t) &\leq -\delta\varphi'(aE(t))E(t) \\ &\quad + n\delta[R_0 + 2C_0^2k_1^2 + C_3'k_2^2R_0]\text{mes}(\Gamma_2)[a\varphi'(aE(t))E(t) - \varphi(aE(t))] \\ &= -n\delta[R_0 + 2C_0^2k_1^2 + C_3'k_2^2R_0]\text{mes}(\Gamma_2)\varphi(aE(t)) \\ &\quad - \frac{\delta}{2}\varphi'(aE(t))E(t), \end{aligned} \quad (4.64)$$

with

$$a = \frac{1}{2n[R_0 + 2C_0^2k_1^2 + C_3'k_2^2R_0]\text{mes}(\Gamma_2)}. \quad (4.65)$$

On the other hand, since $\varphi(s)$ and $\varphi'(s)$ are positive and increasing on $[0, \infty)$, it follows from (4.10) that

$$[1 - \delta C_1'\varphi'(aE(0))]E(t) \leq V(t) \leq [1 + \delta C_1'\varphi'(aE(0))]E(t). \quad (4.66)$$

It therefore follows from (4.64) that

$$\begin{aligned} V'(t) &\leq -\frac{\delta V(t)}{2(1 + \delta C_1'\varphi'(aE(0)))}\varphi'\left(\frac{aV(t)}{1 + \delta C_1'\varphi'(aE(0))}\right) \\ &\quad - n\delta[R_0 + 2C_0^2k_1^2 + C_3'k_2^2R_0]\text{mes}(\Gamma_2)\varphi\left(\frac{aV(t)}{1 + \delta C_1'\varphi'(aE(0))}\right). \end{aligned} \quad (4.67)$$

This is (2.115).

It remains to prove (2.118). We argue by contradiction. Suppose that $E(t)$ does not tend to zero as $t \rightarrow \infty$. Since $E(t)$ is decreasing on $[0, \infty)$, we have

$$E(t) \geq \sigma > 0, \quad \forall t \geq 0, \quad (4.68)$$

and by (4.66), we have

$$V(t) \geq \beta > 0, \quad \forall t \geq 0. \quad (4.69)$$

Thus we have

$$\varphi'\left(\frac{aV(t)}{b}\right) \geq \gamma > 0, \quad \forall t \geq 0. \quad (4.70)$$

It therefore follows from (2.115) that

$$V'(t) \leq -\frac{\delta\gamma}{2b}V(t), \quad \forall t \geq 0. \quad (4.71)$$

which is in contradiction with (4.69). This completes the proof. $\square$

Proof of Corollary 2.9. By Theorem 2.7, it suffices to prove that, for t large enough, (2.115) becomes (2.121). Since h is odd in $[-1, 1]$ and $s^{1/(2q)}h(s^{1/(2q)})$ is convex in $[0, 1]$, we can take for $0 \leq s \leq 1$

$$\varphi(s) = s^{1/(2q)}h(s^{1/(2q)}), \quad (4.72)$$

and then

$$\varphi'(s) = \frac{1}{2q} [s^{(1-2q)/(2q)}h(s^{1/(2q)}) + s^{(1-q)/q}h'(s^{1/(2q)})]. \quad (4.73)$$

On the other hand, by (2.118), there exists $T > 0$ such that

$$V(t) \leq 1, \quad \forall t \geq T. \quad (4.74)$$

Thus, for $t \geq T$, substituting (4.72) and (4.73) into (2.115), we obtain (2.121). $\square$

Proof of Corollary 2.10. If h satisfies (2.95), then we can take

$$\varphi(s) = k_3 s^{(p+1)/(2q)}, \quad \text{for } s \geq 0. \quad (4.75)$$

Thus, (2.115) becomes

$$V'(t) = -K(V(t))^{(p+1)/(2q)}, \quad (4.76)$$

where K is a positive constant independent of V . Hence, the decay properties (2.52) and (2.53) follow from (4.76). $\square$

5. Further comments. In the proof of Theorems 2.2–2.7, inequality (2.25) of Poincaré type plays a key role. This inequality is guaranteed by assumption (2.16). If this assumption does not hold, i.e., $\Gamma_1 = \emptyset$ and $a(x) \equiv 0$, then this inequality is no longer true for all $u \in H^1(\Omega)$. When $\Gamma_1 = \phi$ and $a(x) \equiv 0$, in order to guarantee the coercivity of the energy, it is natural to look for a closed subspace or subset $\mathcal{W}$ of $(H^1(\Omega))^n \times (L^2(\Omega))^n \times L^2(\Omega)$ invariant under the flow generated by the semigroup such that the energy norm (2.20) on $\mathcal{W}$ is equivalent to the usual one induced by $(H^1(\Omega))^n \times (L^2(\Omega))^n \times L^2(\Omega)$.

Naturally, one is tempted to consider the following space of functions with zero average:

$$\mathcal{H}_0 = \left\{ (u, v, \theta) \in (H^1(\Omega))^n \times (L^2(\Omega))^n \times L^2(\Omega) : \int_{\Omega} u(x) dx = \int_{\Omega} v(x) dx = 0 \right\}, \quad (5.1)$$

on which the energy norm is really equivalent to the usual one. Unfortunately, $\mathcal{H}_0$ is not invariant. To see this, we define the function

$$f(t) = \int_{\Omega} u(t) dx. \quad (5.2)$$

We look at the special case where $g(u) = u$. Take sufficiently regular initial $(u^0, u^1, \theta^0) \in \mathcal{H}_0$ such that

$$\int_{\Gamma} m \cdot \nu u^1 d\Gamma \neq 0, \quad (5.3)$$

and let u, θ be the solution of (2.7) corresponding to this initial data. By the continuity of $u'(t)$ with respect to t , we have that

$$\int_{\Gamma} m \cdot \nu u'(t) d\Gamma \neq 0 \quad (5.4)$$

for $t > 0$ small enough. It therefore follows that

$$\begin{aligned} f''(t) &= \int_{\Omega} u''(t) dx \\ &= \int_{\Omega} [\mu \Delta u + (\lambda + \mu) \nabla \operatorname{div} u - \alpha \nabla \theta] dx \\ &= \int_{\Gamma} \left[\mu \frac{\partial u}{\partial \nu} + (\lambda + \mu) \operatorname{div}(u) \nu - \alpha \theta \nu \right] d\Gamma \\ &= - \int_{\Gamma} m \cdot \nu u'(t) d\Gamma \\ &\neq 0. \end{aligned} \quad (5.5)$$

Hence, $\int_{\Omega} u(t) dx$ and $\int_{\Omega} u'(t) dx$ are not always equal to zero along the solution trajectories of (2.7).

On the other hand, we do have the following conserved quantity:

$$\int_{\Omega} u'(t) dx + \int_0^t \int_{\Gamma} m \cdot \nu g(u'(t)) d\Gamma dt \equiv C \text{ (a constant),} \quad (5.6)$$

since, by (2.7), we have

$$\int_{\Omega} u''(t) dx + \int_{\Gamma} m \cdot \nu g(u'(t)) d\Gamma \equiv 0. \quad (5.7)$$

If $g(u)$ is linear, i.e., $g(u) = ku$, then we can easily find an invariant subspace $\mathcal{W}$ as follows:

$$\mathcal{W} = V \times L^2(\Omega), \quad (5.8)$$

where

$$V = \left\{ (u, v) \in (H^1(\Omega))^n \times (L^2(\Omega))^n : \int_{\Gamma} km \cdot \nu u d\Gamma + \int_{\Omega} v dx = 0 \right\}. \quad (5.9)$$

Moreover, the energy norm on $\mathcal{W}$ is equivalent to the usual one. However, for the general nonlinear boundary feedbacks, it is difficult to find such an invariant closed subset. Thus, the case that $\Gamma_1 = \emptyset$ and $a(x) \equiv 0$ with g nonlinear is open.

Acknowledgment. Part of this work was done when the first author visited the Departamento de Matemática Aplicada, Universidad Complutense de Madrid, in May, 1998. He thanks their hospitality and financial support.

REFERENCES

- [1] R. Adams, *Sobolev Spaces*, Academic Press, New York, 1975
- [2] S. Agmon, *Lectures on Elliptic Boundary Value Problems*, D. Van Nostrand Company, Inc., Princeton, NJ, 1965
- [3] F. Alabau and V. Komornik, *Boundary observability, controllability and stabilization of linear elastodynamic systems*, SIAM J. Control Optim. **37**, no. 2, 521–542 (1999)
- [4] V. I. Arnold, *Mathematical Methods of Classical Mechanics*, Springer-Verlag, New York, 1989
- [5] G. Avalos and I. Lasiecka, *Exponential stability of a thermoelastic system with free boundary conditions without mechanical dissipation*, SIAM J. Math. Anal. **29**, 155–182 (1998)
- [6] V. Barbu, *Nonlinear Semigroups and Differential Equations in Banach Spaces*, Noordhoff, Leyden, The Netherlands, 1976
- [7] E. Bisognin, V. Bisognin, G. P. Menzala, and E. Zuazua, *On exponential stability for the Von Kármán equations in the presence of thermal effects*, Math. Methods Appl. Sci. **21**, 393–416 (1998)
- [8] H. Brezis, *Analyse Fonctionnelle: Théorie et Applications*, Masson, Paris, 1983
- [9] J. A. Burns, Z. Y. Liu, and S. Zheng, *On the energy decay of a linear thermoelastic bar*, J. Math. Anal. Appl. **179**, 574–591 (1993)
- [10] G. Chen, *Energy decay estimates and exact boundary value controllability for the wave equation in a bounded domain*, J. Math. Pures Appl. **58**, 249–273 (1979)
- [11] F. Conrad and B. Rao, *Decay of solutions of the wave equation in a star-shaped domain with nonlinear boundary feedback*, Asymptotic Anal. **7**, 159–177 (1993)
- [12] C. M. Dafermos, *On the existence and the asymptotic stability of solutions to the equations of linear thermoelasticity*, Arch. Rational Mech. Anal. **29**, 241–271 (1968)
- [13] R. Dautray and J. L. Lions, *Mathematical Analysis and Numerical Methods for Science and Technology, Vol. 2, Functional and Variational Methods*, Springer-Verlag, Berlin, 1992
- [14] P. Grisvard, *Singularities in Boundary Value Problems*, Masson, Paris, 1992
- [15] P. Grisvard, *Elliptic Problems in Nonsmooth Domains*, Pitman, London, 1985
- [16] P. Grisvard, *Contrôlabilité exacte des solutions de l'équation des ondes en présence de singularités*, J. Math. Pures Appl. **68**, 215–259 (1989)
- [17] S. W. Hansen, *Exponential energy decay in a linear thermoelastic rod*, J. Math. Anal. Appl. **167**, 429–442 (1992)
- [18] M. A. Horn, *Implications of sharp trace regularity results on boundary stabilization of the system of linear elasticity*, J. Math. Anal. Appl. **223**, 126–150 (1998)
- [19] M. A. Horn, *Sharp trace regularity for the solutions of the equations of dynamic elasticity*, J. Math. Systems Estim. Control **8** (2), 217–219 (1998)
- [20] J. U. Kim, *On the energy decay of a linear thermoelastic bar and plate*, SIAM J. Math. Anal. **23**, 889–899 (1992)
- [21] V. Komornik, *Exact Controllability and Stabilization: The Multiplier Method*, John Wiley and Sons, Masson, Paris, 1994
- [22] V. Komornik and B. Rao, *Boundary stabilization of compactly coupled wave equations*, Asymptotic Anal. **14**, 339–359 (1997)
- [23] V. Komornik and E. Zuazua, *A direct method for the boundary stabilization of the wave equation*, J. Math. Pures Appl. **69**, 33–54 (1990)
- [24] A. D. Kovalenko, *Thermoelasticity, Basic Theory and Applications*, Wolters-Noordhoff Publishing, Gröningen, Netherlands, 1969
- [25] J. Lagnese, *Decay of solutions of wave equations in a bounded region with boundary dissipation*, J. Differential Equations **50**, 163–182 (1983)
- [26] J. Lagnese, *Boundary stabilization of linear elastodynamic systems*, SIAM J. Control Optim. **21**, 968–984 (1983)
- [27] J. Lagnese, *Boundary Stabilization of Thin Plates*, SIAM Studies in Applied Mathematics, vol. 10, SIAM Publications, Philadelphia, 1989
- [28] J. Lagnese, *Uniform asymptotic energy estimates for solutions of the equations of dynamic plane elasticity with nonlinear dissipation at the boundary*, Nonlinear Anal. **16**, 35–54 (1991)
- [29] J. Lagnese and G. Leugering, *Uniform stabilization of a nonlinear beam by nonlinear boundary feedback*, J. Differential Equations **91**, 355–388 (1991)
- [30] J. Lagnese and J. L. Lions, *Modelling Analysis and Control of Thin Plates*, Masson, Paris, 1989

- [31] I. Lasiecka, *Stabilization of wave and plate-like equations with nonlinear dissipation on the boundary*, J. Differential Equations **79**, 340–381 (1989)
- [32] I. Lasiecka and D. Tataru, *Uniform boundary stabilization of semilinear wave equations with nonlinear boundary damping*, Differential Integral Equations **6**, 507–533 (1993)
- [33] G. Lebeau and E. Zuazua, *Sur la décroissance non uniforme de l'énergie dans le système de la thermoélasticité linéaire*, C. R. Acad. Sci. Paris Sér. I. Math. **324**, 409–415 (1997)
- [34] G. Leugering, *On boundary feedback stabilization of a viscoelastic membrane*, Dynam. Stability Systems **4**, 71–79 (1989)
- [35] G. Leugering, *On boundary feedback stabilization of a viscoelastic beam*, Proc. Roy. Soc. Edinburgh, Sect. A **114**, 57–69 (1990)
- [36] J. L. Lions and E. Magenes, *Non-homogeneous Boundary Value Problems and Applications*, vols. I and II, Springer-Verlag, New York, 1972
- [37] W. J. Liu, *Partial exact controllability and exponential stability of the higher-dimensional linear thermoelasticity*, ESAIM Contrôle Optim. Calc. Var. **3**, 23–48 (1998)
- [38] W. J. Liu, *Exact Controllability for Some Partial Differential Equations of Evolutional Type*, Ph.D. Thesis, University of Wollongong, Australia, 1997
- [39] W. J. Liu, *The exponential stabilization of the higher-dimensional linear thermoviscoelasticity*, J. Math. Pures Appl. **77**, 355–386 (1998)
- [40] Z. Liu and S. Zheng, *Exponential stability of the semigroup associated with a thermoelastic system*, Quart. Appl. Math. **LI**, 535–545 (1993)
- [41] W. J. Liu and E. Zuazua, *Decay rates for dissipative wave equations*, Ricerche de Matematica **48**, 61–75 (1999)
- [42] G. P. Menzala and E. Zuazua, *Explicit exponential decay rates for solutions of von Kármán's system of thermoelastic plates*, C. R. Acad. Sci Paris Sér. I Math. **324**, 49–54 (1997)
- [43] G. P. Menzala and E. Zuazua, *Energy decay rates for the Von Kármán system of thermoelastic plates*, Differential Integral Equations, to appear.
- [44] K. Narukawa, *Boundary value control of thermoelastic systems*, Hiroshima Math. J. **13**, 227–272 (1983)
- [45] L. A. F. Oliveira, *Exponential decay in thermoelasticity*, Commun. Appl. Anal. **1**, 113–118 (1997)
- [46] J. E. M. Rivera, *Decomposition of the displacement vector field and decay rates in linear thermoelasticity*, SIAM J. Math. Anal. **24**, 390–406 (1993)
- [47] J. E. M. Rivera, *Asymptotic behaviour in n -dimensional thermoelasticity*, Appl. Math. Lett. **10**, 47–53 (1997)
- [48] J. E. M. Rivera and M. L. Oliveira, *Stability in inhomogeneous and anisotropic thermoelasticity*, Boll. Un. Mat. Ital. A A (7) **11**, 115–127 (1997)
- [49] W. Rudin, *Real and Complex Analysis*, second edition, McGraw-Hill, Inc., New York, 1974
- [50] H. K. Wang and G. Chen, *Asymptotic behavior of solutions of the one-dimensional wave equation with a nonlinear boundary stabilizer*, SIAM J. Control Optim. **27**, 758–775 (1989)
- [51] E. Zuazua, *Uniform stabilization of the wave equation by nonlinear boundary feedback*, SIAM J. Control Optim. (2) **28**, 466–477 (1990)