ON THE ZEROS OF ENTIRE FUNCTIONS WITH A MAJORANT OF INFINITE ORDER

S. V. BYKOV AND F. A. SHAMOYAN

To our Teacher Viktor Petrovich Havin on the occasion of his 75th birthday

ABSTRACT. The class of entire functions f satisfying $\ln |f(z)| \leq C_f \lambda(|z|)$, $z \in \mathbb{C}$, where λ is a majorant mentioned in the title, admits a zero set $\{z_j\}$ such that $\{|z_j|\}$ is a uniqueness set for this class.

INTRODUCTION

Let $\mathbb{C}$ denote the complex plane, let $H(\mathbb{C})$ be the set of all entire functions, and let λ be a monotone increasing, positive function on $\mathbb{R}_+ = (0; +\infty)$; we put $\varphi(x) = \ln \lambda(x)$, $x \in \mathbb{R}_+$.

We introduce the following function classes:

- (1) $H(\lambda, +\infty) = \{f \in H(\mathbb{C}) : \ln |f(z)| \leq C_f \cdot \lambda(|z|), z \in \mathbb{C}\},$
- (2) $\tilde{H}(\lambda, +\infty) = \{f \in H(\mathbb{C}) : \ln |f(z)| \leq A_f \cdot \lambda(B_f \cdot |z|), z \in \mathbb{C}\}.$

Here and in what follows, A_f, B_f, C_f are arbitrary positive numbers depending on the function f only. If $\lambda \in C^1(\mathbb{R}_+)$ and $\alpha_\lambda = \lim_{x \rightarrow +\infty} x \frac{\lambda'(x)}{\lambda(x)} < +\infty$, then it is not hard to show that the classes $H(\lambda, +\infty)$ and $\tilde{H}(\lambda, +\infty)$ coincide. However, this is no longer true if $\alpha_\lambda = +\infty$, for example, in the case where

$$\lambda(t) = \underbrace{\exp \exp \cdots \exp}_n(t^\rho), \quad t \in \mathbb{R}_+, \quad \rho \in \mathbb{R}_+.$$

If $\lambda(t) = t^\rho$, $t \in \mathbb{R}_+$, $\rho \in \mathbb{R}_+$, then $H(\lambda, +\infty)$ coincides with the class of functions of finite order and normal type; we denote this class by $H(\rho, +\infty)$.

For $f \in H(\mathbb{C})$, we denote $Z_f \stackrel{\text{def}}{=} \{z \in \mathbb{C} : f(z) = 0\}$. The following property of the zeros of functions of class $H(\rho, +\infty)$ is well known: a sequence $\{z_k\}_{k=1}^{+\infty}$ can be represented as $Z = Z_f$ with $f \in H(\rho, +\infty)$, $f \not\equiv 0$, $\rho \notin \mathbb{N}$, $\rho > 0$, if and only if

- (3) $n(r) = \text{card}\{z_k : |z_k| \leq r\} \leq C_f \cdot r^\rho.$

For $\rho \in \mathbb{N}$, besides (3), the following Lindelöf condition arises (see [1]): there exists $M > 0$ such that

$$(4) \quad \left| \sum_{|z_k| \leq r} \frac{1}{z_k^\rho} \right| \leq M, \quad r \in \mathbb{R}_+.$$

By using (4), it is not difficult to construct a sequence $\{z_k\}_{k=1}^{+\infty} = Z_f$, $f \not\equiv 0$, $f \in H(\rho, +\infty)$, $\rho \in \mathbb{N}$, such that any $g \in H(\rho, +\infty)$ with $Z_g = \tilde{Z}_f$, where $\tilde{Z}_f = \{|z_k|\}_{k=1}^{+\infty}$,

2010 *Mathematics Subject Classification.* Primary 30D15.

Key words and phrases. Entire function, Lindelöf condition, zero set.

Supported by RFBR (grant no. 09-01-97517).

satisfies $g(z) = 0$, $z \in \mathbb{C}$; we mention $\{k^{\frac{1}{\rho}} e^{\frac{i\pi k}{\rho}}\}_{k=1}^{+\infty}$ as an example of such a sequence. In other words, the set $\tilde{Z}_f$ cannot coincide with Z_g for any $g \in H(\rho, +\infty)$, $\rho \in \mathbb{N}$, $g(z) \not\equiv 0$. Thus, the distribution of the arguments of $\{z_k\}_{k=1}^{+\infty}$ is also of importance for the representation $Z = Z_g$, rather than only the growth of the function $n(r)$.

Definition. We say that a set X of entire functions satisfies the *Lindelöf condition* if there exists $f \in X$, $f \not\equiv 0$, $Z_f = \{z_k\}_{k=1}^{+\infty}$, such that if $g \in X$, $Z_g \supseteq \tilde{Z}_f$, where $\tilde{Z}_f = \{|z_k|\}_{k=1}^{+\infty}$, then $g(z) = 0$ for all $z \in \mathbb{C}$.

The above remarks imply that the class $H(\rho, +\infty)$ with $\rho \in \mathbb{N}$ satisfies the Lindelöf condition, while for $\rho \notin \mathbb{N}$ it does not. The following question arises naturally: what happens for the other λ , e.g., for $\lambda(t) = \underbrace{\exp \cdots \exp}_{n}(t^\rho)$, $t, \rho \in \mathbb{R}_+$?

In this paper we show that if $\alpha_\lambda = +\infty$, then the class $H(\lambda, +\infty)$ satisfies the Lindelöf condition and the class $\tilde{H}(\lambda, +\infty)$ does not. In the case where $\alpha_\lambda = +\infty$ we obtain a description of the zero sets of functions in $\tilde{H}(\lambda, +\infty)$. This description is in terms of $\{|z_k|\}_{k=1}^{+\infty}$, i.e., in terms of $n(r)$. At the same time, Theorem 1 shows that the zero sets of functions of class $H(\lambda, +\infty)$ fail to admit a similar description; here the distribution of the arguments of the zeros becomes important.

There are many publications devoted to properties of the root sets for functions in $\tilde{H}(\lambda, +\infty)$; we mention [2, 3] and the references therein. The method applied in those papers is based on Fourier series, and the results do not even characterize the functions λ for which the zero sets can be described in terms of the counting function $n(r)$ only.

§1. FORMULATION OF THE MAIN RESULTS AND PROOF OF AUXILIARY STATEMENTS

In what follows, a monotone increasing positive function λ of class $C^{(1)}(\mathbb{R}_+)$ is called a *weight function* if $\alpha_\lambda = +\infty$.

Theorem 1. Let λ be a weight function of class $C^{(2)}(\mathbb{R}_+)$. Suppose that the function $\psi(x) = \ln \lambda(e^x)$ is convex on $\mathbb{R}_+$ and that

$$(5) \quad \frac{\psi''(x)}{(\psi'(x))^{2-\delta}} = O(1), \quad x \rightarrow +\infty,$$

for some $0 < \delta < 1$. Then the class $H(\lambda, +\infty)$ satisfies the Lindelöf condition.

Theorem 2. Let λ be a weight function of class $C^{(1)}(\mathbb{R}_+)$. Suppose that the function $\varphi(x) = \ln \lambda(x)$ is convex on $\mathbb{R}_+$. The following statements are equivalent:

- 1) a sequence of complex numbers $\{z_k\}_{k=1}^{+\infty}$ is representable in the form Z_f for some $f \in \tilde{H}(\lambda, +\infty)$, $f \not\equiv 0$;
- 2) there exists a positive number C such that

$$\sum_{k=1}^{+\infty} \exp(-\varphi(C|z_k|)) < +\infty.$$

In Theorem 2 it is assumed that $\lambda \in C^{(1)}(\mathbb{R}_+)$. However, the claim remains valid under the assumption that $\lambda \in C(\mathbb{R}_+)$. Indeed, it is easily seen that if $\lambda \in C(\mathbb{R}_+)$ and $\lambda_*(r) = \int_r^{er} \frac{\lambda(t)}{t} dt$, then $\lambda(r) \leq \lambda_*(r) \leq \lambda(er)$, while $\tilde{H}(\lambda, +\infty) = \tilde{H}(\lambda_*, +\infty)$.

The proof of the above theorems is based on several auxiliary statements.

Lemma 1 (see [4]). Let z_n be a sequence of complex numbers converging to infinity and satisfying

$$\sum_{n=1}^{+\infty} \frac{1}{|z_n|^{p_n+1}} < +\infty.$$

Then the Weierstrass product

$$B(z, z_n, p_n) = \prod_{n=1}^{+\infty} \left(1 - \frac{z}{z_n}\right) \cdot \exp\left(\frac{z}{z_n} + \frac{1}{2}\left(\frac{z}{z_n}\right)^2 + \cdots + \frac{1}{p_n}\left(\frac{z}{z_n}\right)^{p_n}\right)$$

converges in $\mathbb{C}$ and obeys the estimate

$$\ln |B(z, z_n, p_n)| \leq \sum_{n=1}^{+\infty} \left|\frac{z}{z_n}\right|^{p_n+1}, \quad z \in \mathbb{C}.$$

Lemma 2 (see [5]). Suppose $\lambda \in C^{(1)}(\mathbb{R}_+)$, $\lambda(x) > 0$, $x \in \mathbb{R}_+$, and $x \frac{\lambda'(x)}{\lambda(x)} \nearrow +\infty$ as $x \rightarrow +\infty$. Then there exists a function $f \in H(\lambda, +\infty)$ with the following properties:

$$\text{a) } \lim_{r \rightarrow +\infty} \frac{\ln M(r, f)}{\lambda(r)} = 1, \quad \text{b) } \lim_{r \rightarrow +\infty} \frac{T(r, f)}{\lambda(r)} = 1, \quad \text{c) } \lim_{r \rightarrow +\infty} \frac{N(r, f)}{\lambda(r)} = 1,$$

where, as usual, $M(r, f) = \max_{|z| \leq r} |f(z)|$,

$$T(r, f) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \ln^+ |f(re^{i\varphi})| d\varphi,$$

and $N(r, f)$ is the Nevanlinna counting function (see [1]):

$$N(r, f) = \int_0^r \frac{n(t) - n(0)}{t} dt,$$

$n(0)$ being the order of the zero of f at $z = 0$.

In the next lemma we estimate the number of real zeros of a function of class $H(\lambda, +\infty)$ on the segment $[0; r]$; this lemma will play an important role in the proof of Theorem 1.

Lemma 3. Let λ be as in Theorem 1. If $g \in H(\lambda, +\infty)$ and $n_R(r, g)$ is the number of positive (real) zeros of g on the segment $[0; r]$, then $n_R(g, r) \leq C_g \cdot \lambda(r)$, $C_g > 0$, $r \in \mathbb{R}_+$.

Proof. Let $\Omega = \{t + i\tau : |\tau| < s(t)\}$, $t \in \mathbb{R}_+$, where $s(t) = \frac{\pi}{\psi'(t)}$ and $\psi(t) = \ln \lambda(e^t)$; there is no loss of generality in assuming that $\psi'(0) > 3$. Obviously, if $f \in H(\lambda, +\infty)$, then $\ln |f(e^z)| \leq C_f \cdot \lambda(e^{\operatorname{Re} z})$, $z \in \mathbb{C}$.

Let $s(t) = \frac{\lambda(e^t)}{\pi \cdot \lambda'(e^t) \cdot e^t}$, $t \in \mathbb{R}_+$; we put $\Omega_+ = \{t + i\tau : t > 0, 0 < \tau < s(t)\}$ and $\Pi_+ = \{u + iv : 0 < v < \frac{\pi}{2}\}$. Let $Z_+(\zeta) = Z(t + i\tau)$ be a conformal mapping onto the strip Π_+ such that $Z_+(iS(0)) = \frac{i\pi}{2}$, $Z_+(0) = 0$, $\lim_{t \rightarrow +\infty} Z_+(t) = +\infty$. By the Riemann-Schwarz principle, the function $Z_+(\zeta)$ admits an analytic continuation Z to the domain

$$\Omega = \{\zeta = t + i\tau : |\tau| < s(|t|), t \in \mathbb{R}\},$$

Z maps Ω onto the strip $\Pi = \{u + iv : -\frac{\pi}{2} < v < \frac{\pi}{2}\}$, and the function $Z_+(t) = Z(t)$, $t \in \mathbb{R}_+$, is strictly monotone increasing on $[0; +\infty)$.

By the Warschawski theorem (the assumptions of it are fulfilled by (5)), we have

$$\begin{aligned} u = \operatorname{Re} Z(\sigma + i\tau) &= \frac{\pi}{2} \cdot \int_0^\sigma \frac{dt}{s(t)} + O(1) = \frac{\pi}{2} \cdot \frac{1}{\pi} \int_0^\sigma \frac{\lambda'(e^t) \cdot e^t}{\lambda(e^t)} dt + O(1) \\ &= \frac{1}{2} \cdot \int_0^\sigma d \ln \lambda(e^t) + O(1) = \frac{1}{2} \cdot \ln \lambda(e^\sigma) + O(1), \quad \sigma \rightarrow +\infty. \end{aligned}$$

Consequently,

$$(6) \quad \ln \lambda(e^\sigma) = 2 \cdot \operatorname{Re} Z(\sigma + i\tau) + O(1),$$

where $u + iv = Z(\sigma + i\tau) \in \Pi$.

We map the right half-plane $\mathbb{C}_+ = \{z \in \mathbb{C} : \operatorname{Re} z > 0\}$ conformally onto Π . Then, if $f \in H(\lambda, +\infty)$, then the function $F(z) = f(e^{Z^{-1}(\ln z)})$ is holomorphic in $\mathbb{C}_+$, and

$$(7) \quad \ln |F(z)| = \ln |f(e^{Z^{-1}(\ln z)})| \leq C_f \cdot \lambda(e^{\operatorname{Re} Z^{-1}(\ln z)}),$$

where Z^{-1} is the function inverse to Z .

Using (6), we see that if $\sigma + i\tau = Z^{-1}(\ln z)$, $Z(\sigma + i\tau) = \ln z = \ln |z| + i \cdot \arg z$, where $|\arg z| < \frac{\pi}{2}$, then $Z^{-1}(\ln z) = Z^{-1}(\ln |z| + i \cdot \arg z) = \sigma + i\tau \in \Omega$. Therefore, $\ln |z| = \operatorname{Re} Z(Z^{-1}(\ln z))$, but, by (6),

$$\frac{1}{2} \cdot \ln \lambda(e^\sigma) = \operatorname{Re} Z(\sigma + i\tau) + O(1) = \ln |z| + O(1) \quad (\sigma \rightarrow +\infty).$$

Thus, using (7), finally we obtain

$$\ln |F(z)| = \ln |f(e^{Z^{-1}(\ln z)})| \leq C_f \cdot \lambda(e^{\operatorname{Re} Z^{-1}(\ln z)}) \leq \widetilde{C}_f \cdot \lambda(e^\sigma) = C_f \cdot |z|^2.$$

Consequently, $\ln |F(z)| \leq C_f \cdot |z|^2$, $z \in \mathbb{C}_+$, and the function $F(z)$ is holomorphic in $\mathbb{C}_+$.

The results pertaining to the zeros of such functions (see [8]) show that the number $\tilde{n}_R(t)$ of real zeros of F on the segment $[1; t]$, $t > 1$, satisfies

$$(8) \quad \tilde{n}_R(t) \leq A_f \cdot t^2.$$

On the other hand, if $F(\rho_k) = 0$, $k = 1, 2, \dots$, then $f(e^{Z^{-1}(\ln \rho_k)}) = 0$, $k = 1, 2, \dots$. This means that $e^{Z^{-1}(\ln \rho_k)} = r_k$, where the r_k , $k = 1, 2, \dots$, are real zeros of f . Since $Z^{-1}(\ln \rho_k) = \ln r_k$, we have

$$(9) \quad \ln \rho_k = Z(\ln r_k).$$

Therefore, $\tilde{n}_R(t) = \operatorname{card}\{\rho_k : \rho_k \leq t\} \leq C_f \cdot t^2$. Since Z^{-1} grows monotonically on $\mathbb{R}_+$, estimate (8) implies that

$$n_R(t) = \operatorname{card}\{\rho_k : Z^{-1}(\ln \rho_k) \leq Z^{-1}(\ln t)\} \leq A_f \cdot t^2.$$

Now, by (9), we have

$$\tilde{n}_R(t) = \operatorname{card}\{r_k : \ln r_k \leq Z^{-1}(\ln t)\} \leq C_f \cdot t^2$$

or

$$\tilde{n}_R(t) = \operatorname{card}\{r_k : r_k \leq e^{Z^{-1}(\ln t)}\} \leq A_f \cdot t^2.$$

Now, we put $Z^{-1}(\ln t) = \rho$, $t \in \mathbb{R}_+$; then $t^2 = 2Z(\rho)$. Therefore,

$$\tilde{n}_R(\rho) = \operatorname{card}\{r_k : r_k \leq e^\rho\} \leq A_f \cdot e^{2Z(\rho)}, \quad \rho \in \mathbb{R}_+.$$

Recalling (6), we see that the last estimate implies that

$$\tilde{n}_R(\rho) = \operatorname{card}\{r_k : r_k \leq e^\rho\} \leq A_f \cdot \lambda(e^\rho) \cdot e^{O(1)} = C_f \cdot \lambda(e^\rho).$$

Putting $e^\rho = r$, $r > 0$, finally we get

$$n_R(r) \leq C_f \cdot \lambda(r),$$

which proves the lemma. $\square$

Lemma 4. *If $\varphi \in C^1(\mathbb{R}_+)$ is convex on $\mathbb{R}_+$ and $\varphi(0) = 0$, then*

$$C\varphi(x) \leq \varphi(Cx)$$

for any number $C \geq 1$.

Proof. Obviously, $\varphi(x) = \int_0^x \varphi'(u) du + \varphi(0) = \int_0^x \varphi'(u) du$, so that $C\varphi(x) = C \int_0^x \varphi'(u) du$. Changing the variables $Cu = t$, we obtain

$$C\varphi(x) = \int_0^{Cx} \varphi'\left(\frac{t}{C}\right) dt, \quad C \geq 1.$$

Since $\varphi'(x)$ is a monotone increasing function, we have $\varphi'\left(\frac{t}{C}\right) \leq \varphi'(t)$ for all t ; consequently, $\int_0^{Cx} \varphi'\left(\frac{t}{C}\right) dt \leq \int_0^{Cx} \varphi'(t) dt$, i.e., $C\varphi(x) \leq \varphi(Cx)$. $\square$

Lemma 5 (see [9]). *Let $\varphi(x)$ be a positive measurable function bounded on every segment $[a; b] \subset \mathbb{R}_+$. For each $K > 1$, there exists a sequence $x_n \rightarrow +\infty$ such that for all x with $x_n \leq x \leq x_n + \frac{1}{[\ln^+ \varphi(x_n)]^K} + \frac{1}{\varphi(x_n)}$ we have*

$$\varphi(x) \leq K \cdot \varphi(x_n).$$

Lemma 6. *Let $Z_f = \{z_k\}_{k=1}^{+\infty}$, $f \in \tilde{H}(\lambda, +\infty)$, $f \not\equiv 0$. Then $n(r) \leq \frac{A_f}{\ln 2} \cdot \lambda(2B_f \cdot r)$, where A_f and B_f are as in (2).*

Proof. Suppose $f \in \tilde{H}(\lambda, +\infty)$, $f(z_k) = 0$, $k = 1, 2, \dots$, $f \not\equiv 0$. Without loss of generality, we may assume that $f(0) = 1$. The Jensen identity yields

$$\begin{aligned} \int_0^r \frac{n(t)}{t} dt &= \frac{1}{2\pi} \int_0^{2\pi} \ln |f(re^{i\varphi})| d\varphi \leq \frac{1}{2\pi} \int_0^{2\pi} A_f \lambda(B_f r) d\varphi \\ &= \frac{1}{2\pi} \cdot 2\pi \cdot A_f \lambda(B_f r) = A_f \cdot \lambda(B_f r), \end{aligned}$$

where $0 < r < +\infty$. Since $n(r)$ grows monotonically, we deduce that

$$n(r) \cdot \ln 2 \leq \int_r^{2r} \frac{n(t)}{t} dt \leq A_f \cdot \lambda(2B_f \cdot r),$$

whence $n(r) \leq \frac{A_f}{\ln 2} \cdot \lambda(2B_f \cdot r)$, as required. $\square$

§2. PROOF OF THE MAIN RESULTS

First, we prove Theorem 1.

Let f be the function of class $H(\lambda, +\infty)$ constructed in Lemma 2, let $\{z_k\}_{k=1}^{+\infty}$ be the set of zeros of f , let $n(r) = \{\text{card } z_k : |z_k| < r\}$, and let

$$N(r) = \int_0^r \frac{n(t)}{t} dt.$$

Without loss of generality, we assume that $f(0) = 1$. By Lemma 2, we have $N(r) \geq C_0 \cdot \lambda(r)$, $r \geq 1$, for some $C_0 > 0$. Suppose that $g \in H(\lambda, +\infty)$, $g(|z_k|) = 0$, $k = 1, 2, \dots$, and $g(z) \not\equiv 0$. Then, by Lemma 3, we have $n_R(r) = n(r)$, where n_R is the set of all positive roots of g on the segment $[0; r]$, and $n_R(r) = n(r) \leq C_1 \cdot \lambda(r)$, where $C_1 > 0$, $r \in \mathbb{R}_+$. We put $\varphi(r) = \frac{N'_+(r)}{N(r)}$, $r \in \mathbb{R}_+$, where $N'_+(r)$ is the right derivative of the function N . Let $\{x_n\}_{n=1}^{+\infty}$ be the sequence built in Lemma 5 starting with φ . Applying that lemma and the identity

$$\ln \frac{N(x'_n)}{N(x_n)} = \int_{x_n}^{x'_n} \frac{N'_+(t)}{N(t)} dt,$$

where $x'_n = x_n + \frac{1}{\varphi(x_n)}$, we obtain the inequality

$$\ln \frac{N(x'_n)}{N(x_n)} \leq K \cdot \varphi(x_n) \cdot (x'_n - x_n) = K \cdot \varphi(x_n) \cdot \frac{1}{\varphi(x_n)} = K,$$

i.e., $N(x'_n) \leq e^K \cdot N(x_n)$. Now, since λ is convex, we get

$$\begin{aligned} \lambda'(x_n) &\leq \frac{1}{x'_n - x_n} \cdot \int_{x_n}^{x'_n} \lambda'(t) dt \leq \frac{1}{x'_n - x_n} \cdot [\lambda(x'_n) - \lambda(x_n)] \\ &\leq \frac{\lambda(x'_n)}{x'_n - x_n} \leq C \cdot \varphi(x_n) \cdot N(x'_n). \end{aligned}$$

Then $\lambda'(x_n) \leq C \cdot e^K \cdot N(x_n) \cdot \varphi(x_n)$, i.e., $\lambda'(x_n) \leq C \cdot e^K \cdot N(x_n) \cdot \frac{N'_+(x_n)}{N(x_n)}$, whence $\lambda'(x_n) \leq C \cdot e^K \cdot N'_+(x_n)$ (note that $x_n \rightarrow +\infty$ as $n \rightarrow +\infty$).

Therefore, $\lambda'(x_n) \leq C \cdot e^K \cdot \frac{n(x_n)}{x_n} \leq C \cdot e^K \cdot \frac{\lambda(x_n)}{x_n}$, so that $\frac{\lambda'(x_n)}{\lambda(x_n)} \cdot x_n \leq C \cdot e^K < +\infty$, which contradicts the assumptions of the theorem. This proves Theorem 1.

Now we start the proof of Theorem 2. We check the implication (1) $\Rightarrow$ (2).

Suppose $f \in \tilde{H}(\lambda, +\infty)$, $f(z_k) = 0$, $k = 1, 2, \dots$, $f \not\equiv 0$. Without loss of generality, we may assume that $\lambda(0) = 1$, i.e., $\varphi(0) = 0$. Lemma 1 shows that if $Z = Z_f$, $Z = \{z_k\}_{k=1}^{+\infty}$ (we assume that $0 < |z_1| \leq |z_2| \leq \dots \leq |z_k| \leq \dots$), then $n(r) \leq A \cdot \lambda(B \cdot r)$, i.e.,

$$(10) \quad n(r) \leq A \cdot \exp \varphi(B \cdot r), \quad r \in \mathbb{R}_+, \quad A \in \mathbb{R}_+, \quad B \in \mathbb{R}_+.$$

We fix $C > 0$. For $R > 0$ we have

$$I_R = \sum_{|z_k| \leq R} \exp(-\varphi(C|z_k|)) = \int_0^R \exp(-\varphi(Ct)) dn(t).$$

Integration by parts yields the formula

$$\begin{aligned} (11) \quad I_R &= n(t) \cdot \exp(-\varphi(Ct)) \Big|_0^R + \int_0^R n(t) \cdot \exp(-\varphi(Ct)) \cdot \varphi'(Ct) dt \\ &= n(R) \cdot \exp(-\varphi(CR)) + C \int_0^R n(t) \cdot \exp(-\varphi(Ct)) \cdot \varphi'(Ct) dt. \end{aligned}$$

Using (10) and choosing $C > B_f$, we see that $I_R \leq A_f \cdot \exp(\varphi(B_f r) - \varphi(Cr)) + C \int_0^R n(t) \cdot \exp(-\varphi(Ct)) \cdot \varphi'(Ct) dt$. Since $\varphi(t)$ is a monotone increasing function on $\mathbb{R}_+$, we have

$$I_R \leq A_f + C \int_0^R n(t) \cdot \exp(-\varphi(Ct)) \cdot \varphi'(Ct) dt \leq A_f + C \int_0^R \exp(\varphi(B_f t) - \varphi(Ct)) \cdot \varphi'(Ct) dt.$$

We take C so large that $\varphi(B_f x) \leq \frac{\varphi(Cx)}{2}$, $x \in \mathbb{R}_+$; then

$$\begin{aligned} I_R &\leq A_f + A_0 \cdot \int_0^R \exp\left(\frac{-\varphi(Cx)}{2}\right) \cdot \varphi'(Cx) dx \\ &= A_f + \frac{A_f}{2} \cdot \int_0^R \exp\left(\frac{-\varphi(Cx)}{2}\right) \cdot d\left(\frac{\varphi(Cx)}{2}\right) \leq \exp\left(\frac{-\varphi(CR)}{2}\right) \leq \text{const}, \end{aligned}$$

i.e., $I_R \leq \text{const}$ for all $R \in \mathbb{R}_+$. Therefore, $\sum_{k=1}^{+\infty} \exp(-\varphi(C|z_k|)) < +\infty$, which establishes the implication (1) $\Rightarrow$ (2).

Now we prove the reverse implication (2) $\Rightarrow$ (1). Let $\{z_k\}_{k=1}^{+\infty}$ be a sequence in $\mathbb{C}$ satisfying (10) and such that $1 < |z_1| \leq |z_2| \leq \dots \leq |z_k| \leq \dots$. There is no loss of generality in assuming that $C = 1$. We construct a function $f \in \tilde{H}(\lambda, +\infty)$, $f \not\equiv 0$, such that $f(z_k) = 0$, $k = 1, 2, \dots$, and $f(z) \not\equiv 0$ for $z \neq z_k$. Put $p_n = \left[\frac{\varphi(|z_n|)}{\ln 2}\right]$, $n = 1, 2, \dots$, where $[a]$ stands for the integral part of a .

We prove that the infinite product

$$B(z, z_n) = \prod_{n=1}^{+\infty} \left(1 - \frac{z}{z_n}\right) \cdot \exp\left(\frac{z}{z_n} + \frac{1}{2}\left(\frac{z}{z_n}\right)^2 + \cdots + \frac{1}{p_n}\left(\frac{z}{z_n}\right)^{p_n}\right)$$

converges and does not vanish identically, and $B \in \tilde{H}(\lambda, +\infty)$. The estimate of Lemma 1 shows that

$$\ln |B(z, z_n)| \leq \sum_{k=1}^{+\infty} \left| \frac{z}{z_n} \right|^{p_n+1}, \quad z \in \mathbb{C}.$$

To estimate the sum obtained, we put $m_n(z) = \left| \frac{z}{z_n} \right|$, $z \in \mathbb{C}$, $n = 1, 2, \dots$. Then

$$I = \sum_{k=1}^{+\infty} \left| \frac{z}{z_n} \right|^{p_n+1} = \sum_{m_n(z) \leq \frac{1}{2}} \left| \frac{z}{z_n} \right|^{p_n+1} + \sum_{m_n(z) > \frac{1}{2}} \left| \frac{z}{z_n} \right|^{p_n+1} = I_1 + I_2.$$

We bound each sum separately. We have

$$\begin{aligned} I_1 &= \sum_{m_n(z) \leq \frac{1}{2}} \left| \frac{z}{z_n} \right|^{p_n+1} \leq \sum_{m_n(z) \leq \frac{1}{2}} \left(\frac{1}{2}\right)^{p_n+1} = \sum_{m_n(z) \leq \frac{1}{2}} \exp(-(p_n+1) \cdot \ln 2) \\ &< e \cdot \sum_{m_n(z) \leq \frac{1}{2}} \exp(-\varphi(|z_n|)) \leq C, \end{aligned}$$

because $\sum_{k=1}^{+\infty} \exp(-\varphi(|z_k|)) < +\infty$.

Passing to the sum I_2 , we have

$$\begin{aligned} I_2 &= \sum_{m_n(z) > \frac{1}{2}} \left| \frac{z}{z_n} \right|^{p_n+1} = \sum_{m_n(z) > \frac{1}{2}} \exp\left((p_n+1) \cdot \ln \left| \frac{z}{z_n} \right|\right) \\ &= \sum_{\frac{1}{2} < m_n(z) < 1} \exp\left((p_n+1) \cdot \ln \left| \frac{z}{z_n} \right|\right) + \sum_{m_n(z) \geq 1} \exp\left((p_n+1) \cdot \ln \left| \frac{z}{z_n} \right|\right) \\ &= I_2^1 + I_2^2. \end{aligned}$$

Since $\ln \left| \frac{z}{z_n} \right| \geq 0$ for $0 < |z_n| \leq |z|$, and $p_n+1 \leq \frac{\varphi(|z_n|)}{\ln 2} + 1 \leq 2\varphi(|z_n|) + 1$, $n = 1, 2, \dots$, we get

$$\begin{aligned} I_2^2 &\leq \sum_{|z_n| \leq |z|} \exp\left((2\varphi(|z_n|) + 1) \cdot \ln \left| \frac{z}{z_n} \right|\right) \leq |z| \cdot \sum_{|z_n| \leq |z|} \exp\left(2\varphi(|z_n|) \cdot \ln \left| \frac{z}{z_n} \right|\right) \\ &\leq |z| \cdot \int_1^{|z|} \exp\left(2\varphi(t) \cdot \ln \frac{|z|}{t}\right) dn(t). \end{aligned}$$

Now we note that the convergence of the series $\sum_{n=1}^{+\infty} \exp(-\varphi(|z_n|)) \leq C < +\infty$ implies that $\sum_{|z_n| \leq R} \exp(-\varphi(|z_n|)) \leq C$. Therefore, $\exp(-\varphi(R)) \cdot n(R) \leq C$. This results in the estimate

$$(12) \quad n(R) \leq C \cdot \exp(\varphi(R)),$$

valid for all $R \in \mathbb{R}_+$. Using (12) and integrating by parts, we obtain

$$I_2^2 \leq |z| \cdot \left(n(|z|) - \int_1^{|z|} \left(\exp(2\varphi(t)) \cdot \ln \frac{|z|}{t} \right) \cdot n(t) \cdot \left(2\varphi'(t) \cdot \ln \frac{|z|}{t} - \frac{2\varphi(t)}{t} \right) dt \right).$$

Recall that φ is a monotone increasing function of class $C^{(1)}(\mathbb{R}_+)$, so that $\varphi'(t) \geq 0$, $t \in \mathbb{R}_+$, and $\ln \frac{|z|}{t} \geq 0$, $1 \leq t \leq |z|$. We continue the last-written estimate:

$$(13) \quad \begin{aligned} I_2^2 &\leq n(|z|) \cdot |z| + 2|z| \cdot \int_1^{|z|} \frac{n(t) \cdot \varphi(t)}{t} \cdot \exp\left(2\varphi(t) \cdot \ln \frac{|z|}{t}\right) dt \\ &\leq 3 \cdot n(|z|) \cdot |z| \cdot \varphi(|z|) \cdot \int_1^{|z|} \exp\left(2\varphi(t) \cdot \ln \frac{|z|}{t}\right) dt. \end{aligned}$$

We denote the last integral in (13) by $\tilde{I}_2^2$. To estimate it, we put $\ln \frac{|z|}{t} = u$, obtaining

$$(14) \quad \tilde{I}_2^2 \leq |z| \cdot \int_0^{\ln |z|} \exp(2\varphi(|z|e^{-u})u) \cdot e^{-u} du \leq |z| \cdot \int_0^{\ln |z|} \exp(2\varphi(|z|e^{-u})u) du.$$

Since φ is convex, we can bound the expression $\varphi(|z|e^{-u})u$ for $u \geq 1$ as follows:

$$\begin{aligned} \varphi(|z|e^{-u})u &= u \int_0^{|z|e^{-u}} \varphi'(t) dt = \int_0^{u|z|e^{-u}} \varphi'\left(\frac{x}{u}\right) dx \\ &\leq \int_0^{|z|} \varphi'\left(\frac{x}{u}\right) dx \leq \int_0^{|z|} \varphi'(x) dx = \varphi(|z|). \end{aligned}$$

Thus,

$$(15) \quad \varphi(|z|e^{-u})u \leq \varphi(|z|)$$

for all $u \geq 1$.

Obviously, if $0 \leq u \leq 1$, then

$$(16) \quad \varphi(|z|e^{-u})u \leq \varphi(|z|).$$

Consequently, $\varphi(|z|e^{-u})u \leq \varphi(|z|)$ for all $u \in \mathbb{R}_+$. Therefore, estimate (14) implies that

$$(17) \quad \tilde{I}_2^2 \leq |z| \cdot \exp(2\varphi(|z|)) \cdot \ln |z|.$$

A combination of (14) and (15) yields

$$I_2^2 \leq 3 \cdot n(|z|) \cdot \varphi(|z|) \cdot |z|^2 \cdot \ln |z| \cdot \exp(2\varphi(|z|)).$$

Since φ is convex and $\varphi'(x)x \rightarrow +\infty$ as $x \rightarrow +\infty$, we can use Lemma 6 to obtain the final estimate:

$$(18) \quad I_2^2 \leq C_1 \cdot \exp(\varphi(C_2 \cdot |z|)).$$

The quantity I_2^1 can be estimated as follows:

$$I_2^1 = \sum_{\frac{1}{2} < m_n(z) < 1} \exp\left((p_n + 1) \cdot \ln \left|\frac{z}{z_n}\right|\right) \leq \sum_{m_n(z) > \frac{1}{2}} 1 \leq \sum_{|z_n| < 2|z|} 1 \leq n(2|z|).$$

Using (12), we conclude that

$$(19) \quad n(2|z|) \leq C \cdot \exp(\varphi(2|z|)).$$

To complete the proof of Theorem 2, now it suffices to combine estimates (18) and (19).

REFERENCES

- [1] B. Ya. Levin, *Distribution of zeros of entire functions*, Gostekhizdat, Moscow, 1956; English transl., Transl. Math. Monogr., vol. 5, Amer. Math. Soc., Providence, RI, 1980. MR0589888 (81k:30011)
- [2] L. A. Rubel, *A survey of a Fourier series method for meromorphic functions*, L'analyse harmonique dans le domaine complexe (Actes Table Ronde Internat., Centre Nat. Recherche Sci., Montpellier, 1972), Lecture Notes in Math., vol. 336, Springer, Berlin, 1973, pp. 51–62. MR0390219 (52:11045)
- [3] L. A. Rubel and B. A. Taylor, *A Fourier series method for meromorphic and entire functions*, Bull. Soc. Math. France **96** (1968), 53–96. MR0243075 (39:4399); MR0265600 (42:509)
- [4] L. De Branges, *Hilbert spaces of entire functions*, Prentice-Hall, Inc., Englewood Cliffs, N.J., 1968. MR0229011 (37:4590)
- [5] J. Clunie and T. Kovari, *On integral functions having prescribed asymptotic growth*. II, Canad. J. Math. **20** (1968), 7–20. MR0227411 (37:2995)
- [6] B. Rodin and S. E. Warschawski, *Extremal length and the boundary behavior of conformal mappings*, Ann. Acad. Sci. Fenn. Ser. A I Math. **2** (1976), 467–500. MR0466516 (57:6394)
- [7] S. E. Warschawski, *On conformal mapping of infinite strips*, Trans. Amer. Math. Soc. **51** (1942), 280–335. MR0006583 (4:9b)
- [8] N. V. Govorov, *The Riemann boundary value problem with an infinite index*, Nauka, Moscow, 1986, pp. 29–41; English transl., Oper. Theory Adv. Appl., vol. 67, Birkhäuser Verlag, Basel, 1994. MR0887532 (88k:30001); MR1269780 (95d:30001)
- [9] W. K. Hayman and F. M. Stewart, *Real inequalities with applications to function theory*, Proc. Cambridge Philos. Soc. **50** (1954), 250–260. MR0061638 (15:857g)

BRYANSK STATE UNIVERSITY, 14 BEZHITSKAYA STREET, BRYANSK 241036, RUSSIA
E-mail address: `b.serecha@mail.ru`

BRYANSK STATE UNIVERSITY, 14 BEZHITSKAYA STREET, BRYANSK 241036, RUSSIA
E-mail address: `shamoyanfa@yandex.ru`

Received 15/APR/2008

Translated by A. PLOTKIN