FAMILIES OF FRACTIONAL CAUCHY TRANSFORMS IN THE BALL

E. S. DUBTSOV

Dedicated to Victor Petrovich Havin on the occasion of his 75th birthday

ABSTRACT. Let B_n denote the unit ball in $\mathbb{C}^n$, $n \geq 1$. Given $\alpha > 0$, let $\mathcal{K}_\alpha(n)$ denote the class of functions defined for $z \in B_n$ by integrating the kernel $(1 - \langle z, \zeta \rangle)^{-\alpha}$ against a complex-valued Borel measure on the sphere $\{\zeta \in \mathbb{C}^n : |\zeta| = 1\}$. The families $\mathcal{K}_\alpha(1)$ of fractional Cauchy transforms have been investigated intensively by several authors. In the paper, various properties of $\mathcal{K}_\alpha(n)$, $n \geq 2$, are studied. In particular, relations between $\mathcal{K}_\alpha(n)$ and other spaces of holomorphic functions in the ball are obtained. Also, pointwise multipliers for the spaces $\mathcal{K}_\alpha(n)$ are investigated.

§1. INTRODUCTION

For $n \geq 1$, put $B_n = \{z \in \mathbb{C}^n : |z| < 1\}$. Let $M(n)$ denote the space of complex-valued Borel measures on the sphere ∂B_n .

1.1. Fractional Cauchy transforms. Let $\alpha > 0$. Given a measure $\mu \in M(n)$, its fractional Cauchy transform of order α is defined by the identity

$$K_\alpha[\mu](z) = \int_{\partial B_n} \frac{1}{(1 - \langle z, \zeta \rangle)^\alpha} d\mu(\zeta), \quad z \in B_n.$$

Here and in what follows we use the principal branch of the logarithm. Put

$$\mathcal{K}_\alpha(n) = \{K_\alpha[\mu] : \mu \in M(n)\}.$$

1.2. Multipliers. Let $\mathcal{H}ol(B_n)$ denote the space of holomorphic functions in the ball B_n . A function $g \in \mathcal{H}ol(B_n)$ is called a multiplier for the family $\mathcal{K}_\alpha(n)$, $\alpha > 0$, if $fg \in \mathcal{K}_\alpha(n)$ for all $f \in \mathcal{K}_\alpha(n)$. Let $\mathfrak{M}_\alpha(n)$ denote the set of all multipliers for $\mathcal{K}_\alpha(n)$.

1.3. Families $\mathcal{K}_\alpha(n)$ and $\mathfrak{M}_\alpha(n)$ as Banach spaces. For $f \in \mathcal{K}_\alpha(n)$, put

$$\|f\|_{\mathcal{K}_\alpha(n)} = \inf \{\|\mu\|_{M(n)} : f = K_\alpha[\mu]\}.$$

Standard arguments show that the above infimum is attained, and that $\mathcal{K}_\alpha(n)$ with the norm $\|\cdot\|_{\mathcal{K}_\alpha(n)}$ is a Banach space. Next, assume that $f = K_\alpha[\rho]$, where ρ is a positive measure. For $\mu \in M(n)$, let $f = K_\alpha[\mu]$; then $\|\rho\| = K_\alpha[\rho](0) = K_\alpha[\mu](0) \leq \|\mu\|$. Therefore, $\|f\|_{\mathcal{K}_\alpha(n)} = \|\rho\|_{M(n)}$.

For $g \in \mathfrak{M}_\alpha(n)$, put

$$\|g\|_{\mathfrak{M}_\alpha(n)} = \sup \{\|fg\|_{\mathcal{K}_\alpha(n)} : \|f\|_{\mathcal{K}_\alpha(n)} \leq 1\}.$$

The closed graph theorem guarantees that $\|g\|_{\mathfrak{M}_\alpha(n)} < \infty$. The space $\mathfrak{M}_\alpha(n)$ with the norm $\|\cdot\|_{\mathfrak{M}_\alpha(n)}$ is a Banach space. Also, note that $\mathfrak{M}_\alpha(n)$ is a Banach algebra.

2010 *Mathematics Subject Classification.* Primary 32A26, 32A37.

Key words and phrases. Fractional Cauchy transform, Bergman–Sobolev space, pointwise multiplier. Supported by RFBR (grant no. 08-01-00358-a) and by the Russian Science Support Foundation.

1.4. Historical remarks. The families $\mathcal{K}_\alpha(n)$ generalize the classical family $\mathcal{K}_n(n)$ of Cauchy integrals. Investigations of the family $\mathcal{K}_1(1)$ as a Banach space were initiated by Havin in [14, 15]. Further results and references are presented in the monograph [9]. The spaces $\mathcal{K}_\alpha(1)$, $\alpha > 0$, were introduced by MacGregor [20]. Various properties of the families $\mathcal{K}_\alpha(1)$ are collected in [17]. In the present paper we focus on the study of the families $\mathcal{K}_\alpha(n)$ with $n \geq 2$. Certain properties of the spaces $\mathcal{K}_n(n)$, $n \in \mathbb{N}$, were proved in the survey [3]. To the best knowledge of the author, for $n \geq 2$ the families $\mathcal{K}_\alpha(n)$ have not been investigated systematically.

1.5. Related problems. Let σ_n denote the normalized Lebesgue measure on the sphere ∂B_n . For $1 \leq p \leq \infty$ and $0 < \alpha < n$, put

$$\mathcal{K}_\alpha^p(n) = \{K_\alpha[f\sigma_n] : f \in L^p(\sigma_n)\}.$$

The boundary behavior of functions belonging to $\mathcal{K}_\alpha^p(1)$ was investigated in [21], where the families $\mathcal{K}_\alpha^p(1)$ were called Dirichlet type spaces. The families $\mathcal{K}_\alpha^p(n)$, $n \in \mathbb{N}$, were investigated in [26]; see also [1, 19, 8].

1.6. Organization of the paper. Embedding properties for $\mathcal{K}_\alpha(n)$ and $\mathfrak{M}_\alpha(n)$ are studied in §2. Relations between $\mathcal{K}_\alpha(n)$ and the classical spaces of holomorphic functions are obtained in §§3 and 5. Also, in §§4 and 5 we study differentiation operators on the spaces $\mathcal{K}_\alpha(n)$. The boundary behavior of the fractional Cauchy transforms is investigated in §6. Finally, some results about the multiplier spaces $\mathfrak{M}_\alpha(n)$ are obtained in §7.

§2. EMBEDDING PROPERTIES

2.1. Embedding properties for families of fractional Cauchy transforms. Below we shall use the following lemma.

Lemma 2.1 (see [7, Lemma 1]). *Assume that $u, v \in B_1$, $\alpha > 0$, $\beta > 0$. Then*

$$(1-u)^{-\alpha}(1-v)^{-\beta} = \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 t^{\alpha-1}(1-t)^{\beta-1}[1-(tu+(1-t)v)]^{-\alpha-\beta} dt.$$

Let $\mathcal{P}(n)$ denote the set of all probability measures on the sphere ∂B_n , $n \in \mathbb{N}$. For $n = 1$, the following result was obtained in [7].

Proposition 2.2. *Put $\mathcal{F}_\alpha(n) = \{K_\alpha[\mu] : \mu \in \mathcal{P}(n)\}$, $n \in \mathbb{N}$.*

- (i) *If $\alpha > 0$ and $\beta > 0$, then $\mathcal{F}_\alpha(n) \cdot \mathcal{F}_\beta(n) \subset \mathcal{F}_{\alpha+\beta}(n)$.*
- (ii) *If $0 < \alpha < \beta$, then $\mathcal{F}_\alpha(n) \subset \mathcal{F}_\beta(n)$.*

Proof. First, we show how to deduce statement (i) from the following property:

$$(2.1) \quad (1 - \langle z, \zeta \rangle)^{-\alpha} (1 - \langle z, \xi \rangle)^{-\beta} \in \mathcal{F}_{\alpha+\beta}(n), \quad z \in B_n,$$

for any fixed points $\zeta, \xi \in \partial B_n$. So, let $f \in \mathcal{F}_\alpha(n)$, and let $g \in \mathcal{F}_\beta(n)$. Then, by the definition,

$$f(z)g(z) = \int_{\partial B_n \times \partial B_n} (1 - \langle z, \zeta \rangle)^{-\alpha} (1 - \langle z, \xi \rangle)^{-\beta} d\rho(\zeta, \xi), \quad z \in B_n,$$

where ρ is a probability measure on the product $\partial B_n \times \partial B_n$. We approximate the measure ρ in the weak* topology by probability measures ρ_j , where every ρ_j is a finite sum of atomic charges. The set $\mathcal{F}_{\alpha+\beta}(n)$ is convex; therefore, on the one hand, we have

$$\int_{\partial B_n \times \partial B_n} (1 - \langle z, \zeta \rangle)^{-\alpha} (1 - \langle z, \xi \rangle)^{-\beta} d\rho_j(\zeta, \xi) \in \mathcal{F}_{\alpha+\beta}(n).$$

On the other hand, without loss of generality we may assume that the convergence

$$\int_{\partial B_n \times \partial B_n} (1 - \langle z, \zeta \rangle)^{-\alpha} (1 - \langle z, \xi \rangle)^{-\beta} d\rho_j(\zeta, \xi) \rightarrow f(z)g(z)$$

as $j \rightarrow \infty$ is uniform on the compact subsets of the ball B_n . It remains to observe that the set $\mathcal{F}_{\alpha+\beta}(n)$ is closed in the topology of uniform convergence on compact subsets of the ball.

Now, we prove property (2.1). Lemma 2.1 guarantees that

$$(2.2) \quad (1 - \langle z, \zeta \rangle)^{-\alpha} (1 - \langle z, \xi \rangle)^{-\beta} = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 t^{\alpha-1} (1-t)^{\beta-1} [1 - \langle z, w(t) \rangle]^{-\alpha-\beta} dt,$$

where $z \in B_n$ and $w(t) = t\zeta + (1-t)\xi$. Since

$$\Gamma(\alpha + \beta)\Gamma(\alpha)^{-1}\Gamma(\beta)^{-1}t^{\alpha-1}(1-t)^{\beta-1}dt$$

is a probability measure on the interval $[0, 1]$, it suffices to show that

$$(2.3) \quad [1 - \langle z, w(t) \rangle]^{-\alpha-\beta} \in \mathcal{F}_{\alpha+\beta}(n), \quad z \in B_n,$$

for every fixed parameter $t \in [0, 1]$. If $t = 0$ or $t = 1$, then property (2.3) holds true by definition. So, assume that $0 < t < 1$. Then $w(t) \in B_n$. Next, note that $[1 - \langle z, \cdot \rangle]^{-\alpha-\beta}$ is a harmonic function in the closed ball $\bar{B}_n$. Hence,

$$(2.4) \quad \frac{1}{[1 - \langle z, w \rangle]^{\alpha+\beta}} = \int_{\partial B_n} \frac{1 - |w|^2}{|w - \eta|^{2n}} \frac{1}{[1 - \langle z, \eta \rangle]^{\alpha+\beta}} d\sigma_n(\eta).$$

If a point $w \in B_n$ is fixed, then $(1 - |w|^2)|w - \eta|^{-2n} d\sigma_n(\eta)$ is a probability measure. Thus, to prove (2.3), it suffices to put $w = w(t)$, $0 < t < 1$. The proof of property (i) is finished.

Let $\beta > \alpha > 0$. Recall that the ball algebra $A(B_n)$ is defined by

$$A(B_n) = \mathcal{H}ol(B_n) \cap C(\bar{B}_n).$$

If $z \in B_n$, then $(1 - \langle z, \cdot \rangle)^{-\beta+\alpha} \in \bar{A}(B_n)$. Hence,

$$1 = \int_{\partial B_n} \frac{1}{(1 - \langle z, \zeta \rangle)^{\beta-\alpha}} d\sigma_n(\zeta), \quad z \in B_n.$$

In other words, $1 \in \mathcal{F}_{\beta-\alpha}(n)$; thus, (i) implies (ii). $\square$

Corollary 2.3. *Let $n \in \mathbb{N}$.*

- (i) *If $\alpha > 0$ and $\beta > 0$, then $\mathcal{K}_\alpha(n) \cdot \mathcal{K}_\beta(n) \subset \mathcal{K}_{\alpha+\beta}(n)$.*
- (ii) *If $0 < \alpha < \beta$, then $\mathcal{K}_\alpha(n) \subset \mathcal{K}_\beta(n)$.*

Proof. Let $f = K_\alpha[\mu]$ and let $g = K_\beta[\rho]$, where $\mu, \rho \in M(n)$ and $\alpha, \beta > 0$. Considering the Jordan decompositions of the measures μ, ρ and applying part (i) of Proposition 2.2, we obtain properties (i) and (ii). $\square$

2.2. Embedding properties for the multiplier families. The proof of the following lemma will be omitted, because it practically coincides with that given in [29] for the case where $n = \alpha = 1$ (see also [16], where the case of $n = 1$, $\alpha > 0$ was considered).

Lemma 2.4. *Assume that $n \in \mathbb{N}$ and $\alpha > 0$. Then the following properties are equivalent:*

- (i) $g \in \mathfrak{M}_\alpha(n)$;
- (ii) $g(z)(1 - \langle z, \zeta \rangle)^{-\alpha} \in \mathcal{K}_\alpha(n)$ for all $\zeta \in \partial B_n$ and

$$(2.5) \quad \sup \left\{ \left\| \frac{g(z)}{(1 - \langle z, \zeta \rangle)^\alpha} \right\|_{\mathcal{K}_\alpha(n)} : \zeta \in \partial B_n \right\} < \infty.$$

For $n = 1$, the following proposition was proved in [16].

Proposition 2.5. *If $n \in \mathbb{N}$ and $0 < \alpha < \beta$, then $\mathfrak{M}_\alpha(n) \subset \mathfrak{M}_\beta(n)$.*

Proof. Let $g \in \mathfrak{M}_\alpha(n)$. By Lemma 2.4, there exists a constant $C > 0$ and there exist measures $\mu_\zeta \in M(n)$, $\zeta \in \partial B_n$, such that $\|\mu_\zeta\| \leq C$ and

$$\frac{g(z)}{(1 - \langle z, \zeta \rangle)^\alpha} = \int_{\partial B_n} \frac{1}{(1 - \langle z, \xi \rangle)^\alpha} d\mu_\zeta(\xi), \quad z \in B_n.$$

Hence,

$$\frac{g(z)}{(1 - \langle z, \zeta \rangle)^\beta} = \int_{\partial B_n} \frac{1}{(1 - \langle z, \xi \rangle)^\alpha} \frac{1}{(1 - \langle z, \zeta \rangle)^{\beta-\alpha}} d\mu_\zeta(\xi).$$

By (2.2) and (2.4), for every pair of points $\zeta, \xi \in \partial B_n$, there exists a probability measure $\rho_{\zeta, \xi} \in M(n)$ such that

$$\frac{1}{(1 - \langle z, \xi \rangle)^\alpha} \frac{1}{(1 - \langle z, \zeta \rangle)^{\beta-\alpha}} = \int_{\partial B_n} \frac{1}{(1 - \langle z, \eta \rangle)^\beta} d\rho_{\zeta, \xi}(\eta).$$

Therefore,

$$\frac{g(z)}{(1 - \langle z, \zeta \rangle)^\beta} = \int_{\partial B_n} \int_{\partial B_n} \frac{1}{(1 - \langle z, \eta \rangle)^\beta} d\rho_{\zeta, \xi}(\eta) d\mu_\zeta(\xi).$$

Fix a point $\zeta \in \partial B_n$. We approximate the measure $\mu = \mu_\zeta$ in the weak* topology by measures μ_k such that $\|\mu_k\| \leq C$ and $\mu_k = \sum_{j=1}^{J(k)} a_{j,k} \delta_{\xi_{j,k}}$, $a_{j,k} \in \mathbb{C}$. Let $\lambda = \lambda_\zeta$ denote an accumulation point of the sequence $\lambda_k = \sum_{j=1}^{J(k)} a_{j,k} \rho_{\zeta, \xi_{j,k}}$ in the weak* topology. Then

$$\frac{g(z)}{(1 - \langle z, \zeta \rangle)^\beta} = \int_{\partial B_n} \frac{1}{(1 - \langle z, \eta \rangle)^\beta} d\lambda_\zeta(\eta),$$

where $\|\lambda_\zeta\| \leq C$. Thus, Lemma 2.4 guarantees that $g \in \mathfrak{M}_\beta(n)$. $\square$

§3. FRACTIONAL CAUCHY TRANSFORMS AND HARDY SPACES

For $f \in \mathcal{H}ol(B_n)$, $0 < p < +\infty$ and $0 < r < 1$, put

$$M_p(f, r) = \left(\int_{\partial B_n} |f(r\zeta)|^p d\sigma_n(\zeta) \right)^{\frac{1}{p}}.$$

Therefore, the classical Hardy space $H^p(B_n)$ is defined by

$$H^p(B_n) = \{f \in \mathcal{H}ol(B_n) : M_p(f, r) \leq C\}.$$

Recall that if f and p are fixed, then the integral mean $M_p(f, r)$ is a monotone increasing function of the variable $r \in (0, 1)$.

For $n \in \mathbb{N}$, let $\mathcal{K}_0(n)$ denote the family of all functions f such that

$$(3.1) \quad f(z) - f(0) = \int_{\partial B_n} \log \frac{1}{1 - \langle z, \zeta \rangle} d\mu(\zeta), \quad z \in B_n,$$

for a measure $\mu \in M(n)$.

Let $f \in \mathcal{K}_\alpha(n)$, $\alpha \geq 0$. In the present section we study the growth of the integral means $M_p(f, r)$ as $r \rightarrow 1-$. For $n = 1$, the corresponding results were obtained in [11].

The following lemma will be used below in numerous situations. The notation $a(z) \approx b(z)$ means that the quotient $a(z)/b(z)$ has a finite positive limit as $|z| \rightarrow 1-$.

Lemma 3.1 (see [23, Proposition 1.4.10]). *Suppose $n \in \mathbb{N}$ and $c \in \mathbb{R}$. Put*

$$I_c(z) = \int_{\partial B_n} \frac{d\sigma_n(\zeta)}{|1 - \langle z, \zeta \rangle|^{n+c}}, \quad z \in \bar{B}_n.$$

- (i) If $c < 0$, then the function $I_c(z)$ is bounded in the closed ball $\bar{B}_n$.
- (ii) If $c > 0$, then $I_c(z) \approx (1 - |z|^2)^{-c}$.
- (iii) Finally,

$$I_0(z) \approx \log \frac{1}{(1 - |z|^2)}.$$

Proposition 3.2. For $\alpha \geq 0$, let $f \in \mathcal{K}_\alpha(n)$. The following estimates hold true as $r \rightarrow 1-$.

If $\alpha = 0$ and $0 < p < +\infty$, then

$$(3.2) \quad M_p^p(f, r) = \mathcal{O}(1).$$

If $0 < \alpha \leq n$ and $0 < p < n/\alpha$, then

$$(3.3) \quad M_p^p(f, r) = \mathcal{O}(1).$$

If $0 < \alpha \leq n$ and $p = n/\alpha$, then

$$(3.4) \quad M_p^p(f, r) = \mathcal{O}\left(\log \frac{1}{1-r}\right).$$

If $p > n/\alpha$ and $p \geq 1$, then

$$(3.5) \quad M_p^p(f, r) = \mathcal{O}\left[\frac{1}{(1-r)^{\alpha p - n}}\right].$$

If $0 < p < 1$ and $\alpha \geq 2n$, then

$$(3.6) \quad M_p^p(f, r) = \mathcal{O}\left[\frac{1}{(1-r)^{(\alpha-n)p}}\right].$$

Proof. It is well known that $\mathcal{K}_n(n) \subset H^p(B_n)$ for all $0 < p < 1$ [23, Theorem 6.2.3]. In other words, estimate (3.3) is fulfilled for $\alpha = n$. Next, let $f = K_\alpha[\mu]$.

Assume that $p \geq 1$. Then the integral Minkowski inequality, Fubini's theorem, and Lemma 3.1 guarantee that

$$\begin{aligned} M_p^p(f, r) &= \int_{\partial B_n} |f(r\xi)|^p d\sigma_n(\xi) \leq \int_{\partial B_n} \int_{\partial B_n} \frac{1}{|1 - \langle r\xi, \zeta \rangle|^{\alpha p}} d\sigma_n(\xi) d|\mu|(\zeta) \\ &= \begin{cases} \mathcal{O}(1) & \text{if } \alpha p < n, \\ \mathcal{O}\left(\log \frac{1}{1-r}\right) & \text{if } \alpha p = n, \\ \mathcal{O}\left[\frac{1}{(1-r)^{\alpha p - n}}\right] & \text{if } \alpha p > n. \end{cases} \end{aligned}$$

This proves estimates (3.4) and (3.5). Property (3.3) is proved for $p \geq 1$. Recall that $H^q(B_n) \subset H^t(B_n)$ for $q > t > 0$. Hence, estimate (3.3) is valid for all $0 < p < n/\alpha$.

If $\gamma > 0$, then

$$-\log 2 \leq \log \frac{1}{|1 - \langle r\xi, \zeta \rangle|} \leq C(\gamma) \frac{1}{|1 - \langle r\xi, \zeta \rangle|^\gamma},$$

where $0 < r < 1$ and $\xi, \zeta \in \partial B_n$. Hence, estimate (3.2) is proved for all $p \in (0, +\infty)$.

Now, assume that $0 < p < 1$ and $\alpha \geq 2n$. Note that $(a + b)^p \leq C(p)(a^p + b^p)$ for $a \geq 0, b \geq 0$. Hence, by the Jordan decomposition theorem, we may assume that μ is a probability measure. Therefore, we obtain

$$\begin{aligned} (3.7) \quad |f(z)| &\leq \int_{\partial B_n} \frac{1}{|1 - \langle z, \zeta \rangle|^\alpha} d\mu(\zeta) \\ &\leq \frac{1}{(1 - |z|)^{\alpha - 2n}} \int_{\partial B_n} \frac{1}{|1 - \langle z, \zeta \rangle|^{2n}} d\mu(\zeta) \\ &= \frac{1}{(1 - |z|)^{\alpha - n}} \int_{\partial B_n} \frac{(1 - |z|^2)^n}{|1 - \langle z, \zeta \rangle|^{2n}} d\mu(\zeta). \end{aligned}$$

Note that

$$u(z) = \int_{\partial B_n} \frac{(1 - |z|^2)^n}{|1 - \langle z, \zeta \rangle|^{2n}} d\mu(\zeta)$$

is a positive $\mathcal{M}$ -harmonic function in the ball. Thus,

$$\int_{\partial B_n} u(r\zeta) d\sigma_n(\zeta) = u(0)$$

for $0 < r < 1$; see [23, Theorem 4.2.4]. Next, since $1/p > 1$, Minkowski's inequality yields

$$(3.8) \quad \left(\int_{\partial B_n} u^p(r\zeta) d\sigma_n(\zeta) \right)^{\frac{1}{p}} \leq \int_{\partial B_n} u^{p \cdot \frac{1}{p}}(r\zeta) d\sigma_n(\zeta) = u(0)$$

for $0 < r < 1$.

Estimates (3.7) and (3.8) imply that

$$M_p^p(f, r) = \int_{\partial B_n} |f(r\zeta)|^p d\sigma_n(\zeta) \leq \frac{1}{(1-r)^{(\alpha-n)p}} u^p(0) = \frac{(\mu(\partial B_n))^p}{(1-r)^{(\alpha-n)p}}.$$

This proves (3.6). $\square$

Corollary 3.3. *Let $n \in \mathbb{N}$.*

- (i) *If $0 \leq \alpha \leq n$, then $\mathcal{K}_\alpha(n) \subset H^p(B_n)$ for $0 < p < n/\alpha$.*
- (ii) *If $f \in H^1(B_n)$, then $f \in \mathcal{K}_n(n)$; moreover, $\|f\|_{\mathcal{K}_n(n)} \leq \|f\|_{H^1(B_n)}$.*
- (iii) $\mathcal{K}_0(n) \subset \mathcal{K}_n(n)$.

Proof. Part (i) is equivalent to estimates (3.2) and (3.3). Next, if $f \in H^1(B_n)$, then $f = K_n[f^*]$, where $f^*(\zeta) = \lim_{r \rightarrow 1-} f(r\zeta)$, $\zeta \in \partial B_n$. Therefore, $\|f\|_{\mathcal{K}_n(n)} \leq \|f\|_{H^1(B_n)}$. Finally, part (iii) follows from (i) and (ii). $\square$

Part (iii) of Corollary 3.3 will be improved in Corollary 5.5. Also, estimate (3.3) can be refined in terms of the weak Hardy spaces $H^{p,\infty}(B_n)$. Let $p > 0$. By definition, $f \in H^{p,\infty}(B_n)$ if $f \in H^q(B_n)$ for some $q > 0$ and

$$\|f\|_{p,\infty}^p = \sup_{t>0} t^p \sigma_n\{\zeta \in \partial B_n : |f^*(\zeta)| > t\} < \infty.$$

Proposition 3.4. *If $0 < \alpha \leq n$, then $\mathcal{K}_\alpha(n) \subset H^{\frac{n}{\alpha},\infty}(B_n)$.*

Proof. The argument below is due to A. B. Aleksandrov. The case of $\alpha = n$ is well known. So, assume that $p = n/\alpha > 1$; there exists a norm $\|\cdot\|_p^*$ on the space $H^{p,\infty}(B_n)$ such that

$$(3.9) \quad \|f\|_{p,\infty} \leq C_1(p) \|f\|_p^* \leq C_2(p) \|f\|_{p,\infty}$$

for all $f \in H^{p,\infty}(B_n)$, for some positive constants $C_1(p)$ and $C_2(p)$ (see, e.g., [6]).

Now, assume that $f \in K_\alpha[\mu]$, $\mu \in M(n)$. We must prove that $f \in H^{p,\infty}(B_n)$. Suppose, without loss of generality, that μ is a probability measure. We approximate the measure μ in the weak* topology by probability measures $\mu_k = \sum_{j=1}^{J(k)} a_{k,j} \delta_{\xi_{k,j}}$, where $a_{k,j} > 0$ and $\delta_{\xi_{k,j}}$ denotes the δ -measure at the point $\xi_{k,j} \in \partial B_n$. Applying estimates (3.9), the triangle inequality, and Proposition 5.1.4 in [23], we obtain

$$\begin{aligned} \|K_\alpha[\mu_k]\|_{p,\infty} &\leq C_1(p) \|K_\alpha[\mu_k]\|_p^* \leq C_1(p) \|(1 - z_1)^{-\alpha}\|_p^* \\ &\leq C_2(p) \|(1 - z_1)^{-\alpha}\|_{p,\infty} \leq C(p, n). \end{aligned}$$

Finally, since $\mu_k \rightarrow \mu$ in the weak* topology, we see that $f \in H^{p,\infty}(B_n)$. The proof is finished. $\square$

§4. FAMILIES $\mathcal{K}_\alpha(n)$ AND DIFFERENTIATION

Let $f \in \mathcal{H}ol(B_1)$, and let $\alpha \geq 0$. It is well known that $f \in \mathcal{K}_\alpha(1)$ if and only if $f' \in \mathcal{K}_{\alpha+1}(1)$. In the present section, we prove similar assertions for $f \in \mathcal{H}ol(B_n)$, $n \in \mathbb{N}$.

4.1. Radial derivatives. Let $f \in \mathcal{H}ol(B_n)$. The radial derivative $\mathcal{R}f$ is defined by the identity

$$\mathcal{R}f(z) = \sum_{j=1}^n z_j \frac{\partial f}{\partial z_j}(z).$$

Recall that

$$(4.1) \quad f(z) - f(0) = \int_0^1 \frac{\mathcal{R}f(tz)}{t} dt, \quad z \in B_n.$$

Proposition 4.1. *Let $f \in \mathcal{H}ol(B_n)$, $n \in \mathbb{N}$. Then $f \in \mathcal{K}_0(n)$ if and only if $\mathcal{R}f \in \mathcal{K}_1(n)$.*

Proof. Suppose that $f \in \mathcal{K}_0(n)$, that is, (3.1) is true. Then

$$\mathcal{R}f(z) = \int_{\partial B_n} \frac{\langle z, \zeta \rangle}{1 - \langle z, \zeta \rangle} d\mu(\zeta) = K_1[\rho](z),$$

where $\rho = \mu - \mu(\partial B_n)\sigma_n$.

To prove the reverse implication, assume that

$$\mathcal{R}f(z) = \int_{\partial B_n} \frac{1}{1 - \langle z, \zeta \rangle} d\rho(\zeta),$$

where $\rho \in M(n)$. Note that $\mathcal{R}f(0) = 0$; thus, $\rho(\partial B_n) = 0$. Applying identity (4.1) and the property $\rho(\partial B_n) = 0$, we obtain

$$\begin{aligned} f(z) - f(0) &= \int_0^1 \int_{\partial B_n} \frac{1}{t(1 - t\langle z, \zeta \rangle)} d\rho(\zeta) dt \\ &= \int_0^1 \int_{\partial B_n} \left(\frac{1}{t} + \frac{\langle z, \zeta \rangle}{1 - t\langle z, \zeta \rangle} \right) d\rho(\zeta) dt \\ &= \int_{\partial B_n} \int_0^1 \frac{\langle z, \zeta \rangle}{1 - t\langle z, \zeta \rangle} dt d\rho(\zeta) \\ &= \int_{\partial B_n} \log \frac{1}{1 - \langle z, \zeta \rangle} d\rho(\zeta), \end{aligned}$$

as required. $\square$

Proposition 4.2. *Assume that $\alpha > 0$ and $f \in \mathcal{H}ol(B_n)$, $n \in \mathbb{N}$. Then $f \in \mathcal{K}_0(n) + \mathcal{K}_\alpha(n)$ if and only if $\mathcal{R}f \in \mathcal{K}_{\alpha+1}(n)$.*

Proof. Let $g \in \mathcal{K}_0(n)$ and $h \in \mathcal{K}_\alpha(n)$, that is,

$$h(z) = \int_{\partial B_n} \frac{1}{(1 - \langle z, \zeta \rangle)^\alpha} d\mu(\zeta), \quad z \in B_n.$$

Straightforward calculations show that

$$\mathcal{R}h(z) = \int_{\partial B_n} \frac{\alpha}{(1 - \langle z, \zeta \rangle)^{\alpha+1}} d\mu(\zeta) - \alpha h(z) \in \mathcal{K}_{\alpha+1}(n)$$

because $\alpha h \in \mathcal{K}_\alpha(n) \subset \mathcal{K}_{\alpha+1}(n)$ by part (ii) of Corollary 2.3. On the other hand, Proposition 4.1 guarantees that $\mathcal{R}g \in \mathcal{K}_1(n) \subset \mathcal{K}_{\alpha+1}(n)$.

It is convenient to split the proof of the reverse implication into two steps.

Step 1. Let $\alpha = m \in \mathbb{N}$. By the hypothesis, we have

$$\mathcal{R}f(z) = \int_{\partial B_n} \frac{1}{(1 - \langle z, \zeta \rangle)^{m+1}} d\mu(\zeta), \quad z \in B_n.$$

Hence,

$$f(z) - f(0) = \int_0^1 \frac{\mathcal{R}f(tz)}{t} dt = \int_0^1 \int_{\partial B_n} \frac{1}{t(1 - t\langle z, \zeta \rangle)^{m+1}} d\mu(\zeta) dt, \quad z \in B_n.$$

If $w \in \mathbb{C}$ and $|w| < 1$, then

$$\frac{1}{t(1 - tw)^{m+1}} = \frac{1}{t} + \sum_{j=1}^{m+1} \frac{w}{(1 - tw)^j}.$$

Since $\mathcal{R}f(0) = 0$, we have $\mu(\partial B_n) = 0$. Putting $w = \langle z, \zeta \rangle$, we obtain

$$\begin{aligned} f(z) - f(0) &= \sum_{j=1}^{m+1} \int_{\partial B_n} \int_0^1 \frac{\langle z, \zeta \rangle}{(1 - t\langle z, \zeta \rangle)^j} dt d\mu(\zeta) \\ &= - \int_{\partial B_n} \log(1 - \langle z, \zeta \rangle) d\mu(\zeta) + \sum_{j=2}^{m+1} \int_{\partial B_n} \frac{1}{(j-1)(1 - \langle z, \zeta \rangle)^{j-1}} d\mu(\zeta). \end{aligned}$$

The embeddings $\mathcal{K}_{j-1}(n) \subset \mathcal{K}_m(n)$, $j = 2, \dots, m$, guarantee that $f \in \mathcal{K}_0(n) + \mathcal{K}_m(n)$.

Step 2. Let $\alpha > 0$, $\alpha \notin \mathbb{N}$. Repeating the arguments of Step 1 and reversing the order of integration, we have

$$\begin{aligned} f(z) - f(0) &= \int_0^1 \int_{\partial B_n} \frac{1}{t(1 - t\langle z, \zeta \rangle)^{\alpha+1}} d\mu(\zeta) dt \\ &= \int_{\partial B_n} \int_0^1 \frac{\langle z, \zeta \rangle}{(1 - t\langle z, \zeta \rangle)^{\alpha+1}} dt d\mu(\zeta) + \int_0^1 \frac{1}{t} \int_{\partial B_n} \frac{1}{(1 - t\langle z, \zeta \rangle)^\alpha} d\mu(\zeta) dt. \end{aligned}$$

The inner integral in the first summand can be calculated explicitly. So, consider the second summand. Put $[\alpha] = m \in \mathbb{N} \cup \{0\}$. Note that $m+1 > \alpha$ and $\mu(\partial B_n) = 0$. Hence, by part (ii) of Corollary 2.3, there exists a measure $\rho \in M(n)$ such that $K_{m+1}[\rho] = K_\alpha[\mu]$. Also, we have $\rho(\partial B_n) = K_{m+1}[\rho](0) = K_\alpha[\mu](0) = \mu(\partial B_n) = 0$. Therefore, reversing the order of integration once again, we obtain

$$f(z) - f(0) = \frac{1}{\alpha} \int_{\partial B_n} \frac{1}{(1 - \langle z, \zeta \rangle)^\alpha} d\mu(\zeta) + \int_{\partial B_n} \int_0^1 \frac{1}{t(1 - t\langle z, \zeta \rangle)^{m+1}} dt d\rho(\zeta).$$

By the definition, the first summand in the above sum belongs to $\mathcal{K}_\alpha(n)$. If $m = 0$, then the proof of Proposition 4.1 shows that the second summand is in the family $\mathcal{K}_0(n)$. Finally, for $m \in \mathbb{N}$, at Step 1 it was proved that the second summand belongs to $\mathcal{K}_0(n) + \mathcal{K}_m(n) \subset \mathcal{K}_0(n) + \mathcal{K}_\alpha(n)$. $\square$

Note that Proposition 4.2 admits refinement (see Corollary 5.6).

4.2. Fractional differential operators of order 1. By definition, we put $R^1 f = f + \mathcal{R}f$ for $f \in \mathcal{H}ol(B_n)$. The operator R^1 is called the fractional differential operator of order 1. Observe that

$$(4.2) \quad f(z) = \int_0^1 R^1 f(tz) dt, \quad z \in B_n.$$

The following proposition can be proved with the help of elementary methods.

Proposition 4.3. *Let $f \in \mathcal{H}ol(B_n)$, $n \in \mathbb{N}$.*

- (i) *If $\alpha > 0$, then the property $f \in \mathcal{K}_\alpha(n)$ implies that $R^1 f \in \mathcal{K}_{\alpha+1}(n)$.*
- (ii) *If $\alpha \geq 1$, then the property $R^1 f \in \mathcal{K}_{\alpha+1}(n)$ implies that $f \in \mathcal{K}_\alpha(n)$.*

Proof. Let $f \in \mathcal{K}_\alpha(n)$, that is, $f = K_\alpha[\mu]$. Then

$$f + \mathcal{R}f = \alpha K_{\alpha+1}[\mu] + (1 - \alpha)f \in \mathcal{K}_{\alpha+1}(n)$$

because $(1 - \alpha)f \in \mathcal{K}_\alpha(n) \subset \mathcal{K}_{\alpha+1}(n)$. The proof of part (i) is complete.

We split the proof of part (ii) into two steps.

Step 1. $\alpha = m \in \mathbb{N}$. By assumption, we have

$$R^1 f(tz) = \int_{\partial B_n} \frac{1}{(1 - t\langle z, \zeta \rangle)^{m+1}} d\mu(\zeta), \quad z \in B_n, \quad t \in [0, 1].$$

Hence, using (4.2), we obtain

$$f(z) = \int_{\partial B_n} \int_0^1 \frac{1}{(1 - t\langle z, \zeta \rangle)^{m+1}} dt d\mu(\zeta), \quad z \in B_n.$$

Let $I(z, \zeta)$ denote the inner integral. If $\langle z, \zeta \rangle = 0$, then $I(z, \zeta) = 1$. If $\langle z, \zeta \rangle \neq 0$, then

$$I(z, \zeta) = \frac{1}{m} \frac{1 - (1 - \langle z, \zeta \rangle)^m}{\langle z, \zeta \rangle (1 - \langle z, \zeta \rangle)^m} = \frac{1}{m} \sum_{j=1}^m (1 - \langle z, \zeta \rangle)^{-j}.$$

Therefore, in both cases we have

$$I(z, \zeta) = \frac{1}{m} \sum_{j=1}^m (1 - \langle z, \zeta \rangle)^{-j}.$$

Consequently,

$$f(z) = \frac{1}{m} \sum_{j=1}^m \int_{\partial B_n} \frac{1}{(1 - \langle z, \zeta \rangle)^j} d\mu(\zeta).$$

Finally, we see that $f \in \mathcal{K}_m(n)$, because $\mathcal{K}_j(n) \subset \mathcal{K}_m(n)$ for $j = 1, \dots, m - 1$.

Step 2. $\alpha > 1$, $\alpha \notin \mathbb{N}$. Repeating the arguments of Step 1, we get

$$f(z) = \int_{\partial B_n} \int_0^1 \frac{1}{(1 - t\langle z, \zeta \rangle)^{\alpha+1}} dt d\mu(\zeta), \quad z \in B_n.$$

We represent the inner integral as a sum of two summands and integrate by parts in the second summand, obtaining

$$\begin{aligned} \int_0^1 \frac{1}{(1 - t\langle z, \zeta \rangle)^{\alpha+1}} dt &= \int_0^1 \frac{1}{(1 - t\langle z, \zeta \rangle)^\alpha} dt + \int_0^1 \frac{t\langle z, \zeta \rangle}{(1 - t\langle z, \zeta \rangle)^{\alpha+1}} dt \\ &= \frac{1}{\alpha} \frac{1}{(1 - \langle z, \zeta \rangle)^\alpha} + \frac{\alpha - 1}{\alpha} \int_0^1 \frac{1}{(1 - t\langle z, \zeta \rangle)^\alpha} dt. \end{aligned}$$

Hence, Fubini's theorem yields

$$f(z) = \frac{1}{\alpha} \int_{\partial B_n} \frac{1}{(1 - \langle z, \zeta \rangle)^\alpha} d\mu(\zeta) + \frac{\alpha - 1}{\alpha} \int_0^1 \int_{\partial B_n} \frac{1}{(1 - t\langle z, \zeta \rangle)^\alpha} d\mu(\zeta) dt.$$

By definition, the first summand in the above sum belongs to $\mathcal{K}_\alpha(n)$.

Consider the second summand. Let $[\alpha] = m \in \mathbb{N}$. Note that $m + 1 > \alpha$. Hence, by part (ii) of Corollary 2.3, there exists a measure $\rho \in M(n)$ such that $K_{m+1}[\rho] = K_\alpha[\mu]$. Therefore, the second summand has the following form:

$$\frac{\alpha - 1}{\alpha} \int_0^1 \int_{\partial B_n} \frac{1}{(1 - \langle tz, \zeta \rangle)^{m+1}} d\rho(\zeta) dt.$$

At Step 1 it was proved that the above function belongs to $\mathcal{K}_m(n) \subset \mathcal{K}_\alpha(n)$. $\square$

To refine Proposition 4.3, we need the following lemma.

Lemma 4.4. *Assume that $\alpha > 0$ and $g(w) = 1 - (1 - w)^\alpha$, $w \in B_1$. Then there exists a measure $\rho \in M(1)$ such that $(1 - w)^{-\alpha} g(w)/w = K_\alpha[\rho](w)$, $w \in B_1$.*

Proof. The definitions imply that $(g(w)/w)'w^2 = g'(w)w - g(w) \in H^1(B_1)$, where $H^1(B_1)$ denotes the Hardy space. Note that $h \in H^1(B_1)$ if $h \in \mathcal{H}ol(B_n)$ and $h(w)w^2 \in H^1(B_1)$. Putting $h(w) = (g(w)/w)'$, we obtain $(g(w)/w)' \in H^1(B_1)$. Hence, $g(w)/w \in \mathfrak{M}_\alpha(1)$ by [16, Theorem 3.5]. Since $(1 - w)^{-\alpha} \in \mathcal{K}_\alpha(1)$, the required measure $\rho \in M(1)$ exists by the definition of the multiplier space $\mathfrak{M}_\alpha(1)$. $\square$

Theorem 4.5. *Assume that $\alpha > 0$ and $f \in \mathcal{H}ol(B_n)$, $n \in \mathbb{N}$. Then $f \in \mathcal{K}_\alpha(n)$ if and only if $R^1 f \in \mathcal{K}_{\alpha+1}(n)$.*

Proof. Suppose that $R^1 f = K_{\alpha+1}[\mu]$, where $\mu \in M(n)$. Applying formula (4.2) and reversing the order of integration, we obtain

$$\begin{aligned} \alpha f(z) &= \int_{\partial B_n} \int_0^1 \frac{\alpha dt}{(1 - t\langle z, \zeta \rangle)^{\alpha+1}} d\mu(\zeta) \\ &= \int_{\partial B_n} \frac{1 - (1 - \langle z, \zeta \rangle)^\alpha}{\langle z, \zeta \rangle (1 - \langle z, \zeta \rangle)^\alpha} d\mu(\zeta), \quad z \in B_n, \end{aligned}$$

where the integrand is assumed to be equal to α if $\langle z, \zeta \rangle = 0$. We put $w = \langle z, \zeta \rangle$ and apply Lemma 4.4. Reversing the order of integration, we have

$$\begin{aligned} \alpha f(z) &= \int_{\partial B_1} \int_{\partial B_n} (1 - \langle z, \lambda \zeta \rangle)^{-\alpha} d\mu(\zeta) d\rho(\lambda) \\ &= \int_{\partial B_1} K_\alpha[\mu_\lambda](z) d\rho(\lambda), \quad z \in B_n, \end{aligned}$$

where $\mu_\lambda(E) = \mu(\bar{\lambda}E)$ for the Borel sets $E \subset \partial B_n$. Note that $K_\alpha[\mu_\lambda](z)$ is a continuous function of the variable $\lambda \in \partial B_1$. Below we show that the last integral is in the family $\mathcal{K}_\alpha(n)$. Without loss of generality, we may assume that ρ is a probability measure. We approximate the measure ρ in the weak* topology by the probability measures $\rho_k = \sum_{j=1}^{J(k)} a_{k,j} \delta_{\lambda_{k,j}}$, where $\delta_{\lambda_{k,j}}$ denotes the δ -measure at $\lambda_{k,j} \in \partial B_1$. By the definition of weak* convergence, we have

$$\sum_{j=1}^{J(k)} a_{k,j} K_\alpha[\mu_{\lambda_{k,j}}](z) \xrightarrow{k \rightarrow \infty} \int_{\partial B_1} K_\alpha[\mu_\lambda](z) d\rho(\lambda), \quad z \in B_n.$$

Next, observe that the sequence $\left\{ \sum_{j=1}^{J(k)} a_{k,j} \mu_{\lambda_{k,j}} \right\}$ is norm bounded. Hence, there exists a subsequence which converges in the weak* topology to a measure $\nu \in M(n)$. Therefore,

$$\sum_{j=1}^{J(k)} a_{k,j} K_\alpha[\mu_{\lambda_{k,j}}](z) \xrightarrow{k \rightarrow \infty} K_\alpha[\nu](z), \quad z \in B_n.$$

Finally, we obtain $f = K_\alpha[\nu/\alpha]$. Now, it suffices to refer to Proposition 4.3. $\square$

Note that Theorem 4.5 extends to the case of $\alpha = 0$ (see Corollary 5.7).

§5. FAMILIES $\mathcal{K}_\alpha(n)$ AND BERGMAN-SOBOLEV SPACES

5.1. Modified operators of fractional differentiation. Consider a pair of parameters $(\beta, t) \in \mathbb{R}^2$ with the property that neither $n - 1 - \beta$ nor $n - 1 - \beta + t$ is a strictly negative integer. We define an operator

$$R^{\beta, t} : \mathcal{H}ol(B_n) \rightarrow \mathcal{H}ol(B_n)$$

as follows. If

$$(5.1) \quad f(z) = \sum_{k=0}^{\infty} f_k(z)$$

is the homogeneous expansion of $f \in \mathcal{H}ol(B_n)$, then

$$(5.2) \quad R^{\beta, t} f(z) = \sum_{k=0}^{\infty} \frac{\Gamma(n - \beta) \Gamma(n + k - \beta + t)}{\Gamma(n - \beta + t) \Gamma(n + k - \beta)} f_k(z).$$

The inverse of $R^{\beta, t}$, denoted by $R_{\beta, t}$, is given by the formula

$$R_{\beta, t} f(z) = \sum_{k=0}^{\infty} \frac{\Gamma(n - \beta + t) \Gamma(n + k - \beta)}{\Gamma(n - \beta) \Gamma(n + k - \beta + t)} f_k(z).$$

Assume that $\beta < n$, $t > 0$, and identity (5.1) is true. Also, suppose that $r \in [0, 1]$ and $z \in B_n$. By the definition of the operator $R^{\beta, t}$, we have

$$\frac{\Gamma(n - \beta + t)}{\Gamma(n - \beta) \Gamma(t)} R^{\beta, t} f(rz) = \sum_{k=0}^{\infty} \frac{r^k f_k(z)}{B(n + k - \beta, t)}.$$

Therefore,

$$(5.3) \quad \frac{\Gamma(n - \beta + t)}{\Gamma(n - \beta) \Gamma(t)} \int_0^1 R^{\beta, t} f(rz) r^{n-\beta-1} (1-r)^{t-1} dr = \sum_{k=0}^{\infty} f_k(z) = f(z).$$

Straightforward calculations imply the following lemma.

Lemma 5.1 (cf. [30, Proposition 1.14]). *Let $n \in \mathbb{N}$. Suppose that neither $n - 1 - \beta$ nor $n - 1 - \beta + t$ is a strictly negative integer. Then*

$$R^{\beta, t} \left(\frac{1}{(1 - \langle z, \zeta \rangle)^{n-\beta}} \right) = \frac{1}{(1 - \langle z, \zeta \rangle)^{n-\beta+t}}$$

for all points $z \in B_n$ and $\zeta \in \partial B_n$.

5.2. A condition sufficient for membership in $\mathcal{K}_\alpha(n)$ with $\alpha > n$. For $n = 1$, the following fact was proved in [13].

Proposition 5.2. *Suppose $n \in \mathbb{N}$ and $\alpha > n$. If $g \in \mathcal{H}ol(B_n)$ and*

$$\int_0^1 \int_{\partial B_n} |g(r\zeta)| (1-r)^{\alpha-n-1} d\sigma_n(\zeta) dr = V < +\infty,$$

then $g \in \mathcal{K}_\alpha(n)$ and $\|g\|_{\mathcal{K}_\alpha(n)} \leq C(\alpha, n, V) = (\alpha - n) \cdots (\alpha - 1) V / (n - 1)!$.

Proof. Put $t = \alpha - n$ and $f = R_{0, t} g$. Then $g = R^{0, t} f$ and

$$f(z) = \frac{\Gamma(n + t)}{\Gamma(n) \Gamma(t)} \int_0^1 g(rz) r^{n-1} (1-r)^{t-1} dr$$

by (5.3). If $0 < s < 1$, then

$$\begin{aligned} \frac{\Gamma(n)\Gamma(t)}{\Gamma(n+t)} \int_{\partial B_n} |f(s\zeta)| d\sigma_n(\zeta) &\leq \int_0^1 \int_{\partial B_n} |g(sr\zeta)| d\sigma_n(\zeta) (1-r)^{t-1} dr \\ &\leq \int_0^1 \int_{\partial B_n} |g(r\zeta)| d\sigma_n(\zeta) (1-r)^{t-1} dr \\ &= V < +\infty. \end{aligned}$$

Therefore, $f \in H^1(B_n) \subset \mathcal{K}_n(n)$ and $\|f\|_{\mathcal{K}_n(n)} \leq \|f\|_{H^1(B_n)} \leq C(\alpha, n, V)$. In other words,

$$f(z) = \int_{\partial B_n} \frac{1}{(1 - \langle z, \zeta \rangle)^n} d\mu(\zeta), \quad z \in B_n,$$

where $\|\mu\|_{M(n)} \leq C(\alpha, n, V)$. Thus, with the help of Lemma 5.1, we obtain

$$g(z) = R^{0,t}f(z) = \int_{\partial B_n} \frac{1}{(1 - \langle z, \zeta \rangle)^{n+t}} d\mu(\zeta), \quad z \in B_n.$$

Hence, $\|g\|_{\mathcal{K}_\alpha(n)} \leq C(\alpha, n, V)$. □

5.3. Bergman–Sobolev spaces. Suppose $j \in \mathbb{N}$ and $q > 0$. By definition, the Bergman–Sobolev space $A_{q,j}^1(B_n)$ consists of all functions $f \in \mathcal{H}ol(B_n)$ for which

$$\int_0^1 \int_{\partial B_n} |R^j f(r\zeta)| (1-r)^{q-1} d\sigma_n(\zeta) dr < +\infty,$$

where $R^j = (R^1)^j$. For $n = 1$, the following assertion coincides with [13, Lemma 2].

Proposition 5.3. *Suppose $n \in \mathbb{N}$, $j \in \{1, \dots, n\}$, and $\alpha > n - j$.*

- (i) *If $f \in A_{\alpha-n+j,j}^1(B_n)$, then $f \in \mathcal{K}_\alpha(n)$.*
- (ii) *Let $\beta > \alpha$. Then $\mathcal{K}_\alpha(n) \subset A_{\beta-n+j,j}^1(B_n)$.*

Proof. We prove part (i). Since $\alpha + j > n$, Proposition 5.2 shows that $R^j f \in \mathcal{K}_{\alpha+j}(n)$. Since $\alpha > 0$, repeated application of Theorem 4.5 guarantees that $f \in \mathcal{K}_\alpha(n)$.

Now, we turn to part (ii). Let $f \in \mathcal{K}_\alpha(n)$, $\alpha > n - j$. By Theorem 4.5, we have $R^j f \in \mathcal{K}_{\alpha+j}(n)$; that is,

$$R^j f(z) = \int_{\partial B_n} \frac{1}{(1 - \langle z, \zeta \rangle)^{\alpha+j}} d\mu(\zeta), \quad z \in B_n,$$

for some measure $\mu \in M(n)$. Therefore, if $0 \leq r < 1$, then

$$\int_{\partial B_n} |R^j f(r\xi)| d\sigma_n(\xi) \leq \int_{\partial B_n} \int_{\partial B_n} \frac{1}{|1 - r\langle \xi, \zeta \rangle|^{\alpha+j}} d\sigma_n(\xi) d|\mu|(\zeta).$$

Since $\alpha + j > n$, Lemma 3.1 yields

$$\int_{\partial B_n} |R^j f(r\xi)| d\sigma_n(\xi) \leq C\|\mu\|_{M(n)}(1-r)^{n-\alpha-j}.$$

Now, assume that $\beta > \alpha$. Then the above inequality guarantees that

$$\int_0^1 \int_{\partial B_n} |R^j f(r\xi)| (1-r)^{\beta-n+j-1} d\sigma_n(\xi) dr \leq C\|\mu\| \int_0^1 (1-r)^{\beta-\alpha-1} < +\infty.$$

In other words, $f \in A_{\beta-n+j,j}^1(B_n)$. □

5.4. Bergman–Sobolev spaces and $\mathcal{K}_0(n)$.

Lemma 5.4. *Let $n \in \mathbb{N}$. Then $\mathcal{K}_0(n) \subset A_{\varepsilon,n}^1(B_n)$ for any $\varepsilon > 0$.*

Proof. Let $f \in \mathcal{K}_0(n)$. Recall that $\mathcal{K}_0(n) \subset \mathcal{K}_n(n)$ by part (iii) of Corollary 3.3. Next, $\mathcal{R}^j f \in \mathcal{K}_j(n) \subset \mathcal{K}_n(n)$ for $j = 1, 2, \dots, n$ by Propositions 4.1, 4.2 and Corollary 2.3. Hence, $R^n f = (I + \mathcal{R})^n f \in \mathcal{K}_n(n)$; that is,

$$R^n f(z) = \int_{\partial B_n} \frac{1}{(1 - \langle z, \zeta \rangle)^n} d\mu(\zeta), \quad z \in B_n,$$

for some measure $\mu \in M(n)$. Hence, if $0 \leq r < 1$, then

$$\int_{\partial B_n} |R^n f(r\xi)| d\sigma_n(\xi) \leq \int_{\partial B_n} \int_{\partial B_n} \frac{1}{|1 - r\langle \xi, \zeta \rangle|^n} d\sigma_n(\xi) d|\mu|(\zeta).$$

By Lemma 3.1, we have

$$\int_{\partial B_n} |R^n f(r\xi)| d\sigma_n(\xi) \leq C \log \frac{e}{1-r}.$$

The above estimate implies that

$$\int_0^1 \int_{\partial B_n} |R^n f(r\xi)| (1-r)^{\varepsilon-1} d\sigma_n(\xi) dr \leq C \int_0^1 (1-r)^{\varepsilon-1} \log \frac{e}{1-r} < +\infty,$$

as required. $\square$

5.5. An embedding property for $\mathcal{K}_0(n)$. It is well known that $\mathcal{K}_0(1) \subset \mathcal{K}_\alpha(1)$ for all $\alpha > 0$.

Corollary 5.5. *Let $n \in \mathbb{N}$. Then $\mathcal{K}_0(n) \subset \mathcal{K}_\alpha(n)$ for all $\alpha > 0$.*

Proof. Lemma 5.4 guarantees that $\mathcal{K}_0(n) \subset A_{\alpha,n}^1(B_n)$, $\alpha > 0$. It remains to note that $A_{\alpha,n}^1(B_n) \subset \mathcal{K}_\alpha(n)$ by part (i) of Proposition 5.3. $\square$

A. B. Aleksandrov observed that Corollary 5.5 can also be deduced from the embedding $\mathcal{K}_0(1) \subset \mathcal{K}_\alpha(1)$ for all $\alpha > 0$.

5.6. The families $\mathcal{K}_\alpha(n)$, $\alpha \geq 0$, and differentiation. We have the following refinement of Proposition 4.2.

Corollary 5.6. *Suppose $n \in \mathbb{N}$ and $\alpha \geq 0$. Then $f \in \mathcal{K}_\alpha(n)$ if and only if $\mathcal{R}f \in \mathcal{K}_{\alpha+1}(n)$.*

Proof. We apply Propositions 4.1, 4.2 and Corollary 5.5. $\square$

The next corollary refines Theorem 4.5.

Corollary 5.7. *Suppose $n \in \mathbb{N}$ and $\alpha \geq 0$. Then $f \in \mathcal{K}_\alpha(n)$ if and only if $R^1 f \in \mathcal{K}_{\alpha+1}(n)$.*

Proof. The case of $\alpha > 0$ is covered by Theorem 4.5. So, assume that $f \in \mathcal{K}_0(n)$. Corollary 5.5 guarantees that $f \in \mathcal{K}_1(n)$. On the other hand, $\mathcal{R}f \in \mathcal{K}_1(n)$ by Corollary 5.6. Therefore, $R^1 f = f + \mathcal{R}f \in \mathcal{K}_1(n)$.

To prove the reverse implication, assume that $R^1 f \in \mathcal{K}_1(n)$. Since $\mathcal{K}_1(n) \subset \mathcal{K}_2(n)$, we have $R^1 f \in \mathcal{K}_2(n)$. Hence, $f \in \mathcal{K}_1(n)$ by Theorem 4.5. Therefore, $\mathcal{R}f = R^1 f - f \in \mathcal{K}_1(n)$. So, $f \in \mathcal{K}_0(n)$ by Corollary 5.6. $\square$

5.7. Modified Bergman–Sobolev spaces and $\mathcal{K}_\alpha(n)$. Suppose $n \in \mathbb{N}$, $j \in \mathbb{N}$, and $q > 0$. In some applications, it is convenient to use the modified Bergman–Sobolev space $\tilde{A}_{q,j}^1(B_n)$ in place of $A_{q,j}^1(B_n)$. By definition, the space $\tilde{A}_{q,j}^1(B_n)$ consists of functions $f \in \mathcal{H}ol(B_n)$ such that

$$\|f\|_{\tilde{A}_{q,j}^1(B_n)} = \sum_{|m| \leq j-1} \left| \frac{\partial^m f}{\partial z^m}(0) \right| + \int_0^1 \int_{\partial B_n} |\mathcal{R}^j f(r\zeta)| (1-r)^{q-1} d\sigma_n(\zeta) dr < +\infty,$$

where $m = (m_1, \dots, m_n) \in \mathbb{Z}_+^n$ and $|m| = m_1 + \dots + m_n$. The space $\tilde{A}_{q,j}^1(B_n)$ with the norm $\|\cdot\|_{\tilde{A}_{q,j}^1(B_n)}$ is a Banach space (cf. [30, Proposition 6.2]).

Proposition 5.8. Suppose $n \in \mathbb{N}$, $j \in \{1, \dots, n\}$, and $\alpha > n - j$.

- (i) The embedding $\tilde{A}_{\alpha-n+j,j}^1(B_n) \subset \mathcal{K}_\alpha(n)$ holds true. If $f \in \tilde{A}_{\alpha-n+j,j}^1(B_n)$, then $\|f\|_{\mathcal{K}_\alpha(n)} \leq C \|f\|_{\tilde{A}_{\alpha-n+j,j}^1(B_n)}$, where the constant $C > 0$ does not depend on f .
- (ii) Let $\beta > \alpha$. Then $\mathcal{K}_\alpha(n) \subset \tilde{A}_{\beta-n+j,j}^1(B_n)$.

Proof. Assume that $f \in \tilde{A}_{\alpha-n+j,j}^1(B_n)$. We have

$$\int_0^1 \int_{\partial B_n} |\mathcal{R}^j f(r\zeta)| (1-r)^{\alpha+j-n-1} d\sigma_n(\zeta) dr < +\infty.$$

Since $\alpha + j > n$, Proposition 5.2 yields $\mathcal{R}^j f \in \mathcal{K}_{\alpha+j}(n)$. Repeated application of Corollary 5.6 shows that $f \in \mathcal{K}_\alpha(n)$.

Now, consider the operator $I : \tilde{A}_{\alpha-n+j,j}^1(B_n) \rightarrow \mathcal{K}_\alpha(n)$ defined by $If = f$. If a functional sequence converges in the space $\tilde{A}_{\alpha-n+j,j}^1(B_n)$ or in the space $\mathcal{K}_\alpha(n)$, then this sequence converges uniformly on the compact subsets of the ball B_n . Hence, the graph of the operator I is closed. So, by the closed graph theorem,

$$\|f\|_{\mathcal{K}_\alpha(n)} \leq C \|f\|_{\tilde{A}_{\alpha-n+j,j}^1(B_n)}$$

for all $f \in \tilde{A}_{\alpha-n+j,j}^1(B_n)$. Part (i) is proved.

Part (ii). Let $f \in \mathcal{K}_\alpha(n)$, $\alpha > n - j$. We have $\mathcal{R}^j f \in \mathcal{K}_{\alpha+j}(n)$ by Proposition 4.2. To finish the argument, it suffices to repeat the proof of part (ii) of Proposition 5.3. $\square$

5.8. Fractional Cauchy transforms and inner functions. Assume that $n \in \mathbb{N}$, $j \in \mathbb{N}$, and $q > 0$. The space $\tilde{A}_{q,j}^2(B_n)$ consists of all functions $f \in \mathcal{H}ol(B_n)$ such that

$$\int_0^1 \int_{\partial B_n} |\mathcal{R}^j f(r\zeta)|^2 (1-r)^{q-1} d\sigma_n(\zeta) dr < +\infty.$$

Lemma 5.9. Suppose $n \in \mathbb{N}$, $j \in \{1, \dots, n\}$, and $\beta > \alpha > n - j$. Then

$$H^\infty(B_n) \cap \mathcal{K}_\alpha(n) \subset \tilde{A}_{\beta-n+2j,j}^2(B_n).$$

Proof. Let $f \in H^\infty(B_n) \cap \mathcal{K}_\alpha(n)$. Since $f \in H^\infty(B_n)$, we have

$$f(z) = \int_{\partial B_n} \frac{f^*(\zeta)}{(1 - \langle z, \zeta \rangle)^n} d\sigma_n(\zeta), \quad z \in B_n.$$

Therefore,

$$\mathcal{R}f(z) = \int_{\partial B_n} \frac{n \langle z, \zeta \rangle f^*(\zeta)}{(1 - \langle z, \zeta \rangle)^{n+1}} d\sigma_n(\zeta), \quad z \in B_n.$$

Hence, using Lemma 3.1, we obtain

$$|\mathcal{R}f(z)| \leq \int_{\partial B_n} \frac{C}{|1 - \langle z, \zeta \rangle|^{n+1}} d\sigma_n(\zeta) \leq \frac{C}{(1 - |z|)}, \quad z \in B_n.$$

Arguing similarly, we verify that

$$(5.4) \quad |\mathcal{R}^j f(z)| \leq \frac{C}{(1 - |z|)^j}, \quad z \in B_n.$$

Since $f \in \mathcal{K}_\alpha(n)$, Proposition 4.2 guarantees that $\mathcal{R}^j f \in \mathcal{K}_{\alpha+j}(n)$. Since $\alpha + j > n$ and $\beta > \alpha$, part (ii) of Proposition 5.8 yields

$$\int_0^1 \int_{\partial B_n} |\mathcal{R}^j f(r\zeta)| (1-r)^{\beta-n+j-1} d\sigma_n(\zeta) dr < \infty.$$

Hence, applying inequality (5.4), we get

$$\int_0^1 \int_{\partial B_n} |\mathcal{R}^j f(r\zeta)| (1-r)^j \cdot |\mathcal{R}^j f(r\zeta)| (1-r)^{\beta-n+j-1} d\sigma_n(\zeta) dr < +\infty,$$

as required. $\square$

Recall that a nonconstant function $f \in H^\infty(B_n)$ is said to be *inner* if $|f^*| = 1$ σ_n -a.e. On the one hand, the families $\mathcal{K}_\alpha(n)$ with $\alpha \geq n$ contain all inner functions because $H^\infty(B_n) \subset H^1(B_n) \subset \mathcal{K}_n(n) \subset \mathcal{K}_\alpha(n)$. On the other hand, the following assertion is true for $n \geq 2$.

Proposition 5.10. *Let $n \geq 2$. If $0 \leq \alpha < n - 1/2$, then the family $\mathcal{K}_\alpha(n)$ contains no inner functions.*

Proof. Let $f \in \mathcal{K}_\alpha(n)$ be an inner function. By Corollary 5.5 and part (ii) of Corollary 2.3, there is no loss of generality in assuming that $\alpha \in (n - 1, n - 1/2)$.

Suppose $\rho \in [1/2, 1)$ and $f_\rho(\zeta) = f(\rho\zeta)$, $\zeta \in \partial B_n$. For $\zeta \in \partial B_n$ and $\lambda \in B_1$, put $f_\zeta(\lambda) = f(\lambda\zeta)$. Observe that $\mathcal{R}f(\lambda\zeta) = \lambda f'_\zeta(\lambda)$; hence,

$$(5.5) \quad |f^*(\zeta) - f_\rho(\zeta)| \leq 2 \int_\rho^1 |\mathcal{R}f(r\zeta)| dr$$

provided that $f^*(\zeta)$ is well defined. Applying estimate (5.5), Hölder's inequality, and Fubini's theorem, we obtain

$$\begin{aligned} \|f_\rho - f^*\|_{L^2(\partial B_n)}^2 &\leq C \int_{\partial B_n} \left(\int_\rho^1 |\mathcal{R}f(r\zeta)| (1-r)^{\frac{1}{4}} (1-r)^{-\frac{1}{4}} dr \right)^2 d\sigma_n(\zeta) \\ &\leq C(1-\rho)^{\frac{1}{2}} \int_{\partial B_n} \int_\rho^1 |\mathcal{R}f(r\zeta)|^2 (1-r)^{\frac{1}{2}} dr d\sigma_n(\zeta) \\ &= o(1-\rho)^{\frac{1}{2}} \quad \text{as } \rho \rightarrow 1 \end{aligned}$$

by Lemma 5.9. Tamm [27] proved that no inner function can satisfy the estimate $\|f_\rho - f^*\|_{L^2(\partial B_n)}^2 = o(1-\rho)^{\frac{1}{2}}$ as $\rho \rightarrow 1-$ if $n \geq 2$. This contradiction finishes the proof of the proposition. $\square$

§6. BOUNDARY BEHAVIOR

6.1. Directions of the maximal radial growth. The definition of the norm in the space $\mathcal{K}_\alpha(n)$, $\alpha > 0$, implies that

$$|f(z)| \leq \frac{\|f\|_{\mathcal{K}_\alpha(n)}}{(1-|z|)^\alpha}, \quad z \in B_n.$$

Proposition 6.1 shows that this maximal growth is possible for an at most countable set of radial directions. For $n = 1$, this was proved in [11].

Suppose $\xi \in \partial B_n$ and $C > 1$. Recall that the corresponding Korányi domain $D_C(\xi)$ is defined by

$$D_C(\xi) = \{z \in B_n : |1 - \langle z, \xi \rangle| < C(1 - |z|)\}.$$

Proposition 6.1. *Suppose $n \in \mathbb{N}$, $\mu \in M(n)$, $\alpha > 0$, and $f = K_\alpha[\mu]$. If $\xi \in \partial B_n$, then*

$$\lim_{\substack{z \rightarrow \xi \\ z \in D_C(\xi)}} (1 - \langle z, \xi \rangle)^\alpha f(z) = \mu(\{\xi\})$$

for all $C > 1$.

Proof. Let $\zeta, \xi \in \partial B_n$. We have

$$\lim_{\substack{z \rightarrow \xi \\ z \in B_n}} \frac{(1 - \langle z, \xi \rangle)^\alpha}{(1 - \langle z, \zeta \rangle)^\alpha} = \delta_{\zeta, \xi},$$

where $\delta_{\zeta, \xi} = 0$ for $\zeta \neq \xi$, and $\delta_{\xi, \xi} = 1$. If $z \in D_C(\xi)$, then

$$\left| \frac{1 - \langle z, \xi \rangle}{1 - \langle z, \zeta \rangle} \right|^\alpha \leq \frac{|1 - \langle z, \xi \rangle|^\alpha}{(1 - |z|)^\alpha} \leq C^\alpha.$$

Hence, if $f = K_\alpha[\mu]$, then

$$\lim_{\substack{z \rightarrow \xi \\ z \in D_C(\xi)}} (1 - \langle z, \xi \rangle)^\alpha f(z) = \lim_{\substack{z \rightarrow \xi \\ z \in D_C(\xi)}} \int_{\partial B_n} \frac{(1 - \langle z, \xi \rangle)^\alpha}{(1 - \langle z, \zeta \rangle)^\alpha} d\mu(\zeta) = \mu(\{\xi\})$$

by the dominated convergence theorem. $\square$

6.2. Radial behavior of functions in $\mathcal{K}_\alpha(n)$ with $\alpha > n$. Recall that the Bloch space $\mathfrak{B}(B_n)$ consists of the functions $f \in \mathcal{H}ol(B_n)$ such that

$$|\mathcal{R}f(z)| \leq \frac{C}{1 - |z|}, \quad z \in B_n,$$

for some constant $C > 0$ (see [30, Chapter 3] for equivalent definitions).

Proposition 6.2. *Suppose $n \in \mathbb{N}$ and $\alpha > n$. Then $\mathfrak{B}(B_n) \subset \mathcal{K}_\alpha(n)$.*

Proof. Let $f \in \mathfrak{B}(B_n)$, that is, $|\mathcal{R}f(r\zeta)| \leq C(1 - r)^{-1}$ for all $\zeta \in \partial B_n$ and $r \in [0, 1]$. The assumption $\alpha > n$ guarantees that

$$\int_0^1 \int_{\partial B_n} |\mathcal{R}f(r\zeta)| (1 - r)^{(\alpha+1)-n-1} d\sigma_n(\zeta) dr < +\infty.$$

Since $\alpha + 1 > n$, we have $\mathcal{R}f \in \mathcal{K}_{\alpha+1}(n)$ by Proposition 5.2. Finally, applying Corollary 5.6, we obtain $f \in \mathcal{K}_\alpha(n)$. $\square$

Corollary 6.3. *Let $n \in \mathbb{N}$. Then there exists a function $f \in \bigcap_{\alpha > n} \mathcal{K}_\alpha(n)$ such that the finite limit $\lim_{r \rightarrow 1-} f(r\zeta)$ fails to exist for all $\zeta \in \partial B_n$.*

Proof. Ullrich [28] constructed a function $f \in \mathfrak{B}(B_n)$ with no finite radial limits. It remains to apply Proposition 6.2. $\square$

6.3. Exceptional sets and Hausdorff contents: definitions. Let $0 < \tau \leq 1$, and let $\zeta \in \partial B_n$. Put

$$D_{\tau,C}(\zeta) = \{z \in B_n : |1 - \langle z, \zeta \rangle| < C(1 - |z|)^\tau\},$$

where $C > 1$ for $\tau = 1$, and $C > 0$ for $\tau \in (0, 1)$. Note that the sets $D_{1,C}(\zeta)$ coincide with the Korányi domains $D_C(\zeta)$, which were considered in Subsection 6.1. Also, note that the order of contact between the set $D_{\tau,C}(\zeta)$ and the sphere ∂B_n increases as the parameter τ decreases.

Consider a function $f : B_n \rightarrow \mathbb{C}$. By definition, the set $E_{\tau,C}(f)$ consists of all $\zeta \in \partial B_n$ for which the function $f(z)$ fails to have a limit at the point ζ as z approaches ζ through $D_{\tau,C}(\zeta)$. The sets $E_{1,C}(f)$ are said to be *exceptional*; the sets $E_{\tau,C}(f)$, $0 < \tau < 1$, are *tangentially exceptional*.

To estimate the size of the sets $E = E_{\tau,C}(f)$ for $f \in \mathcal{K}_\alpha(n)$, it is natural to apply the nonisotropic Hausdorff contents, which are defined as follows:

$$(6.1) \quad H_m(E) = H_m(E, \partial B_n) = \inf \left\{ \sum_k \delta_k^m : E \subset \bigcup_k Q(\zeta_k, \delta_k) \right\},$$

where $\zeta_k \in \partial B_n$ and $Q(\zeta, \delta) = \{\xi \in \partial B_n : |1 - \langle \zeta, \xi \rangle| < \delta\}$.

6.4. Hardy–Sobolev spaces. Assume that a function $f \in \mathcal{H}ol(B_n)$ has homogeneous expansion $f = \sum_k f_k$. Then the fractional derivative of order $\beta > 0$ is defined by the formula

$$R^\beta f(z) = \sum_k (k+1)^\beta f_k(z), \quad z \in B_n.$$

For $\beta > 0$ and $0 < p < \infty$, the Hardy–Sobolev spaces are defined by

$$H_\beta^p(B_n) = \left\{ f \in \mathcal{H}ol(B_n) : \sup_{0 < r < 1} \int_{\partial B_n} |R^\beta f(r\zeta)|^p d\sigma_n(\zeta) < \infty \right\}.$$

Theorem 6.4 (Ahern and Cohn [1]). *Let $n - \beta p \geq 0$. If $f \in H_\beta^p(B_n)$, then*

$$H_{n-\beta p+\varepsilon}(E(f)) = 0$$

for all $\varepsilon > 0$.

Theorem 6.5 (Cascante and Ortega [8, Corollary 2.1]; see also [26]). *Assume that $n \in \mathbb{N}$, $0 < p < \infty$, $0 < \beta < n/p$, $0 < \tau < 1$, and $m = (n - \beta p)/\tau$. If $f \in H_\beta^p(B_n)$, then $H_m(E_{\tau,C}(f)) = 0$ for all $C > 0$.*

6.5. Exceptional sets. By Corollary 6.3, there exists a function $f \in \bigcap_{\alpha > n} \mathcal{K}_\alpha(n)$ such that $E_{1,C}(f) = \partial B_n$ for all $C > 1$. On the other hand, if $0 \leq \alpha \leq n$, then $\mathcal{K}_\alpha(n) \subset H^p(B_n)$ for all $0 < p < 1$. Hence, if $f \in \mathcal{K}_\alpha(n)$, $0 \leq \alpha \leq n$, then $\sigma_n(E_{1,C}(f)) = 0$ for all $C > 1$. This observation can be refined in the case where $0 \leq \alpha < n$. For the spaces $\mathcal{K}_\alpha(1)$, $0 \leq \alpha < 1$, similar refinements in terms of the Bessel capacities were obtained in [12].

Proposition 6.6. *Suppose $n \in \mathbb{N}$ and $\alpha \in (0, n)$. If $f \in \mathcal{K}_\alpha(n)$, then $H_{\alpha+\varepsilon}(E_{1,C}(f)) = 0$ for all $\varepsilon > 0$ and $C > 1$.*

Proof. Let $n - \alpha > \varepsilon > 0$. Then $\mathcal{K}_\alpha(n) \subset A_{\alpha+\varepsilon,n}^1(B_n) \subset H_{n-\alpha-\varepsilon}^1(B_n)$ by Proposition 5.3 and by [4, Theorem 5.12], respectively. It remains to apply Theorem 6.4. $\square$

The following example corresponds to Proposition 6.6.

Proposition 6.7. *Suppose $n \in \mathbb{N}$, $\alpha \in (0, n)$, $0 \leq m < \alpha$, $E \subset \partial B_n$ is a compact set, $H_m(E) = 0$, and $C > 1$. Then $E = E_{1,C}(f)$ for some function $f \in \mathcal{K}_\alpha(n)$.*

Proof. Fix $p_0 > 1$ such that $m = n - (n - \alpha)p_0$. We have $H_m(E) = 0$; hence, Theorem 1.2 in [1] guarantees that $E = E_{1,C}(f)$ for some function $f \in H_{n-\alpha}^{p_0}(B_n)$. Next, recall that

$$\mathcal{K}_\alpha^{p_0}(n) = \{K_\alpha[g\sigma_n] : g \in L^{p_0}(\sigma_n)\}.$$

By [8, Theorem 2.1], we have $H_{n-\alpha}^{p_0}(B_n) = \mathcal{K}_\alpha^{p_0}(n)$. It remains to note that $\mathcal{K}_\alpha^{p_0}(n) \subset \mathcal{K}_\alpha(n)$. $\square$

6.6. Tangentially exceptional sets. For the spaces $\mathcal{K}_\alpha(1)$, $0 \leq \alpha < 1$, tangentially exceptional sets were investigated in [10]. In the case of an arbitrary dimension $n \in \mathbb{N}$, the following assertion holds true.

Proposition 6.8. *Suppose $n \in \mathbb{N}$, $\alpha \in (0, n)$, $\tau \in (\alpha/n, 1)$, and $m > \alpha/\tau$. If $f \in \mathcal{K}_\alpha(n)$, then $H_m(E_{\tau,C}(f)) = 0$ for all $C > 0$.*

Proof. We have $\mathcal{K}_\alpha(n) \subset H_{n-\alpha-\varepsilon}^1(B_n)$ for $n - \alpha > \varepsilon > 0$. It remains to apply Theorem 6.5. $\square$

The following example corresponds to Proposition 6.8.

Proposition 6.9. *Suppose $n \in \mathbb{N}$, $\alpha \in (0, n)$, $\tau \in [\alpha/n, 1)$, $m < \alpha/\tau$, and $C > 0$. If $E \subset \partial B_n$ is a compact set and $H_m(E) = 0$, then $E = E_{\tau,C}(f)$ for some function $f \in \mathcal{K}_\alpha(n)$.*

Proof. Since $m < \alpha/\tau$, there exists $p_0 > 1$ such that

$$m = (n - (n - \alpha)p_0)/\tau.$$

Since $p_0 > 1$ and $H_m(E) = 0$, we have $E = E_{\tau,C}(f)$ for some function $f \in \mathcal{K}_\alpha^{p_0}(n)$; see [26, Remark after Corollary 3.11]. It remains to note that $\mathcal{K}_\alpha^{p_0}(n) \subset \mathcal{K}_\alpha(n)$. $\square$

§7. MULTIPLIERS

In this section, we study the spaces $\mathfrak{M}_\alpha(n)$ that consist of multipliers for the families $\mathcal{K}_\alpha(n)$. Note that multipliers for the Hardy–Sobolev spaces were investigated in the recent papers [22, 5] and [25].

7.1. Necessary conditions. For $n = 1$, the results of this subsection were obtained in [16].

Lemma 7.1. *Suppose $n \in \mathbb{N}$, $\alpha > 0$, and $f \in \mathfrak{M}_\alpha(n)$. Then $f \in H^\infty(B_n)$; moreover, $\|f\|_{H^\infty(B_n)} \leq \|f\|_{\mathfrak{M}_\alpha(n)}$.*

Proof. Fix a point $\zeta \in \partial B_n$ and fix a constant K for which $\|f\|_{\mathfrak{M}_\alpha(n)} < K$. Since $\|(1 - \langle \cdot, \zeta \rangle)^{-\alpha}\|_{\mathcal{K}_\alpha(n)} = 1$, there exists a measure $\mu_\zeta \in M(n)$ such that $\|\mu_\zeta\|_{M(n)} < K$ and

$$\frac{f(z)}{(1 - \langle z, \zeta \rangle)^\alpha} = \int_{\partial B_n} \frac{1}{(1 - \langle z, \xi \rangle)^\alpha} d\mu_\zeta(\xi).$$

In other words,

$$f(z) = \int_{\partial B_n} \left(\frac{1 - \langle z, \zeta \rangle}{1 - \langle z, \xi \rangle} \right)^\alpha d\mu_\zeta(\xi).$$

Put $z = r\zeta$, $0 \leq r < 1$. Then

$$|f(r\zeta)| \leq \int_{\partial B_n} \left(\frac{1 - r}{1 - r|\langle \zeta, \xi \rangle|} \right)^\alpha d\mu_\zeta(\xi) \leq \|\mu_\zeta\|_{M(n)} < K.$$

Since the point $\zeta \in \partial B_n$ and the constant $K > \|f\|_{\mathfrak{M}_\alpha(n)}$ are arbitrary, we have $\|f\|_{H^\infty(B_n)} \leq \|f\|_{\mathfrak{M}_\alpha(n)}$. $\square$

Lemma 7.2. *Suppose $n \in \mathbb{N}$, $\alpha > 0$, and $f \in \mathfrak{M}_\alpha(n)$. Then $f \in \mathcal{K}_\alpha(n)$; moreover, $\|f\|_{\mathcal{K}_\alpha(n)} \leq \|f\|_{\mathfrak{M}_\alpha(n)}$.*

Proof. Put $I(z) = 1$ for $z \in B_n$. Recall that $I = K_\alpha[\sigma_n]$. Since σ_n is a positive measure, we have $\|I\|_{\mathcal{K}_\alpha(n)} = \|\sigma_n\| = 1$. The assumption $f \in \mathfrak{M}_\alpha(n)$ implies that $f = fI \in \mathcal{K}_\alpha(n)$. Moreover, $\|f\|_{\mathcal{K}_\alpha(n)} = \|fI\|_{\mathcal{K}_\alpha(n)} \leq \|f\|_{\mathfrak{M}_\alpha(n)}\|I\|_{\mathcal{K}_\alpha(n)} = \|f\|_{\mathfrak{M}_\alpha(n)}$. $\square$

Proposition 7.3. *Let $\alpha > 0$. Then there exists a constant C_α such that, for any $f \in \mathfrak{M}_\alpha(n)$ and any $n \in \mathbb{N}$, the radial variation in the direction $\xi \in \partial B_n$ satisfies the estimate $V(f, \xi) \leq C_\alpha \|f\|_{\mathfrak{M}_\alpha}$ for all $\xi \in \partial B_n$.*

Proof. Suppose $f \in \mathfrak{M}_\alpha(n)$, $\xi \in \partial B_n$, and $\varepsilon > 0$. By the definition of the space $\mathfrak{M}_\alpha(n)$, there exists a measure μ_ξ such that $\|\mu_\xi\|_{M(n)} \leq \|f\|_{\mathfrak{M}_\alpha(n)} + \varepsilon$ and

$$f(z) = \int_{\partial B_n} \frac{(1 - \langle z, \xi \rangle)^\alpha}{(1 - \langle z, \zeta \rangle)^\alpha} d\mu_\xi(\zeta), \quad z \in B_n.$$

Let $\lambda \in B_1$. We put $z = \lambda\xi$ and $f_\xi(\lambda) = f(\lambda\xi)$. Then

$$\begin{aligned} f_\xi(\lambda) &= \int_{\partial B_n} \frac{(1 - \lambda)^\alpha}{(1 - \lambda\langle \xi, \zeta \rangle)^\alpha} d\mu_\xi(\zeta), \\ f'_\xi(\lambda) &= \alpha \int_{\partial B_n} \frac{(1 - \lambda)^{\alpha-1}(\langle \xi, \zeta \rangle - 1)}{(1 - \lambda\langle \xi, \zeta \rangle)^{\alpha+1}} d\mu_\xi(\zeta). \end{aligned}$$

Now, set $\lambda = r$, $0 \leq r < 1$. Fubini's theorem guarantees that

$$(7.1) \quad \int_0^1 |f'_\xi(r)| dr \leq \alpha \int_{\partial B_n} \int_0^1 \frac{(1 - r)^{\alpha-1} |1 - \langle \xi, \zeta \rangle|}{|1 - r\langle \xi, \zeta \rangle|^{\alpha+1}} dr d|\mu_\xi|(\zeta).$$

Put $w = \langle \xi, \zeta \rangle$ and $b = |1 - \langle \xi, \zeta \rangle|$. Observe that

$$|1 - rw|^2 = (1 - r)(1 - r|w|^2) + r|1 - w|^2 \geq (1 - r)^2 + r^2b^2.$$

Hence,

$$\int_0^1 \frac{(1 - r)^{\alpha-1}b}{|1 - r\langle \xi, \zeta \rangle|^{\alpha+1}} dr \leq \int_0^1 \frac{(1 - r)^{\alpha-1}b}{((1 - r)^2 + r^2b^2)^{(\alpha+1)/2}} dr := I(\alpha, b).$$

If $b = 0$, then the proof is finished. Otherwise, as was shown in the proof of Theorem 2.6 in [16], we have $I(\alpha, b) \leq C_\alpha < \infty$. Therefore, estimate (7.1) implies that $V(f, \xi) \leq C_\alpha \|f\|_{\mathfrak{M}_\alpha(n)}$. The proof of the proposition is finished. $\square$

Recall that a bounded nonconstant function $f \in \text{Hol}(B_n)$ is said to be *inner* if $|f^*| = 1$ σ_n -a.e. A complete description of the inner functions belonging to $\mathfrak{M}_1(1)$ was obtained in [18]. For arbitrary $\alpha > 0$, the inner functions in the family $\mathfrak{M}_\alpha(1)$ were investigated in [13]. For $n \geq 2$, the answer to the corresponding question follows from Proposition 7.3.

Corollary 7.4. *Assume that $n \geq 2$ and $\alpha > 0$. If f is an inner function in the ball B_n , then $f \notin \mathfrak{M}_\alpha(n)$.*

Proof. For $n \geq 2$, it is well known that any inner function has no radial limits on a dense subset of the sphere ∂B_n ; see [24, §1]. $\square$

7.2. Sufficient conditions. Recall that $A(B_n)$ denotes the ball algebra. If $f \in A(B_n)$, then put $f^* = f|_{\partial B_n}$. The complex modulus of continuity of the function f^* is defined by

$$\omega_{\mathbb{C}}(f^*, \delta) = \sup_{\zeta, \xi \in \partial B_n} \{|f^*(\zeta) - f^*(\xi)| : d(\zeta, \xi) \leq \delta\},$$

where $\delta \in (0, 2]$ and $d(\zeta, \xi) = |1 - \langle \zeta, \xi \rangle|$. Note that $d(\zeta, \xi)$ is a quasimetric on the sphere.

Let ν_n denote the Lebesgue measure on B_n normalized by the condition $\nu_n(B_n) = 1$. Suppose that $n \geq 2$. For a Borel function $f : B_1 \rightarrow \mathbb{C}$, it is well known that the identity

$$(7.2) \quad \int_{\partial B_n} f(\langle \xi, \zeta \rangle) d\sigma_n(\xi) = (n-1) \int_{B_1} f(z)(1-|z|^2)^{n-2} d\nu_1(z)$$

is fulfilled if its left-hand side or right-hand side is well defined.

An analog of the following statement for $n = 1$ was proved in [29].

Proposition 7.5. *Let $\alpha \geq n \geq 2$. Suppose that a function $g \in A(B_n)$ satisfies the following condition:*

$$(7.3) \quad (n-1) \int_{B_1} \frac{(1-|z|^2)^{n-2} \omega_{\mathbb{C}}(g^*, |1-z|)}{|1-z|^n} d\nu_1(z) = C_{\omega} < +\infty.$$

Then $g \in \mathfrak{M}_{\alpha}(n)$.

Proof. By Lemma 2.4 and Proposition 2.5, it suffices to verify that the function g satisfies condition (2.5) with $\alpha = n$.

Fix a point $\zeta \in \partial B_n$. Then

$$\frac{g(w)}{(1-\langle w, \zeta \rangle)^n} = \frac{g(w) - g(\zeta)}{(1-\langle w, \zeta \rangle)^n} + \frac{g(\zeta)}{(1-\langle w, \zeta \rangle)^n} := h_1(w) + h_2(w), \quad w \in B_n.$$

Observe that $\|h_2\|_{\mathcal{K}_n(n)} \leq \|g\|_{A(B_n)}$.

On the other hand, we have $g(\cdot) - g(\zeta) \in H^{\infty}(B_n)$ and $(1-\langle \cdot, \zeta \rangle)^{-n} \in H^p(B_n)$ for all $0 < p < 1$. Hence, $h_1 \in H^p(B_n)$ for all $0 < p < 1$. Let $h_1^*(\xi) = \lim_{r \rightarrow 1-} h_1(r\xi)$, $\xi \in \partial B_n$, $\xi \neq \zeta$. Formula (7.2) and condition (7.3) show that $\|h_1^*\|_{L^1(\partial B_n)} \leq C_{\omega}$. Therefore, $h_1 \in H^1(B_n)$; also, we obtain $\|h_1\|_{\mathcal{K}_n(n)} \leq \|h_1\|_{H^1(B_n)} \leq C_{\omega}$. This proves (2.5). $\square$

Assume that $0 < \beta < 1$. By definition, the standard Lipschitz space $\Lambda^{\beta}(\partial B_n)$ consists of all functions $f : \partial B_n \rightarrow \mathbb{C}$ such that

$$|f(\zeta) - f(\xi)| \leq C_f |\zeta - \xi|^{\beta}$$

for all $\zeta, \xi \in \partial B_n$. In other words, $\Lambda^{\beta}(\partial B_n)$ are the Lipschitz spaces with respect to the Euclidean metric on the sphere. Elements of these spaces can be used to construct examples of functions belonging to families $\mathfrak{M}_{\alpha}(n)$ with $\alpha \geq n$.

Corollary 7.6. *Suppose $n \geq 2$, $0 < \beta < 1/2$, and $f \in \Lambda^{\beta}(\partial B_n)$. Then $K_n[f] \in \mathfrak{M}_{\alpha}(n)$ for $\alpha \geq n$.*

Proof. Let $f \in \Lambda^{\beta}(\partial B_n)$. It is well known that $\omega_{\mathbb{C}}(K_n[f]^*, \delta) = \mathcal{O}(\delta^{\beta})$ (see [2]). Hence, condition (7.3) is satisfied. $\square$

Explicit examples of functions belonging to $\mathfrak{M}_{\alpha}(n)$, $\alpha \geq 0$, can be obtained with the help of the following assertion.

Proposition 7.7. *Suppose $n \in \mathbb{N}$, $j \in \{0, 1, \dots, n\}$, and $\alpha \geq n - j$. Also, suppose that the derivative $\mathcal{R}^j g \in A(B_n)$ satisfies condition (7.3). Then $g \in \mathfrak{M}_{\alpha}(n)$.*

Proof. We argue by induction. As the base of induction, we take the case of $j = 0$. Then the claim is true by Proposition 7.5.

Now, let $k \in \{0, 1, \dots, n-1\}$. Assume that the required implication is proved for $j = k$.

Suppose $j = k + 1 \in \{1, \dots, n\}$ and $f \in \mathcal{K}_\alpha(n)$, $\alpha \geq n - j$. We must prove that $fg \in \mathcal{K}_\alpha(n)$. Since $\alpha \geq n - j \geq 0$, Corollary 5.6 guarantees that $fg \in \mathcal{K}_\alpha(n)$ if and only if $\mathcal{R}(fg) \in \mathcal{K}_{\alpha+1}(n)$. Note that

$$\mathcal{R}(fg) = f\mathcal{R}g + g\mathcal{R}f.$$

The function $\mathcal{R}^{j-1}(\mathcal{R}g) \in A(B_n)$ satisfies condition (7.3). We have $\alpha + 1 \geq n - (j - 1)$; hence, $\mathcal{R}g \in \mathfrak{M}_{\alpha+1}(n)$ by the inductive hypothesis. Since $f \in \mathcal{K}_\alpha(n) \subset \mathcal{K}_{\alpha+1}(n)$, we obtain $f\mathcal{R}g \in \mathcal{K}_{\alpha+1}(n)$ by the definition of the multiplier space.

On the other hand, $f \in \mathcal{K}_\alpha(n)$ if and only if $\mathcal{R}f \in \mathcal{K}_{\alpha+1}(n)$. Next, the function $\mathcal{R}^{j-1}g \in A(B_n)$ satisfies condition (7.3). By the inductive hypothesis, we have $g \in \mathfrak{M}_{\alpha+1}(n)$; thus, $g\mathcal{R}f \in \mathcal{K}_{\alpha+1}(n)$.

Therefore, $\mathcal{R}(fg) \in \mathcal{K}_{\alpha+1}(n)$ and $fg \in \mathcal{K}_\alpha(n)$. The induction step is finished, and the proposition is proved. $\square$

By using the family $\mathcal{K}_0(n)$, other examples of multipliers can be obtained.

Proposition 7.8. *Suppose $n \in \mathbb{N}$, $0 < \alpha < n$, and $g \in \mathcal{K}_0(n) \cap \mathfrak{M}_n(n)$. Then $g \in \mathfrak{M}_\alpha(n)$.*

Proof. We argue by induction. As the base of induction, we take the statement that $g \in \mathfrak{M}_\alpha(n)$ for $\alpha \geq n$, which is true by assumption.

Now, suppose that $j \in \{0, 1, \dots, n-1\}$. Assume that the required implication is valid for $\alpha \geq n - j$. We prove that $g \in \mathfrak{M}_\alpha(n)$ if $\alpha > 0$ and $\alpha \geq n - j - 1$.

Let $f \in \mathcal{K}_\alpha(n)$. We must check that $fg \in \mathcal{K}_\alpha(n)$.

The condition $g \in \mathcal{K}_0(n)$ implies that $\mathcal{R}g \in \mathcal{K}_1(n)$. Since $\alpha > 0$, we obtain $f\mathcal{R}g \in \mathcal{K}_\alpha(n) \cdot \mathcal{K}_1(n) \subset \mathcal{K}_{\alpha+1}(n)$. On the other hand, $\mathcal{R}f \in \mathcal{K}_{\alpha+1}(n)$ because $f \in \mathcal{K}_\alpha(n)$. Next, we have $g \in \mathfrak{M}_{n-j}(n)$ by the inductive hypothesis. Also, $\mathfrak{M}_{n-j}(n) \subset \mathfrak{M}_{\alpha+1}(n)$ because $n - j \leq \alpha + 1$. Thus, $g\mathcal{R}f \in \mathfrak{M}_{\alpha+1}(n) \cdot \mathcal{K}_{\alpha+1}(n) \subset \mathcal{K}_{\alpha+1}(n)$.

Therefore, $\mathcal{R}(fg) = f\mathcal{R}g + g\mathcal{R}f \in \mathcal{K}_{\alpha+1}(n)$ and $fg \in \mathcal{K}_\alpha(n)$ by Corollary 5.6. The induction step is finished. The proof of the proposition is complete. $\square$

Corollary 7.9. *Suppose $n \in \mathbb{N}$ and $0 < \alpha < n$. Also, suppose that a function $g \in \mathcal{K}_0(n) \cap A(B_n)$ satisfies condition (7.3). Then $g \in \mathfrak{M}_\alpha(n)$.*

Proof. It suffices to apply Propositions 7.5 and 7.8. $\square$

ACKNOWLEDGMENT

The author is grateful to A. B. Aleksandrov for suggested improvements and useful comments.

REFERENCES

- [1] P. Ahern and W. Cohn, *Exceptional sets for Hardy–Sobolev functions*, $p > 1$, Indiana Univ. Math. J. **38** (1989), no. 2, 417–453. MR0997390 (90m:32013)
- [2] P. Ahern and R. Schneider, *A smoothing property of the Henkin and Szegő projections*, Duke Math. J. **47** (1980), no. 1, 135–143. MR0563371 (80i:32015)
- [3] A. B. Aleksandrov, *Function theory in the ball*, Itogi Nauki i Tekhniki, Sovrem. Probl. Mat. Fund. Naprav., vol. 8, VINITI, Moscow, 1985, pp. 115–190; English transl., Encyclopaedia Math. Sci., vol. 8, Springer-Verlag, Berlin, 1994, pp. 107–178. MR0850487 (88b:32002)
- [4] F. Beatrous and J. Burbea, *Holomorphic Sobolev spaces on the ball*, Diss. Math. (Rozprawy Mat.) **276** (1989), 60 pp. MR1010151 (90k:32010)
- [5] ———, *On multipliers for Hardy–Sobolev spaces*, Proc. Amer. Math. Soc. **136** (2008), no. 6, 2125–2133. MR2383518 (2009c:32007)
- [6] J. Bergh and J. Löfström, *Interpolation spaces. An introduction*, Grundlehren Math. Wiss., Bd. 223, Springer-Verlag, Berlin, 1976. MR0482275 (58:2349)

- [7] L. Brickman, D. J. Hallenbeck, T. H. MacGregor, and D. R. Wilken, *Convex hulls and extreme points of families of starlike and convex mappings*, Trans. Amer. Math. Soc. **185** (1973), 413–428. MR0338337 (49:3102)
- [8] C. Cascante and J. M. Ortega, *Tangential-exceptional sets for Hardy–Sobolev spaces*, Illinois J. Math. **39** (1995), no. 1, 68–85. MR1299649 (95j:32009)
- [9] J. A. Cima, A. L. Matheson, and W. T. Ross, *The Cauchy transform*, Math. Surveys Monogr., vol. 125, Amer. Math. Soc., Providence, RI, 2006. MR2215991 (2006m:30003)
- [10] D. J. Hallenbeck, *Tangential limits of Cauchy–Stieltjes transforms*, Complex Variables Theory Appl. **33** (1997), no. 1–4, 129–136. MR1624915 (99b:31001)
- [11] D. J. Hallenbeck and T. H. MacGregor, *Growth and zero sets of analytic families of Cauchy–Stieltjes integrals*, J. Anal. Math. **61** (1993), 231–259. MR1253443 (95d:30078)
- [12] ———, *Radial limits and radial growth of Cauchy–Stieltjes transforms*, Complex Variables Theory Appl. **21** (1993), no. 3–4, 219–229. MR1276578 (95d:30077)
- [13] D. J. Hallenbeck, T. H. MacGregor, and K. Samotij, *Fractional Cauchy transforms, inner functions and multipliers*, Proc. London Math. Soc. (3) **72** (1996), no. 1, 157–187. MR1357091 (96j:30056)
- [14] V. P. Havin, *On analytic functions representable by an integral of Cauchy–Stieltjes type*, Vestnik Leningrad. Univ. Mat. Mekh. Astronom. **1958**, vyp. 1, 66–79. (Russian) MR0095256 (20:1762)
- [15] ———, *Relations between certain classes of functions regular in the unit circle*, Vestnik Leningrad. Univ. Mat. Mekh. Astronom. **1962**, vyp. 1, 102–111. (Russian) MR0152660 (27:2635)
- [16] R. A. Hirschweiler and T. H. MacGregor, *Multipliers of families of Cauchy–Stieltjes transforms*, Trans. Amer. Math. Soc. **331** (1992), no. 1, 377–394. MR1120775 (93g:30057)
- [17] ———, *Fractional Cauchy transforms*, Chapman & Hall/CRC Monogr. Surveys in Pure Appl. Math., vol. 136, Chapman & Hall/CRC, Boca Raton, FL, 2006. MR2189294 (2007b:30001)
- [18] S. V. Hruščev and S. A. Vinogradov, *Inner functions and multipliers of Cauchy type integrals*, Ark. Mat. **19** (1981), no. 1, 23–42. MR0625535 (83c:30027)
- [19] V. G. Krotov, *Estimates for maximal operators that are connected with boundary behavior and their applications*, Trudy Mat. Inst. Steklov. **190** (1989), 117–138; English transl., Proc. Steklov Inst. Math. **1992**, no. 1, 123–144. MR1005342 (90h:42030)
- [20] T. H. MacGregor, *Analytic and univalent functions with integral representations involving complex measures*, Indiana Univ. Math. J. **36** (1987), no. 1, 109–130. MR0876994 (87m:30037)
- [21] A. Nagel, W. Rudin, and J. H. Shapiro, *Tangential boundary behavior of functions in Dirichlet-type spaces*, Ann. of Math. (2) **116** (1982), no. 2, 331–360. MR0672838 (84a:31002)
- [22] J. M. Ortega and J. Fàbrega, *Multipliers in Hardy–Sobolev spaces*, Integral Equations Operator Theory **55** (2006), no. 4, 535–560. MR2250162 (2007f:46034)
- [23] W. Rudin, *Function theory in the unit ball of $\mathbf{C}^n$* , Grundlehren Math. Wiss., Bd. 241, Springer-Verlag, New York–Berlin, 1980. MR0601594 (82i:32002)
- [24] ———, *New constructions of functions holomorphic in the unit ball of $\mathbf{C}^n$* , CBMS Regional Conf. Ser. in Math., vol. 63, Published for the Conference Board of the Mathematical Sciences, Washington, DC, Amer. Math. Soc., Providence, RI, 1986. MR0840468 (87f:32013)
- [25] P. Ryan and M. Stoll, *Hardy–Sobolev spaces and algebras of holomorphic functions on the unit ball in $\mathbf{C}^n$* , Complex Var. Elliptic Equ. **53** (2008), no. 6, 565–584. MR2421818 (2009b:46111)
- [26] J. Sueiro, *Tangential boundary limits and exceptional sets for holomorphic functions in Dirichlet-type spaces*, Math. Ann. **286** (1990), no. 4, 661–678. MR1045395 (91b:32008)
- [27] M. Tamm, *Sur l’image par une fonction holomorphe bornée du bord d’un domaine pseudoconvexe*, C. R. Acad. Sci. Paris Sér. I Math. **294** (1982), no. 16, 537–540. MR0679938 (84g:32008)
- [28] D. C. Ullrich, *A Bloch function in the ball with no radial limits*, Bull. London Math. Soc. **20** (1988), no. 4, 337–341. MR0940289 (89k:32009)
- [29] S. A. Vinogradov, M. G. Goluzina, and V. P. Khavin, *Multipliers and divisors of Cauchy–Stieltjes type integrals*, Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) **19** (1970), 55–78; English transl., Sem. in Math. Steklov Math. Inst., Leningrad, vol. 19, Consultants Bureau, New York–London, 1972, pp. 29–42. MR0291471 (45:562)
- [30] K. Zhu, *Spaces of holomorphic functions in the unit ball*, Grad. Texts in Math., vol. 226, Springer-Verlag, New York, 2005. MR2115155 (2006d:46035)

ST. PETERSBURG BRANCH, STEKLOV MATHEMATICAL INSTITUTE, FONTANKA 27, ST. PETERSBURG 191023, RUSSIA

E-mail address: dubtsov@pdmi.ras.ru

Received 23/NOV/2008

Translated by THE AUTHOR