REDUCIBILITY OF FUNCTION PAIRS IN $H_{\mathbb{R}}^{\infty}$

RAYMOND MORTINI

ABSTRACT. A short proof of a result by Brett Wick on the reducibility of function pairs in $H_{\mathbb{R}}^{\infty}$ is presented, and some unusual properties of the solutions to the associated Bézout equations are unveiled.

§1. THE PROBLEM

Let $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ be the unit disk, and let H^{∞} be the (complex) algebra of all bounded holomorphic functions on $\mathbb{D}$. Moreover, let $H_{\mathbb{R}}^{\infty}$ be the (real) algebra of all bounded holomorphic functions in $\mathbb{D}$ that satisfy the symmetry condition $f(z) = \overline{f(\bar{z})}$ for $z \in \mathbb{D}$. It is well known that $H_{\mathbb{R}}^{\infty}$ is the set of functions in H^{∞} with real Fourier coefficients, and that $H_{\mathbb{R}}^{\infty}$ coincides with the set of all functions in H^{∞} that are real-valued on $] -1, 1[$. This algebra gained a lot of interest the last years in view of its appearance in control theory (see [13, 14, 9]). It is a direct consequence of the corona theorem for H^{∞} that if (f, g) is a function pair in $H_{\mathbb{R}}^{\infty}$ with $\delta := \inf_{z \in \mathbb{D}} |f(z)| + |g(z)| > 0$, then there exist $p, q \in H_{\mathbb{R}}^{\infty}$ such that $pf + qg = 1$ (see [14]). On the other hand, in contrast to the H^{∞} -setting (see [12]), it is not always possible to choose the first factor p to be invertible in $H_{\mathbb{R}}^{\infty}$. In fact, as was already mentioned in [13], if $f(z) = z$ and $g(z) = 1 - z^2$, then each function of the form $F := z + h(1 - z^2)$ with $h \in H_{\mathbb{R}}^{\infty}$ must have a zero within $] -1, 1[$, because F is real-valued on $] -1, 1[$, $F(-1) < 0$ and $F(1) > 0$.

It is therefore a natural problem to characterize the pairs (f, g) in $H_{\mathbb{R}}^{\infty}$ that are reducible. Recall that a pair (f, g) in a commutative, unital Banach algebra A is said to be *invertible* if $af + bg = 1$ for some $a, b \in A$ and that (f, g) is *reducible* if there exists $h \in A$ such that $f + hg$ is invertible in A . In other words, (f, g) is reducible if there exists a solution (u, h) of the Bézout equation $uf + hg = 1$ such that the first factor u is invertible.

Motivated by his earlier result [13] for the algebra $A_{\mathbb{R}}(\mathbb{D})$ of functions in $H_{\mathbb{R}}^{\infty}$ that admit continuous extensions to the Euclidean closure of $\mathbb{D}$, in a series of papers [14, 15] Brett Wick developed the following theorem.

Theorem W (see [15]). *Let $(f, g) \in (H_{\mathbb{R}}^{\infty})^2$ be a function pair satisfying $\|f\|_{\infty} \leq 1$, $\|g\|_{\infty} \leq 1$ and $|f| + |g| \geq \delta > 0$. Suppose that there exists $\varepsilon > 0$ such that f has constant sign on the set $\{x \in] -1, 1[: |g(x)| < \varepsilon\}$. Then there exists a constant $C = C(\delta, \varepsilon)$ depending only on δ and ε such that the Bézout equation*

$$uf + hg = 1$$

admits a solution $(u, h) \in (H_{\mathbb{R}}^{\infty})^2$ with u invertible and

$$\Delta(u, u^{-1}, h) := \|u\|_{\infty} + \|u^{-1}\|_{\infty} + \|h\|_{\infty} \leq C(\delta, \varepsilon).$$

Note that the hypothesis on the sign is satisfied automatically whenever $|g|$ is bounded away from zero on $] -1, 1[$ (in that case for ε we can take any number strictly smaller than

2010 *Mathematics Subject Classification*. Primary 30H05, 46H15, 30J10, 30H80.

Key words and phrases. Corona theorem, Bézout equation, Blaschke product.

$\inf_{x \in]-1, 1[} |g(x)|$. Also, if (f, g) is reducible in $H_{\mathbb{R}}^{\infty}$, then the sign condition is fulfilled necessarily (see below).

The intention of our paper is two-fold. First, in §2 we give a new short proof of the reducibility part of this theorem. Our proof will be based on the maximal ideal space theory for H^{∞} and on Treil's theorem [12] on the reducibility of pairs in H^{∞} . The original proof by Wick involved the $\bar{\partial}$ -calculus and depended on the methods developed by Treil to solve Bézout equations.

Then, in §3, we present a new proof of Theorem **W** with norm estimates that depends only on the corresponding result by Treil [12], but is otherwise quite elementary.

In §4, we provide some counterexamples to the original statement of Wick's result [14]. First, we give examples of invertible pairs (f, g) that show that, in general, the weaker hypothesis of f having constant sign on the set of real zeros of g is not sufficient in order to get a solution of the Bézout equation

$$uf + hg = 1, \quad u \text{ invertible in } H_{\mathbb{R}}^{\infty}.$$

We mention that in [9, Proposition 2.8] it was shown that if g is a Blaschke product having only real and simple zeros, then the hypothesis of f being positive on the set of real zeros of g is indeed sufficient to guarantee the reducibility of the pair (whenever (f, g) is invertible).

Then we show that there exist reducible pairs (f_n, g_n) of functions in the unit ball of $H_{\mathbb{R}}^{\infty}$ with $|f_n| + |g_n| \geq \delta > 0$ such that, quite surprisingly,

$$\sup_n \Delta(u_n, u_n^{-1}, h_n) = \infty$$

for any solutions (u_n, h_n) of the equation $uf_n + hg_n = 1$ with u invertible in $H_{\mathbb{R}}^{\infty}$.

This shows that in Wick's Theorem **W** the parameter ε plays an essential role. To the best of our knowledge, this is the first occurrence of a corona-type theorem for a uniform algebra where the uniform boundedness of solutions fails.

§2. A MAXIMAL IDEAL SPACE VIEW OF WICK'S RESULT

Let $M(H^{\infty})$ be the spectrum (or maximal ideal) space of H^{∞} ; that is, $M(H^{\infty})$ is the space of all (nonzero) multiplicative linear functionals on H^{∞} endowed with the trace of the weak-* topology $\sigma((H^{\infty})^*, H^{\infty})$, also called the Gelfand topology. Identifying each point z_0 in $\mathbb{D}$ with the evaluation functional $\Phi_{z_0} : f \mapsto f(z_0)$, we can identify $\mathbb{D}$ with a subset of $M(H^{\infty})$. By Carleson's famous corona theorem, $\mathbb{D}$ is dense in $M(H^{\infty})$. The Gelfand transform $\hat{f} : M(H^{\infty}) \rightarrow \mathbb{C}$ of an element $f \in H^{\infty}$ is given by $\hat{f}(m) = m(f)$. It can be viewed as a unique continuous extension of f from $\mathbb{D}$ to $M(H^{\infty})$. As usual, we shall identify $\hat{f}$ with f .

A corona pair in $A = H^{\infty}$ (or $A = H_{\mathbb{R}}^{\infty}$) is a pair (f, g) of functions in A such that $\inf_{z \in \mathbb{D}} |f(z)| + |g(z)| \geq \delta > 0$. The zero set of $f \in H^{\infty}$ is given by $Z(f) = \{m \in M(H^{\infty}) : f(m) = 0\}$. Moreover, let $Z_{\mathbb{D}}(f) = \{z \in \mathbb{D} : f(z) = 0\}$. It is well known that if b is an interpolating Blaschke product, then $Z(b)$ is the closure of $Z_{\mathbb{D}}(b)$ (see [3, p. 379]); in general, for arbitrary Blaschke products B , the set $\overline{Z_{\mathbb{D}}(B)}$ is much smaller than $Z(B)$ (see [4] for a study of the inner functions Θ for which $Z(\Theta) = \overline{Z_{\mathbb{D}}(\Theta)}$).

In the sequel, we may identify $M(H^{\infty})$ with $M(H_{\mathbb{R}}^{\infty})$ (because H^{∞} is the complexification of $H_{\mathbb{R}}^{\infty}$; see [6, pp. 37–38]), and we look upon the functions in $H_{\mathbb{R}}^{\infty}$ as being defined on $M(H^{\infty})$.

A prominent role in our approach to Wick's result will be played by the closure E of the interval $] -1, 1[$ in $M(H^{\infty})$. If $g \in H_{\mathbb{R}}^{\infty}$, then the set of *extended real zeros* of g is the set $Z(g) \cap E$.

The following topological observation will give a relationship between the hypotheses in Theorem **W** and our approach (see [9, p. 178]).

Observation 2.1. Let $(f, g) \in H_{\mathbb{R}}^{\infty}$ be a corona pair. The following assertions are equivalent:

- (1) there exists $\varepsilon > 0$ such that f has constant sign on the set

$$\{x \in]-1, 1[: |g(x)| < \varepsilon\};$$

- (2) f has constant sign on the set of extended real zeros of g .

Proof. Suppose that $|f| + |g| \geq \delta > 0$.

(1) $\implies$ (2). We may assume that $\varepsilon < \delta/2$. Then $|f(z)| > \delta/2$ whenever $|g(z)| < \varepsilon$. Say that f is positive at each $x \in]-1, 1[$ with $|g(x)| < \varepsilon$. Let $m_1, m_2 \in Z(g) \cap E$. Choose a net z_{α} in $]-1, 1[$ that converges to m_1 . Hence, $g(z_{\alpha}) \rightarrow 0$. In particular, there is α_0 such that $|g(z_{\alpha})| < \varepsilon$ for all $\alpha > \alpha_0$, whence $f(z_{\alpha}) > \delta/2$. Therefore, $f(m_1) \geq \delta/2 > 0$. By the same argument, $f(m_2) \geq \delta/2 > 0$. Thus, f has constant sign on the set of extended real zeros of g .

(2) $\implies$ (1). Since $|f| + |g| \geq \delta > 0$, by (2) we may assume that $f > \delta/2$ on $Z(g) \cap E$. Now,

$$Z(g) \cap E = \bigcap_{n=1}^{\infty} \{m \in E : |g(m)| \leq 1/n\}.$$

The finite intersection property for compact spaces (here E) implies that there is n_0 such that

$$\{m \in E : |g(m)| < 1/n_0\} \subseteq \{m \in E : f(m) > \delta/2\}.$$

Thus, $f(x) > 0$ at each point $x \in]-1, 1[$ with $|g(x)| \leq 1/n_0$. □

The following lemma is due to Corach and Suárez [1, 2].

Lemma 2.2 (see [1, p. 636] and [2, p. 608]). *Let A be a real or complex commutative unital Banach algebra with unit element denoted by 1.*

a) *Suppose that $(f, g) \in A^2$ is an invertible pair. If (f_n) converges to f in the norm and if the pairs (f_n, g) are reducible, then (f, g) is also reducible.*

b) *For $g \in A$, the set*

$$R(g) = \{f \in A : (f, g) \text{ is reducible}\}$$

is open-closed inside

$$I(g) = \{f \in A : (f, g) \text{ is invertible}\}.$$

c) *If $\phi : [0, 1] \rightarrow I(g)$ is a continuous curve and $(\phi(0), g)$ is reducible, then $(\phi(1), g)$ is reducible.*

Note that part a) follows from the identity

$$f_n + u_n g = f(x f_n + y g) + (y(f_n - f) + u_n)g,$$

where $f_n + u_n g \in A^{-1}$ and $x f + y g = 1$ (see [10] or [8]).

One of our tools will be the following result from [7]. Recall that a compact set $C \subseteq M(A)$ is said to be A -convex if C coincides with its A -convex hull

$$\tilde{C} = \{m \in M(A) : |\hat{f}(m)| \leq \max_C |\hat{f}| \text{ for all } f \in A\}.$$

Here we represent the short proof for the reader's convenience. In the sequel, let f^* be the function defined by $f^*(z) = \overline{f(\bar{z})}$.

Theorem 2.3 (see [7]). *The set E is H^{∞} and $H_{\mathbb{R}}^{\infty}$ -convex.*

Proof. Let $m \in M(H^\infty) \setminus E$. Since H^∞ is separating (see [11, p. 242]), there exists $f \in H^\infty$ such that $f(m) = 0$ and $f \neq 0$ on E . We may assume that $\|f\|_\infty \leq 1$. Let $g = ff^*$. Then $g \in H_{\mathbb{R}}^\infty$ and g is real-valued on E . By the definition of g , actually we have $1 \geq g \geq 0$ on E . Since $f \neq 0$ on E and $f^* = \bar{f}$ on E , we see that $\sigma := \min_E g > 0$. Now, let $h = 1 - g$. Then $h \in H_{\mathbb{R}}^\infty \subseteq H^\infty$ and $h(m) = 1$, whence

$$\max_E |h| = \max_E h \leq 1 - \sigma < |h(m)|.$$

Thus, m does not belong to the $H_{\mathbb{R}}^\infty$ -convex closure of E . Therefore, E is $H_{\mathbb{R}}^\infty$ -convex as well as H^∞ -convex. $\square$

Now we are ready to give the announced short proof of the reducibility part of Wick's result. It is essentially based on Treil's result that in H^∞ every corona pair is reducible [12].

Theorem 2.4 (Treil's theorem). *Let $(f, g) \in (H^\infty)^2$ be a corona pair satisfying*

$$\|f\|_\infty \leq 1, \quad \|g\|_\infty \leq 1, \quad \text{and} \quad |f| + |g| \geq \delta > 0.$$

Then there exist $h \in H^\infty$ and a constant $\kappa = \kappa(\delta)$ depending only on δ such that $v := f + hg$ is invertible in H^∞ and

$$\|h\|_\infty + \|v^{-1}\|_\infty \leq \kappa(\delta).$$

Theorem 2.5 (see [14, 15]). *Let (f, g) be a corona pair in $H_{\mathbb{R}}^\infty$. Then (f, g) is reducible if and only if f has constant sign on the set of extended real zeros of g .*

Proof. First, we prove the “only if” part. Let $u = f + hg$ be invertible in $H_{\mathbb{R}}^\infty$. Here $h \in H_{\mathbb{R}}^\infty$. Suppose that there exist $m_1, m_2 \in Z(g) \cap E$ such that $f(m_1) < 0$ and $f(m_2) > 0$. In particular $u(m_1) < 0$ and $u(m_2) > 0$. Since E is a connected set, the continuity of the (real-valued) function u on E implies that u has a zero on E . Hence, u is no longer invertible. (A finer analysis actually would yield a zero of u in $] -1, 1[$.)

To prove the sufficiency of the sign condition, consider the (real) trace-algebra $A = \overline{H_{\mathbb{R}}^\infty|_E}$, where the closure is taken with respect to the supremum norm in the space $C(E, \mathbb{R})$ of all real-valued continuous functions on E . Since E is $H_{\mathbb{R}}^\infty$ -convex by Theorem 2.3, the spectrum $M(A)$ of A can be identified with E (see [6, p. 125]). By our hypothesis, we may assume that $f > 0$ on $Z(g) \cap E$. This implies that for every $t \geq 0$ the pairs $(f|_E + t, g|_E)$ are invertible in A . Since $f|_E + N$ is invertible in A for large $N \in \mathbb{N}$, the pair $(f|_E + N, g|_E)$ is reducible in A . By Lemma 2.2, this implies that $(f|_E, g|_E)$ is also reducible in A . This means that there exists $q \in A$ such that $f + qg \neq 0$ on E . Approximating q uniformly on E by a function $h \in H_{\mathbb{R}}^\infty$, we see that $v := f + hg \in H_{\mathbb{R}}^\infty$ and that v does not vanish on E . In particular, v has no real zeros. Now, (v, g) is also a corona pair. Therefore, by [9, Proposition 2.6]¹ (or Proposition 3.1 below), (v, g) is reducible in $H_{\mathbb{R}}^\infty$. Thus, there exists $H \in H_{\mathbb{R}}^\infty$ so that

$$(f + hg) + Hg = f + (h + H)g$$

is invertible in $H_{\mathbb{R}}^\infty$. In other words, (f, g) is reducible in $H_{\mathbb{R}}^\infty$. $\square$

§3. WICK'S THEOREM WITH NORM ESTIMATES

In this section we present a new proof of Theorem **W** with norm estimate. Our proof depends only on Treil's result, i.e., Theorem 2.4, but does not need the quite delicate construction from [12].

For this, we shall need the following extensions of [9, Proposition 2.2] and [9, Proposition 2.6]. The proofs work exactly in the same way as those in [9]: at each occurrence

¹Here Treil's result is used.

of Treil's theorem, we additionally must take into account his norm estimates. Upon the referee's request, we present here the details.

Proposition 3.1. *Let (f, g) be a corona pair in $H_{\mathbb{R}}^{\infty}$ with $\|f\|_{\infty} \leq 1$, $\|g\|_{\infty} \leq 1$, and $|f| + |g| \geq \delta > 0$. Then the following assertions hold.*

- (1) *There exist a constant $C(\delta)$ (depending only on δ) and functions $u, h \in H_{\mathbb{R}}^{\infty}$ with u invertible such that $1 = uf^2 + hg$ and*

$$\|u\|_{\infty} + \|u^{-1}\|_{\infty} + \|h\|_{\infty} \leq C(\delta).$$

- (2) *Suppose that f has no real zeros. Then, again, there exist a constant $C(\delta)$ (depending only on δ) and functions $u, h \in H_{\mathbb{R}}^{\infty}$ with u invertible such that $1 = uf + hg$ and*

$$\|u\|_{\infty} + \|u^{-1}\|_{\infty} + \|h\|_{\infty} \leq C(\delta).$$

Proof. (1) By Theorem 2.4, there exist a constant κ depending only on δ , and $k \in H^{\infty}$ such that $v := f + kg$ is invertible in H^{∞} and $\|k\|_{\infty} + \|v^{-1}\|_{\infty} \leq \kappa$. In particular, $v^* = f + k^*g$ is invertible in H^{∞} . Multiplying the two equations, we see that

$$f^2 + g(f(k + k^*) + kk^*g) = vv^*$$

is invertible in H^{∞} . But the factors $k + k^*$ and kk^* belong to $H_{\mathbb{R}}^{\infty}$. Thus, with $u = (vv^*)^{-1}$ and $h = (f(k + k^*) + kk^*g)(vv^*)^{-1}$ we get $1 = uf^2 + hg$ and

$$\|u\|_{\infty} + \|u^{-1}\|_{\infty} + \|h\|_{\infty} \leq C(\delta).$$

(2) Write $f = BF$ where F is zero-free and B is the Blaschke product associated with the zeros of f . Since the zeros of B are symmetric with respect to the real axis, B and F are in $H_{\mathbb{R}}^{\infty}$. Without loss of generality, we may assume that $F > 0$ on $] -1, 1[$. Moreover, $F = G^2$ for some $G \in H_{\mathbb{R}}^{\infty}$ (see [9, p. 180]). Thus, (B, g) and (G, g) are invertible pairs in $(H_{\mathbb{R}}^{\infty})^2$ satisfying $|B| + |g| \geq \delta$ and $|G| + |g| \geq \delta$. By (1), there exists a constant $\kappa_1 > 0$ independent of δ such that

$$w := G^2 + k_1g = F + k_1g$$

is invertible in $H_{\mathbb{R}}^{\infty}$ for some $k_1 \in H_{\mathbb{R}}^{\infty}$ with $\|k_1\|_{\infty} + \|w^{-1}\|_{\infty} \leq \kappa_1$.

It remains to prove the same property for the pair (B, g) . Let C be the Blaschke product formed with the zeros of B on the upper-half plane, and let C^* be defined by $C^*(z) := \overline{C(\bar{z})}$. Since f (and with it, B) has no real zeros, we see that $B = CC^*$. Since (C, g) is an invertible pair in $(H^{\infty})^2$ with $|C| + |g| \geq \delta$, Treil's Theorem 2.4 yields the existence of $k_2 \in H^{\infty}$ and $\kappa_2 > 0$ depending only on δ such that $v := C + k_2g$ is invertible in H^{∞} and $\|k_2\|_{\infty} + \|v^{-1}\|_{\infty} \leq \kappa_2$. But $v^* = C^* + k_2^*g$. Multiplying the two equations, we get

$$CC^* + ((Ck_2^* + C^*k_2) + k_2k_2^*g) = vv^*.$$

Thus, with

$$K = (Ck_2^* + C^*k_2) + k_2k_2^*g \in H_{\mathbb{R}}^{\infty},$$

the function $\tilde{w} := vv^* = B + K$ is invertible in $H_{\mathbb{R}}^{\infty}$ and

$$\|K\|_{\infty} + \|\tilde{w}^{-1}\|_{\infty} \leq \kappa(\delta)$$

for some constant $\kappa(\delta)$.

To summarize, we have shown that

$$R := w\tilde{w} = (F + k_1g)(B + K) = BF + g(KF + k_1B + k_1Kg)$$

is invertible in $H_{\mathbb{R}}^{\infty}$ and that $R = f + hg$, where $h := KF + k_1B + k_1Kg$ and $\|h\|_{\infty} + \|R^{-1}\|_{\infty} \leq c(\delta)$.

It is now easy to deduce the claim. $\square$

Now we are ready to prove Theorem **W**, which we restate here, for the sake of a better exposition.

Theorem 3.2. *Let $(f, g) \in (H_{\mathbb{R}}^{\infty})^2$ be a corona pair satisfying $\|f\|_{\infty} \leq 1$, $\|g\|_{\infty} \leq 1$, and $|f| + |g| \geq \delta > 0$. Suppose that there exists $\varepsilon > 0$ such that f has constant sign on the set $\{x \in]-1, 1[: |g(x)| < \varepsilon\}$. Then there exists a constant $C = C(\delta, \varepsilon)$ depending only on δ and ε such that the Bézout equation*

$$uf + hg = 1$$

admits a solution $(u, h) \in (H_{\mathbb{R}}^{\infty})^2$ with u invertible and

$$\Delta(u, u^{-1}, h) := \|u\|_{\infty} + \|u^{-1}\|_{\infty} + \|h\|_{\infty} \leq C(\delta, \varepsilon).$$

Proof. The idea is to replace the function g by g^2 , so that for large N , the function $f + Ng^2$ is positive on the set of all $x \in]-1, 1[$ where $|g(x)|$ is small.

So, suppose that $f(x) > 0$ for all $x \in]-1, 1[$ with $|g(x)| < \varepsilon$. We may assume that $\varepsilon < \delta/2$. Then $|f| + |g| \geq \delta > 0$ implies that $f(x) > \delta/2$ for those x . Let

$$N = \max \left\{ \frac{4}{\delta}, \frac{1}{\varepsilon^2} \right\}.$$

Then $N = \frac{1}{\varepsilon^2}$. We claim that $f + Ng^2$ has no zeros on $]-1, 1[$. In fact, if $|g(x)| \leq \varepsilon$, then

$$|f(x) + Ng^2(x)| = f(x) + Ng^2(x) > \delta/2 > 0.$$

If $|g(x)| > \varepsilon$, then $g^2(x) > \varepsilon^2$, whence

$$|f(x) + Ng^2(x)| \geq Ng^2(x) - |f(x)| > N\varepsilon^2 - 1 \geq 0.$$

Let $F := \frac{f + Ng^2}{1 + N}$. Then (F, g) is a corona pair. Next we show that

$$(3.1) \quad |F| + |g| \geq \frac{\delta}{4N}.$$

Indeed, if $|g| \geq \sqrt{\frac{\delta}{4N}}$, then we are done. Now, let $|g| \leq \sqrt{\frac{\delta}{4N}}$. By the choice of N , we have

$$\sqrt{\frac{\delta}{4N}} \leq \frac{\delta}{4}.$$

Hence, since $|f| + |g| \geq \delta$, we see that $|f| \geq \frac{3}{4}\delta$. Accordingly,

$$|F| \geq \frac{|f| - N|g|^2}{N + 1} \geq \frac{\frac{3}{4}\delta - N\frac{\delta}{4N}}{1 + N} = \frac{\frac{\delta}{2}}{1 + N} \geq \frac{\delta}{4N}.$$

This establishes (3.1).

Since F has no real zeros, we may apply Proposition 3.1 to obtain functions u and h in $H_{\mathbb{R}}^{\infty}$ such that u is invertible,

$$1 = uF + hg,$$

and

$$\|u\|_{\infty} + \|u^{-1}\|_{\infty} + \|h\|_{\infty} \leq C(\delta/4N) = C((\varepsilon^2\delta)/4) =: C.$$

Thus,

$$1 = \frac{u}{1 + N}f + \left(\frac{Nug}{1 + N} + h \right)g$$

is the solution we were looking for; simply put

$$C(\delta, \varepsilon) = 2C + \left(1 + \frac{1}{\varepsilon^2} \right) C.$$

□

A natural question is whether it is possible to get bounds on the solution triple (u, u^{-1}, h) of the Bézout equation $uf + hg = 1$, u invertible in $H_{\mathbb{R}}^{\infty}$, that depend only on

$$\delta = \inf_z (|f(z)| + |g(z)|).$$

For example, this is the case under the hypotheses of Proposition 3.1. The answer is provided in §5.

§4. THE SIGN CONDITION ON $] -1, 1[$ IS NOT SUFFICIENT

In [9, Proposition 2.8] it was shown that the pair (f, B) is reducible in $H_{\mathbb{R}}^{\infty}$ whenever B is a Blaschke product having only real zeros (of multiplicity one) and $f \in H_{\mathbb{R}}^{\infty}$ is such that f is positive on the real zeros of B and $|f| + |B| \geq \delta > 0$.

If B is replaced by an outer function g or a singular inner function, then g has no zeros at all in $\mathbb{D}$ and the hypothesis of f being positive at the real zeros of g is true by convention. We show that in that case reducibility is not necessarily forced by this vacuous condition.

Example 4.1. Let $f(z) = z$ and $g(z) = \exp(-\frac{1+z^2}{1-z^2})$. Then g has no zeros in $] -1, 1[$, but nevertheless (f, g) is not reducible in $H_{\mathbb{R}}^{\infty}$.

Proof. Let $h \in H_{\mathbb{R}}^{\infty}$ be given, and set $L(z) = z + h(z)g(z)$. Since $\lim_{r \rightarrow 1} g(r) = 0$ and $\lim_{x \rightarrow -1} g(x) = 0$, we see that $\lim_{r \rightarrow 1} L(r) = 1$ and $\lim_{x \rightarrow -1} L(x) = -1$. Thus, being real valued on $] -1, 1[$, L has a zero within $] -1, 1[$, so that L is not invertible. $\square$

We could have worked also with the pair $(z, 1 - z^2)$. But the example above has the additional property that g has no zeros in $\mathbb{C}$ at all (apart from the essential singularities at -1 and 1).

Here is yet another example; this time we give an example such that the second member g does have real zeros (note that conditions involving empty sets are sometimes arguable). It is based on the fact that f is positive on the real zeros of g if and only if between two real zeros of g there is an even number of zeros of f (or none). In the following, let $\mathbb{N} = \{0, 1, 2, \dots\}$.

Example 4.2. Let B be the interpolating Blaschke product with the zeros $z_n = 1 - 2^{-n}$ ($n = 1, 2, \dots$), let B_1 be the subproduct of B with the zero set

$$\{z_n : n = 3j + 2 \text{ or } n = 3(j + 1), j \in \mathbb{N}\},$$

that is, $Z_{\mathbb{D}}(B_1) = \{z_2, z_3, z_5, z_6, \dots\}$, and let B_2 be the subproduct of B with the zero set

$$\{z_n : n = 3j + 1, j \in \mathbb{N}\}.$$

We introduce two functions f and g by

$$f(z) = zB_1(z), \quad g(z) = B_2(z) \exp\left(-\frac{1-z}{1+z}\right).$$

We are going to show that (f, g) is a corona pair, f has constant sign on the real zeros of g , but (f, g) is not reducible in $H_{\mathbb{R}}^{\infty}$.

Proof. Since $B = B_1B_2$ is an interpolating Blaschke product, from [5, p. 208] we see that the ideal generated by B_1 and B_2 in H^{∞} equals H^{∞} . Hence B_1 and B_2 have no common zeros on the spectrum of H^{∞} . Moreover, since the zeros of $\exp(-\frac{1-z}{1+z})$ in $M(H^{\infty})$ are in the fiber over the point $z = -1$, we conclude that f and g have no zeros in common on $M(H^{\infty})$. Thus (f, g) is a corona pair in H^{∞} , and consequently also in $H_{\mathbb{R}}^{\infty}$.

By construction, between two distinct real zeros of B_2 there are exactly two zeros of f . Thus f has constant sign on the real zeros of g . Now all the zeros of B_1 are larger than the first zero z_1 of g . Recall that B_1 has the form

$$(4.1) \quad B_1(x) = \prod_j \frac{a_j - x}{1 - a_j x}.$$

Hence, $f(z_1) > 0$, and so this constant sign is the positive sign. Moreover, B_1 is bounded away from zero on the zeros of B_2 (because (B_1, B_2) is a corona pair). So, we can conclude that for given $h \in H_{\mathbb{R}}^{\infty}$, the function $W(z) = zB_1(z) + h(z)g(z)$ satisfies

$$\liminf_n W(z_{3n+1}) = \liminf_n z_{3n+1} B_1(z_{3n+1}) > 0.$$

On the other hand, B_1 is unimodular and continuous at -1 and so, by (4.1), $B_1(-1) = 1 > 0$. Thus,

$$\lim_{x \rightarrow -1} W(x) = -1.$$

Again we see that W has a zero in $] -1, 1[$. □

§5. NON-UNIFORMLY BOUNDED SOLUTIONS

In contrast to the situation in H^{∞} (see [12]), we will show here that the reducibility of sequences of corona pairs (f_n, g_n) in the unit ball of $H_{\mathbb{R}}^{\infty}$ with $|f_n| + |g_n| \geq \delta > 0$ does not imply the existence of uniformly bounded solutions of the Bézout equation $u_n f_n + h_n g_n = 1$ with u_n invertible, where the bounds of the norms of u_n, u_n^{-1} and h_n depend only on δ . Recall that

$$\Delta(u, u^{-1}, h) = \|u\|_{\infty} + \|u^{-1}\|_{\infty} + \|h\|_{\infty}.$$

Example 5.1. Put $f(z) = z$ and $g(z) = \exp(-\frac{1+z^2}{1-z^2})$. We choose a sequence (r_n) of positive numbers converging to 1 ($0 < r_n < 1$); let $f_n(z) = f(r_n z)$ and $g_n(z) = g(r_n z)$. Then the pairs (f_n, g_n) are reducible in $H_{\mathbb{R}}^{\infty}$ and satisfy $|f_n| + |g_n| \geq \delta > 0$, but

$$\sup_n \Delta(u_n, u_n^{-1}, h_n) = \infty$$

for any solutions $(u_n, h_n) \in (H_{\mathbb{R}}^{\infty})^2$ of the equation $u_n f_n + h_n g_n = 1$ with u_n invertible in $H_{\mathbb{R}}^{\infty}$.

Proof. Suppose to the contrary that there exist solutions (u_n, h_n) satisfying

$$\sup_n \Delta(u_n, u_n^{-1}, h_n) < \infty.$$

Using a normal families argument, we can find subsequences of u_n and h_n converging locally uniformly to some functions u and h in $H_{\mathbb{R}}^{\infty}$. Since $|u_n|$ is bounded away from zero by a constant independent of n , the limit function u is also invertible. Thus,

$$uf + hg = 1,$$

but this contradicts Example 4.1. □

We can also create counterexamples to this norm-estimate problem by working with finite Blaschke products. Note that this is the class of functions Brett Wick worked with in [14].

Example 5.2. There exists a sequence (B_n) of finite Blaschke products in $H_{\mathbb{R}}^{\infty}$ such that the pairs (z, B_n) are reducible in $H_{\mathbb{R}}^{\infty}$ and satisfy $|z| + |B_n| \geq \eta > 0$, but

$$\sup_n \Delta(u_n, u_n^{-1}, h_n) = \infty$$

for any solutions $(u_n, h_n) \in (H_{\mathbb{R}}^{\infty})^2$ of $u_n z + h_n B_n = 1$ with u_n invertible in $H_{\mathbb{R}}^{\infty}$.

Proof. Let $g^*(z) = \overline{g(\bar{z})}$. Note that a function $g \in H^{\infty}$ belongs to $H_{\mathbb{R}}^{\infty}$ if and only if $g = g^*$. Let $f(z) = z, g(z) = (1 - z^2)/2$. By the Carathéodory theorem (see [3, p. 6]), we can choose a sequence of finite Blaschke products b_n converging locally uniformly to g in $\mathbb{D}$. If b_n has a real zero, we move it vertically a little to the upper half disk. Since $g = g^*$, $B_n := b_n b_n^*$ tends locally uniformly to g^2 . Moreover, $B_n \in A(\mathbb{D})_{\mathbb{R}}$ and has no real zeros. Obviously, (z, g^2) is a corona pair, say $|z| + |g^2| \geq \delta > 0$, where we may choose $\delta < 1$. Since B_n converges to g^2 , uniformly on $|z| \leq 1/2$, we see that $|z| + |B_n| \geq \delta/2$ on $|z| \leq 1/2$ whenever n is large. For $1/2 \leq |z| \leq 1$ we obtain $|z| + |B_n| \geq 1/2 \geq \delta/2$.

Thus, the (z, B_n) are corona pairs with $|z| + |B_n| \geq \delta/2$. Using Lemma 2.2, we show that these pairs are reducible in $A(\mathbb{D})_{\mathbb{R}}$. Indeed, since $B_n > 0$ on $[-1, 1]$, the pairs $(z + t, B_n)$ are invertible for any $t \geq 0$. For $t = 2$, e.g., $z + 2$ is an invertible function in $A(\mathbb{D})_{\mathbb{R}}$; hence, $(z + 2, B_n)$ is reducible. Thus, by Lemma 2.2, (z, B_n) is also reducible.

So, let

$$1 = u_n z + h_n B_n,$$

where u_n is invertible in $A_{\mathbb{R}}(\mathbb{D})$.

As above, no bound on the norms of u_n, u_n^{-1} and h_n is possible, because otherwise $1 = uz + hg^2$ for some $u \in H_{\mathbb{R}}^{\infty}$, with u invertible and $h \in H_{\mathbb{R}}^{\infty}$. But the pair (z, g^2) is not reducible in $H_{\mathbb{R}}^{\infty}$, because for every $k \in H_{\mathbb{R}}^{\infty}$ the function $z + k(1 - z^2)^2$ takes the value -1 at -1 and 1 at 1 , so that it admits a zero on $] -1, 1[$. $\square$

Examples 5.1 and 5.2 show that for fixed δ , the constants $C(\delta, \varepsilon)$ in Wick's theorem **W** may tend to infinity as $\varepsilon \rightarrow 0$. This is the case, e.g., if $\varepsilon := \varepsilon_n = \frac{1}{2} \min_{[-1, 1]} B_n$, where B_n is the finite Blaschke product in Example 5.2. (Indeed, fix $\varepsilon' > 0$ and choose r so close to 1 that $g^2(r) < \varepsilon'$. Since $B_n(r)$ converges to $g^2(r)$, we see that $\varepsilon_n \leq B_n(r) < \varepsilon'$ for n sufficiently large. Hence, $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$.)

REFERENCES

- [1] G. Corach and F. D. Suárez, *Stable rank in holomorphic function algebras*, Illinois J. Math. **29** (1985), 627–639. MR0806470 (87b:46056)
- [2] ———, *On the stable range of uniform algebras and H^{∞}* , Proc. Amer. Math. Soc. **98** (1986), 607–610. MR0861760 (87m:46103)
- [3] J. B. Garnett, *Bounded analytic functions*, Pure Appl. Math., vol. 96, Acad. Press, New York–London, 1981. MR0628971 (83g:30037)
- [4] P. Gorkin, R. Mortini, and N. Nikolski, *Norm controlled inversions and a corona theorem for H^{∞} -quotient algebras*, J. Funct. Anal. **255** (2008), 854–876. MR2433955 (2009h:30061)
- [5] K. Hoffman, *Banach spaces of analytic functions*, Reprint of the 1962 original, Dover Publ., Inc., New York, 1988. MR1102893 (92d:46066)
- [6] S. H. Kulkarni and B. V. Limaye, *Real function algebras*, Monogr. Textbooks Pure Appl. Math., vol. 168, Marcel Dekker, New York, 1992. MR1197884 (93m:46059)
- [7] R. Mortini, *The covering dimension of a distinguished subset of the spectrum $M(H^{\infty})$ of H^{∞} and the algebra of real symmetric and continuous functions on $M(H^{\infty})$* , Topology Appl. **159** (2012), 900–910. MR2868890
- [8] R. Mortini and R. Rupp, *A note on some uniform algebra generated by smooth functions in the plane*, J. Funct. Spaces Appl. Article ID 905650 (2012), 15 pages.
- [9] R. Mortini and B. Wick, *The Bass and topological stable ranks of $H_{\mathbb{R}}^{\infty}(\mathbb{D})$ and $A_{\mathbb{R}}(\mathbb{D})$* , J. Reine Angew. Math. **636** (2009), 175–191. MR2572249 (2010m:46079)
- [10] ———, *Spectral characteristics and stable ranks for the Sarason algebra $H^{\infty} + C$* , Michigan Math. J. **59** (2010), 395–409. MR2677628 (2011j:46085)
- [11] F. D. Suárez, *Čech cohomology and covering dimension for the H^{∞} maximal ideal space*, J. Funct. Anal. **123** (1994), 233–263. MR1283028 (95g:46100)
- [12] S. Treil, *The stable rank of the algebra H^{∞} equals 1*, J. Funct. Anal. **109** (1992), 130–154. MR1183608 (93h:46076)
- [13] B. Wick, *A note about stabilization in $A_{\mathbb{R}}(\mathbb{D})$* , Math. Nachr. **282** (2009), 912–916. MR2530887 (2010d:46065)
- [14] ———, *Stabilization in $H_{\mathbb{R}}^{\infty}(\mathbb{D})$* , Publ. Mat. **54** (2010), 25–52. MR2603587 (2011c:46112)

- [15] ———, *Corrigenda*. “Stabilization in $H_{\mathbb{R}}^{\infty}(\mathbb{D})$ ”, Publ. Mat. **55** (2011), 251–260. MR2779583 (2012a:46090)

DÉPARTEMENT DE MATHÉMATIQUES, LMAM, UMR 7122, UNIVERSITÉ PAUL VERLAINE, ÎLE DU SAULCY, F-57045 METZ, FRANCE

E-mail address: `mortini@math.univ-metz.fr`

Received 28/JUN/2010

Originally published in English