AN ESTIMATE OF THE RATE OF CONVERGENCE TO THE NORMAL LAW

UDC 519.21

I. YO. POLYAK

ABSTRACT. The paper contains an estimate of the rate of convergence to the normal law for nonidentically distributed random variables.

A generalization of a V. M. Zolotarev result for the case of nonidentically distributed random variables is obtained in [1]. This paper continues the line of research initiated in [1] on applications of pseudomoments to the rate of convergence to the normal law. However below we use other pseudomoments different from those used in [1].

Let $\xi_1, \xi_2, \dots, \xi_n, \dots$ be a sequence of independent random variables such that $M\xi_i = 0$ and $\text{Var}\xi_i = \sigma_i^2$. Let $F_i(x)$ and $f_i(t)$ be the distribution function and characteristic function of the random variable ξ_i , $i \geq 1$, respectively. Denote by $\Phi_n(x)$ the distribution function of the random variable $(\xi_1 + \xi_2 + \dots + \xi_n)/B_n$ and put $B_n^2 = \sigma_1^2 + \dots + \sigma_n^2$, $\bar{\sigma}_i = \min(1; \sigma_i)$, and $\bar{B}_n^2 = \bar{\sigma}_1^2 + \dots + \bar{\sigma}_n^2$. Further let $\Phi(x)$ be the standard normal distribution function, $\rho_n = \sup_x |\Phi_n(x) - \Phi(x)|$, and

$$\begin{aligned}\lambda_{ni}^{(1)} &= \int_{|x| \leq B_n} \max(1, |x|^3) \left| d \left(F_i(x) - \Phi \left(\frac{x}{\sigma_i} \right) \right) \right|, \\ \lambda_{ni}^{(2)} &= \int_{|x| \geq B_n} x^2 \left| d \left(F_i(x) - \Phi \left(\frac{x}{\sigma_i} \right) \right) \right|, \\ \Lambda_n^{(1)} &= \frac{1}{B_n^2} \sum_{i=1}^n \lambda_{ni}^{(1)}, \quad \Lambda_n^{(2)} = \frac{1}{B_n^2} \sum_{i=1}^n \lambda_{ni}^{(2)}, \quad b_n = \min(\bar{\sigma}_1, \dots, \bar{\sigma}_n).\end{aligned}$$

Theorem. For all $n \geq 1$,

$$(1) \quad \rho_n \leq C \left(\frac{1}{B_n} \Lambda_n^{(1)} + \Lambda_n^{(2)} \right) b_n^{-3},$$

where C is a universal constant.

Proof. Let n be fixed and $c \in (0; 1)$ be a certain constant; the exact value of c will be chosen below. Assume that $\Lambda_n^{(2)} \leq c$ and note that estimate (1) is obvious otherwise.

For $n = 1$ we have

$$\begin{aligned}\rho_1 &= \sup_x |F_1(x\sigma) - \Phi(x)| = \sup_x |F_1(x) - \Phi(x/\sigma_1)| = \sup_x \left| \int_{-\infty}^{+\infty} d(F_1(x) - \Phi(x/\sigma_1)) \right| \\ &\leq \int_{-\infty}^{+\infty} |d(F_1(x) - \Phi(x/\sigma_1))| \leq \lambda_{n1}^{(1)} + \frac{1}{\sigma_1^2} \lambda_{n1}^{(2)},\end{aligned}$$

and inequality (1) follows. Now let $n \geq 2$.

Put $\phi_n(t) = \prod_{i=1}^n f_i(t/B_n)$. We apply the following inequality:

$$(2) \quad \rho_n \leq \min \left\{ \max \left(\int_0^X \left| \phi_n(t) - e^{-t^2/2} \right| \frac{dt}{t}, \frac{6\sqrt{2}}{X} \right); X > 0 \right\}$$

(see [2, p. 377]) with $X = TB_n$, $T = c^{5/2}\Lambda_n^{-1}$, and $\Lambda_n = \max(\Lambda_n^{(1)}; \Lambda_n^{(2)})$. Let $T_1 = \sqrt{c}$, $T_2 = \min(T_1, T)$. Using the inequalities

$$\left| \phi_n(t) - e^{-t^2/2} \right| \leq \sum_{i=1}^n \left| \omega_i \left(\frac{t}{B_n} \right) \right| |\psi_{ni}(t)|,$$

where $\omega_i(t) = f_i(t) - \exp\{-t^2\sigma_i^2/2\}$, $\psi_{ni}(t) = \prod_{k=1}^{i-1} \exp\{-t^2\sigma_k^2/2B_n^2\} \prod_{k=i+1}^n f_k(t/B_n)$, and

$$|\omega_i(t)| \leq \min \left(\frac{|t|^3}{6} \lambda_{ni}^{(1)} + 2t^2 \lambda_{ni}^{(2)}; \lambda_{ni}^{(1)} + \frac{1}{B_n^2} \lambda_{ni}^{(2)} \right),$$

we get

$$\begin{aligned} |f_k(t)| &\leq \left| \exp\{-t^2\sigma_k^2/2\} + \omega_k(t) \right| \leq \exp\{-t^2\sigma_k^2/2\} + \frac{|t|^3}{6} \lambda_{nk}^{(1)} + 2t^2 \lambda_{nk}^{(2)} \\ &\leq \exp\{-t^2\bar{\sigma}_k^2/2\} + t^2 \left(\frac{c^{5/2}}{6} + 2c \right) \leq \exp\{-c_1 t^2 \bar{\sigma}_k^2\} \end{aligned}$$

for $|t| \leq T_2$, while

$$|\psi_{ni}(t)| \leq e \cdot \exp \left\{ -t^2 \frac{\bar{B}_n^2}{B_n^2} c_1 \right\}$$

for $|t| \leq T_2 B_n$, where $c_1 = (1 - e^{-c/2})/c - 13c/6$ and c is chosen such that $c_1 > 0$. Then (see inequality (12) in [1])

$$(3) \quad \left| \phi_n(t) - e^{-t^2/2} \right| \leq e \left(\frac{|t|^3}{6} \Lambda_n^{(1)} + 2t^2 \Lambda_n^{(2)} \right) \exp \left\{ -t^2 \frac{\bar{B}_n^2}{B_n^2} c_1 \right\}.$$

We represent the integral in (2) as follows:

$$(4) \quad \begin{aligned} I &= \int_0^{TB_n} \left| \phi_n(t) - e^{-t^2/2} \right| \frac{dt}{t} \\ &= \int_0^{T_2 B_n} \left| \phi_n(t) - e^{-t^2/2} \right| \frac{dt}{t} + \int_{T_2 B_n}^{TB_n} \left| \phi_n(t) - e^{-t^2/2} \right| \frac{dt}{t} = I_1 + I_2. \end{aligned}$$

Inequality (3) implies

$$(5) \quad I_1 = \int_0^{T_2 B_n} \left| \phi_n(t) - e^{-t^2/2} \right| \frac{dt}{t} \leq C^{(1)} \left(\frac{1}{\bar{B}_n} \Lambda_n^{(1)} + \Lambda_n^{(2)} \right).$$

The integral I_2 is nonzero if $T_2 = T_1$, in which case $\Lambda_n^{(1)} \leq c^2$. Thus

$$\begin{aligned} \left| f_i \left(\frac{t}{B_n} \right) \right| &\leq \exp \{ -t^2 \sigma_i^2 / (2B_n^2) \} + \left| \omega_i \left(\frac{t}{B_n} \right) \right| \\ &\leq \exp \left\{ \left(e^{-c/2} - 1 \right) \bar{\sigma}_i^2 + \lambda_{ni}^{(1)} + \frac{1}{B_n^2} \lambda_{ni}^{(2)} \right\}, \\ \prod_{\substack{k=1 \\ k \neq i, j}}^n \left(\exp \{ -t^2 \sigma_k^2 / (2B_n^2) \} + \left| \omega_k \left(\frac{t}{B_n} \right) \right| \right) &\leq e^{2+c} \exp \left\{ - \left(\frac{1 - e^{-c/2}}{c} - c \right) c \bar{B}_n^2 \right\} \end{aligned}$$

for $T_2 B_n \leq |t| \leq T B_n$. Moreover by inequality (13) in [1],

$$\begin{aligned}
 \left| \phi_n(t) - e^{-t^2/2} \right| &\leq \sum_{i=1}^n \left| \omega_i \left(\frac{t}{B_n} \right) \right| \left| \sum_{\substack{j=1 \\ j \neq i}}^n e^{-t^2 \sigma_j^2 / (2B_n^2)} \prod_{\substack{k=1 \\ k \neq i, j}}^n \left(e^{-t^2 \sigma_k^2 / (2B_n^2)} + \left| \omega_k \left(\frac{t}{B_n} \right) \right| \right) \right| \\
 &\quad + \prod_{k=1}^n \left| \omega_k \left(\frac{t}{B_n} \right) \right| \\
 &\leq \sum_{i=1}^n \left| \omega_i \left(\frac{t}{B_n} \right) \right| e^{-t^2 b_n^2 / (2B_n^2)} (n-1) e^{2+c} \exp \{ -c_3 c \overline{B}_n^2 \} \\
 &\quad + \prod_{k=1}^n \left(\lambda_{nk}^{(1)} + \frac{1}{B_n^2} \lambda_{nk}^{(2)} \right),
 \end{aligned}$$

where $c_3 = (1 - e^{-c/2})/c - c > 0$.

Thus

$$\begin{aligned}
 I_2 &= \int_{T_2 B_n}^{T B_n} \left| \phi_n(t) - e^{-t^2/2} \right| \frac{dt}{t} \\
 &\leq (n-1) e^{2+c} \exp \{ -c_3 c \overline{B}_n^2 \} \sum_{i=1}^n \left(\lambda_{ni}^{(1)} + \frac{1}{B_n^2} \lambda_{ni}^{(2)} \right) \int_{T_2 B_n}^{T B_n} e^{-t^2 b_n^2 / (2B_n^2)} \frac{dt}{t} \\
 (6) \quad &\quad + \prod_{k=1}^n \left(\lambda_{nk}^{(1)} + \frac{1}{B_n^2} \lambda_{nk}^{(2)} \right) \int_{T_2 B_n}^{T B_n} \frac{dt}{t} \\
 &\leq C^{(3)} \left(\frac{\Lambda_n^{(1)}}{\overline{B}_n} + \Lambda_n^{(2)} \right) b_n^{-3}
 \end{aligned}$$

and (1) follows from (2) and (4)–(6). $\square$

BIBLIOGRAPHY

1. P. V. Slyusarchuk and I. Yo. Polyak, *A generalization of a V. M. Zolotarev result*, Naukovyi Visnyk Uzhgorod. Univer., Ser. Matem. **3** (1997), 184–189. (Ukrainian)
2. V. M. Zolotarev, *A modern theory of summation of independent random variables*, “Nauka”, Moscow, 1986. (Russian) MR 89a:60003

CHAIR OF MATHEMATICAL ANALYSIS, DEPARTMENT OF MATHEMATICS, UZHGOROD NATIONAL UNIVERSITY, PIDGIRNA STREET 46, UZHGOROD, UKRAINE

E-mail address: poliak_ivan@mail.ru

Received 20/NOV/2002

Translated by OLEG KLESOV