GENERALIZED SOLUTIONS OF A HYPERBOLIC EQUATION WITH A φ -SUB-GAUSSIAN RIGHT HAND SIDE

UDC 519.21

B. V. DOVGAI

ABSTRACT. A hyperbolic equation with a φ -sub-Gaussian right hand side is considered. Conditions for the existence of a generalized solution of such an equation are found.

1. INTRODUCTION

This paper is devoted to studying generalized solutions of the wave equation with zero initial and boundary conditions and with a φ -sub-Gaussian right hand side. We find sufficient conditions for the existence of a generalized solution expressed in terms of convergence of a certain deterministic series and in terms of the covariance function of the random field presented on the right hand side of the equation.

2. GENERALIZED SOLUTION OF A BOUNDARY PROBLEM FOR THE DETERMINISTIC CASE

Let $T > 0$ be a constant, let $p(x)$ be a continuously differentiable function on the interval $[a, b]$ such that $p(x) > 0$, let $\rho(x)$ and $q(x)$ be continuous functions on the interval $[a, b]$ such that $\rho(x) > 0$ and $q(x) \geq 0$, and let $f(x, t)$, $x \in [a, b]$, $t \in [0, T]$, be a continuous function.

Consider the boundary problem for the nonhomogeneous hyperbolic equation on the interval $[a, b]$:

$$(1) \quad \frac{\partial}{\partial x} \left(p(x) \frac{\partial u}{\partial x} \right) - q(x)u - \rho(x) \frac{\partial^2 u}{\partial t^2} = -\rho(x)f(x, t),$$

$$x \in [a, b], \quad t \in [0, T],$$

$$(2) \quad u|_{t=0} = 0, \quad \frac{\partial u}{\partial t} \Big|_{t=0} = 0,$$

$$(3) \quad u|_{x=a} = 0, \quad u|_{x=b} = 0.$$

The corresponding Sturm–Liouville problem is written as follows:

$$\frac{d}{dx} \left(p \frac{dX}{dx} \right) - qX + \lambda \rho X = 0,$$

$$X(a) = X(b) = 0.$$

Let $X_n(x)$ be the orthonormal eigenfunctions with respect to the weight ρ and let λ_n be the corresponding eigenvalues for this problem. We assume that λ_n are taken in

2000 *Mathematics Subject Classification.* Primary 60H15, 35L20, 35R60.

Key words and phrases. Generalized solutions, hyperbolic equation, φ -sub-Gaussian stochastic process.

ascending order. In view of the restrictions imposed on the functions p , q , and ρ , all eigenvalues are positive, and zero is not an eigenvalue.

Put $\mu_n = \sqrt{\lambda_n}$.

Denote by $u_n(x, t)$ a solution of the problem (1)–(3) where a partial Fourier sum of the function f substitutes the function f itself on the right hand side of equation (1), namely

$$f_n(x, t) = \sum_{k=1}^n f_k(t) X_k(x)$$

replaces f . Here $f_k(t) = \int_a^b f(x, t) X_k(x) \rho(x) dx$. Thus

$$u_n(x, t) = \sum_{k=1}^n X_k(x) \frac{1}{\mu_k} \int_0^t \sin \mu_k(t - u) f_k(u) du.$$

Definition 1. Following [11], a function $u(x, t)$ is called a generalized solution of the problem (1)–(3) if there is a subsequence $u_{n_k}(x, t)$, $k \geq 1$, such that $u_{n_k}(x, t) \rightarrow u(x, t)$ as $k \rightarrow \infty$ in the norm of the space $C([a, b] \times [0, T])$.

3. SUFFICIENT CONDITIONS FOR THE EXISTENCE OF A GENERALIZED SOLUTION

Let $T > 0$ be a constant; let $p(x)$ and $\rho(x)$, $x \in [0, \pi]$, be twice continuously differentiable functions such that $p(x) > 0$ and $\rho(x) > 0$; let $q(x)$, $x \in [0, \pi]$, be a continuously differentiable function such that $q(x) \geq 0$; and let $\xi(x, t)$, $x \in [0, \pi]$, $t \in [0, T]$, be an almost surely continuous random field.

Consider the following problem:

$$(4) \quad \frac{\partial}{\partial x} \left(p(x) \frac{\partial u}{\partial x} \right) - q(x)u - \rho(x) \frac{\partial^2 u}{\partial t^2} = -\rho(x)\xi(x, t), \quad x \in [0, \pi], \quad t \in [0, T],$$

$$(5) \quad u|_{t=0} = 0, \quad \left. \frac{\partial u}{\partial t} \right|_{t=0} = 0,$$

$$(6) \quad u|_{x=0} = 0, \quad u|_{x=\pi} = 0.$$

The corresponding Sturm–Liouville problem is as follows:

$$\begin{aligned} \frac{d}{dx} \left(p \frac{dX}{dx} \right) - qX + \lambda \rho X &= 0, \\ X(0) &= X(\pi) = 0. \end{aligned}$$

Let $X_n(x)$ be the orthonormal eigenfunctions with respect to the weight ρ and let λ_n be the corresponding eigenvalues of the problem. We assume that λ_n are taken in ascending order. In view of the assumptions imposed on p , q , and ρ , all eigenvalues are positive, and zero is not an eigenvalue.

Put $\mu_n = \sqrt{\lambda_n}$.

Below is the definition of a generalized condition compatible with Definition 1.

Definition 2. A random field $u(x, t)$, $x \in [0, \pi]$, $t \in [0, T]$, is called a generalized solution of the problem (4)–(6) if

$$(7) \quad u(x, t) = \sum_{n=1}^{\infty} X_n(x) \frac{1}{\mu_n} \int_0^t \sin \mu_n(t - u) \zeta_n(u) du,$$

where

$$\zeta_n(t) = \int_0^\pi \xi(x, t) X_n(x) \rho(x) dx$$

and the series on the right hand side of (7) converges in probability in the norm of the space $C([0, \pi] \times [0, T])$.

Remark 1. If a series of random variables converges in probability, then there exists a subsequence of partial sums that converges almost surely. Thus Definition 2 is indeed compatible with Definition 1.

Theorem 1. *Let $\xi(x, t)$ be a strictly φ -sub-Gaussian random field that is continuous with probability one. Assume that $p > 1$ and $\varphi(x) = |x|^p$ for $|x| > 1$. If the following double series:*

$$\sum_{k,m=1}^{\infty} \frac{(\ln^{\alpha} k \ln^{1-\alpha} m)^{2\delta}}{km} C_{k,m}$$

converges for some $\alpha \in [0, 1]$ and $\delta > 1 - 1/p$, where

$$C_{k,m} = \sup_{0 \leq s \leq T, 0 \leq t \leq T} |\mathbb{E} \zeta_k(s) \zeta_m(t)|,$$

then $u(x, t)$ defined by (7) is a generalized solution of the problem (4)–(6).

Proof. It is known [11] that there exists a constant $C_X > 0$ such that

$$|X_n(x)| \leq C_X$$

for all $x \in [0, \pi]$. Moreover, in the case under consideration,

$$\mu_n = \frac{n\pi}{l} + O\left(\frac{1}{n}\right),$$

$$X_n(x) = \frac{1}{\sqrt[4]{p(x)\rho(x)}} \sqrt{\frac{2}{l}} \sin\left(\frac{n\pi}{l} \int_0^x \sqrt{\frac{\rho(u)}{p(u)}} du\right) + \frac{K_n(x)}{n}$$

(see [11]), where

$$l = \int_0^\pi \sqrt{\rho(u)/p(u)} du,$$

$|K_n(x)| \leq C'$ for all $n \geq 1$ and $x \in [0, \pi]$, and

$$\begin{aligned} u(x, t) &= \frac{1}{\sqrt[4]{p(x)\rho(x)}} \sqrt{\frac{2}{l}} \sum_{n=1}^{\infty} \frac{1}{\mu_n} \sin\left(\frac{n\pi}{l} \int_0^x \sqrt{\frac{\rho(u)}{p(u)}} du\right) \int_0^t \sin \mu_n(t-u) \zeta_n(u) du \\ &\quad + \sum_{n=1}^{\infty} \frac{1}{n\mu_n} K_n(x) \int_0^t \sin \mu_n(t-u) \zeta_n(u) du, \\ \sum_{n=1}^{\infty} \frac{1}{n\mu_n} K_n(x) \int_0^t \sin \mu_n(t-u) \zeta_n(u) du &\leq C' C_X T \pi \max_{\substack{x \in [0, \pi] \\ t \in [0, T]}} \{|\xi(x, t)\rho(x)|\} \sum_{n=1}^{\infty} \frac{1}{n\mu_n} < \infty \end{aligned}$$

with probability 1. Thus the series

$$\sum_{n=1}^{\infty} \frac{1}{n\mu_n} K_n(x) \int_0^t \sin \mu_n(t-u) \zeta_n(u) du$$

converges with probability 1 in the norm of the space $C([0, \pi] \times [0, T])$, and therefore it converges in probability.

Further,

$$\begin{aligned} & \sum_{k,n=1}^{\infty} \frac{1}{\mu_n \mu_k} \frac{1}{\sqrt{p(x)\rho(x)}} \frac{2}{l} \sin \left(\frac{n\pi}{l} \int_0^x \sqrt{\frac{\rho(u)}{p(u)}} du \right) \sin \left(\frac{k\pi}{l} \int_0^x \sqrt{\frac{\rho(u)}{p(u)}} du \right) \\ & \quad \times \int_0^t \int_0^t \sin \mu_n(t-u) \sin \mu_k(t-v) \mathbb{E} \zeta_n(u) \zeta_k(v) du dv \\ & \leq C'' \sum_{k,n=1}^{\infty} \frac{1}{\mu_n \mu_k} C_{n,k} < \infty. \end{aligned}$$

Let

$$R_{k,m}(t, s) = \int_0^t \int_0^s \sin \mu_k(t-u) \sin \mu_m(s-v) \mathbb{E} \zeta_k(u) \zeta_m(v) dv du.$$

Similarly to the proof of Lemma 5 of the paper [5], we obtain

$$\begin{aligned} & \mathbb{E} \left| \sum_{k=1}^n \frac{1}{\mu_k} \sin \left(\frac{k\pi}{l} \int_0^x \sqrt{\frac{\rho}{p}} du \right) \int_0^t \sin \mu_k(t-u) \zeta_k(u) du \right. \\ & \quad \left. - \sum_{k=1}^n \frac{1}{\mu_k} \sin \left(\frac{k\pi}{l} \int_0^y \sqrt{\frac{\rho}{p}} du \right) \int_0^s \sin \mu_k(s-u) \zeta_k(u) du \right|^2 \\ & = \sum_{k,m=1}^n \frac{1}{\mu_k \mu_m} \left[\sin \left(\frac{k\pi}{l} \int_0^x \sqrt{\frac{\rho}{p}} du \right) \sin \left(\frac{m\pi}{l} \int_0^x \sqrt{\frac{\rho}{p}} du \right) R_{k,m}(t, t) \right. \\ & \quad - \sin \left(\frac{k\pi}{l} \int_0^x \sqrt{\frac{\rho}{p}} du \right) \sin \left(\frac{m\pi}{l} \int_0^y \sqrt{\frac{\rho}{p}} du \right) R_{k,m}(t, s) \\ & \quad - \sin \left(\frac{k\pi}{l} \int_0^y \sqrt{\frac{\rho}{p}} du \right) \sin \left(\frac{m\pi}{l} \int_0^x \sqrt{\frac{\rho}{p}} du \right) R_{k,m}(s, t) \\ & \quad \left. + \sin \left(\frac{k\pi}{l} \int_0^y \sqrt{\frac{\rho}{p}} du \right) \sin \left(\frac{m\pi}{l} \int_0^y \sqrt{\frac{\rho}{p}} du \right) R_{k,m}(s, s) \right] \\ & \leq 4T^2 \sum_{k,m=1}^{\infty} \frac{C_{k,m}}{\mu_k \mu_m} \left(\ln^{2\delta} \left(e^{2\delta} + \frac{m\pi C_1}{2l} \right) + \ln^{2\delta} \left(e^{2\delta} + \frac{\mu_m}{2} \right) + A \right)^{\alpha} \\ & \quad \times \left(\ln^{2\delta} \left(e^{2\delta} + \frac{k\pi C_1}{2l} \right) + \ln^{2\delta} \left(e^{2\delta} + \frac{\mu_k}{2} \right) + A \right)^{1-\alpha} \cdot |\ln h|^{-2\delta}, \end{aligned}$$

where $h = \max\{|t-s|, |x-y|\}$ and where $C_1 > 0$ and $A > 0$ are some constants such that $|\sqrt{\rho(x)/p(x)}| \geq C_1$ for all $x \in [0, \pi]$ and

$$\frac{|t-s|}{2T} \leq \frac{A}{|\ln |t-s||^{2\delta}}$$

for sufficiently small $|t-s|$. □

4. CONDITIONS FOR THE EXISTENCE OF A GENERALIZED SOLUTION EXPRESSED IN TERMS OF THE COVARIANCE FUNCTION

Let $B(x, y, t, s) = \mathbb{E} \xi(x, t) \xi(y, s)$ for $(x, y, t, s) \in [0, \pi]^2 \times [0, T]^2$. Assume that

$$B(0, y, t, s) = B(\pi, y, t, s) = 0, \quad y \in [0, \pi], \quad t \in [0, T], \quad s \in [0, T],$$

$$B(x, 0, t, s) = B(x, \pi, t, s) = 0, \quad x \in [0, \pi], \quad t \in [0, T], \quad s \in [0, T].$$

Fix a pair $(t, s) \in [0, T]^2$ and extend $B(x, y, t, s)$ as a function of x and y to the whole plane $\mathbb{R}^2$ in such a way that the extension is a periodic function of period 2π in x and in y and

$$B(-x, y, t, s) = -B(x, y, t, s) = B(x, -y, t, s).$$

Such an extension exists in view of the assumptions imposed above on the function B .

Consider a particular case of the problem (4)–(6), namely let $\rho(x) = p(x)$ in (4).

We extend the functions $p(x)$ and $q(x)$ to the whole axis such that the extension is periodic with period 2π and

$$p(x) = p(-x), \quad q(x) = q(-x).$$

We also assume that the extension of $p(x)$ is twice continuously differentiable, while that of $q(x)$ is continuously differentiable.

Theorem 2. *Let $\xi(x, t)$ be a strictly φ -sub-Gaussian almost surely continuous random field whose covariance function $B(x, y, t, s)$ is continuous. Assume that $p > 1$ and $\varphi(x) = |x|^p$ for $x > 1$. If*

$$\sup_{\substack{y \in [0, \pi], \\ s \in [0, T]}} \int_{-\pi}^{\pi} \left| B(x, y, t, s) \sqrt{p(x)} - B(x + \beta, y, t, s) \sqrt{p(x + \beta)} \right| dx \leq \frac{C}{|\ln \beta|^{2\gamma}}$$

for a sufficiently small constant $\beta > 0$ and some $C > 0$ and $\gamma > 2 - 1/p$, then $u(x, t)$ defined by (7) is a generalized solution of the problem (4)–(6).

Remark 2. The above condition holds if, for example, the derivatives $\partial B / \partial x$ and $p'(x)$ are continuous.

Proof. We show that the assumptions of the theorem imply that the series

$$\sum_{k, m=1}^{\infty} \frac{((\ln k)^\alpha (\ln m)^{(1-\alpha)})^{2\delta}}{km} C_{k, m}$$

converges for some constants $\alpha \in [0, 1]$ and $\delta > 1 - 1/p$, where

$$C_{k, m} = \sup_{0 \leq t, s \leq T} |\mathbb{E} \zeta_k(t) \zeta_m(s)|.$$

Let $\alpha = \frac{1}{2}$. It is sufficient to show that

$$C_{k, m} \leq \frac{C}{(\ln k \ln m)^\gamma}, \quad C > 0, \quad \gamma > 2 - \frac{1}{p}$$

for all sufficiently large k and m . In the case under consideration,

$$X_n(x) = \frac{1}{\sqrt{p(x)}} \sqrt{\frac{2}{\pi}} \sin nx + \frac{K_n(x)}{n}, \quad |K_n(x)| \leq L$$

(see [11]). Put $B^*(x, y, t, s) = p(x)p(y)B(x, y, t, s)$.

Similarly to the proof of Theorem 2 of the paper [6] we obtain

$$\begin{aligned}
|\mathbb{E} \zeta_k(t) \zeta_m(s)| &= \left| \int_0^\pi \int_0^\pi X_k(x) X_m(y) B^*(x, y, t, s) dx dy \right| \\
&\leq \left| \frac{2}{\pi} \int_0^\pi \int_0^\pi B^*(x, y, t, s) \frac{1}{\sqrt{p(x)p(y)}} \sin kx \sin my dx dy \right| \\
&\quad + \left| \frac{\sqrt{2}}{m\sqrt{\pi}} \int_0^\pi \int_0^\pi B^*(x, y, t, s) \frac{1}{\sqrt{p(x)}} \sin kx K_m(y) dx dy \right| \\
&\quad + \left| \frac{\sqrt{2}}{k\sqrt{\pi}} \int_0^\pi \int_0^\pi B^*(x, y, t, s) \frac{1}{\sqrt{p(y)}} \sin my K_k(x) dx dy \right| \\
&\quad + \left| \frac{1}{km} \int_0^\pi \int_0^\pi B^*(x, y, t, s) K_k(x) K_m(y) dx dy \right| \\
&\leq \frac{1}{2\pi} \left(\int_0^\pi \int_{-\pi}^\pi \left| B^*(x, y, t, s) \frac{1}{\sqrt{p(x)}} - B^*\left(x + \frac{\pi}{k}, y, t, s\right) \frac{1}{\sqrt{p(x + \frac{\pi}{k})}} \right| \right. \\
&\quad \left. \times \frac{1}{\sqrt{p(y)}} |\sin kx \sin my| dx dy \right)^{1/2} \\
&\quad \times \left(\int_0^\pi \int_{-\pi}^\pi \left| B^*(y, x, s, t) \frac{1}{\sqrt{p(y)}} - B^*\left(y + \frac{\pi}{m}, x, s, t\right) \frac{1}{\sqrt{p(y + \frac{\pi}{m})}} \right| \right. \\
&\quad \left. \times \frac{1}{\sqrt{p(x)}} |\sin kx \sin my| dy dx \right)^{1/2} \\
&\quad + \frac{\sqrt{2}}{4m\sqrt{\pi}} \int_0^\pi \int_{-\pi}^\pi \left| B^*(x, y, t, s) \frac{1}{\sqrt{p(x)}} - B^*\left(x + \frac{\pi}{k}, y, t, s\right) \frac{1}{\sqrt{p(x + \frac{\pi}{k})}} \right| \\
&\quad \times |\sin kx K_m(y)| dx dy \\
&\quad + \frac{\sqrt{2}}{4k\sqrt{\pi}} \int_0^\pi \int_{-\pi}^\pi \left| B^*(y, x, s, t) \frac{1}{\sqrt{p(y)}} - B^*\left(y + \frac{\pi}{m}, x, s, t\right) \frac{1}{\sqrt{p(y + \frac{\pi}{m})}} \right| \\
&\quad \times |\sin my K_k(x)| dy dx \\
&\quad + \frac{1}{km} \int_0^\pi \int_0^\pi |B^*(x, y, t, s) K_k(x) K_m(y)| dx dy.
\end{aligned}$$

Since

$$\begin{aligned}
&\left| B^*(x, y, t, s) \frac{1}{\sqrt{p(x)}} - B^*\left(x + \frac{\pi}{k}, y, t, s\right) \frac{1}{\sqrt{p(x + \frac{\pi}{k})}} \right| \\
&= \left| B(x, y, t, s) \frac{\sqrt{p(x)}}{p(y)} - B\left(x + \frac{\pi}{k}, y, t, s\right) \frac{\sqrt{p(x + \frac{\pi}{k})}}{p(y)} \right| \\
&\leq \left| B(x, y, t, s) \sqrt{p(x)} - B\left(x + \frac{\pi}{k}, y, t, s\right) \sqrt{p\left(x + \frac{\pi}{k}\right)} \right| \cdot \frac{1}{\min_{0 \leq y \leq \pi} p(y)},
\end{aligned}$$

there exist constants $C_i > 0$, $i = 0, \dots, 4$, such that

$$|\mathbb{E} \zeta_k(t) \zeta_m(s)| \leq \frac{C_1}{\left| \ln \frac{\pi}{k} \ln \frac{\pi}{m} \right|^\gamma} + \frac{C_2}{m \left| \ln \frac{\pi}{k} \right|^{2\gamma}} + \frac{C_3}{k \left| \ln \frac{\pi}{m} \right|^{2\gamma}} + \frac{C_4}{km} \leq \frac{C_0}{(\ln k \ln m)^\gamma}$$

for sufficiently large k and m . $\square$

5. CONCLUDING REMARKS

We consider generalized solutions of a hyperbolic equation with zero initial and boundary conditions and find sufficient conditions for the existence of a generalized solution of the problem (4)–(6).

BIBLIOGRAPHY

1. V. V. Buldygin and Yu. V. Kozachenko, *On a question of the applicability of the Fourier method for solving problems with random boundary conditions*, Random Processes in Problems of Mathematical Physics, Akad. Nauk Ukrain. SSR, Inst. Mat., Kiev, 1979, pp. 4–35. (Russian) MR587658 (83i:35080)
2. E. Beisenbaev and Yu. V. Kozachenko, *Uniform convergence in probability of random series, and solution of boundary value problems with random initial conditions*, Teor. Veroyatnost. i Mat. Statist. **21** (1979), 9–23; English transl. in Theory Probab. Math. Statist. **21** (1980), 9–23. MR550238 (81g:60036)
3. E. Barrasa de la Krus and Yu. V. Kozachenko, *Boundary-value problems for equations of mathematical physics with strictly Orlicz random initial conditions*, Random Oper. Stochastic Equations **3** (1995), no. 3, 201–220. MR1354813 (96h:60009)
4. V. V. Buldygin and Yu. V. Kozachenko, *Metric Characterization of Random Variables and Random Processes*, TViMS, Kiev, 1998; English transl., American Mathematical Society, Providence, RI, 2000. MR1743716 (2001g:60089)
5. B. V. Dovgay and Yu. V. Kozachenko, *The condition for application of Fourier method to the solution of nonhomogeneous string oscillation equation with φ -sub-Gaussian right hand side*, Random Oper. Stochastic Equations **13** (2005), no. 3, 281–296. MR2165325 (2006h:60042)
6. B. V. Dovgay and Yu. V. Kozachenko, *The properties of solution of nonhomogeneous string oscillation equation with φ -sub-Gaussian right side*, Random Oper. Stochastic Equations **17** (2009), no. 3, 221–241. MR2574102 (2010k:35281)
7. M. A. Krasnosel'skiĭ and J. B. Rutitskiĭ, *Convex Functions and Orlicz Spaces*, Fizmatgiz, Moscow, 1958; English transl., P. Noordhoff Ltd., Groningen, 1961. MR0126722 (23:A4016)
8. Yu. V. Kozachenko and G. I. Slivka, *Justification of the Fourier method for a hyperbolic equation with random initial conditions*, Teor. Imovir. Mat. Stat. **69** (2003), 63–78; English transl. in Theory Probab. Math. Statist. **69** (2004), 67–83. MR2110906 (2005k:60127)
9. Yu. V. Kozachenko and Yu. A. Kovalchuk, *Boundary value problems with random initial conditions, and functional series from $\text{Sub}_\varphi(\Omega)$* , Ukrain. Mat. Zh. **50** (1998), no. 4, 504–515; English transl. in Ukrainian Math. J. **50** (1999), no. 4, 572–585. MR1698149 (2000f:60029)
10. R. Giuliano Antonini, Yu. Kozachenko, and T. Nikitina, *Spaces of φ -sub-Gaussian random variables*, Rend. Accad. Naz. Sci. XL Mem. Mat. Appl. **XXVII** (2003), no. 1, 95–124. MR2056414 (2005f:60036)
11. G. N. Polozhiĭ, *Equations of Mathematical Physics*, Vysshaya shkola, Moscow, 1964. (Russian)

DEPARTMENT OF OPERATIONS RESEARCH, FACULTY FOR CYBERNETICS, NATIONAL TARAS SHEVCHENKO UNIVERSITY, GLUSHKOV AVENUE 6, KYIV 03127, UKRAINE
E-mail address: bogdov@gmail.com

Received 20/MAY/2009

Translated by N. SEMENOV