ON THE PROJECTIVE AXIOMS OF GEOMETRY*

BY

ELIAKIM HASTINGS MOORE

Introduction.

In the present paper I wish to consider the axioms called by HILBERT† (1899) *the axioms of connection and of order* (I 1–7, II 1–5 of HILBERT's list), and called by SCHUR‡ (1901) *the projective axioms* of geometry.

HILBERT states (l. c., pp. 3, 21) that his body of axioms consists of independent axioms, that is, that no one of the axioms is logically deducible from the remaining axioms. This statement, accepted by HÖLDER§ and SOMMER,|| is not accepted by SCHUR,¶ who holds that HILBERT's axioms I 3, 4, 5 are consequences of the axioms I 1–2, 7, II 1–5. For reasons explained below, this last remark cannot be accepted as valid.

There are, however, redundancies** in HILBERT's axioms I–II, the two axioms

* Presented to the Society December 28, 1901. Received for publication December 31, 1901.

† HILBERT, *Grundlagen der Geometrie, Festschrift zur Feier der Enthüllung des Gauss-Weber-Denkmal in Göttingen*, Leipzig, 1899.

‡ SCHUR, *Ueber die Grundlagen der Geometrie*, *Mathematische Annalen*, vol. 55 (1901), pp. 265–292.

§ HÖLDER, *Anschauung und Denken in der Geometrie*, Leipzig, 1900; cf. p. 35.

|| SOMMER, *Hilbert's Foundations of Geometry*, *Bulletin*, vol. 6 (1900), p. 292.

¶ loc. cit., pp. 266, 267, 271.

** As to the independence of the axioms I 1–7, II 1–5 HILBERT (l. c., p. 21) refers to his Göttingen lectures of the winter semester 1898–99, as reported by DR. VON SCHAPER and manifolded. By the kindness of DR. BOSWORTH-FOCKE I have seen a copy of this report. Proofs are given that (1) I 2 is independent of I 1; (2) I 5 of I 1–4, 6–7; (3) I 6 of I 1–5, 7; (4) II 4 of I 1–7, II 1–3; (5) II 3 of I 1–7, II 1–2, 4; (6) II 5 of I 1–7, II 1–4.—In contravention of the remark of SCHUR, I prove below further that (7) I 3 is independent of I 1–2, 4–7, II 1–5; (8) I 4 of I 1–2, 6–7, II 1–5; (9) I 5 of I 1–3, 6–7, II 1–5.

Each of these proofs involves the exhibition of a particular geometry, in which, e. g., in case (9) the relations I 1–3, 6–7, II 1–5 hold while the relation I 5 does not hold. This exhibition is in terms of a science supposed to exist without self-contradictions, and the proposition of independence is contingent on the correctness of this supposition. As science of reference one prefers the usual analysis of real numbers or a science (e. g., the analysis of complex numbers; euclidean n -dimensional geometry) whose existence is deducible from that of analysis.

Such proofs were used by PEANO in investigations on the foundations of arithmetic and geometry (l. c., p. 62) and they play a brilliant part in the work of HILBERT.

I 4, II 4 being deducible from I 1-3, 5, 7, II 1-3, 5. I establish these two conclusions in § 5 of this paper.*

In § 1 I give a simplified arrangement of the projective axioms of n -dimensional geometry ($n \geq 2$). I use as elements the point and the line and the segment of a line, the line and the segment being certain sets of points, and give convenient definitions of the plane or 2-space, and the k -spaces in general ($0 \leq k \leq n$); the properties of the k -spaces are developed in § 3.

It is convenient here briefly to characterize the systems of projective axioms of 3-dimensional geometry given by PASCH † (1882), PEANO ‡ (1889, 1894). INGRAMI § (1899) and HILBERT (1899).

PASCH undertook to make pure geometry in a strict sense a purely deductive (abstract) science based on certain assumed (abstract) notions subject to certain assumed relations. He introduced as basal notions for geometry the point, the linear-segment (*gerade Strecke*) and the planar-segment (*ebene Fläche*), in terms of which the line and the plane were defined. The linear-segment and the planar-segment are certain sets of points.

PEANO, following PASCH, retained the point and the linear-segment; but he defined the line and the plane by means of these elements alone. His system, as simplified by INGRAMI, is given by SCHUR (l. c., p. 267 ff.)

HILBERT has as basal notions the point, the line, the segment of a line (connected with the notion *between*) and the plane. The line and the plane are by the Hilbert axioms (I 1, 7₁; I 3, 7₂) connected with certain sets of points (lying on the line; the plane), in such a way that we may (and for simplicity we do) identify them with those sets of points. For convenience of reference I set down the Hilbert axioms I 1-7, II 5; the axioms II 1, 2, 3 are in effect the axioms 2, 5, 3 of § 1 of this paper, while axiom II 4 is quoted in the theorem of § 2.

I 1, 2, 7₁. *Two distinct points A, B determine in every case a line g ; we set $AB = g$ or $BA = g$. Any two distinct points of a line determine this line; that is, if $AB = g$ and $AC = g$, and $B \neq C$, then $BC = g$. On every line there are at least two points.*

I 3, 4, 5, 7₂. *Three non-collinear points A, B, C determine in every case a plane π ; we set $ABC = \pi$. Any three non-collinear points of a plane determine this plane. If two points A, B of a line g lie in a plane π , then every point of the line g lies in π . On every plane there are at least three non-collinear points.*

* This paper has been prepared in connection with my current Chicago seminar-course on the foundations of geometry and analysis, and queries and remarks of members of this course, in particular, of Mr. O. VEULEN, have been a source of much stimulus.

† PASCH, *Vorlesungen über neuere Geometrie*, Leipzig, 1882.

‡ PEANO, *I principii di Geometria*, Torino, 1889 (cited in the following paper), *Sui fondamenti della Geometria*, *Rivista di Matematica*, vol. 4 (1894), pp. 51-90.

§ INGRAMI, *Elementi di Geometria per le scuole secondarie superiori*, Bologna, 1899 (citation of SCHUR).

I 7₃, 6. *There exist at least [one plane and] * four non-coplanar points. If two planes α, β have a point A in common, then they have at least a second point B in common.*

II 5. *Let A, B, C be three non-collinear points and g a line lying in the plane ABC and containing no one of the points A, B, C ; if the line g contains a point within the segment AB , then it always contains either a point within the segment AC or a point within the segment BC .*

Here every one of two (three) points which determine a line (plane) is defined as a point of or on or in or lying in the line (plane).—In interpretation we agree: (1) I 2 has the full meaning of its original statement, viz.: If $AB = g$ and $CD = g$, and $B \neq D$, then $BD = g$. (2) I 4 has the specification: If $A_1A_2A = \pi$, $B_1B_2B = \pi$ and $C_1C_2C = \pi$, and A, B, C are non-collinear, then $ABC = \pi$. (3) In I 3, if $ABC = \pi$, then $BAC = \pi$ and $ACB = \pi$. (4) The determinations in I 1, 3 are unique determinations.

Clearly the body of axioms of a system depends essentially upon the choice of the basal notions of the system. In this connection a remark is pertinent with respect to one's attitude concerning the foundations of geometry. I suppose that if geometry † is taken to be a natural science—the science or a science of the space in which or according to which we live—it would, as is contended by PASCH and PEANO, be undesirable to introduce the line as a basal notion. The linear-segment seems to be a more fundamental notion. But we may discriminate between that part of geometry which establishes a body of postulates based as directly as may be on spatial experience or intuition, and that part which consists in the organization of the science on the basis of the accepted body of axioms; and so we understand that it may in the development of the theory be convenient to replace the body of primary notions and relations by another body of notions and relations, less fundamental, but, with respect to the deductive geometry, more convenient. I suppose that with this thought HILBERT introduced the line and the plane as basal notions in his abstract geometry.—It is understood that greater generality would be obtained by introducing the axioms as valid, as one says, for a limited region of space. In this connection reference is made to KLEIN's introduction ‡ of the ideal elements of projective geometry without the use of the parallel axiom, and to the remarks of PASCH (pp. 4, 18, 126), of PEANO (p. 75) and of SCHUR (pp. 267, 274).

As has been stated, SCHUR has remarked that in the body of projective axioms of HILBERT the axioms I 3, 4, 5 are redundant. This criticism is, however,

* The bracketed addition to the original form of I 7₃ is a necessary addition; it is an implication of the remarks with which HILBERT introduces the axioms I.

† For $n = 3$.—In case $n > 3$ the geometry is perhaps essentially abstract.

‡ *Mathematische Annalen*, vol. 6, p. 134.—Cf. SCHUR, *ibid.*, vol. 39, p. 113, and vol. 55, p. 274.

incorrect. For if in the ordinary euclidean 3-dimensional space we take the points, the lines, the segments of lines, and two intersecting spheres as the points, the lines, the segments of lines, and the (only) planes of a new geometry one has a geometry in which I 1-2, 6-7, II 1-5 are satisfied, while all the statements I 3, 4, 5 are invalid. Thus I 3, 4, 5 are not logical consequences of the remaining axioms I, II. Indeed, one may prove* that (1) I 3 is independent of I 1-2, 4-7, II 1-5; (2) I 5 of I 1-3, 6-7, II 1-5. These two remarks together with the fact (already stated) that I 4 is deducible from I 1-3, 5, 7, II 1-3, 5 furnish a satisfactory account of the rôles of the axioms I 3, 4, 5 in the Hilbert system I, II.

The remark of SCHUR was connected with the fact that, as PEANO pointed out, the plane is capable of explicit definition. Apparently SCHUR directly identified the Hilbert plane with the Peano plane. But in a particular Hilbert geometry the plane postulated by HILBERT is to be identified with the plane defined by PEANO certainly only by the mediation (to some extent) of the axioms I 3-7, II 5 by which alone the Hilbert plane is conditioned. Pursuing this thought I find (§ 5) that by replacing in the Hilbert system axioms I 3, 5, 7, II 5 by axioms considerably milder one is still able to make this identification and thus to prove the axioms in their original form, as used by HILBERT.

Evidently, with PEANO and SCHUR, we may better define the plane explicitly than thus, by the mediation of axioms, implicitly. The implicit definitions of the k -spaces in n -dimensional geometry (cf. HILBERT, loc. cit., p. 71) would be still more undesirable. The explicit definitions of § 1 seem to be simpler than those of § 4, the analogues of the Peano-Ingrami definitions.

To indicate briefly the arrangement of this paper :

In § 1 I formulate a body of seven projective axioms of n -dimensional geometry ($n \geq 2$); these for $n = 3$ correspond with HILBERT's axioms I, II as follows :

$$(1; 2; 3; 4; 5; \underline{6}; \underline{7}) = (I\ 1, 2, 7_1; II\ 1; II\ 3; \underline{II\ 5}; II\ 2; \underline{I\ 7_2, 7_3}; \underline{I\ 6});$$

the axioms I 3, 4, 5 are omitted, the plane (and the k -spaces, in general) being explicitly defined; in the cases underscored the correspondence is only approximate; in all cases the effort has been to give clear and precise expression to the meaning intended to be conveyed.—Axiom 4, the triangle-transversal axiom, is funda-

* A geometry for case (1) is the ordinary euclidean 3-dimensional geometry, with the omission of one plane; one for case (2) is the ordinary euclidean 2-dimensional geometry with the modification that the general plane ABC is the set of points, O apart, of the lines AO, BO, CO , the point O being the center of the circle inscribed in the triangle ABC .

One has the feeling that, even on the assumption that the Hilbert plane ABC is defined explicitly as identical with the Peano plane ABC , the criticism of SCHUR remains incorrect, in view of the fact that the plane order axiom II 5 relates to a line supposed to lie in the plane and is accordingly in this respect milder than PEANO's two triangle axioms fundamental in the theory of the Peano plane. A geometry showing the correctness of this conjecture I have not found.

mental in the theory of the k -spaces; it replaces (its two corollaries 3, 4) the two triangle axioms introduced by PEANO as fundamental for the theory of the plane, which replaced PASCH's plane order axiom, retained by HILBERT as axiom II 5. —Axiom 7, the axiom of limitation, in agreement with SCHUR (loc. cit., p. 271), I think desirably replaces HILBERT's I 6.

HILBERT's axiom II 4 is in § 2 proved as a theorem depending on axioms 1–6, in particular on axiom 4, the triangle-transversal axiom.

In § 3 the properties of the k -spaces are developed on the basis of the definitions of § 1 and the axioms 1–6. Comparison with another type of definition is made in § 4.

In § 5 I prove that in HILBERT's system with the body of axioms I 1–3, 5, 7, II 1–3, 5, or with the milder body of axioms I 1, 2, 3', 5', 7₁, 7₂', 7₃', II 1–3, 5' one may identify the Hilbert plane with the plane defined in § 1, and that the axioms 1–6 ($n = 3$) of § 1 hold. Then it follows (by §§ 2, 3, 4) that the Hilbert axioms I 4, II 4 are deducible from the Hilbert axioms I 1–3, 5, 7, II 1–3, 5.

§ 1. *The projective axioms of n -dimensional geometry.*

In an abstract deductive geometry (of space of n dimensions, $n \geq 2$) we consider a set of things called *points*, having properties fully* determined by the body of postulates or axioms laid down as the basis of the geometry. The space under consideration, call it *the fundamental space*, is the set of all points under consideration.

The projective axioms involve certain sets of points, called *lines*, and certain sets of collinear points, called *segments* (of lines). These lines and segments receive definition only implicitly by the mediation of the axioms.

The projective axioms may be grouped as follows:

1–4: *Axioms of conditioned existence and of definition;*

5–6: *Axioms of absolute existence;*

7: *Axiom of limitation.*

The reader will understand that the designations attached to the various groups of axioms are intended to indicate somewhat clearly, although roughly, what seem to be the principal functions of the various axioms in the body of axioms.

AXIOM 1: THE LINE. *Two distinct points A, B determine uniquely a set of points, the line AB . The points A, B belong to or lie on the line AB . A line is a set of points thus determined by certain two and indeed by any† two distinct points belonging to it.*

*In so far as they pertain to the arbitrary geometry of the system of geometries in question.

†This statement may be replaced by these three: *The line AB contains and thus is identical with the line BA . The line AB contains the line AC , where C is any point not A of the line AB . The line AC contains the point B .*

AXIOM 2: THE SEGMENT OF A LINE. *Two distinct points A, B of a line determine uniquely a set of points distinct from A, B and lying on the line; this set of points is the segment AB , with extremities A, B . The segment AB contains and thus is identical with the segment BA . A segment is a set of points thus determined by two distinct points, its extremities.*

Def. 1. A point C of the segment AB is said to lie between or to separate* the extremities A, B of the segment, in notation ACB or BCA .—The notation indicates the order-relation of the three points.

AXIOM 3: THREE COLLINEAR POINTS. *Of three distinct collinear points A, B, C one and only one lies on the segment determined by the other two: of the three order-relations, ABC, BCA, CAB one and but one† holds.*

AXIOM 4: THE TRIANGLE* WITH TRANSVERSAL LINE. *A line which cuts‡ one side of a triangle externally and another side internally cuts the third side internally.*

In other words (cf. fig. 1):

AXIOM 4. *If A, B, C are three non-collinear points and if the line $\frac{DF}{DE}$ cuts the lines $\frac{BC}{BC}, \frac{AB}{CA}$ in points $\frac{D}{D}, \frac{F}{E}$ respectively, where $\frac{BCD}{BCD}$ and $\frac{AFB}{CEA}$, then it cuts the line $\frac{CA}{AB}$ in a point $\frac{E}{F}$ such that $\frac{CEA}{AFB}$.*

FIG. 1.

FIG. 2.

Notation. We denote the axioms into which axiom 4 has just been analyzed by the notations§

$$\begin{matrix} 4_1 \\ 4_2 \end{matrix} \quad \text{or} \quad \begin{matrix} 4_{11}, 4_{12} \\ 4_{21}, 4_{22} \end{matrix}.$$

* Similarly, a set Γ of points having a point C in common with the segment AB is said to separate A and B .

† Here, as Mr. VEULEN has remarked, it is obviously sufficient to state that the relation ACB does not hold if the relation ABC does hold.

‡ For brevity I use, without explanation, a few geometric terms whose meaning in the abstract geometry in question is supposed to be sufficiently clear.

§ Similar self-explanatory partitional suffix-notations will be used elsewhere in this paper.

Remark. Axiom 4_{22} is a logical consequence* of axioms 4_{11} , 4_{12} and 4_{21} together with axioms 1–3.

Remark. The following seven theorems depend so immediately on axiom 4 by the mediation of the preceding axioms that I insert them here as corollaries of axiom 4.

Cor. 1. (In the notation of the second statement of axiom 4.)—*The three points D, E, F are distinct and such that DEF . The intermediate point E lies on the side adjacent to the point D lying on its side externally.*—One applies axiom 4_1 to the transversal AC of the triangle BDF .

Cor. 2. *If a line cuts two sides of a triangle externally and (cyclically) similarly, then it cuts the third side externally but dissimilarly, and the point of intersection with the third side lies between the points of intersection with the other two sides.*—That is (cf. fig. 2), in analogous notation: *If $\frac{BCD}{BCD}$ and $\frac{CAE}{ABF}$, then $\frac{BAF}{ACE}$ and $\frac{DFE}{DEF}$.*—One applies axiom 4_2 and cor. 1 to the transversal $\frac{BA}{AC}$ of the triangle $\frac{CDE}{BDF}$.—In this case the two statements are equivalent.

FIG. 3.

FIG. 4.

Cor. 3. *The two segments AD, BE from two vertices A, B of a triangle ABC to two points D, E of the respectively opposite segments BC, CA have in common a point G .*—One applies axiom 4_1 and cor. 1 to the transversal AD of the triangle BCE . Cf. fig. 3.

Cor. 4. *The point G being any point of the segment AD from the vertex A of a triangle ABC to a point D of the opposite segment BC , there exists a*

* By use of axioms 4_{21} and 3 we see that it is sufficient to prove that the relations ABF and BAF are impossible, and here we use besides axiom 4_1 itself its corollary 7 (of the text) which is independent of axiom 4_2 .

The relation ABF is impossible. For by 4_1 the transversal FEE of the triangle ABC would yield the relation BDC .

Further, the relation BAF is impossible. For, in conjunction with DEF , it would imply that the line AEC meets the triangle BDF in inner points of its three sides, which by cor. 7 is impossible; in conjunction with DFE , it would imply, by 4_1 for the triangle CED with the transversal BAF , the relation CAE ; and similarly, in conjunction with EDF , it would lead to the relation ACE .

point E of the segment CA such that the segment BE contains the point G .—One applies axiom 4_2 and cor. 1 to the transversal BG of the triangle ACD .

Cor. 5. In a triangle ABC the line AD , containing the vertex A and a point D of the opposite segment BC , and the segment EF , from a point E of the segment CA to a point F of the segment AB , have in common a point* H .—One applies axiom 4_1 to the line AD as a transversal first of the triangle BCE and then of the triangle BEF . Cf. fig. 4.

Cor. 6. In a triangle ABC the line AH , containing the vertex A and a point H of the segment EF , from a point E of the segment CA to a point F of the segment AB , contains a point D of the segment BC .—One applies axiom 4_2 to the line AH as a transversal first of the triangle BEF and then of the triangle BCE .

Cor. 7. No transversal meets the three sides of a triangle internally.—If three points D, E, F on the sides BC, CA, AB of the triangle ABC were collinear, then, for instance, E would lie between D and F , and further, however, by cor. 5, the line AD would meet the segment EF in a point—in fact, the point D —lying between E and F .

Def. 2 ($k = 2$). Three distinct points A_1, A_2, B_0 are (in the order specified) independent if B_0 does not lie in the line A_1A_2 .

Def. 3 ($k = 2$). Three independent points A_1, A_2, B_0 determine uniquely a plane, or 2-space, $A_1A_2B_0$. The plane $A_1A_2B_0$ is the aggregate set of all points A of the line A_1A_2 , of all points C separated (cf. def. 1) from B_0 by the line A_1A_2 , and of all points B separated from at least one of these points C by the line A_1A_2 .

Def. 2 ($k = 3$). Four points A_1, A_2, A_3, B_0 are independent if A_1, A_2, A_3 are independent and B_0 does not lie in the plane $A_1A_2A_3$.

Def. 3 ($k = 3$). Four independent points A_1, A_2, A_3, B_0 determine uniquely a 3-space $A_1A_2A_3B_0$. The 3-space $A_1A_2A_3B_0$ is the aggregate set of all points A of the 2-space $A_1A_2A_3$, of all points C separated from B_0 by the 2-space $A_1A_2A_3$, and of all points B separated from at least one of these points C by the 2-space $A_1A_2A_3$.

Def. 2 ($k = k$). The $k + 1$ points $A_1, \dots, A_k, B_0$ are independent if the first k points $A_1, \dots, A_k$ are independent and the last point B_0 does not lie in the $(k - 1)$ -space $A_1 \dots A_k$.

Def. 3 ($k = k$). † A set of $k + 1$ independent points $A_1, \dots, A_k, B_0$ de-

* It is to be noted that on the basis of axioms 1-5, the point H lies not only on the segment EF but also on the segment AD .—Denoting by G the point of intersection of the line AD and the segment BE , we have, by cor. 1, AGD and AHG , from which, by the theory of the order-relations of four collinear points (§2) we have indeed $AHID$.

† Denoting a point as a 0-space, we define similarly the notions: Two independent points A_1, B_0 ; the 1-space A_1B_0 . Then within a fundamental space, which contains a 2-space, by the theory of §2, the 1-space A_1B_0 and the line A_1B_0 are identical. This remark makes general, for positive integral values of k , the notions of $k + 1$ independent points and of a k -space.—Evidently it does not involve an explicit definition of the notion line.

termines uniquely a k -space $A_1 \cdots A_k B_0$; the k -space $A_1 \cdots A_k B_0$ is the aggregate set of all points A of the $(k-1)$ -space $A_1 \cdots A_k$, of all points C separated from B_0 by the $(k-1)$ -space $A_1 \cdots A_k$, and of all points B separated from at least one of these points C by the $(k-1)$ -space $A_1 \cdots A_k$.

AXIOM 5. *On the line AB determined by two distinct points A, B there exists a point C on the segment AB and there exists a point D distinct from A such that the segment AD contains B .*

AXIOM 6. *There exists in the fundamental space a set of $n+1$ independent points $A_1, \dots, A_n, A_{n+1}$ (where n is any particular integer greater than 1); and thus there exists a set of $k+1$ independent points $A_1, \dots, A_k, A_{k+1}$ for every integer k ($1 \leq k \leq n$).*

Cor. *The fundamental space contains k -spaces ($1 \leq k \leq n$).*

AXIOM 7. *There exists in the fundamental space* no $(n+1)$ -space.*

Cor. *The fundamental space contains no set of $n+2$ independent points.*

In the system of axioms, axioms 6 and 7 may be condensed into the following

AXIOM. *The fundamental space is an n -space.*

§ 2. *The order-relations of four collinear points. Deduction from axioms 1-4, 5₁, 6 ($n \geq 2$).*

In the usual geometry one will admit the validity of the following four statements concerning the order-relations of four distinct collinear points A, B, C, D .

1. *If CAB and ABD , then CAD and CBD .*
2. *If CAB and ADB , then CAD and CDB .*
3. *If CAB and DAB , then either CDA or DCA ; either CDB or DCB ; if CDA , then CDB ; and if CDB , then CDA .*
4. *If ACB and ADB , then either ACD or ADC ; either CDB or DCB ; if ACD , then CDB ; and if CDB , then ACD .*

Concerning these four statements we establish the fundamental

THEOREM. *The statements 1, 2, 3, 4 are evidently equivalent to Hilbert's linear order axiom II 4:*

To any four distinct points of a line the notation A, B, C, D may always be assigned in such a way that ABC, ABD, ACD, BCD . Furthermore, these statements are theorems of a geometry in which axioms 1-4, 5₁ and 6 for $n \geq 2$ hold.

We prove, in the first place, by use of the axioms 2, 3 and 5₁, that the

* That in an n -space there exists no set of $n+2$ independent points and thus no $(n+1)$ -space is a theorem. (Cf. § 3.)

statements are consequences* of the single statement 2_1 , which, in modified notation, is:

2_1 . If ABC and ACD , then BCD .

One observes readily that statements 3 and 4 follow from statements 1 and 2.

Now statement 2_2 follows from 2_1 , for we have CAB and ADB , and if we had DBC , then by 2_1 from DBC and BAC we should have DBA in contradiction with ADB ; and if we had BCD , then by 2_1 from DCB and CAB we should have DCA , and, at the same time, from ADB and DCB we should have ADC ; these two conclusions are contradictory; hence we have, as stated, CDB .

Next the statements 1_1 and 1_2 are evidently equivalent, and we prove the truth of 1_1 on the assumption that 2_1 and consequently 2_2 are valid.

Changing the notation, we have ABC and BCD and are to prove ABD . If we had ADB , then by 2_1 from CBA and BDA we should have CBD , which is impossible. We must prove that DAB is impossible. Suppose we had DAB with ABC and BCD . If we had ACD , then by 2_2 from DCA and CBA we should have DBA , which is impossible. If we had DAC , then by 2_2 from DAC and ABC we should have DBC , which is impossible. Hence we should have CDA , and so the four points A, B, C, D would have the relations ABC, BCD, CDA, DAB ; this set of four relations we denote by the notation $(ABCD)$ cyclic.

Accordingly we have at least proved that if ABC and BCD , then either ABD or $(ABCD)$ cyclic, and so likewise either DCA or $(DCBA)$ cyclic; that is, if ABC and BCD , then either ABD and ACD or else $(ABCD)$ cyclic.

We consider the case $(ABCD)$ cyclic. By axiom 5₁ there is in this case on the line $ABCD$ a point O different from A, B, C, D and such that DOA . By 2 from CDA, DOA, DAB we have CDO, COA, DOB, OAB .

Further, from BCD and CDO we have, in accordance with the alternative above proved, either BCO and BDO or else $(BCDO)$ cyclic. The former of these alternatives contradicts the result DOB obtained previously. Thus we have $(BCDO)$ cyclic, and from it DOB, OBC .

Similarly, from OAB and ABC we have either OAC and OBC or else $(OABC)$ cyclic. The latter of these alternatives is contradictory to the result OBC just obtained, while the former is contradictory to the result COA obtained earlier. Thus, by the intervention of the point O , we recognize that the case $(ABCD)$ cyclic is impossible. Hence 1 follows from 2_1 .

Accordingly we have proved that the four statements 1-4 depend, by use of the axioms 2, 3 and 5₁, upon the single statement 2_1 :

* Mr. VEBLEN has found that HILBERT's axiom II 4 is deducible from the two statements:—
If ABC and ACD , then ABD . If ABD and ACD , then ABC or CBD .

If ABC and ACD , then BCD .

We are to prove further, by the use of axioms 1, 6 in addition to those previously mentioned, that this statement 2₁ depends upon axiom 4, the triangle-transversal axiom.

This will be proved in three parts:

(α) *The relations ABC , ACD , CDB cannot hold simultaneously.*

(β) *If ABC and ACD , then ABD .*

(γ) *The relations ABC , ACD , DBC cannot hold simultaneously.*

Here (β) depends upon (α) and is a lemma for (γ), while (α) and (γ) together yield the theorem in question.

In proof of (α): We take a point E , not on the line $ABCD$; we draw the distinct lines EC , EB ; and on the segment EC we select a point F . Then, in accordance with axiom 4 and its first corollary as applied to the transversal AF of the triangle EBC , in view of the relations ABC and EFC , we see that the line AF cuts the segment EB in a point G for which EGB and AGF . In the triangle EBC the segment FD cutting the two segments EC , BC (in view of the relation CDB) is itself cut by the line CG , in accordance with cor. 5 of axiom 4, in a point H for which FHD . Then (in view of the relation ACD) the line CGH cuts the three sides of the triangle ADF in three inner points. But this, by cor. 7 of axiom 4, is impossible. Accordingly statement (α) is proved.

In proof of (β): The case ABC , ACD , BAD (that is, DCA , DAB , ABC) is excluded by (α), and the case ABC , ACD , ADB leads by (α) to this impossibility that no one of the three points B , C , D is between the other two. Accordingly statement (β) is proved.

In proof of (γ): We take a point E , not on the line $ABCD$; we draw the line EC ; and on the segment EC we select a point F . Then, in view of EFC and ACD , the transversal AF cuts the side ED of the triangle ECD in a point G , where EGD and AFG .—Further, in view of ABC and AFG , the transversal GB of the triangle ACF cuts the segment CF in a point H ; for this point H , in view of CHF and CFE , we have by (β) the relation CHE . Then the transversal BGH , in view of DBC , DGE and CHE , meets the three sides of the triangle DCE internally, and this is impossible.

§3. *Properties of the k -spaces of the fundamental space.*

We suppose that axioms 1–6 with $n \geq 2$ hold for the fundamental space under consideration in §3. By the first statement of the following theorem it appears that a k -space of the fundamental space is a fundamental k -space in the sense of §1, the axioms 1–7 holding with k as n ; those points, lines, segments of the fundamental space, which lie in the k -space, being directly identified with the points, lines, segments of the fundamental k -space.

THEOREM.—*In a fundamental space for which the axioms 1–6 with $n \geq 2$ are valid, the sets of independent points and the k -spaces have the following properties:*

1. *Any two distinct points of a k -space determine a line or 1-space lying entirely in it.*
2. *A k -space is determined by any set of $k + 1$ independent points lying in it.*
3. *A k -space contains no set of $k + 2$ independent points.*
4. *Any $k' + 1$ independent points of a k -space ($0 \leq k' \leq k$) determine a k' -space lying entirely in it.*
5. *If $k + 1$ points are independent when taken in a certain order, they are independent when taken in any order.*
6. *A $(k - 1)$ -space lying in a k -space separates the points of the k -space which do not lie in the $(k - 1)$ -space into two sets or parts in such a way that two points of ^{the same part are not} _{opposite parts are} separated by the $(k - 1)$ -space.*
7. *In a k -space, a p -space and a q -space ($p + q \geq k$) having a common point have in common precisely an r -space, where r is a definite integer such that $r \geq p + q - k$ and (by 3) $0 \leq r \leq p$, $0 \leq r \leq q$.*

In view of the theorem of § 2 this theorem is true for the case $k = 1$, a 0-space being a point. Its truth for the general k may be proved by the induction from k to $k + 1$. The general step of induction will however be clearer if the proof for the case $k = 2$ is given by itself.

For the case $k = 2$ we consider three independent points A_1, A_2, B_0 . We denote by a the line $A_1 A_2$; by $\{A\}$ the set of points A of the line a ; by $\{C\}$ the set of points C separated from B_0 by a ; with respect to a particular point C , by $\{B_C\}$ the set of points B_C separated from the point C by a ; by $\{\{B_C\}\}$ the aggregate of all such sets $\{B_C\}$. In these notations the plane $\pi = A_1 A_2 B_0$ is by def. 2 of § 1 a certain set $\pi = \{P\}$ of points P , viz:

$$\{P\} = \{A\} + \{C\} + \{\{B_C\}\}.$$

The proof of the statements of the theorem may easily be made to depend on the following three remarks:

The plane $\pi = A_1 A_2 B_0 = a B_0$ contains all the lines of the type AB_0 .

The two planes $\pi = A_1 A_2 B_0$ and $\pi' = A_1 A_2 B'_0$, where B'_0 is any point P of π not in a , are identical.

The two planes $\pi = A_1 A_2 B_0$ and $\pi' = A_1 B_0 A_2$ are identical.

In proof of the first remark the theorem of § 2 is at once applicable.

In proof of the second remark we have in mind axioms 4 and 5 and the theorem of § 2. In the plane $\pi = A_1 A_2 B_0 = a B_0$, the line a does not separate two points C . Further the sets $\{B_{C_1}\}$, $\{B_{C_2}\}$ connected with two distinct points C_1, C_2 of the set $\{C\}$ are identical; this is true if the line $C_1 C_2$ contains a

point A , and likewise, if the line C_1C_2 contains the point B_0 , and further if the two lines B_0C_1 , B_0C_2 are distinct, in view of the fact that on the line B_0C_2 there is a point C_3 of the set $\{C\}$ such that the line C_1C_3 contains a point A . Thus the various sets $\{B_c\}$ are identical with one another; we set $\{B_c\} = \{B\}$ and have

$$\{P\} = \{A\} + \{C\} + \{B\} = \{A\} + \{B\} + \{C\}.$$

The sets $\{B\}$, $\{C\}$ have no points in common, since two points C_1, C_2 or B_1, B_2 are not separated by a , while two points B, C are separated by a . The line a separates the plane π into the two parts $\{B\}$, $\{C\}$. It is now easy to see that the plane $\pi = A_1A_2B_0$ is identical with the plane $\pi' = A_1A_2B'_0$, if B'_0 is any particular point C , and so, by repetition of the argument, also if B'_0 is any particular point B . Accordingly the second remark is proved.

From the second remark a point P of the plane $\pi = A_1A_2B_0 = aB_0$ is a point A of the line $a = A_1A_2$, or a point A' of line $a' = A_1B_0$, or a point P such that for certain two distinct points A_3 of a , A'_3 of a' the relation A'_3A_3P holds; in the last case selecting a point A_4 of a such that $A_3A_1A_4$, we find a point A'_4 of a' such that $A_4A'_4P$. Accordingly the planes $A_1A_2B_0$ and $A_1B_0A_2$ are identical.

We proceed to prove the truth of the theorem in question for the case $k = m + 1$ on the assumption of its validity for the case $k = m$.

We consider the $(m + 1)$ -space $\Sigma = A_1A_2 \cdots A_m A_{m+1} B_0 = aB_0$. One proves as before (but with the use of statement 1 of the theorem for $k = m$) that, in notations analogous to those previously used,

$$\Sigma = \{S\} = \{A\} + \{B\} + \{C\},$$

and that the $(m + 1)$ -space $\Sigma = aB_0$ is identical with the $(m + 1)$ -space $\Sigma' = aB'_0$, where B'_0 is any point of Σ not in a .

Further, one sees that the $(m + 1)$ -space Σ contains the m -space a , that all lines joining points S to points A lie in Σ , and that the m -space a divides the space Σ into two parts as specified in statement 6.

By statement 2 for the m -spaces, the space Σ may then be obtained from any $m + 1$ independent points of a together with any point S of Σ not in a .

Statements 1 to 6 will follow easily from the remarks already made, when we prove that the two $^*(m + 1)$ -spaces,

$$\Sigma = A_1 \cdots A_m A_{m+1} B_0, \quad \Sigma' = A_1 \cdots A_m B_0 A_{m+1}$$

are identical, and in proof of this last statement it is convenient to modify the notations. The $(m - 1)$ -space $A_1 \cdots A_m$ is denoted by $d = D_1 \cdots D_m$; the

* One uses statement 3 for $k = m$.

points A_{m+1} , B_0 are denoted by A_0 , A'_0 ; and the lines $D_m A_0$, $D_m A'_0$ are denoted by l , l' . We have then

$$a = dA_0 = dl, \quad a' = dA'_0 = dl',$$

$$\Sigma = aA'_0 = al', \quad \Sigma' = a'A_0 = a'l.$$

We see that the statement is proved when we prove that every point S of Σ is a point S' of Σ' . This is true evidently for a point S which is a point A of a or a point A' of a' or, by the discussion of the third remark for $k = 2$, a point of the plane $D_n A_0 A'_0 = D_n A'_0 A_0 = ll' = l'l$.—We consider any other point S of Σ . Such a point S is collinear with certain points A of a and L' of l' , where $L'AS$; and in its turn A , lying in $a = dl$ and lying neither in d nor in l , is collinear with certain points D of d and L of l , where LDA ; and accordingly, by axiom 4, S is collinear with the point L and a point A' of the segment $L'D$ of the space $a = dl'$; this relation $LA'S$ indicates that the point S is a point S' of $\Sigma' = a'l$.—This completes the proof of statements 1 to 6 for the general k .

It remains to prove the validity of statement 7. We suppose that in the k -space Σ a p -space α and a q -space β ($p + q \geq k$) have a common point C_0 . Then they have in common precisely an r -space γ , where $0 \leq r \leq p$, $0 \leq r \leq q$. We are to show that $r \geq p + q - k$.

We take γ as determined by the $r + 1$ linearly independent points $C_0, \dots, C_r$; and α as determined by these points with $p - r$ points $A_1, \dots, A_{p-r}$; and β as determined by these points with $q - r$ points $B_1, \dots, B_{q-r}$. If β lies in α , then $r = q$; hence, since $k \geq p$, we have indeed $r \geq p + q - k$.

We suppose then that β does not lie in α ; thus there is a point B_1 . Then the $(p + 1)$ -space αB_1 has in common with the q -space β precisely the $(r + 1)$ -space γB_1 . For let B' be any point common to αB_1 and β . If B' is of α , then it is of γ and thus of γB_1 . If it is not of α , then in the $(p + 1)$ -space αB_1 it lies either on the opposite side of α from B_1 , in which case the segment $B_1 B'$ contains a point A of α which is, however, likewise a point B , and thus is a point C of γ , and thus B' is of the line CB_1 and hence of the $(r + 1)$ -space γB_1 ; or it is on the same side of α as B_1 , in which case a similar result will follow by the mediation of a point B'_1 lying on the line through B_1 and the point C_0 and on the opposite side of α from the point B_1 . So we see that if the spaces α and β have γ in common, then the spaces αB_1 and β have precisely γB_1 in common. Hence αB_1 does not contain B_2 . Similarly, $\alpha B_1 B_2$ and β have precisely $\gamma B_1 B_2$ in common, and $\alpha B_1 B_2$ does not contain B_3 . Thus it appears finally that the points $C_0, \dots, C_r, A_1, \dots, A_{p-r}, B_1, \dots, B_{q-r}$ are $1 + p + q - r$ independent points of the k -space Σ , and accordingly we have $p + q - r \leq k$, that is, $r \geq p + q - k$, which was to be proved.

§ 4. *The figure in a k -space determined by $k + 1$ independent points.*

In a k -space a $(k + 1)$ -gon $A_1 \cdots A_{k+1}$ whose $k + 1$ vertices A_g are independent points determines by its $k + 1$ $(k - 1)$ -space faces $a_1 \cdots a_{k+1}$ a certain partition of the k -space.

To every point P of the k -space there is assigned a certain $(k + 1)$ -partite character; every partial character of P is $+$, 0 , or $-$; the g th partial character is 0 if P lies in a_g , $\pm$ if P and A_g lie on ^{the same side}_{or opposite sides} of a_g . Thus, for instance, A_1 has the character $(+, 0, \cdots, 0)_{k \ 0's}$. From the g th partial characters of two points P_1, P_2 one infers at once whether the segment $P_1 P_2$ has lying in a_g no point, one point, or all its points.

Points having the same character are classed together and form one of the parts in the partition in question. There are 3^{k+1} characters; of these, however, only those containing at least one partial character $+$ are characters of points. This is provable by induction. One notes, for instance, that the $(k + 1)$ -partite characters of points of the k -space $a_{k+1} = A_1 \cdots A_k$ are the k -partite characters of these points as points of a_{k+1} with the adjunction of 0 as $(k + 1)$ th character.

The points with character $(+, \cdots, +)_{k+1 \ +}$ constitute the interior* of the $(k + 1)$ -gon, while similarly the points with no partial character $-$ constitute, as variously grouped, the interiors of the various faces of various dimensions of the $(k + 1)$ -gon. If P is a point of the interior of the $(k + 1)$ -gon, the line $A_g P$ cuts the interior of the face a_g in a point A'_g such that P lies between A_g and A'_g .

We speak of complementary or opposite faces of the $(k + 1)$ -gon.† If a point P lies on no face and also not in the interior of the $(k + 1)$ -gon, then (in only one way) there are two points Q, R lying in the interior of opposite faces of the $(k + 1)$ -gon and such that Q lies on the segment PR . The character of P is made up of say q $+$'s and r $-$'s ($q + r = k + 1$; $q > 0, r > 0$), and, by convenient determination of the notation, it is $(+, \cdots, +, -, \cdots, -)_{q \ +'; r \ -'}$; then the points Q, R have respectively the characters

$$(+, \cdots, +, 0, \cdots, 0)_{q \ +'; r \ 0's}, \quad (0, \cdots, 0, +, \cdots, +)_{q \ 0's; r \ +'}.$$

The definition of the k -space analogous to the Peano-Ingrami definition‡ of the plane and of the 3-space is the following:

A $(k + 1)$ -gon $A_1 \cdots A_{k+1}$ lying in no $(k - 1)$ -space determines as its interior the set of all points lying on the segments connecting its vertices to points lying in the interior of the respectively opposite k -gon faces; and it determines as the k -space $A_1 \cdots A_{k+1}$ the set of all points lying either on one of the $(k - 1)$ -space

* For $k = 0$: a point constitutes simultaneously a 0-space, a 1-gon, the interior of a 1-gon.

† For example: ($k = 2$) in a triangle a vertex and opposite side are complementary faces; ($k = 3$) in a tetrahedron a vertex and opposite face are complementary faces, and two opposite edges are complementary faces.

‡ Cf. SCHUB, loc. cit., p. 268.

faces of the $(k + 1)$ -gon or on a line intersecting the interiors of two opposite faces of the $(k + 1)$ -gon.

This definition and that of § 1 are equivalent. Although in this definition the $k + 1$ determining points enter symmetrically, the definition of § 1 is apparently much simpler.

§ 5. *The Hilbert plane.*

We consider a Hilbert geometry in which the basal elements are points, lines, segments of lines and Hilbert planes subject to the Hilbert axioms I 1, 2, 7₁, II 1, 2, 3, which do not involve the Hilbert plane, and furthermore subject to the following axioms I 7'₃, 7'₂, 3', 5', II 5', which do involve and furnish the implicit definition of the Hilbert plane. These axioms I 7'₃, 7'₂, 3', 5', II 5' are (by use of I 1, 2, 7₁, II 1, 2, 3) deducible from the Hilbert axioms I 7₃, 7₂, 3, 5, II 5.

I 7'₃, 7'₂. *There exists a Hilbert plane. Every Hilbert plane π has a notation ABC with respect to certain* three non-collinear points A, B, C . In the notation ABC there is no question of order of the three points A, B, C .*

I 3'. *Three non-collinear points A, B, C determine* a set of points: a Hilbert plane ABC , to which they belong.*

I 5'. *With respect to a Hilbert plane ABC every line joining any point A' of the segment BC to a point D distinct from A' and lying in the plane lies wholly in the plane.*

II 5'. (*The plane order axiom*). *With respect to a Hilbert plane ABC every line which lies in the plane and which contains a point of the segment BC contains also A or B or C or a point of the segment CA or a point of the segment AB .*

In this geometry we prove first the validity, as theorems, of the axioms 1-5 and 6 ($n = 2$) of § 1; from them follow by § 2 the validity as a theorem of the Hilbert axiom II 4, by § 3, 4 the theory of the plane as defined in § 1, and the identity of this plane with that defined by PEANO.

Now we consider any Hilbert plane ABC with notation connected with three non-collinear points A, B, C . It contains the points A, B and C , the line BC (in view of the existence of a point A' of the segment BC), the lines CA and AB , and all lines of the types AA', BB', CC' where A', B', C' are points of the respective segments BC, CA, AB . Hence the Hilbert plane ABC contains the Peano plane ABC . Further, if D is a point of the Hilbert plane ABC not on one of the lines BC, CA, AB , then the line $A'D$ (where A' is a point of the segment BC) contains the point A or a point B' of the segment CA or a point C' of the segment AB . The line $A'D$ quâ $A'A$ or $A'B'$ or $A'C'$

* It is not implied that a plane π has such notation ABC in a definite way, nor that a notation ABC denotes a definite plane.

lies entirely in the plane ABC (of § 1). Hence this plane ABC contains the Hilbert plane ABC .

Accordingly the plane ABC (of § 1), the Peano plane ABC and the Hilbert plane ABC are identical. Then by § 3 follows the validity as theorems for the Hilbert plane of the Hilbert axioms I 3, 4, 5, 7₂, II 5 in their fulness.

Hence *the Hilbert axioms I 4, II 4 are logically deducible from the Hilbert axioms I 1-3, 5, 7, II 1-3, 5.*—If we had wished to exhibit merely this dependence it would have been simpler not to introduce the axioms I 7₂, 7₃, 3', 5', II 5'

THE UNIVERSITY OF CHICAGO,

December 31, 1901.