

# ON THE INTEGRATION OF THE HOMOGENEOUS LINEAR DIFFERENCE EQUATION OF SECOND ORDER\*

BY

WALTER B. FORD.

## I. *Introduction.*

1. It is proposed in the present paper to study the behavior for large positive integral values of  $x$  of the general solution of the equation

$$(1) \quad a_0(x)u(x+2) + a_1(x)u(x+1) + a_2(x)u(x) = 0,$$

the coefficients  $a_0(x)$ ,  $a_1(x)$ ,  $a_2(x)$  being given functions (real or complex) of  $x$  and subject only to conditions relative to the form taken by the expression

$$4 \frac{a_0(x)a_2(x+1)}{a_1(x)a_1(x+1)} \rightarrow 1$$

when  $x$  is very large.† The general results are summarized in Theorem I, after which application is made to the following special equation considered by HORN:‡

$$P_0(x)u(x+2) + x^k P_1(x)u(x+1) + x^{2k} P_2(x)u(x) = 0,$$

where  $k$  is any integer, positive, negative or zero and  $P_0(x)$ ,  $P_1(x)$ ,  $P_2(x)$  are developable asymptotically (or as convergent series) in the form

$$P_\lambda(x) = a_\lambda + \frac{b_\lambda}{x} + \frac{c_\lambda}{x^2} + \dots + \frac{p_\lambda + \omega_\lambda(x)}{x^n} \quad (\lim_{x \rightarrow \infty} \omega_\lambda(x) = 0).$$

The results obtained for this type of equation are stated in Theorem II and are shown to be in accord with those obtained by HORN. Also certain results addi-

---

\* Presented to the Society (Chicago) April 17, 1908.

† Papers relating to this subject but placing more restrictive conditions upon the coefficients have been published by HORN, *Mathematische Annalen*, vol. 53 (1900), pp. 117-192, and by the present author, *Annali di Matematica*, series 3, vol. 13 (1907), pp. 313-328. For a summary of literature concerning equations of form (1) see BARNES, *Messenger of Mathematics*, vol. 34 (1904), p. 53.

‡ *Loc. cit.*, p. 190. For simplicity we shall here confine ourselves to equations of order 2, whereas HORN considers similar equations of any order. A generalization of our results to any order is immediate, in view of the general character of the investigations contained in the above mentioned memoir in the *Annali* upon which the present paper depends.

tional to his are to be here found, and application of these is made in § 11 to the study of the behavior of Legendre's function of the first kind  $P_n$  for large values of  $n$ .

## II. Reduction of the Equation.

2. We shall first show that in case  $a_1(x) \neq 0$ ,  $a_2(x) \neq 0$  for all  $x \geq \alpha = \text{constant}$ ,\* the study of equation (1) may be made to depend upon that of a certain linear difference equation of the type

$$(2) \quad \Delta^2 y(x) + a(x)y(x+2) = 0,$$

in which  $a(x)$  depends only upon  $a_0$ ,  $a_1$  and  $a_2$  and is defined for all  $x \geq \alpha$ .

In fact, if one places  $u(x) = t(x)y(x)$ , equation (1) becomes

$$a_0(x)t(x+2)y(x+2) + a_1(x)t(x+1)y(x+1) + a_2(x)t(x)y(x) = 0$$

or, since

$$y(x+1) = y(x+2) - \Delta y(x+1),$$

$$y(x) = y(x+2) - 2\Delta y(x+1) + \Delta^2 y(x),$$

the same equation becomes

$$(3) \quad a_2(x)t(x)\Delta^2 y(x) - [a_1(x)t(x+1) + 2a_2(x)t(x)]\Delta y(x+1) \\ + [a_0(x)t(x+2) + a_1(x)t(x+1) + a_2(x)t(x)]y(x+2) = 0.$$

Let us now choose the heretofore undetermined function  $t(x)$  in such a manner that the second coefficient of (3) vanishes:

$$(4) \quad a_1(x)t(x+1) + 2a_2(x)t(x) = 0.$$

The function  $t(x)$  thus becomes determined,† except for a constant factor, by the equation

$$(5) \quad t(x) = \prod_{x_1=\alpha}^{x_1=x-1} \left( -\frac{2a_2}{a_1} \right)_{x_1},$$

since  $a_1(x) \neq 0$  for  $x \geq \alpha$ . Moreover, from the hypothesis  $a_2(x) \neq 0$  it follows that  $t(x) \neq 0$  and hence also  $a_2(x)t(x) \neq 0$ .

Consequently, equation (3) may be reduced to the form

$$(6) \quad \Delta^2 y(x) + [4(a_0/a_1)_x(a_2/a_1)_{x+1} - 1]y(x+2) = 0 \quad (x \geq \alpha),$$

i. e., to the desired form (2).

3. We shall now make the assumption that  $4(a_0/a_1)_x(a_2/a_1)_{x+1} - 1$  for large values of  $x$  has the form  $-\nu^2 - \phi(x) - \theta(x)$ , where  $\nu$  is a constant (real or

\* Throughout the paper it will be understood, unless otherwise stated, that  $x \geq \alpha =$  a sufficiently large constant.

† See BOOLE's *Finite Differences*, Chap. IX, § 6.

complex) and  $\nu(\nu^2 - 1) \neq 0$ , where also  $\phi(x)$  is a function of  $x$  of the form  $\mu/x^{1+\delta}$ , in which  $\mu$  and  $\delta$  are constants with  $0 < \delta \leq \frac{1}{2}$ , and where  $\theta(x)$  is any function of  $x$  such that the series

$$\sum_{x_1=x}^{x_1=\infty} |\theta(x_1)|$$

converges. For example, the function  $\theta(x)$  may be taken as any function of the form  $g(x) \cdot \tau(x)/x$ , where  $|g(x)| < c = \text{const.}$  and  $\tau(x)$  is one of the functions

$$\frac{1}{x^p}, \frac{1}{(\log x)^{1+p}}, \frac{1}{\log x (\log_2 x)^{1+p}}, \dots \quad (p > 0).$$

In particular, we observe that the above hypotheses are realized whenever the original equation (1) has coefficients  $a_0(x)$ ,  $a_1(x)$ ,  $a_2(x)$  which are developable either in the form of convergent power series in  $1/x$  ( $x \geq \alpha$ ) or, more generally, as asymptotic series in  $1/x$ .

Equation (6) thus becomes

$$(7) \quad \Delta^2 y(x) - [\nu^2 + \phi(x) + \theta(x)]y(x+2) = 0.$$

In the study of this equation we shall now make use of Theorem II of the paper entitled *Sur les équations linéaires aux différences finies*,\* using without further remark the results and notation there found.

In the present instance let us choose the auxiliary functions  $z_1(x)$ ,  $z_2(x)$  as follows, for reasons which will appear presently:

$$(8) \quad z_1(x) = \prod_{x_1=\alpha}^{x_1=x-1} (1 + \nu + \phi(x_1)/2\nu), \quad z_2(x) = \prod_{x_1=\alpha}^{x_1=x-1} (1 - \nu - \phi(x_1)/2\nu).$$

Then

$$(9) \quad \Delta z_1(x) = (\nu + \phi(x)/2\nu)z_1(x), \quad \Delta^2 z_1(x) = (\nu^2 + \phi(x) + \xi_1(x))z_1(x),$$

where

$$\xi_1(x) = \frac{1+\nu}{2\nu} \Delta \phi(x) + \frac{\phi(x+1)\phi(x)}{4\nu^2}.$$

From our hypothesis respecting  $\phi(x)$  it appears directly that  $\xi_1(x)$  has the character of one of the functions  $\theta(x)$  mentioned above.

Likewise we have

$$\Delta z_2(x) = -(\nu + \phi(x)/2\nu)z_2(x), \quad \Delta^2 z_2(x) = (\nu^2 + \phi(x) + \xi_2(x))z_2(x),$$

where  $\xi_2(x)$  has the properties of  $\xi_1(x)$  just mentioned and is obtained from it by changing  $\nu$  into  $-\nu$ .

\* *Annali di Matematica*, loc. cit., p. 301. The expression  $\sum_{x_1=x+1}^{x_1=\infty} |u_m(x_1)|$  there occurring should be replaced by  $\sum_{m=1}^{m=\infty} |u_m(x_1)|$ .

Equations (8) and (9) together with those just noted give the following values for the expressions  $A(x)$ ,  $Q(x)$ ,  $f_1(x)$ ,  $f_2(x)$ ,  $\bar{q}(x, x_1)$ ,  $\Phi(x, x_1)$ ,  $\Psi(x, x_1)$ , and  $F(x)$  which occur in the statement of the above mentioned Theorem II:

$$(10) \quad Q(x) = -2(\nu + \phi(x)/2\nu)z_1(x)z_2(x),$$

$$(11) \quad A(x) = c_2z_1(x) - c_1z_2(x),$$

$$(12) \quad \begin{aligned} f_1(x) &= [-\nu^2 - \phi(x) - \theta(x)]z_1(x) + \Delta^2z_1(x) = [\xi_1(x) - \theta(x)]z_1(x), \\ f_2(x) &= [\xi_2(x) - \theta(x)]z_2(x), \end{aligned}$$

$$(13) \quad \bar{q}(x, x_1) = \begin{vmatrix} z_1(x) & [\xi_1(x_1) - \theta(x_1)]z_1(x_1) \\ z_2(x) & [\xi_2(x_1) - \theta(x_1)]z_2(x_1) \end{vmatrix},$$

$$(14) \quad \begin{aligned} \Phi(x, x_1) &= \frac{1}{2(\nu + \phi(x_1 + 1)/2\nu)} \\ &\times \left[ \frac{[\xi_2(x_1) - \theta(x_1)]z_1(x_1 + 1)}{[1 - \nu - \phi(x_1)/2\nu]z_1(x_1 + 1)} - \frac{[\xi_1(x_1) - \theta(x_1)]z_2(x_1 + 1)}{[1 + \nu + \phi(x_1)/2\nu]z_2(x_1 + 1)} \right]. \end{aligned}$$

Moreover, since  $\lim_{x_1 \rightarrow x} \phi(x_1) = 0$ ,  $\nu(1 - \nu^2) \neq 0$ , while  $\theta(x_1)$ ,  $\xi_1(x_1)$ ,  $\xi_2(x_1)$  have the properties above mentioned, we now see that

$$(15) \quad \Phi(x, x_1) = \frac{z_1(x + 1)}{z_1(x_1 + 1)} \left[ s_1(x_1) + s_2(x_1) \frac{z_1(x_1 + 1)z_2(x + 1)}{z_1(x + 1)z_2(x_1 + 1)} \right] |\theta(x_1)|,$$

where  $|s_1(x_1)|$  and  $|s_2(x_1)|$  are less than some assignable constant, and  $\theta(x)$ , though not identical with the  $\theta(x)$  of (7), has the properties before described of that function.

Finally, we have

$$(16) \quad \Psi(x, x_1) = A(x_1 + 1)\Phi(x, x_1)$$

and

$$(17) \quad F(x) = \frac{-1}{Q(x + 1)} [A(x + 1) + u_1(x) + u_2(x) + \cdots + u_n(x) + \cdots].$$

### III. The special case $|1 + \nu| = |1 - \nu|$ .

4. Let us suppose in the first place that  $|1 + \nu| = |1 - \nu|$  and that  $|1 + \nu + \phi(x)/2\nu| = |1 - \nu - \phi(x)/2\nu| = \rho_x$ , ( $x \geq \alpha$ ). For this case  $|z_1(x)| = |z_2(x)|$ . Whence if we place

$$(18) \quad G(x) = \prod_{n=\alpha}^{n=x} \rho_n$$

we may write

$$(19) \quad \Phi(x, x_1) = \omega_1(x, x_1) |\theta(x_1)| \frac{G(x)}{G(x_1)} \quad (x \geq \alpha, x_1 > x),$$

where  $|\omega_1(x, x_1)| < \Omega_1 = \text{const.}$

Similarly we have

$$(20) \quad A(x+1) = G(x) [c_1 r_1(x) + c_2 r_2(x)] \quad (x \geq \alpha),$$

where  $|r_1(x)| = |r_2(x)| = 1$ . Whence, by (16), it follows that

$$(21) \quad \Psi(x, x_1) = \omega_2(x, x_1) |\theta(x_1)| G(x) \quad (x \geq \alpha, x_1 > \alpha),$$

where  $|\omega_2(x, x_1)| < \Omega_2 = \text{const.}$

Equations (19) and (21) having been obtained, we turn to consider the series

$$(22) \quad |u_1(x)| + |u_2(x)| + \cdots + |u_m(x)| + \cdots$$

Referring to the definition of the term  $u_m(x)$  given in the before mentioned Theorem II, we have in the present instance

$$(23) \quad \Phi(x, x_1) \Phi(x_1, x_2) \cdots \Phi(x_{m-2}, x_{m-1}) \Psi(x_{m-1}, x_m) = G(x) P(x, x_1, x_2, \cdots, x_m),$$

where for  $x_m \geq x_{m-1} \geq x_{m-2} \cdots \geq x_1 \geq x \geq \alpha$  we may write

$$(24) \quad |P(x, x_1, x_2, \cdots, x_m)| < \Omega_1^{m-1} \Omega_2 |\theta(x_1) \theta(x_2) \cdots \theta(x_m)|.$$

Relations (23) and (24) now enable us to show that the three conditions (a), (b), (c) of the theorem are here fulfilled. Condition (a) is fulfilled inasmuch as we have

$$\sum_{x_1=x+1}^{x_1=\infty} \sum_{x_2=x_1+1}^{x_2=\infty} \cdots \sum_{x_m=x_{m-1}+1}^{x_m=\infty} |\Phi(x, x_1) \Phi(x_1, x_2) \cdots \Phi(x_{m-2}, x_{m-1}) \Psi(x_{m-1}, x_m)| < G(x) \Omega_1^{m-1} \Omega_2 W(x),$$

where

$$W(x) = \sum_{x_1=x+1}^{x_1=\infty} |\theta(x_1)| \sum_{x_2=x_1+1}^{x_2=\infty} |\theta(x_2)| \cdots \sum_{x_m=x_{m-1}+1}^{x_m=\infty} |\theta(x_m)|,$$

and this expression has a meaning by virtue of our hypotheses concerning the function  $\theta(x)$ .

As to condition (b), let us put

$$(25) \quad \theta_1(x) = \sum_{x_1=x+1}^{x_1=\infty} |\theta(x_1)|.$$

Since  $\theta_1(x)$  becomes arbitrarily small for all values of  $x$  sufficiently large, the term  $u_m(x)$  takes the form

$$(26) \quad u_m(x) = G(x) V_m(x),$$

where

$$(27) \quad |V_m(x)| < \Omega_1^{m-1} \Omega_2 |\theta_1(x)|^m.$$

Whence, if  $\alpha$  be sufficiently large, we shall have

$$(28) \quad |u_m(x)| < G(x) k^m, \quad k = \text{const.} < 1 \quad (x \geq \alpha).$$

Thus the series (22) converges for  $x \geq \alpha$  to a value  $U(x)$  such that

$$(29) \quad U(x) = \gamma(x) G(x), \quad \gamma(x) < \frac{k}{1-k}.$$

By virtue of (19) and (20) we may now write

$$|\Phi(x, x_1)| U(x_1) = \beta_1(x, x_1) G(x),$$

where

$$\beta_1(x, x_1) < \Omega_3 |\theta(x_1)| \quad (\Omega_3 = \text{const.}, x \geq \alpha, x_1 \geq \alpha).$$

Thus the series

$$\sum_{x_1=x+1}^{x_1=\infty} |\Phi(x, x_1)| U(x_1) \quad (x \geq \alpha)$$

converges.

Finally, the expressions  $F(x_1) f_r'(x_1)$ , ( $r = 1, 2$ ), of condition (c) are to be considered. We obtain in the first place from (17) and (29)

$$(30) \quad F(x) = - \left( \frac{A(x+1)}{Q(x+1)} + \frac{\beta_2(x) G(x)}{Q(x+1)} \right) \quad (|\beta_2(x)| < \gamma(x) < \frac{k}{1-k}).$$

But from (10) and (11) we have

$$(31) \quad \frac{A(x+1)}{Q(x+1)} = - \frac{\nu}{2\nu^2 + \phi(x+1)} \left( \frac{c_2}{z_2(x+1)} - \frac{c_1}{z_1(x+1)} \right) = \frac{\beta_3(x)}{G(x)} \quad (x \geq \alpha),$$

where  $|\beta_3(x)| < \beta_3 = \text{const.}$

Similarly we obtain

$$(32) \quad \frac{G(x)}{Q(x+1)} = \frac{\beta_4(x)}{G(x)} \quad (|\beta_4(x)| < \beta_4 = \text{const.}).$$

Whence follows

$$F(x) = \frac{\beta_5(x)}{G(x)} \quad (|\beta_5(x)| \leq \beta_5 = \text{const.}).$$

Moreover, we have from (12),

$$f_r'(x) = \beta_6(x, r) G(x) |\theta(x)| \quad (r = 1, 2),$$

where  $|\beta_6(x, r)| < \beta_6 = \text{const.}$

Thence, noting that  $\rho_a \neq 0$ , we have

$$F(x) f_r'(x) = \beta_7(x, r) |\theta(x)| \quad (|\beta_7(x, r)| < \beta_7 = \text{const.}),$$

and, therefore, the expressions

$$\sum_{x_1=x+1}^{x_1=\infty} F(x_1) f_r'(x_1) \quad (r = 1, 2; x \geq \alpha),$$

have a meaning. Thus all the conditions of Theorem II become fulfilled.

Moreover, the function  $\gamma(x)$  of (29) has the properties of the function  $\theta_1(x)$

defined in (25); i. e.,  $\lim_{x=\infty} \gamma(x) = 0$ . Consequently from (30), (31) and (32) we have

$$(33) \quad Y(x) = \frac{\nu}{2\nu^2 + \phi(x+1)} \left( \frac{c_2}{z_2(x+1)} - \frac{c_1}{z_1(x+1)} + \frac{\delta(x)}{G(x)} \right)$$

where  $x \geq \alpha$ ,  $\lim_{x=\infty} \delta(x) = 0$ .

Upon applying now the result of Theorem II we find that the general solution of (7), under the present hypotheses respecting  $\nu$ ,  $\phi(x)$ , and  $\theta(x)$ , will have the following form when  $x \geq \alpha$  for sufficiently large  $\alpha$ :

$$y(x) = \frac{k_1(1 + \epsilon_1(x))}{z_1(x-1)} + \frac{k_2(1 + \epsilon_2(x))}{z_2(x-1)} \quad (\lim_{x=\infty} \epsilon_1(x) = \lim_{x=\infty} \epsilon_2(x) = 0),$$

$k_1, k_2$  being arbitrary constants. Since also

$$\frac{z_1(x)}{z_1(x-1)} = 1 + \eta_1(x), \quad \frac{z_2(x)}{z_2(x-1)} = 1 + \eta_2(x) \quad (\lim_{x=\infty} \eta_1(x) = \lim_{x=\infty} \eta_2(x) = 0),$$

this may be thrown into the form

$$(34) \quad y(x) = \frac{k_1(1 + \epsilon_1(x))}{z_1(x)} + \frac{k_2(1 + \epsilon_2(x))}{z_2(x)} \quad (\lim_{x=\infty} \epsilon_1(x) = \lim_{x=\infty} \epsilon_2(x) = 0).$$

#### IV. Derivation of a first integral for the general case.

5. We turn now to consider the cases in which  $|1 + \nu| \neq |1 - \nu|$ . Placing  $\rho_x = |1 + \nu + \phi(x)/2\nu|$  as before and also placing  $\sigma_x = |1 - \nu - \phi(x)/2\nu|$  and

$$(35) \quad H(x) = \prod_{n=\alpha}^{n=x} \sigma_n,$$

we shall have

$$\frac{z_2(x+1)}{z_1(x+1)} = \lambda_3(x, x_1) \frac{H(x)}{H(x_1)} \quad (|\lambda_3(x, x_1)| = 1).$$

Thus, from (15) we obtain

$$\Phi(x, x_1) = \frac{H(x)}{H(x_1)} \left( s_1(x_1) \frac{G(x)H(x_1)}{G(x_1)H(x)} + s_2(x_1) \right) |\theta(x_1)|.$$

Then

$$\frac{G(x)H(x_1)}{G(x_1)H(x)} = \left( \frac{\sigma_{x_1}}{\rho_{x_1}} \right) \left( \frac{\sigma_{x_1-1}}{\rho_{x_1-1}} \right) \cdots \left( \frac{\sigma_{x+1}}{\rho_{x+1}} \right).$$

Let us now suppose  $x_1 > x \geq \alpha$  and let us consider  $\nu$  to be that square root of  $\nu^2$  in (7) for which  $|1 + \nu| > |1 - \nu|$ . Then the factors in the right hand member of our last equation will each be less than 1 and we shall be able to write

$$(36) \quad \Phi(x, x_1) = \omega_3(x, x_1) |\theta(x_1)| \frac{H(x)}{H(x_1)} \quad (|\omega_3(x, x_1)| < \Omega_3 = \text{const.}).$$

As regards the function

$$\Psi(x, x_1) = A(x_1 + 1) \Phi(x, x_1) = [c_2 z_1(x_1 + 1) - c_1 z_2(x_1 + 1)] \Phi(x, x_1),$$

let us take in the present instance  $c_2 = 0$ . Then

$$A(x + 1) = c_1 r_1(x) H(x) \quad (|r(x)| = 1),$$

so that

$$(37) \quad \Psi(x, x_1) = \omega_4(x, x_1) H(x) |\theta(x_1)| \quad (|\omega_4(x, x_1)| < \Omega_4 = \text{const.}).$$

We may now proceed as in the former case. Thus we have in the first place

$$(38) \quad u_m(x) = H(x) V_m(x) \quad (x \geq a),$$

where  $V_m(x)$  has the properties indicated in (27), whence also

$$(39) \quad |u_m(x)| < H(x) k^m \quad (k = \text{const.} < 1; x \geq a),$$

and

$$Y(x) = \frac{\nu}{2\nu^2 + \phi(x+1)} \left( \frac{-c_1}{z_1(x+1)} + \frac{\delta_1(x)}{G(x)} \right) \quad \left( \lim_{x \rightarrow \infty} \delta_1(x) = 0 \right).$$

Thus in place of (34) we now have the solution

$$(40) \quad y_1(x) = \frac{k_1 (1 + \epsilon_1(x))}{z_1(x)} \quad (x \geq a, \lim_{x \rightarrow \infty} \epsilon_1(x) = 0).$$

### V. Derivation of the second integral.

6. Having obtained but a particular solution  $y_1(x)$  of (7) when  $|1 + \nu| \neq |1 - \nu|$ , we proceed in the present section to obtain a second solution and therewith the general solution.

For this purpose equation (7) may be written in the form

$$(41) \quad (1 - \nu^2 - \phi(x) - \theta(x)) y(x+2) - 2y(x+1) + y(x) = 0.$$

If now  $u(x)$  and  $v(x)$  be any two linearly independent solutions of the equation

$$(42) \quad a_0(x) u(x+2) + a_1(x) u(x+1) + a_2(x) u(x) = 0,$$

we have, after placing for brevity  $a_0(x) = a_0$ , etc., and  $u(x+1) = u_1$ ,  $u(x+2) = u_2$ , etc.:

$$a_0(u_2 v_1 - u_1 v_2) + a_2(u_0 v_1 - u_1 v_0) = 0$$

and hence

$$\frac{u_2 v_1 - u_1 v_2}{u_1 v_0 - u_0 v_1} = \frac{a_2}{a_0}.$$

Therefore, for any fixed integer  $\alpha$  we obtain

$$u(\alpha + n)v(\alpha + n - 1) - u(\alpha + n - 1)v(\alpha + n) = c \prod_{p=\alpha}^{p=\alpha+n-2} \frac{a_2(p)}{a_0(p)} \quad (n=2, 3, \dots),$$

where  $c$  is a constant as regards  $n$ . Whence

$$\frac{u(\alpha + n)}{v(\alpha + n)} - \frac{u(\alpha + 1)}{v(\alpha + 1)} = c \sum_{m=2}^{m=n} \frac{1}{v(\alpha + m)v(\alpha + m - 1)} \prod_{p=\alpha}^{p=\alpha+m-2} \frac{a_2(p)}{a_0(p)}.$$

Thus, if  $v(x)$  be looked upon as a *known* solution of (42), the solution  $u(x)$  may be expressed in the following form for all values of  $x = \alpha + n$ ,  $\alpha$  being a constant:

$$(43) \quad u(x) = c_1 v(x) + c_2 v(x) S_n$$

where

$$S_n = \sum_{m=1}^{m=n} \frac{1}{v(\alpha + m)v(\alpha + m - 1)} \prod_{p=\alpha}^{p=\alpha+m-2} \frac{a_2(p)}{a_0(p)}.$$

We proceed to study the properties of  $S_n$  for large values of  $n$  in the case of the special equation (41), for which one solution  $v(x) \equiv y_1(x)$ , given by (40), is known.

Thus we have in the present instance

$$(44) \quad \frac{1}{v(\alpha + m)} = k_1(1 + \epsilon_m)z_1(\alpha + m) = k_1(1 + \epsilon_m) \prod_{p=\alpha}^{p=\alpha+m-1} \left(1 + \nu + \frac{\phi(p)}{2\nu}\right) \quad (\lim_{m \rightarrow \infty} \epsilon_m = 0),$$

and from (41)

$$\frac{a_2(p)}{a_0(p)} = \frac{1 + \eta_p}{\left(1 + \nu + \frac{\phi(p)}{2\nu}\right) \left(1 - \nu - \frac{\phi(p)}{2\nu}\right)},$$

where

$$\eta_p = \frac{\phi(p) - \left[\frac{\phi(p)}{2\nu}\right]^2}{1 - \nu^2 - \phi(p) - \theta(p)}.$$

Thence,

$$(45) \quad \prod_{p=\alpha}^{p=\alpha+m-2} \frac{a_2(p)}{a_0(p)} = \frac{\prod_{p=\alpha}^{p=\alpha+m-2} (1 + \eta_p)}{\prod_{p=\alpha}^{p=\alpha+m-2} \left(1 + \nu + \frac{\phi(p)}{2\nu}\right) \left(1 - \nu - \frac{\phi(p)}{2\nu}\right)}.$$

Moreover, by virtue of our hypotheses upon  $\phi(x)$  and  $\theta(x)$ , the series

$$\sum_{p=\alpha}^{p=\infty} |\eta_p|$$

is convergent, and hence it follows that

$$(46) \quad \prod_{p=\alpha}^{p=\alpha+m-2} (1 + \eta_p) = k_2(1 + \epsilon_m) \quad (k_2 = \text{const.}, \lim_{m \rightarrow \infty} \epsilon_m = 0).$$

Upon making use of (44), (45) and (46) we find that the  $m$ th term of  $S_n$  is of the form

$$k_2(1 + \epsilon_m) \frac{\prod_{p=a}^{p=a+m-1} \left(1 + \nu + \frac{\phi(p)}{2\nu}\right)}{\prod_{p=a}^{p=a+m-2} \left(1 - \nu - \frac{\phi(p)}{2\nu}\right)} = \frac{k'_2(1 + \epsilon'_m)z_1(\alpha + m)}{z_2(\alpha + m)} \quad (\lim_{m \rightarrow \infty} \epsilon'_m = 0).$$

The expression  $S_n$  may therefore be put into the form

$$(47) \quad S_n = k'_2 \frac{z_1(\alpha + n)}{z_2(\alpha + n)} S'_n,$$

where

$$S'_n = \sum_{m=1}^{m=n} \frac{z_1(\alpha + m)z_2(\alpha + n)}{z_1(\alpha + n)z_2(\alpha + m)} (1 + \epsilon'_m),$$

which we may now show directly to be of the form  $k_3 + \epsilon_n$  where  $k_3 = \text{const.}$  and  $\lim_{n \rightarrow \infty} \epsilon_n = 0$ . In fact, we may write

$$S'_n = \left( \sum_{m=1}^{m=n-p-1} + \sum_{m=n-p}^{m=n} \right) \frac{z_1(\alpha + m)z_2(\alpha + n)}{z_1(\alpha + n)z_2(\alpha + m)} (1 + \epsilon_m),$$

where the first sum appearing on the right evidently approaches a limit by virtue of (8) when  $n = \infty$ , while by taking  $p$  sufficiently large the second sum may be made less in absolute value than any pre-assigned positive quantity  $\epsilon$ , whatever  $n$  may be ( $n > p + 1$ ), since

$$\left| \frac{z_1(\alpha + m)z_2(\alpha + n)}{z_1(\alpha + n)z_2(\alpha + m)} \right| = \frac{\sigma_{a+n}\sigma_{a+n-1} \cdots \sigma_{a+m-1}}{\rho_{a+n}\rho_{a+n-1} \cdots \rho_{a+m-1}} < q^{n-m+2} \quad (q = \text{const.} < 1).$$

Availing ourselves of the form just established for  $S'_n$  and recalling the definition of  $v(\alpha + n)$ , we see from (47) that

$$S_n = \frac{k(1 + \epsilon_n)}{v(\alpha + n)z_2(\alpha + n)} = \frac{k(1 + \epsilon(x))}{v(x)z_2(x)} \quad (\lim_{n \rightarrow \infty} \epsilon_n = \lim_{x \rightarrow \infty} \epsilon(x) = 0).$$

Placing this value in (43) and recalling that the expression  $c_1 v(x) = c_1 y_1(x)$  there appearing is itself a solution of (41), we obtain the following second solution linearly independent of  $y_1(x)$ :

$$y_2(x) = \frac{k_2(1 + \epsilon_2(x))}{\tilde{z}_2(x)} \quad (x \geq \alpha, \lim_{x \rightarrow \infty} \epsilon_2(x) = 0).$$

Our results may now be summarized in the following general theorem:

*Theorem I: Given the equation*

$$a_0(x)u(x+2) + a_1(x)u(x+1) + a_2(x)u(x) = 0$$

*whose coefficients (real or complex) are defined for all positive integral values of  $x$  sufficiently large ( $x \geq \alpha$ ), and satisfy for such values the conditions:*

(1)  $a_0(x)$ ,  $a_1(x)$ ,  $a_2(x)$  never vanish.

(2) The expression

$$4 \frac{a_0(x)a_2(x+1)}{a_1(x)a_1(x+1)} - 1$$

has the form  $-\nu^2 - \mu/x^{1+\delta} - \theta(x)$  where  $\mu$ ,  $\nu$ ,  $\delta$  are constants such that  $0 < \delta \leq \frac{1}{2}$  and  $\nu(1 - \nu^2) \neq 0$  and where  $\theta(x)$  is any function of  $x$  such that the series

$$\sum_{x_1=x}^{x_1=\infty} |\theta(x_1)| \quad (x \geq \alpha)$$

converges.

Then if  $|1 + \nu| \neq |1 - \nu|$ , there are two particular solutions of the equation having the forms

$$u_1(x) = \frac{t(x)(1 + \epsilon_1(x))}{z_1(x)}, \quad u_2(x) = \frac{t(x)(1 + \epsilon_2(x))}{z_2(x)} \quad (\lim_{x \rightarrow \infty} \epsilon_1(x) = \lim_{x \rightarrow \infty} \epsilon_2(x) = 0),$$

in which

$$t(x) = \prod_{x_1=\alpha}^{x_1=x-1} \left( \frac{-2a_2}{a_1} \right)_{x_1}, \quad z_1(x) = \prod_{x_1=\alpha}^{x_1=x-1} \left\{ 1 + \nu + \frac{\phi(x_1)}{2\nu} \right\}, \quad z_2(x) = \prod_{x_1=\alpha}^{x_1=x-1} \left\{ 1 - \nu - \frac{\phi(x_1)}{2\nu} \right\}.$$

When  $|1 + \nu| = |1 - \nu|$  the result continues to hold true provided that

$$\left| 1 + \nu + \frac{\phi(x)}{2\nu} \right| = \left| 1 - \nu - \frac{\phi(x)}{2\nu} \right| \text{ for } x \geq \alpha.$$

## VI. Application to the equation considered by Horn.\*

7. We proceed to apply the above theorem to the study of the solutions  $u(x)$  of the equation,

$$(48) \quad P_0(x)u(x+2) + x^k P_1(x)u(x+1) + x^{2k} P_2(x)u(x) = 0,$$

where  $k$  is any integer, positive, negative or zero and  $P_0(x)$ ,  $P_1(x)$  and  $P_2(x)$  are either convergent series for all sufficiently large values of  $x$  or are developable asymptotically in the form

$$P_\lambda(x) = a_\lambda + \frac{b_\lambda}{x} + \frac{c_\lambda}{x^2} + \dots + \frac{p_\lambda + \omega_\lambda(x)}{x^n} \quad (\lim_{x \rightarrow \infty} \omega_\lambda(x) = 0).$$

For this purpose we begin by making the transformation  $u(x) = [\Gamma(x)]^k y(x)$ . Thus equation (48) takes the form

$$(49) \quad \left( a_0 + \frac{b_0}{x} + \frac{c_0}{x^2} + \dots \right) y(x+2) + \left( a'_1 + \frac{b'_1}{x} + \frac{c'_1}{x^2} + \dots \right) y(x+1) + \left( a'_2 + \frac{b'_2}{x} + \dots \right) y(x) = 0,$$

\* Loc. cit., p. 190.

in which the coefficients of  $y(x+1)$  and  $y(x)$  are of the form just indicated for  $P_\lambda(x)$ . In particular,

$$(50) \quad a'_1 = a_1, \quad b'_1 = b_1 - a_1 k, \quad a'_2 = a_2, \quad b'_2 = b_2 - a_2 k.$$

For convenience, let us drop the primes in (49) and take as object of study the equation

$$(51) \quad \left(a_0 + \frac{b_0}{x} + \dots\right)y(x+2) + \left(a_1 + \frac{b_1}{x} + \dots\right)y(x+1) + \left(a_2 + \frac{b_2}{x} + \dots\right)y(x) = 0,$$

in which the coefficients are of the type  $P_\lambda(x)$ .

Equation (4) becomes in the present instance

$$(52) \quad \left(a_1 + \frac{b_1}{x} + \frac{c_1}{x^2} + \dots\right)t(x+1) + 2\left(a_2 + \frac{b_2}{x} + \frac{c_2}{x^2} + \dots\right)t(x) = 0.$$

Place  $t(x) = x^g(1+h/x)v(x)$  where  $g$  and  $h$  are constants yet to be determined. Equation (52) then takes the form

$$\left(1 + \frac{h}{x+1}\right)\left(a_1 + \frac{b_1}{x} + \dots\right)v(x+1) + 2\left(1 + \frac{1}{x}\right)^{-g}\left(1 + \frac{h}{x}\right)\left(a_2 + \frac{b_2}{x} + \dots\right)v(x) = 0,$$

or, upon developing  $(1+1/x)^{-g}$  by the binomial theorem,

$$(53) \quad \left(a_1 + (b_1 + ha_1)\frac{1}{x} + (c_1 + hb_1 - ha_1)\frac{1}{x^2} + \dots\right)v(x+1) + 2\left(a_2 + (b_2 - ga_2 + ha_2)\frac{1}{x} + (c_2 - gb_2 + hb_2 + \frac{1}{2}g(g+1)a_2 - gha_2)\frac{1}{x^2} + \dots\right)v(x) = 0.$$

Let us now choose the undetermined constants  $g, h$  so that the term in  $1/x$  in each coefficient of (53) vanishes. In case  $a_1 \neq 0, a_2 \neq 0$ , these two conditions determine  $g$  and  $h$  as follows:

$$g = \frac{a_1 b_2 - a_2 b_1}{a_1 a_2}, \quad h = -\frac{b_1}{a_1}.$$

The constants  $g, h$  having been thus determined, we may now apply directly to equation (53) the results embodied in Theorem III of my previous paper\* and write for sufficiently large  $\alpha$

$$v(x) = c_1 \left(-\frac{2a_2}{a_1}\right)^x (1 + \epsilon_1(x)) \quad (c_1 = \text{const. } x \geq \alpha, \lim_{x \rightarrow \infty} \epsilon_1(x) = 0),$$

and therefore also

$$(54) \quad t(x) = c_1 x^{\frac{a_1 b_2 - a_2 b_1}{a_1 a_2}} \left(-\frac{2a_2}{a_1}\right)^x (1 + \epsilon_2(x)) \quad (\lim_{x \rightarrow \infty} \epsilon_2(x) = 0).$$

\* Loc. cit., p. 313.

We turn now to consider the forms taken by  $z_1(x)$ ,  $z_2(x)$  for the equation (51). In the first place, let us construct the expression

$$\frac{4a_0(x)a_2(x+1)}{a_1(x)a_1(x+1)} - 1$$

referred to in condition (b) of the preceding theorem. Since

$$\frac{a_0(x)}{a_1(x)} = \frac{a_0}{a_1} + \frac{a_1 b_0 - a_0 b_1}{a_1^2} \left( \frac{1}{x} \right) + \dots,$$

$$\frac{a_2(x+1)}{a_1(x+1)} = \frac{a_2}{a_1} + \frac{a_1 b_2 - a_2 b_1}{a_1^2} \left( \frac{1}{x+1} \right) + \dots,$$

we obtain

$$4 \frac{a_0(x)a_2(x+1)}{a_1(x)a_1(x+1)} - 1 = -\nu^2 - \frac{\mu}{x+1} - \theta(x),$$

in which

$$(55) \quad \nu = \frac{\sqrt{a_1^2 - 4a_0a_2}}{a_1},$$

$$(56) \quad \mu = 4 \frac{2a_0a_2b_1 - a_1a_2b_0 - a_0a_1b_2}{a_1^3},$$

and  $\theta(x)$  vanishes to at least the second order when  $x = \infty$ .

Thus, for the function  $z_1(x)$  we have

$$(57) \quad z_1(x) = \prod_{x_1=\alpha}^{x_1=x} \left( 1 + \nu + \frac{\mu}{2\nu x_1} \right) = g_1(1+\nu)^{x+1} \frac{\Gamma\left(x+1 + \frac{\mu}{2\nu(1+\nu)}\right)}{\Gamma(x+1)}$$

where  $g_1$  is a constant and  $\mu$  and  $\nu$  are given by (56) and (55). Similarly,

$$(58) \quad z_2(x) = g_2(1-\nu)^{x+1} \frac{\Gamma\left(x+1 - \frac{\mu}{2\nu(1-\nu)}\right)}{\Gamma(x+1)}$$

Let us next consider what conditions (a) and (b) of Theorem I become in the present instance. Since

$$\nu(\nu^2 - 1) = \frac{4a_0a_2}{a_1^3} \sqrt{a_1^2 - 4a_0a_2},$$

they will be satisfied if  $a_0a_1a_2(a_1^2 - 4a_0a_2) \neq 0$ . Moreover, the roots of the quadratic equation  $a_0\lambda^2 + a_1\lambda + a_2 = 0$  are

$$\lambda_1 = (1/2a_0)(-a_1 + \sqrt{a_1^2 - 4a_0a_2}), \quad \lambda_2 = (1/2a_0)(-a_1 - \sqrt{a_1^2 - 4a_0a_2}),$$

so that

$$1 + \nu = -\frac{2a_2}{a_1\lambda_1}, \quad 1 - \nu = -\frac{2a_2}{a_1\lambda_2}.$$

Also we have

$$\frac{\mu}{2\nu(1+\nu)} = \frac{\mu'}{a_0 a_1 (a_1 \lambda_2 + 2a_2)}, \quad \frac{-\mu}{2\nu(1-\nu)} = \frac{\mu'}{a_0 a_1 (a_1 \lambda_1 + 2a_2)},$$

where  $\mu' = -\mu a_1^3/4 = a_0 a_1 b_2 + a_1 a_2 b_0 - 2a_0 a_2 b_1$ .

Noting that  $a_1^2 - 4a_0 a_2 \neq 0$  whenever  $\lambda_1 \neq \lambda_2$ , we see that if in equation (51) the roots  $\lambda_1, \lambda_2$  of the quadratic  $a_0 \lambda^2 + a_1 \lambda + a_2 = 0$  are of unequal modulus and  $a_0 a_1 a_2 \neq 0$ , there are two solutions of the same equation which, when considered for all positive integral values of  $x$  sufficiently large, take the forms

$$(59) \quad y_1(x) = x^g \frac{\Gamma(x) \lambda_1^x (1 + \epsilon_1(x))}{\Gamma\left(x + \frac{p}{q\lambda_2 + r}\right)}, \quad y_2(x) = x^g \frac{\Gamma(x) \lambda_2^x (1 + \epsilon_2(x))}{\Gamma\left(x + \frac{p}{q\lambda_1 + r}\right)},$$

wherein  $\lim_{x \rightarrow \infty} \epsilon_1(x) = \lim_{x \rightarrow \infty} \epsilon_2(x) = 0$ , and the constants have the values

$$g = \frac{a_1 b_2 - a_2 b_1}{a_1 a_2}, \quad p = a_0 a_1 b_2 + a_1 a_2 b_0 - 2a_0 a_2 b_1, \quad q = a_0 a_1^2, \quad r = 2a_0 a_1 a_2.$$

Moreover, for the cases in which  $\lambda_1, \lambda_2$  are distinct but of equal modulus we see from (57) and (58) that the result just obtained will continue to hold true, by Theorem I, provided that for all  $x \geq \alpha$  we have

$$|x + \mu/2\nu(1 + \nu)| = |x - \mu/2\nu(1 - \nu)|,$$

i. e.  $|x + c_1| = |x + c_2|$  where

$$c_1 = \frac{p}{q\lambda_2 + r}, \quad c_2 = \frac{p}{q\lambda_1 + r}.$$

But if this latter condition is satisfied, it is evident that  $c_1$  and  $c_2$  are conjugate imaginaries, and conversely. Thus the result already obtained when  $|\lambda_1| \neq |\lambda_2|$  will continue to hold true when  $|\lambda_1| = |\lambda_2|$ , provided that  $\lambda_1 \neq \lambda_2$ , and either  $c_1, c_2$  are conjugate imaginaries or  $c_1 = c_2 = 0$ .

We note also that the form of the solutions (59) may be somewhat simplified by making use of the well known asymptotic relation

$$\Gamma(x) \sim \sqrt{2\pi} e^{-x} x^{x-1/2}.$$

Thus for any constant  $l$  we have

$$\frac{\Gamma(x+l)}{\Gamma(x)} \sim e^{-l} (x+l)^l \left(1 + \frac{l}{x}\right)^{x-l} \sim e^{-l} x^l \left(1 + \frac{l}{x}\right)^x \sim x^l,$$

so that relations (59) may be replaced by

$$(60) \quad y_1(x) = x^{h_1} \lambda_1^x (1 + \epsilon_1(x)), \quad y_2(x) = x^{h_2} \lambda_2^x (1 + \epsilon_2(x)),$$

wherein

$$h_1 = g - c_1 = \frac{a_1 b_2 - a_2 b_1}{a_1 a_2} - \frac{p}{q\lambda_2 + r}, \quad h_2 = \frac{a_1 b_2 - a_2 b_1}{a_1 a_2} - \frac{p}{q\lambda_1 + r}.$$

8. Besides the cases already considered in which  $a_0 a_1 a_2 \neq 0$  it is deserving of note that whenever  $a_0 a_2 \neq 0$ ,  $a_1 = b_1 = 0$ , the nature of the solutions of (51) for large positive values of  $x$  may still be found by the application of known results. For after making the transformation

$$y(x) = x^g \left( 1 + \frac{h}{x} \right) v(x)$$

the same equation takes the form

$$A_0(x)v(x+2) + A_1(x)v(x+1) + A_2(x)v(x) = 0,$$

where

$$A_0(x) = a_0 + (b_0 + ha_0)\frac{1}{x} + \dots,$$

$$A_1(x) = \frac{c_1}{x^2} + \dots,$$

$$A_2(x) = a_2 + (b_2 - 2ga_2 + ha_2)\frac{1}{x} + \dots$$

Then by choosing  $h$  and  $g$  so that the coefficients of  $1/x$  here appearing vanish, i. e.,

$$g = \frac{a_0 b_2 - a_2 b_0}{2a_0 a_2}, \quad h = -\frac{b_0}{a_0},$$

we may at once apply to equation (51) the Theorem III of the aforesaid memoir. As the roots of the equation  $a_0 \lambda^2 + a_2 = 0$  are unequal but of equal modulus, we conclude that there are two solutions under the present hypotheses having respectively the forms

$$(61) \quad y_1(x) = x^g \lambda_1^x (1 + \epsilon_1(x)), \quad y_2(x) = x^g \lambda_2^x (1 + \epsilon_2(x)) \quad \left( \lim_{x \rightarrow \infty} \epsilon_1(x) = \lim_{x \rightarrow \infty} \epsilon_2(x) = 0 \right),$$

where

$$g = (a_0 b_2 - a_2 b_0) / 2a_0 a_2.$$

9. Returning now to the original equation (48) and recalling that

$$u(x)[\Gamma(x)]^k y(x) = \left( \frac{x}{e} \right)^{kx} x^{-ik} (2\pi)^{ik} y(x) (1 + \epsilon(x)),$$

where  $\lim_{x \rightarrow \infty} \epsilon(x) = 0$ , and making use of equations (50) we reach in summary the following theorem:

**THEOREM II.** *Given the equation*

$$P_0(x)u(x+2) + x^k P_1(x)u(x+1) + x^{2k} P_2(x)u(x) = 0,$$

where  $k$  is any integer, positive, negative or zero, while  $P_0(x)$ ,  $P_1(x)$ ,  $P_2(x)$

are convergent series for sufficiently large values of  $x$ , or are any functions of  $x$  developable asymptotically in the form

$$P_{\lambda}(x) = a_{\lambda} + \frac{b_{\lambda}}{x} + \frac{c_{\lambda}}{x^2} + \dots + \frac{p_{\lambda} + \omega_{\lambda}(x)}{x^n} \quad (\lim_{x \rightarrow \infty} \omega_{\lambda}(x) = 0).$$

CASE I. If  $a_0 a_1 a_2 \neq 0$  and if the roots  $\lambda_1, \lambda_2$  of the quadratic equation  $a_0 \lambda^2 + a_1 \lambda + a_2 = 0$  are of unequal modulus, there are two solutions of the given equation which, when considered for all positive integral values of  $x$  sufficiently large, take the respective forms

$$u_1(x) = \left(\frac{x}{e}\right)^{kx} x^{\rho + \sigma_1 \lambda_1^x} (1 + \epsilon_1(x)), \quad u_2(x) = \left(\frac{x}{e}\right)^{kx} x^{\rho + \sigma_2 \lambda_2^x} (1 + \epsilon_2(x))$$

$$(\lim_{x \rightarrow \infty} \epsilon_1(x) = 0, \lim_{x \rightarrow \infty} \epsilon_2(x) = 0).$$

wherein  $\rho, \sigma_1, \sigma_2$  are constants defined as follows:

$$\rho = \frac{a_1 b_2 - a_2 b_1}{a_1 a_2} - \frac{k}{2},$$

$$\sigma_i = \frac{2a_0 a_2 b_1 - a_0 a_1 b_2 - a_1 a_2 b_0 - k a_0 a_1 a_2}{a_0 a_1 (a_1 \lambda_i + 2a_2)} \quad (i = 1, 2).$$

Moreover, the same result holds true whenever the roots  $\lambda_1, \lambda_2$  are unequal but of equal modulus, provided that  $\sigma_1$  and  $\sigma_2$  are conjugate imaginaries, including the case in which  $\sigma_1 = \sigma_2 = 0$ .

CASE II. If  $a_0 a_2 \neq 0, a_1 = b_1 = 0$  and if we represent by  $\lambda_1, \lambda_2$  the roots of the quadratic equation  $a_0 \lambda^2 + a_2 = 0$ , there are two solutions of the given equation which when considered for values of  $x$  sufficiently large take the forms

$$u_1(x) = \left(\frac{x}{e}\right)^{kx} x^{\rho \lambda_1^x} (1 + \epsilon_1(x)), \quad u_2(x) = \left(\frac{x}{e}\right)^{kx} x^{\rho \lambda_2^x} (1 + \epsilon_2(x))$$

$$(\lim_{x \rightarrow \infty} \epsilon_1(x) = \lim_{x \rightarrow \infty} \epsilon_2(x) = 0),$$

where  $\rho$  is defined by the relation

$$\rho = \frac{a_0 b_2 - a_2 b_0}{2a_0 a_2} - \frac{k}{2}.$$

10. The results obtained under Case I of the above theorem are in accord with those obtained by HORN.\* To show this we evidently need to show merely that our values of  $\rho + \sigma_1, \rho + \sigma_2$  are equal respectively to the quantities  $\rho_1, \rho_2$  employed by HORN, and defined† by the relation

$$\rho_1 = -\frac{k(a_1 + 4a_0 \lambda_1)}{2(a_1 + 2a_0 \lambda_1)} - \frac{b_2 + b_1 \lambda_1 + b_0 \lambda_1^2}{a_1 \lambda_1 + 2a_0 \lambda_1^2},$$

with a similar formula for  $\rho_2$  obtained by replacing  $\lambda_1$  by  $\lambda_2$ .

\* Loc. cit., p. 192.

† Loc. cit., p. 191, footnote.

Now in the sum  $\rho + \sigma_1$  the coefficient of  $\frac{1}{2}k$  is  $-1 - 2a_2/(a_1\lambda_2 + 2a_2)$ . But the coefficient of  $-\frac{1}{2}k$  in  $\rho_1$  may be written in the form

$$-1 - \frac{2a_0\lambda_1}{a_1 + 2a_0\lambda_1} = -1 - \frac{2a_0\lambda_1\lambda_2}{a_1\lambda_1 + 2a_0\lambda_1\lambda_2},$$

which becomes the same as the coefficient of  $-\frac{1}{2}k$  in  $\rho + \sigma_1$  since  $\lambda_1\lambda_2 = a_2/a_0$ . It remains only to show that

$$\frac{a_1b_2 - a_2b_1}{a_1a_2} + \frac{2a_0a_2b_1 - a_0a_1b_2 - a_1a_2b_0}{a_0a_1(a_1\lambda_2 + 2a_2)} = -\frac{b_2 + b_1\lambda_1 + b_0\lambda_1^2}{a_1\lambda_1 + 2a_0\lambda_1^2}.$$

This reduces to an identity by virtue of the relations

$$a_1\lambda_2 + 2a_2 = \frac{a_2}{a_0\lambda_1}(a_1 + 2a_0\lambda_1), \quad a_0\lambda_1^2 + a_1\lambda_1 + a_2 = 0.$$

Similarly, by using  $\lambda_2$  instead of  $\lambda_1$  we obtain  $\rho + \sigma_2 = \rho_2$ .

It is to be observed that HORN's work concerns only case I.

11. As an illustration of the application of the preceding theorem, let us consider the equation

$$(62) \quad \left(1 + \frac{2}{x}\right)u(x+2) - \left(2 + \frac{3}{x}\right)zu(x+1) + \left(1 + \frac{1}{x}\right)u(x) = 0,$$

one of whose solutions is Legendre's function of the first kind  $P_x(z)$ . For simplicity, we shall confine ourselves to real values of  $z$ .

Here we have  $k=0$ ,  $a_0=1$ ,  $a_1=-2z$ ,  $a_2=1$ ,  $b_0=2$ ,  $b_1=-3z$ ,  $b_2=1$ . Whence the roots  $\lambda_1, \lambda_2$  are those of the quadratic  $\lambda^2 - 2z\lambda + 1 = 0$ ; i. e.,  $\lambda_1 = z + \sqrt{z^2 - 1}$ ,  $\lambda_2 = z - \sqrt{z^2 - 1}$ . Thus, we shall have  $|\lambda_1| \neq |\lambda_2|$  if  $|z| > 1$ , while we shall have  $|\lambda_1| = |\lambda_2|$  but  $\lambda_1 \neq \lambda_2$  if  $|z| < 1$ . Applying Case I of Theorem II, observing that  $a_0a_1a_2 \neq 0$  when  $z \neq 0$  and that in the present instance  $\rho = -\frac{1}{2}$ ,  $\sigma_1 = \sigma_2 = 0$ , we find that for all real values of  $z$  except  $z=0$  the general solution of the above equation, when considered for all positive integral values of  $x$  sufficiently large takes the form

$$u(x) = \frac{1}{\sqrt{x}} \left[ k_1(z + \sqrt{z^2 - 1})^x (1 + \epsilon_1(x)) + k_2(z - \sqrt{z^2 - 1})^x (1 + \epsilon_2(x)) \right]$$

$$(\lim_{x \rightarrow \infty} \epsilon_1(x) = \lim_{x \rightarrow \infty} \epsilon_2(x) = 0).$$

$k_1, k_2$  being arbitrary constants.

Moreover, precisely the same result holds when  $z=0$ , as appears directly by applying Case II of the same theorem.

If, in particular,  $-1 < z < 1$ , we may place  $z = \cos \xi$  and write  $(z + \sqrt{z^2 - 1})^x = \cos x\xi + i \sin x\xi$ . The solution  $u(x)$  then takes the form

$$u(x) = \frac{1}{\sqrt{x}} \left[ k_1(1 + \epsilon_1(x)) \cos x\xi + k_2(1 + \epsilon_2(x)) \sin x\xi \right],$$

where  $k_1, k_2, \epsilon_1, \epsilon_2$  have the properties already mentioned. Moreover, upon determining two constants  $\lambda, \mu$  from the equations  $\lambda \sin \mu = k_1, \lambda \cos \mu = k_2$ , and inserting for the constants  $k_1, k_2$  these expressions, we obtain for  $u(x)$  the form

$$u(x) = \frac{\lambda}{\sqrt{x}} [\sin(x\xi + \mu) + \epsilon(x)] \quad \left( \lim_{x \rightarrow \infty} \epsilon(x) = 0 \right)$$

where  $\lambda$  and  $\mu$  are arbitrary constants.

This last result for the special case in which  $u(x) = P_x(z)$  agrees with other well known results respecting the behavior of Legendre's function of the first kind for large values of  $x$ . Previous investigations upon the subject, however, appear to have been from the standpoint of the *differential* equation satisfied by  $P_x(z)$  rather than from that of the *difference* equation (62).\*

---

\* Cf. DINI, *Studi sulle equazioni differenziali*, Annali di Matematica, ser. 3, vol. 3 (1899), p. 178.