CONCERNING APPROACHABILITY OF SIMPLE CLOSED AND OPEN CURVES*

BY

JOHN ROBERT KLINE

Schoenflies† was the first to formulate the converse of the fundamental theorem of C. Jordan‡ that a simple closed curve§ lying wholly within a plane decomposes the plane into an inside and an outside region. The statement of this converse theorem is as follows: *Suppose K is a closed, bounded set of points lying in a plane S and that $S - K = S_1 + S_2$, where S_1 and S_2 are point-sets such that (1) every two points of S_i ($i = 1, 2$) can be joined by an arc lying entirely in S_i (2) every arc joining a point of S_1 to a point of S_2 contains at least one point of K (3) if O is a point of K and P is a point not belonging to K , then P can be joined to O by an arc that has no point except O in common with K . Every point-set that satisfies these conditions is a simple closed curve.* Schoenflies used metrical hypotheses in his proof. Lennes gave a proof of this converse theorem using straight lines.|| R. L. Moore pointed out that a proof similar in large part to that of Lennes can be carried through with the use of arcs and closed curves on the basis of his system of postulates Σ_3 , thus furnishing a non-metrical proof of the converse theorem.¶

In all these proofs the condition numbered three, the condition of approachability (*erreichbarkeit*) plays a fundamental rôle. It is the purpose of the present paper to study the effect of substituting for the condition of approach-

* Presented to the Society, April, 1918.

† Cf. A. Schoenflies, *Ueber einen grundlegenden Satz der Analysis Situs*, Nachrichten der Göttinger Gesellschaft der Wissenschaften, 1902, p. 185.

‡ C. Jordan, *Cours d'Analyse*, 2d ed., Paris, 1893, p. 92.

§ If A and B are distinct points, then a *simple continuous arc* from A to B is defined by Lennes as a bounded, closed, connected set of points containing A and B , but containing no proper connected subset containing both A and B , Cf. N. J. Lennes *Curves in non-metrical analysis situs with an application in the calculus of variations*, American Journal of Mathematics, vol. 33 (1911), p. 308. A *simple closed curve* is a set of points composed of two arcs AXB and AYB having no point in common other than A and B . Hereafter in this paper "arc" and "closed curve" will be considered synonymous with "simple continuous arc" and "simple closed curve," respectively.

|| Cf. N. J. Lennes, loc. cit., § 5.

¶ Cf. R. L. Moore, *On the foundations of plane analysis situs*, these Transactions, vol. 17 (1916), p. 59.

ability, the condition that the set is "connected in kleinem."* The results obtained are embodied in the following theorem:

THEOREM A. Suppose K is a closed plane point-set, S is the set of all points of the plane, while $S - K = S_1 + S_2$, where S_1 and S_2 are two mutually exclusive domains† such that every point of K is a common boundary point of S_1 and S_2 . Then a necessary and sufficient condition that K be either a simple closed curve or an open curve‡ is that K be connected in kleinem.

That the condition stated in Theorem A is necessary is evident. I will proceed to show that it is sufficient. Suppose K is a connected in kleinem set satisfying the conditions stipulated in Theorem A. Then the following lemmas hold true:

LEMMA A. Every arc joining a point of S_1 to a point of S_2 contains a point of K .

Proof. Suppose it were possible to draw an arc from a point P_1 of S_1 to a point P_2 of S_2 that contains no point of K . Then let us divide the arc $P_1 P_2$ into two sets, M_1 and M_2 , where M_1 is the set of all points of $P_1 P_2$ that belong to S_1 , while M_2 is the set of all points of $P_1 P_2$ which belong to S_2 . As $P_1 P_2$ is a connected point-set either M_1 contains a limit point of M_2 or M_2 contains a limit point of M_1 .

Case I. A point F of M_1 is a limit point of M_2 . As F is a point of the domain S_1 , there exists a region containing F and lying entirely in S_1 . As S_1 and S_2 are mutually exclusive domains, this region contains no point of M_2 . Hence F cannot be a limit point of M_2 .

Case II. A point G of M_2 is a limit point of M_1 . This is impossible as in Case I.

Hence we are led to a contradiction if we suppose our lemma false.

LEMMA B. The set K is connected.

Proof. Suppose K were not connected. Then it could be divided into two mutually exclusive sets K_1 and K_2 , neither of which contains a limit point of the other one. Let P_i ($i = 1, 2$) denote a point of K_i . Put about P_i a circle R_i having P_i as center and such that R_i and its interior lie entirely

* Cf. Hans Hahn, *Ueber die allgemeinste ebene Punktmenge die stetiges Bild einer Strecke ist*, Jahresbericht der Deutschen Mathematiker Vereinigung, vol. 23 (1914), pp. 318-22. According to Hahn, a set of points C is said to be connected in kleinem if, whenever P is a point of C , ϵ a positive number and K a circle of radius $1/\epsilon$ with center at P , then there exists within K and with center at P , another circle $K_{\epsilon, P}$ such that if X is a point of C within $K_{\epsilon, P}$ then X and P lie together in some connected subset of C that lies entirely within K .

† A domain is a connected set of points M such that if P is a point of M , then there is a region that contains P and lies in M .

‡ An open curve is defined by R. L. Moore as a closed, connected, set of points M such that if P is a point of M , then $M - P$ is the sum of two mutually exclusive connected point-sets, neither of which contains a limit point of the other.

without R_{i+1} .^{*} As K is connected in kleinem, there exists a circle $\bar{R}$, lying within R_i and with center at P_i such that if X_i is a point of K within $\bar{R}_i$, then X_i and P_i lie on some connected subset of K lying within R_i . It may easily be shown that X_i can be joined to P_i by a simple continuous arc of K lying entirely within R_i .[†] As every point of K is a common boundary point of S_1 and S_2 , then there exists within $\bar{R}_i$ a point M_{ij} ($j = 1, 2$) belonging to S_j . As S_j is a domain, then there exists a simple continuous arc $M_{1j} K_j M_{2j}$ lying entirely in S_j . Join M_{ij} to P_i by a simple continuous arc $M_{ij} L_{ij} P_i$ lying entirely within $\bar{R}_i$ and let G_{ij} denote the first point of K on the arc $M_{ij} L_{ij} P_i$ following M_{ij} . Then we may join G_{i1} to G_{i2} by an arc $G_{i1} F_i G_{i2}$ belonging to K and lying entirely within R_i . The point-set $G_{11} M_{11}$ (on $M_{11} L_{11} P_1$) + $M_{11} K_1 M_{21}$ + $M_{21} G_{21}$ (on $M_{21} L_{21} P_2$) contains as a subset a simple continuous arc $G_{11} H_1 G_{21}$ lying except for its endpoints entirely in S_1 , while the set $G_{12} M_{12}$ (on $M_{12} L_{12} P_1$) + $M_{12} K_2 M_{22}$ + $M_{22} G_{22}$ (on $M_{22} L_{22} P_2$) contains as a subset a simple continuous arc $G_{12} H_2 G_{22}$ lying except for its endpoints entirely in S_2 . We then have a closed curve $G_{11} F_1 G_{12} H_2 G_{22} - F_2 G_{21} H_1 G_{11}$ such that the arcs $G_{11} F_1 G_{12}$ and $G_{21} F_2 G_{22}$ lie entirely on K and within R_1 and R_2 , respectively, while $G_{11} H_1 G_{21}$ [‡] and $G_{12} H_2 G_{22}$ belong to S_1 and S_2 , respectively.

All points of $G_{11} F_1 G_{12}$ belong to K_1 . For suppose a point H of $G_{11} F_1 G_{12}$ belonged to K_2 . As H is joined to G_{11} , which in turn can be joined to P_1 by an arc of K lying entirely within R_1 , it follows that H can be joined to P_1 by an arc HFP_1 of K lying entirely within R_1 . Let $[\bar{H}_1]$ denote the set of all points of HFP_1 belonging to K_1 while $[\bar{H}_2]$ denotes the set of all points of HFP_1 belonging to K_2 . Clearly neither of these sets contains a limit point of the other. Hence the arc HFP_1 is not a connected point-set. Hence the supposition that H belongs to K_2 has led to a contradiction. In like manner, all points of $G_{21} F_2 G_{22}$ belong to K_2 .

The interior of $G_{11} F_1 G_{12} H_2 G_{22} F_2 G_{21} H_1 G_{11}$ must contain at least one point of K . For suppose it does not contain a point of K . Then the interior of $G_{11} F_1 G_{12} H_2 G_{22} F_2 G_{21} H_1 G_{11}$ is a subset of $S_1 + S_2$. Suppose it contains a point H of S_1 . Then H can be joined to H_2 by an arc HXH_2 lying except for H_2 entirely within $G_{11} F_1 G_{12} H_2 G_{22} F_2 G_{21} H_1 G_{11}$.[§] Let $[W_1]$ denote the set of all points of HXH_2 belonging to S_1 while $[W_2]$ denotes the set of all points of HXH_2 which are points of S_2 . Clearly neither of these sets contains

^{*} It is understood that subscripts are reduced modulo 2,

[†] Cf. R. L. Moore, *A theorem concerning continuous curves*, Bulletin of the American Mathematical Society, vol. 23 (1917). While Professor Moore's theorem states that every two points of a continuous curve can be joined by a simple continuous arc lying entirely on the given continuous curve, it is clear that his methods suffice to prove the above stronger statement.

[‡] If AXB is an arc, then the symbol AXB will denote $AXB - A - B$,

[§] Cf. R. L. Moore, *Foundations of plane analysis situs*, loc. cit., Theorem 39, pp. 153-5.

a limit point of the other. Hence the arc HXH_2 is not a connected point-set. In like manner the supposition that there is within $G_{11} F_1 G_{12} H_2 G_{22} F_2 - G_{21} H_1 G_{11}$, a point of S_2 leads to a contradiction. Hence $G_{11} F_1 G_{12} H_2 G_{22} F_2 - G_{21} H_1 G_{11}$ must enclose a point of K .

Let $[V_2]$ denote the set of all points V_2 such that either (1) V_2 is a point of $G_{21} F_2 G_{22}$, or (2) V_2 is a point such that there exists a closed connected set $V_2 XF'_2$ belonging to K and lying within or on $G_{11} F_1 G_{12} H_2 G_{22} F_2 G_{21} H_1 G_{11}$, and such that F'_2 is a point of $G_{21} F_2 G_{22}$. As K is connected in kleinem it may easily be proved that $[V_2]$ is a closed set. It is also true that all points of $[V_2]$ belong to K_2 . Hence no point of $G_{11} F_1 G_{12}$ either belongs to or is a limit point of $[V_2]$. It may also be proved with the use of the in kleinem property that no point of $[V_2]$ is a limit point of a set of points of K lying within $G_{11} F_1 G_{12} H_2 G_{22} F_2 G_{21} H_1 G_{11}$ and containing no point of V_2 . There exists an arc $H_1 YH_2$ such that (1) $\overline{H_1 YH_2}$ is a subset of the interior of $G_{11} F_1 G_{12} H_2 G_{22} F_2 G_{21} H_1 G_{11}$ and (2) $\overline{H_1 YH_2}$ contains no points of $[V_2]$.^{*} Let $[V_1]$ denote the set of all points of K within or on the closed curve, $H_1 YH_2 G_{22} F_2 G_{21} H_1$, not belonging to $[V_2]$. The set $[V_1]$ is closed. Put about each point of $[V_1]$ a circle lying entirely within $G_{11} F_1 G_{12} H_2 G_{22} F_2 - G_{21} H_1 G_{11}$ and containing within it or on its boundary no point of $[V_2]$. By the Heine-Borel Property, there exists a finite number of circles of the above set, $C_1, C_2, \dots, C_n$, covering $[V_1]$. With the use of Theorems 41, 42, 43, and 44 of Professor Moore's Foundations we may easily obtain from the set $C_1, C_2, \dots, C_n$ and the closed curve $G_{11} F_1 G_{12} H_2 YH_1 G_{11}$, a new closed curve $G_{11} F_1 G_{12} H_2 ZH_1 G_{11}$, where the arc $H_1 G_{11} F_1 G_{12} H_2$ of the new closed curve $G_{11} F_1 G_{12} H_2 ZH_1 G_{11}$ is the arc $H_1 G_{11} F_1 G_{12} H_2$ of $G_{11} F_1 G_{12} H_2 YH_1 G_{11}$ and where $H_2 ZH_1$ is free from points of K and lies within $G_{11} F_1 G_{12} H_2 G_{22} F_2 - G_{21} H_1 G_{11}$. But then we have a point of S_1 joined to a point of S_2 by an arc containing no point of K . Thus the supposition that K is not connected, leads to a contradiction.

LEMMA C. *If K contains one simple closed curve J , then all points of K belong to J .*

Proof. Suppose Lemma C is not true. Then K contains a closed curve J and at least one point P not on J . Two cases may arise:

Case I. P is within J . As every point of K is a common boundary point of S_1 and S_2 , the interior of J contains a point P_1 of S_1 and a point P_2 of S_2 . The exterior of J cannot contain a point $\bar{P}_1$ of S_1 . For suppose it did. Then any arc from P_1 to $\bar{P}_1$ would contain a point of J and hence a point of K , contrary to the fact that S_1 is a domain. In like manner no point of S_2 can be in the exterior of J . Hence the exterior of J must be a subset of K , while

^{*}Cf. my paper, *A definition of sense on plane closed curves in non-metrical analysis situs*, *Annals of Mathematics*, vol. XIX (1918), Theorem D, pp. 188-9.

S_1 and S_2 are subsets of the interior of J . But this is impossible because no point without J is a limit point of a set of points lying entirely within J thus making it impossible that every point of K be a common boundary point of S_1 and S_2 . Hence the supposition that P is within J has led to a contradiction.

Case II. P is without J . Case II may be proved impossible by an argument similar to that used in Case I.

An immediate consequence of Lemma C is that if K is not a simple closed curve, then there is but one K -arc from a point A of K to a distinct point B of K .

LEMMA D. *The set K does not contain three arcs OP_1 , OP_2 , and OP_3 , no two of which have a common point other than O .*

Proof. Suppose Lemma D were false. Then there would exist three arcs OP_1 , OP_2 , and OP_3 , no two of which have a point in common other than O . Put about P_i ($i = 1, 2, 3$) a circle C_i such that the point-set $OP_{i+1}^* + OP_{i+2}$ is a subset of the exterior of C_i and such that C_i has no point in common with $C_{i+1} + C_{i+2}$. As K is connected in kleinem, there exists within C_i and with center at P_i , another circle C_{P_i, c_i} such that if X_i is a point of K within C_{P_i, c_i} , then there is an arc from X_i to P_i every point of which is a point of K and which lies entirely within C_i .† As all points of K are limit points of both S_1 and S_2 , C_{P_i, c_i} must contain at least one point $P_{i, 1}$ of S_1 . As S_1 is a domain, there is an arc $P_{11}P_{21}$ from P_{11} to P_{21} all points of which belong to S_1 . Join $P_{i, 1}$ to P_i by an arc $P_{i, 1}P_i$ lying entirely within C_{P_i, c_i} and let X_i denote the first point of the arc $P_{i, 1}P_i$ after $P_{i, 1}$, which belongs to K . There exists an arc X_iP_i from X_i to P_i belonging to K and lying entirely within C_i . Let P'_i denote the first point of the arc X_iP_i which is on OP_i . The point-set $P'_1X_1 + X_1P_{11} + P_{11}P_{21} + P_{21}X_2 + X_2P'_2$ contains as a subset an arc $P'_1F_1P'_2$ such that (1) $P'_1F_1P'_2$ has no point in common with $OP_1 + OP_2 + OP_3$, (2) all points of $P'_1F_1P'_2$ belong to either K or S_1 , (3) at least one point, F_1 , of S_1 is a point of $P'_1F_1P'_2$. By methods similar to those just employed, we may construct an arc $Q'_1H_2Q'_2$ from a point Q'_1 of OP_1 to a point Q'_2 of OP'_2 such that (1) Q'_i is on OP_i between P'_i and O , (2) all points of $Q'_1H_2Q'_2$ belong to either S_2 or K , (3) except for Q'_1 and Q'_2 , $Q'_1H_2Q'_2$ has no point in common with $P'_1FP'_2 + OP_1 + OP_2 + OP_3$, (4) at least one point H_2 of $Q'_1H_2Q'_2$ belongs to S_2 . Two cases may arise:

Case I. $Q'_1H_2Q'_2$ is entirely within $OP'_1F_1P'_2O$. Then the interior of $OP'_1F_1P'_2O = Q'_1H_2Q'_2 +$ the interior of $OQ'_1H_2Q'_2O +$ the interior of $P'_1F_1P'_2Q'_2H_2Q'_1P'_1$. The point-set $OP_3 + P_3$ is either entirely within or entirely without $OQ'_1H_2Q'_2O$.

(a) Suppose $OP_3 + P_3$ is entirely within $OQ'_1H_2Q'_2O$. Then $OQ'_1H_2Q'_2O$

* It is understood throughout this argument that subscripts are reduced modulo 3.

† See an earlier footnote.

must enclose at least one point L of S_1 . But then an arc from L to F_1 must contain at least one point of $OQ'_1 H_2 Q'_2 O$. Hence, as $OQ'_1 H_2 Q'_2 O$ is a subset of $K + S_2$, no such arc LF_1 can lie entirely in S_1 , contrary to the fact that S_1 is a domain.

(b) Suppose $\overline{OP_3} + P_3$ is entirely without $OQ'_1 H_2 Q'_2 O$. It follows that $\overline{OP_3} + P_3$ is entirely without $OP'_1 F_1 P'_2 O$. Then the exterior of $OP'_1 F_1 P'_2 O$ contains at least one point M of S_2 . Then any arc from M to H_2 must contain at least one point of $OP'_1 F_1 P'_2 O$ and hence at least one point not in S_2 . But this is contrary to the fact that S_2 is a domain.

Thus in Case I we are led to a contradiction.

Case II. $Q'_1 H_2 Q'_2$ is without $OP'_1 F_1 P'_2 O$. We may show that Case II is impossible by methods similar to those used in Case I.

LEMMA E. *If O is a point of K and P is a point of S_i ($i = 1, 2$) then there exists at least one arc OP such that $\overline{OP} + P$ is a subset of S_i .*

Proof. Two conceivable cases may arise.

Case I. There exist points A_1 and A_2 of K [$A_1 \neq O \neq A_2$] such that O is a point of the arc $A_1 OA_2$ belonging to K . By the same methods as were used in the preceding lemma we may construct an arc $A'_1 F_1 A'_2$ such that (1) on $A_1 OA_2$ the order $A_1 A'_1 OA'_2 A_2$ holds, (2) $A'_1 F_1 A'_2$ is a subset of $S_1 + K$, (3) at least one point F_1 of $A'_1 F_1 A'_2$ is a point of S_1 , (4) no point of $A'_1 F_1 A'_2$ belongs to $A_1 OA_2$. The point O is not a limit point of $K - A'_1 OA'_2$. For suppose it were. Then it would be a sequential limit point of a set of points $P_1, P_2, \dots$, every one of which belongs to $K - A'_1 OA'_2$. Put about O as center a circle M such that A'_1 and A'_2 are both without M . As K is connected in kleinem there exists another circle $\bar{M}$ lying within M and having its center at O such that if X is a point of K within $\bar{M}$, then X and O can be joined by an arc of K lying entirely within M . Let $\bar{P}$ denote that point of the set $P_1, P_2, \dots$ of lowest subscript which lies within $\bar{M}$, while $\bar{P}O$ denotes an arc of K from $\bar{P}$ to O lying entirely within M . Let O' denote the first point of $\bar{P}O$ which is on $A'_1 OA'_2$. Then the set K contains three arcs $A'_1 O'$, $A'_2 O'$, and PO' , no two of which have a point in common other than O' . But this is contrary to Lemma D. Hence O cannot be a limit point of $K - A'_1 OA'_2$. There exists a closed curve G enclosing O but enclosing no points of $A'_1 F_1 A'_2 + [K - A'_1 OA'_2]$. Then there exist two closed curves J'_1 and J'_2 such that (1) every point of J'_1 or J'_2 belongs either to G or to $A'_1 F_1 A'_2 OA'_1$ (2) O is on J'_1 and on J'_2 (3) every point within J'_1 is within $A'_1 F_1 A'_2 OA'_1$ while every point within J'_2 is without $A'_1 F_1 A'_2 OA'_1$ (4) every point within either J'_1 or J'_2 is within G .^{*} It is clear that either the interior of J'_1 or the interior of J'_2 is a subset of S_1 while the interior of the other of these two closed curves is a subset of S_2 . Let J_1 denote that one whose interior is a subset of S_1 while

^{*} Cf. R. L. Moore, *Foundations*, Theorem 43, pp. 156-7.

J_2 denotes the one whose interior is a subset of S_2 . Let E denote a point within J_1 , while P_1 is any other point of S_1 . There exists an arc EO such that $EO - O$ is a subset of the interior of J_1 .* As S_1 is a domain, there is an arc EP_1 lying entirely in S_1 . The point-set $EO + EP_1$ contains as a subset an arc from P_1 to O lying except for O entirely in S_1 . In like manner we may show that any point P_2 of S_2 can be joined to O by an arc lying except for O entirely in S_2 .

Case II. There do not exist two distinct points A_1 and A_2 of K such that O is on an arc of K from A_1 to A_2 . Let A denote a point of K different from O while ARO denotes an arc of K from A to O . By an argument similar to that employed in Case I we may show that if O were a limit point of $K - ARO$, then either there would exist three arcs AR' , $R'O$, and $R'P$, no two of which have a point in common other than R' or there would exist a point A' , ($A \neq A' \neq O$) such that O is an arc of K from A' to A . But the first of these possibilities contradicts Lemma *D* while the second is contrary to the hypothesis of Case II. Hence O cannot be a limit point of $K - ARO$. Put about O a circle C that neither contains or encloses any point of $K - ARO$. Let $P_1, P_2, \dots$ denote a set of points of S_1 approaching O as their sequential limit point. It is possible to pass at least one simple continuous arc† through $ARO + P_1 + P_2 + \dots$. Let $P'_1 ORA$ denote one such arc. If the interval OP'_1 of the arc $P'_1 ORA$ does not lie entirely within C , let P' denote the first point which it has in common with C . Otherwise let P' denote P'_1 . Let $\bar{P}$ denote that point of the set $P_1, P_2, \dots$ of lowest subscript lying on OP' . It is clear that the sub-arc $\bar{OP}$ of $P'_1 ORA$ lies, except for O , entirely in $\bar{S}_1$. Let F_1 denote any other point of S_1 . Join F_1 to $\bar{P}$ by an arc lying entirely in S_1 . Then the point-set $\bar{OP} + \bar{PF}_1$ contains as a subset an arc from O to F_1 , lying except for O entirely in S_1 .

In like manner we may show that if F_2 is a point of S_2 , F_2 can be joined to O by an arc lying except for O entirely in S_2 .

LEMMA F. *A necessary and sufficient condition that K be bounded, is that either S_1 or S_2 be bounded.*

Proof. *The condition is necessary.* Let us suppose that K is bounded while neither S_1 nor S_2 is bounded. As K is bounded, there is a circle C such that all points of K are within C . As S_1 and S_2 are unbounded, there is a point P_1 of S_1 and a point P_2 of S_2 without C . Join P_1 and P_2 by an arc lying entirely without C . By Lemma *A*, this arc must contain a point of K . But all points of K are within C . Hence we are led to a contradiction if we suppose our condition is not necessary.

* Cf. R. L. Moore, *Foundations*, Theorem 39, pp. 153-5.

† Cf. R. L. Moore and J. R. Kline, *On the most general closed point-set through which it is possible to pass a simple continuous arc*, *Annals of Mathematics*, vol. XX (1919), pp. 218-23.

The condition is sufficient. For suppose S_1 is bounded while K is unbounded. Since S_1 is bounded, there exists a circle C enclosing S_1 . Since K is unbounded, it contains a point P without C . The point P cannot be a limit point of S_1 . But this is contrary to hypothesis.

Proof of Theorem A. Two cases may arise:

Case I. K is bounded. Then, by Schoenflies' Theorem and the preceding lemmas, it follows that K is a simple closed curve.

Case II. K is unbounded. It follows, by Lemma F that neither S_1 nor S_2 is bounded. Then K is an open curve. For a proof of this statement see my paper, "The converse of the theorem concerning the division of a plane by an open curve."*

UNIVERSITY OF PENNSYLVANIA,
PHILADELPHIA, PA.

* Cf. these Transactions, vol. 18 (1917), pp. 177-184.