ON CYCLIC NUMBERS OF ONE-DIMENSIONAL COMPACT SETS*

BY

W. A. WILSON

1. An interesting class of one-dimensional compact sets is that of cyclically connected sets. P. Alexandroff† has defined the cyclic numbers of such sets and obtained important properties. It is the main purpose of this article to obtain a theorem on the divisor of a sequence and a generalization of Alexandroff's addition theorem. The proofs are based on a modification of Alexandroff's definition of cyclic number, which seems to the writer to furnish a more natural approach for those whose main interest is the theory of point sets. The work is confined to compact metric spaces.

2. A finite set of closed sets $\{d_i\}$, of which no three have common points and none is a sub-set of the union of the others, is called a *canonical set* of cells. If the set M can be expressed as the union of a canonical set of cells, each of which has a diameter less than some positive σ , we say that these cells constitute a *canonical σ -covering* of M . If M is compact and one-dimensional at most, it is well known that there is a canonical σ -covering of M for each positive σ . If M is contained in the canonical set $S = \{d_i\}$ and $M \cdot d_i \neq 0$ for each i , we call S a *canonical enclosure*.

If the cell d_i has points in common with w_i other cells, w_i is called the *order* of d_i . A *component* of the covering is a maximal sub-set of the cells which cannot be separated into two sets such that no cell of one set contains points of the other set.

If p is the number of components,

$$m = \frac{1}{2} \sum_1^n (w_i - 2) + p = \frac{1}{2} \sum_1^n w_i - n + p$$

is called the *cyclic number* of the covering or enclosure.‡ The number I defined by

$$I = \frac{1}{2} \sum_1^n (w_i - 2) = m - p$$

* Presented to the Society, September 9, 1931; received by the editors July 11, 1931.

† *Über kombinatorische Eigenschaften allgemeiner Kurven*, Mathematische Annalen, vol. 96, pp. 512-554.

‡ It is apparent from §11 of the article referred to above that m is the same as the $s(L)$ of the corresponding linear complex used by Alexandroff.

is called the *index* of the covering or enclosure. The numbers m and I will be used sometimes when the covering is not canonical. In this connection it should be noted that, if the covering satisfies the definition of being canonical, except for the existence of one or more cells each of which is a sub-set of some other cell, the canonical covering resulting from the deletion of these cells has the same cyclic number as the original one. Such a covering may be called almost canonical.

In reckoning m , p , and I , void cells are not counted. Thus, if no two cells have common points, $I = -n = -p$ and $m = 0$. At times we use $w(d_i)$, $m(S)$, $p(S)$, or $I(S)$ for definiteness.

3. THEOREM. *Let M be a compact set imbedded in a compact space Z and $\{d_i\}$ be a canonical σ -covering or σ -enclosure. Then there is an $\epsilon > 0$ such that, if e_i is the set of points of Z whose distances from d_i are not greater than ϵ , then $e_i \cdot e_j \neq 0$ if and only if $d_i \cdot d_j \neq 0$, and the set of cells $\{e_i\}$ is a canonical σ -enclosure of M of the same cyclic and component numbers as $\{d_i\}$.*

There is a positive δ less than one-third the least distance between any pair of disjoint cells $\{d_i\}$. Since the given set is canonical and the number of cells is finite, there is a positive η such that every d_i contains a point whose distance from the union of the remaining cells is greater than η . The conclusion is then valid if ϵ is less than both δ and η , and 2ϵ is less than σ minus the upper bound of the diameters of the cells $\{d_i\}$. For on account of the first statement $e_i \cdot e_j \neq 0$ if and only if $d_i \cdot d_j \neq 0$. On account of the second statement no e_i is contained in the union of the others. Finally the enclosure is a σ -enclosure because the diameter of e_i exceeds that of d_i by at most 2ϵ .

4. THEOREM. *Let A and B be canonical sets of cells such that the point sets which they form are disjoint. Then $m(A+B) = m(A) + m(B)$ and $I(A+B) = I(A) + I(B)$.*

COROLLARY. *The cyclic number or index of a canonical set of cells is the sum of the cyclic numbers or indices, respectively, of the components.*

5. THEOREM. *Let $A = \{d_i\}$ be a canonical set of cells. To each d_i let there correspond a closed sub-set e_i and let B be a canonical set of the cells $\{e_i\}$. Then $m(B) \leq m(A)$.*

First let A be connected and its cells arranged in such an order that the first n form a connected set A_n . Let B_n be the sub-set of B corresponding to cells of A_n and assume that $m(A_n) \geq m(B_n)$.

If e_{n+1} is void, it is obvious that $m(A_{n+1}) \geq m(B_{n+1})$. If e_{n+1} is not void, let it meet c components of B_n in a total of k cells. It is easily seen that

$I(B_{n+1}) = I(B_n) + k - 1$ and $p(B_{n+1}) = p(B_n) - c + 1$; hence $m(B_{n+1}) = m(B_n) + k - c$. Now d_{n+1} may meet more than k cells of A_n ; hence $m(A_{n+1}) \geq m(A_n) + k - 1$. Clearly $k - 1 \geq k - c$, unless $c = 0$. In this case $k = 0$, but on the other hand d_{n+1} meets at least one cell of A_n . Thus in every case the addition of e_{n+1} and d_{n+1} increases $m(A_n)$ by at least as much as $m(B_n)$ is increased. Hence the theorem is true by induction.

The case that A is not connected follows at once by §4.

COROLLARY 1. *Let $A = \{d_i\}$ be a canonical set of cells and $B \subset A$. Then $m(B) \leq m(A)$.*

For it is obvious that B is canonical and this corollary is the special case of the theorem where $e_i = d_i$ or e_i is void.

COROLLARY 2. *If in addition to the hypotheses of the above theorem or corollary we know that $p(B) \geq p(A)$, then $I(B) \leq I(A)$.*

COROLLARY 3. *Let M be a compact set and $\{d_i\}$ be a canonical enclosure of M of cyclic number m . Let $e_i = M \cdot d_i$. Then there is a sub-set of the cells $\{e_i\}$ which is a canonical covering of M of cyclic number not greater than m .*

For the deletion of any cell e_i which is a sub-set of the union of the other cells leaves a covering of M . If this is done successively, we finally reach a sub-set of the cells $\{e_i\}$ which is a canonical covering of M and the theorem is then applicable.

6. If $A = \{d_i\}$ and $B = \{e_k\}$ are two canonical sets of cells which have common points and $A + B$ is canonical, a discussion of $A + B$ requires the assignment of a meaning to $A \cdot B$. By $A \cdot B$ we shall mean the set of cells $\{f_r\}$ which are either common cells of A and B or divisors of a cell of A not in B and a cell of B not in A . If f_r is a cell of the first type and $f_s = d_i \cdot e_k$ is one of the second, $f_r \cdot f_s = 0$, because $A + B$ is canonical. In like manner no two cells of the second type can have common points. Thus $A \cdot B$ is canonical and each component is either a cell of the second type or a connected set of cells of the first type. It is also a simple matter to show by induction that, with the above definitions,

$$p(A + B) + p(A \cdot B) - p(A) - p(B) \geq 0.$$

7. THEOREM. *Let A , B , and $A + B$ be canonical collections of cells. Then*

$$I(A + B) = I(A) + I(B) - I(A \cdot B).$$

Let B_0 be the set of cells common to A and B ; let there be n other cells of B and let B_k be the union of B_0 and the first k of these cells $\{e_i\}$. Suppose that the theorem is true for $A + B_k$; i.e.,

$$(1) \quad I(A + B_k) = I(A) + I(B_k) - I(A \cdot B_k).$$

This is obviously true when $k=0$, since $A \cdot B_0 = B_0$.

Let e_{k+1} meet c cells of B_0 , a cells of $A - B_0$, and b cells of $B_k - B_0$. As the sets are canonical, this gives at once

$$(2) \quad I(A + B_{k+1}) = I(A + B_k) + a + b + c - 1;$$

$$(3) \quad I(B_{k+1}) = I(B_k) + b + c - 1.$$

By §6 we see that the addition of e_{k+1} adds a isolated cells to $A \cdot B_k$; hence

$$(4) \quad I(A \cdot B_{k+1}) = I(A \cdot B_k) - a.$$

Relations (2), (3), and (4) make (1) true for $k+1$. Thus the theorem is true by induction.

COROLLARY 1. *If $A+B$ is a canonical collection of cells,*

$$m(A+B) = m(A) + m(B) - m(A \cdot B) + p(A+B) - p(A) - p(B) + p(A \cdot B).$$

COROLLARY 2. *If $A+B$ is a canonical collection of cells,*

$$m(A+B) \geq m(A) + m(B) - m(A \cdot B).$$

Remark. The above theorem and the one in §14 are variants of the index formulas derived by S. Straszewicz (*Über die Zerschneidung der Ebene durch abgeschlossene Mengen*, Fundamenta Mathematicae, vol. 7, pp. 159–187) for plane sets. The index used here is not, however, the same. This theorem is easily generalized to cover the case where the set of cells is not canonical and may indeed be taken as a basis to define the cyclic number.

8. A compact metric set M is called *at least m cyclic* if for a positive σ small enough every canonical σ -covering of M has its cyclic number at least m . It is called *at most m cyclic* if for every positive σ there is some canonical σ -covering of cyclic number not more than m . If M is imbedded in a compact space, we can of course substitute the word enclosure for covering without altering the sense of these definitions.

We say that M is *m cyclic* if it is both at most and at least m cyclic; i.e., if for some $\sigma > 0$ every canonical σ -covering has its cyclic number at least m and for every $\sigma > 0$ some canonical σ -covering has the cyclic number m . If for every integer m , M is at least m cyclic, we say that it is ∞ cyclic. It is a simple matter to show from these definitions that every (at most) one-dimensional compact set has a definite cyclic number, finite or infinite. It is also clear from page 523 of the reference given in §1 that m is one less than Alexandroff's κ .

9. It is readily seen with the aid of §5, Corollary 3, that, if M and N are one-dimensional compact sets and $M \subset N$, the cyclic number of M is not more

than that of N . (Cf. Alexandroff, loc. cit., p. 523.) Hence the divisor of a decreasing sequence of at most m cyclic one-dimensional compact sets is at most m cyclic, and every set of this character contains a set irreducible with respect to these properties. But the first of these statements is seen to become false and the second doubtful, if the word *most* is replaced by *least*, and so the question of the cyclic number of the divisor of a sequence requires further investigation.

Similarly, the relation between the cyclic number of a set and those of its components is obvious when the component number is finite, but requires a proof when the component number is infinite. We begin with the second question.

10. THEOREM. *Let M be a compact one-dimensional set. If its cyclic number is finite, it is the sum of the cyclic numbers of those components having positive cyclic numbers.*

We know by §7 that for every partition of M into r disjoint closed sets $\{M_i\}$, $m(M) = \sum_i m(M_i)$. There is then a largest $r \leq m$ such that for each i , $m(M_i) \geq 1$.

Now no such M_i can contain more than one component K_i of positive cyclic number. If M_i is decomposed into disjoint closed sets P_i and Q_i , where $K_i \subseteq P_i$, it is clear that $m(P_i) = m(M_i)$ and $m(Q_i) = 0$. If this can be done for each i so that $K_i = P_i$, the theorem is evidently proved.

In the other event there is a decreasing sequence $\{P_{ij}\}$ of closed sub-sets of M_i having the component K_i as its divisor, such that $m(P_{ij}) = m(M_i)$ for each j , $Q_{ij} = M_i - P_{ij}$ is closed, and $m(Q_{ij}) = 0$. Let σ be so small that every canonical σ -covering of M_i has its cyclic number at least $m(M_i)$. Let $\delta < \sigma/2$ and let $A = \{d_n\}$ be a canonical δ -covering of K_i of cyclic number $m(K_i)$. By §3 we can take $\epsilon < \sigma/6$ and so small that, if e_n is the set of points of M_i whose distances from d_n are not greater than ϵ , the set of cells $\{e_n\}$ is a canonical σ -enclosure of some P_{ij} of cyclic number $m(K_i)$, since K_i is the divisor of $\{P_{ij}\}$. Setting $f_n = e_n \cdot P_{ij}$, there is by §5, Corollary 3, a sub-set B of cells $\{f_n\}$ which is a canonical σ -covering of P_{ij} of cyclic number not greater than $m(K_i)$. Let $C = \{g_k\}$ be a canonical σ -covering of Q_{ij} of cyclic number 0. Then $B + C$ is a canonical σ -covering of M_i of cyclic number not more than $m(K_i)$. Hence in this case also $m(K_i) = m(M_i)$, and the theorem is proved.

11. THEOREM. *Let M be an ∞ cyclic compact one-dimensional set. Then either some component of M is ∞ cyclic or an infinity of components have positive cyclic numbers.*

We first show that the assumption that all the components are 0 cyclic

leads to a contradiction. There is a dyadic decomposition* $M = \sum M_{i_1} = \sum M_{i_1 i_2}$, $= \sum M_{i_1 i_2 i_3}$, etc., into disjoint closed sets, such that each component K of M is the divisor of some monotone descending sequence $\{M_n\}$, where M_n is some $M_{i_1 i_2 \dots i_n}$.

Take $\sigma > 0$. There is a canonical $(\sigma/2)$ -covering of K , $K = U[d_r]$, whose cyclic number is 0. By §3 we can take $\delta < \sigma/6$ and so small that, if e_r is the set of points of M whose distances from d_r are not greater than δ , the set of cells $\{e_r\}$ is a canonical σ -enclosure of K of cyclic number 0. For some n_k , this enclosure contains every M_n , $n \geq n_k$, of the sequence whose divisor is K . That is, for each K there is an n_k , such that there is a canonical σ -covering of cyclic number 0 of every M_n , $n \geq n_k$, of the sequence whose divisor is K .

We now prove that n_k has an upper bound for all components $\{K\}$. If this is not true, it follows by well known theorems that there is a sequence $\{K_u\}$ of components such that $n_{k_u} \rightarrow \infty$ and a K containing the upper closed limiting set of $\{K_u\}$, as $u \rightarrow \infty$. But for u large enough, K_u and K lie in the same M_{n_k} ; whence $n_{k_u} \leq n_k$.

Let then n be the upper bound of n_k . There is then a canonical σ -covering of each M_n of cyclic number 0. As these are disjoint and their number is finite, we have a canonical σ -covering of M of cyclic number 0. Since σ was arbitrary, we see that M is 0 cyclic, contrary to hypothesis. Hence not all the components of M are 0 cyclic.

Now let M have s components $\{K_i\}$, each of cyclic number m_i , and let every other component be 0 cyclic. Taking any $\sigma > 0$, let $K = U[d_{1,k}]$ be a canonical $(\sigma/2)$ -covering of K_1 of cyclic number m_1 . Choose a positive δ so small that, if $e_{1,k}$ is the set of points of M whose distances from $d_{1,k}$ are not more than δ , then the set of cells $\{e_{1,k}\}$ is a canonical σ -enclosure of K_1 of cyclic number m_1 . We can set $M = P_1 + Q_1$, where P_1 and Q_1 are disjoint closed sets, $K_1 \subseteq P_1$, $\sum_2^s K_i \subseteq Q_1$, and P_1 is contained in the enclosure $\{e_{1,k}\}$. Hence by §5, Corollary 3, this enclosure generates a canonical σ -covering of P_1 of cyclic number at most m_1 .

In the same way Q_1 is the sum of disjoint closed sets P_2 and Q_2 , such that $K_2 \subseteq P_2$, $\sum_3^s K_i \subseteq Q_2$, and there is a canonical σ -covering of P_2 of cyclic number at most m_2 . Finally we reach $Q_s = M - \sum_1^s P_i$. This has no component of positive cyclic number and so there is a canonical σ -covering of Q_s of cyclic number 0. Thus there is for any σ a canonical σ -covering of M of cyclic number at most $m' = \sum_1^s m_i$. This is contrary to the hypothesis that M is ∞ cyclic, and so the theorem is proved.

Remark. We can combine the results of these two sections in the state-

* F. Hausdorff, *Mengenlehre*, p. 131.

ment that the cyclic number of any compact one-dimensional set is the sum of the cyclic numbers of its components, if we understand that the sum of any number of zeros is zero and that of any number of infinities is infinity. The cardinal number of the cyclic components may be that of any closed set.

12. LEMMA. *Let M and M' be compact one-dimensional sets lying in a compact space Z , $M' \subseteq M$, and M' have p' components, p' finite. Let every canonical σ -enclosure of M have its cyclic number at least m and for some $\delta < \sigma/2$ let there be a canonical δ -covering of M' of cyclic number m' and component number p' . Then there is an $\epsilon > 0$ and an $\eta > 0$ such that, if $E = \{e_k\}$ is any canonical η -enclosure of M and B is a sub-set of E for which $M' \subset B \subset V_\epsilon(M')^*$, while the other cells of E form a set C containing no point of M' , then*

$$m(E) - m(B) + p(B) \geq m - m' + p'.$$

Let $D = \{d_i\}$ be the canonical δ -covering referred to above. Let ϵ be so small that, if the set of points of Z whose distances from d_i are not greater than 2ϵ is taken as a cell, the set of such cells forms a canonical σ -enclosure of M' of cyclic number m' and component number p' . (See §3.) Let $\epsilon < \sigma/6$ and $\eta < \epsilon/2$. Let $E = \{e_k\}$ be any canonical η -enclosure of M and B be a sub-set of E for which $M' \subset B \subset V_\epsilon(M')$ and the other cells form a set C such that $M' \cdot C = 0$. Then each cell e_k of B contains a point of $V_\epsilon(d_i)$ for some i .

Let Γ_i be the union of the cells of B containing points of $V_\epsilon(d_i)$. Then $\Gamma_i \cdot \Gamma_j \neq 0$ if and only if $d_i \cdot d_j \neq 0$, and $\{\Gamma_i\}$ is a canonical σ -enclosure of M' of cyclic number m' and component number p' .

For each i let Δ_i be the union of the cells $\{e_k\}$ contained in Γ_i , but in no $\Gamma_j, j > i$. The set A of non-void cells $\{\Delta_i\}$ is a canonical enclosure of M' since $\Delta_i \subseteq \Gamma_i, m(A) \leq m'$ by §5, and $p(A) \geq p'$. It is easy to see that $A + C$ is a canonical σ -enclosure of M , and we must prove the inequality given in the statement of the theorem.

We know that

- (1) $m(A + C) - m(A) + p(A) = m(C) - m(A \cdot C) + p(A + C) - p(C) + p(A \cdot C);$
- (2) $m(B + C) - m(B) + p(B) = m(C) - m(B \cdot C) + p(B + C) - p(C) + p(B \cdot C).$

Since C has no cells in common with A or B , $m(A \cdot C) = m(B \cdot C) = 0$. As $A + C$ is obtained by combining certain cells of $B + C$ into new cells, $p(B + C) \geq p(A + C)$. Each component of $A \cdot C$ is the divisor of a cell e_k of C and a cell Δ_i of A and is therefore the sum of the divisors of e_k and one or more cells $\{e_i\}$ contained in Δ_i . As no two cells $\{\Delta_i\}$ contain common cells $\{e_i\}$, it follows that $p(B \cdot C) \geq p(A \cdot C)$.

* This notation signifies the set of points of Z whose distances from M' are less than ϵ .

Thus the left member of (2) is greater than or equal to that of (1). Since $m(A+C) \geq m$ and $p(A) \geq p'$, while $m(A) \leq m'$, this proves the theorem.

COROLLARY 1. *Under the hypotheses of the above theorem there is an $\eta > 0$ such that, if $E = \{e_k\}$ is any canonical η -enclosure of M and B is the sub-set of cells containing points of M' , then $m(E) - m(B) \geq m - m'$.*

For in this case $p(B) = p'$.

COROLLARY 2. *Let M and M' be compact one-dimensional sets with cyclic numbers m and m' , let $p' = p(M')$ be finite, and let $M' \subset M$. Then there is an $\epsilon > 0$ and an $\eta > 0$, such that, if $E = \{e_k\}$ is any canonical η -enclosure of M and B is a sub-set of E for which $M' \subset B \subset V_\epsilon(M')$, while the other cells of E form a set C containing no point of M' , then $m(E) - m(B) + p(B) \geq m - m' + p'$.*

For it is easy to show that the hypotheses of the theorem apply here.

COROLLARY 3. *Under the hypotheses of Corollary 2 there is an $\eta > 0$ such that, if $E = \{e_k\}$ is any canonical η -enclosure of M and B is the sub-set of cells containing points of M' , then $m(E) - m(B) \geq m - m'$.*

For again $p(B) = p'$.

13. THEOREM. *Let $\{M_i\}$ be a decreasing sequence of compact one-dimensional sets, each of finite cyclic number m , and let M be the divisor of the sequence. Then m is the cyclic number of M .*

By §10 each M_i contains a sub-set N_i consisting of the components of M_i whose cyclic numbers are positive and $m(N_i) = m$. As there can be at most m components in each N_i and, if $A \subset B$, $m(A) \leq m(B)$, we can assume without loss of generality that each N_i consists of p components $\{K_{ij}\}$, each of cyclic number $m_j \geq 1$, where $p \leq m$ and $m = \sum_1^p m_j$. Moreover, the sequence $\{N_i\}$ is monotone decreasing. Let N be its divisor.

It is easily seen from §8 that for each i there is a greatest σ_i such that every canonical σ_i -covering of N_i has its cyclic number at least m . If the lower bound σ of σ_i is positive, we proceed as follows. Let $m' = m(N)$ and $\{d_n\}$ be a canonical covering of N of cyclic number m' and norm $\sigma/2$. Taking ϵ small enough, we know that, if the set of points of N_1 whose distances from the cell d_n are not greater than ϵ is taken as a cell, then the set of such cells forms a canonical σ -enclosure of N and N_i , for i large enough, of cyclic number m' . This enclosure generates a canonical σ -covering of N_i of cyclic number at most m' , by §5. This is a contradiction unless $m' = m$.

Let ρ be less than the distance between any two components of N_1 , and therefore of any N_i . If the lower bound of σ_i is 0, we may assume that every

$\sigma_i < \rho/3$ and that every $\sigma_i > 3\sigma_{i+1}$, since this is true for a partial sequence. If δ is slightly less than $\sigma_i/2$, it is greater than σ_{i+1} . By definition of σ_{i+1} there is a canonical δ -covering of N_{i+1} of cyclic number not greater than $m-1$ and of component number p . Then by §12 there is an η_i such that every canonical η_i -covering of N_i has its cyclic number at least 1 larger than that of the subset of cells containing points of N_{i+1} . Let r be any integer and η be less than any η_i , $i \leq r$.

Let S_1 be an η -covering of N_1 and T_2 be the set of cells of S_1 containing points of N_2 . By §5, Corollary 3, there is a set S_2 of cells which are divisors of N_2 and cells of T_2 , and which forms a canonical η -covering of N_2 of cyclic number not greater than $m(T_2)$. Then by §12,

$$(1) \quad m(S_1) \geq m(T_2) + 1 \geq m(S_2) + 1.$$

Defining T_3 and S_3 in like manner, we have

$$(2) \quad m(S_2) \geq m(S_3) + 1.$$

Finally we reach

$$(r) \quad m(S_r) \geq m(S_{r+1}) + 1.$$

Relations (1), (2), $\dots$, (r) give

$$m(S_1) \geq r.$$

Hence for every integer r , $m(N_1) \geq r$, which is impossible by the hypothesis. Therefore the lower bound of σ_i cannot be zero and $m(N) = m$. Since $N \subseteq M$ and $M \subseteq M_i$ for every i , $m(M) = m$, which was to be proved.

COROLLARY. *Let M be a one-dimensional compact set of finite cyclic number m . Then N contains a set K irreducible with respect to the properties of being closed and of cyclic number m .*

14. THEOREM. *Let A and B be one-dimensional compact sets, each having a finite number of components and a finite cyclic number. Let $A \cdot B$ have a finite number of components. Then*

$$(1) \quad m(A + B) = m(A) + m(B) - m(A \cdot B) + p(A + B) - p(A) - p(B) + p(A \cdot B).$$

There is a σ so small that every canonical σ -enclosure of A , B , $A \cdot B$, or $A + B$ has the same component number and at least as great a cyclic number as the respective set. Let $D = \{d_i\}$ be a canonical $(\sigma/2)$ -covering of $A \cdot B$ of cyclic number $m(A \cdot B)$. Let $\epsilon < \sigma/6$ and so small that §3 is valid for the covering D and the set $A \cdot B$ considered as imbedded in the space $A + B$. Let $\eta < \epsilon$ and so small that §12, Corollary 1, is valid for A and $A \cdot B$, and for B and

$A \cdot B$. Let $E = \{e_k\}$ be a canonical η -covering of A of cyclic number $m(A)$; let E'' denote the set of cells of E which contain points of $A \cdot B$, and E' the others. Similarly define $F = \{f_k\}$, F'' , and F' with respect to B and $A \cdot B$. Then E'' and F'' have the same cyclic and component numbers as $A \cdot B$.

Let Δ'_i be the union of the cells $\{e_k\}$ containing points of d_i but of no d_j , $j > i$. Let Δ''_i be the union of the cells $\{f_k\}$ containing points of d_i but of no d_j , $j > i$, and $\Delta_i = \Delta'_i + \Delta''_i$. Then $G = \{\Delta_i\}$ is a canonical σ -enclosure of $A \cdot B$, $m(G) = m(A \cdot B)$, and $p(G) = p(A \cdot B)$, by the choice of ϵ , η , and σ .

Now $G + E'$ is a canonical σ -enclosure of A , G and E' have no common cells, $p(G + E') = p(A)$, and $m(G + E') \geq m(A)$. Compare the equations

$$\begin{aligned} m(G + E') &= m(G) + m(E') + p(G + E') - p(G) - p(E') + p(G \cdot E'); \\ m(E) &= m(E'') + m(E') + p(E'' + E') - p(E'') - p(E') + p(E'' \cdot E'). \end{aligned}$$

It is seen as in the proof of §12 that $p(G \cdot E') \leq p(E'' \cdot E')$. As the other pairs of corresponding terms on the right are equal, this gives $m(G + E') \leq m(E)$; whence $m(G + E') = m(A)$. Likewise, $G + F'$ is a canonical σ -enclosure of B , $m(G + F') = m(B)$, and $p(G + F') = p(B)$.

Consider the canonical σ -covering $E' + G + F'$ of $A + B$. Since $E' \cdot F' = 0$, we have at once from the above results and §7, Corollary 1,

$$m(E' + G + F') = m(A) + m(B) - m(A \cdot B) + p(A + B) - p(A) - p(B) + p(A \cdot B).$$

This shows that $m(A + B)$ is not greater than the right side of (1).

Now for any covering D of $A + B$ and any sub-set C we let C_D be the set of cells containing points of C . We first note that $A_D \cdot B_D$ consists solely of cells common to A_D and B_D . For, if a cell d of A_D meets a cell d' of B_D and d contains no point of B , $d \subset A$, and d' contains a point of A . Then $d' \subset A_D$ and so $d \cdot d'$ is deleted by our original definition of the divisor of two canonical collections of cells.

For any $\epsilon > 0$ there is a $\delta > 0$ so that for any canonical δ -covering $A \cdot B \subset A_D \cdot B_D \subset V_\epsilon(A \cdot B)$ and no other cell of $(A + B)_D$ contains points of $A \cdot B$. As $A \cdot B$ is a sub-set of A , there is an $\epsilon > 0$ and an $\eta > 0$ for which §12, Corollary 2, is valid.

Also, for some $\sigma > 0$ and less than both δ and η , every canonical σ -covering of $A + B$ is such that $p(A + B)_D = p(A + B)$, $p(A_D) = p(A)$, $p(B_D) = p(B)$, $m(A_D) \geq m(A)$, and $m(B_D) \geq m(B)$, and for some such covering $m(A + B)_D = m(A + B)$. By §7,

$$\begin{aligned} m(A + B)_D &= m(A_D) + m(B_D) - m(A_D \cdot B_D) \\ &\quad + p(A + B)_D - p(A_D) - p(B_D) + p(A_D \cdot B_D). \end{aligned}$$

By §12, Corollary 2, $m(A_D) - m(A_D \cdot B_D) + p(A_D \cdot B_D) \geq m(A) - m(A \cdot B) + p(A \cdot B)$. Hence the above equation shows that $m(A+B)$ is greater than or equal to the right hand side of (1). As we have already shown that it is not greater, the theorem is proved.

15. If we define the index of a set A by $I(A) = m(A) - p(A)$, the above addition formula becomes $I(A+B) = I(A) + I(B) - I(A \cdot B)$, which is the same as the formula of Straszewicz and is readily generalized. For completeness the following rather obvious result is added.

THEOREM. *Let A and B be compact one-dimensional sets, let each have a finite set of components, and let $A \cdot B$ have an infinite set of components. Then $m(A+B)$ is infinite.*

For some component K of A and some component L of B , $K \cdot L$ has an infinite set of components. Then $K \cdot L$ is the sum of r disjoint closed sets for any integer r . Hence for some $\sigma > 0$ every canonical σ -covering $M+N$ of $K+L$, where M denotes the set of cells containing points of K and N those containing points of L , is such that $M \cdot N$ has at least r components. By §7,

$$m(M+N) = m(M) + m(N) - m(M \cdot N) \\ + p(M+N) - p(M) - p(N) + p(M \cdot N).$$

Obviously, $m(M) \geq 0$, $m(N) - m(M \cdot N) \geq 0$, and $p(M+N) = p(M) = p(N) = 1$. Hence $m(M+N) \geq r-1$. As r was any integer, $m(K+L)$ is infinite; and so, a fortiori, $m(A+B)$ is infinite.

YALE UNIVERSITY,
NEW HAVEN, CONN