ON A CANONICAL CONFORMAL MAPPING OF J. L. WALSH

BY
JAMES A. JENKINS

1. In a recent paper [3] Walsh has given a type of canonical conformal mapping for plane domains of finite connectivity which may be regarded as the culmination of results of de la Vallée Poussin and Julia on canonical conformal mappings associated with level curves of polynomials and rational functions. On the one hand Walsh allows a class of transcendental functions replacing the rational functions of the earlier works. On the other the striking features of his paper are, first, the completeness of the representation obtained as compared with the results of de la Vallée Poussin and Julia and, second, the fact that with an appropriate interpretation the canonical mapping for a given domain is unique. It is of considerable interest that Walsh's results can be obtained by use of the method of quadratic differentials, indeed by a suitable development of considerations given in a paper [2] by the author and D. C. Spencer. The exposition of these ideas is the purpose of the present paper.

2. THEOREM 1. *Let D be a domain on the z -sphere with boundary components $B_1, B_2, \dots, B_\mu; C_1, C_2, \dots, C_\nu, \mu\nu \neq 0$. Then there exists a conformal mapping of D onto a domain Δ on the Z -sphere where Δ is defined by*

$$(1) \quad 1 < |T(Z)| < e^{1/\tau}, \quad T(Z) \equiv \frac{A(Z - a_1)^{M_1}(Z - a_2)^{M_2} \dots (Z - a_\mu)^{M_\mu}}{(Z - b_1)^{N_1}(Z - b_2)^{N_2} \dots (Z - b_\nu)^{N_\nu}},$$

$$\sum_{j=1}^{\mu} M_j = \sum_{j=1}^{\nu} N_j = 1, \quad \tau > 0.$$

The exponents M_j and N_j are positive but need not be rational. The locus $|T(Z)| = 1$ consists of μ mutually disjoint Jordan curves, respective images of the B_j , which separate Δ from the a_j ; the locus $|T(Z)| = e^{1/\tau}$ consists of ν mutually disjoint Jordan curves, respective images of the C_j , which separate Δ from the b_j .

In his corresponding statement [3, p. 129] Walsh includes the remark that if the B_j, C_j are Jordan curves the mapping can be extended to a homeomorphism between $\bar{D}$ and $\bar{\Delta}$. As appears in his proof this follows at once from standard results on boundary correspondence when Theorem 1 is established.

In [2, §8] is treated a canonical conformal mapping of a multiply-connected domain obtained from a regular positive quadratic differential given on the domain. (In that paper negative quadratic differentials were used but

Received by the editors November 10, 1956.

this is inessential.) If one uses a quadratic differential derived from the harmonic measure of the set of curves C_j with respect to the domain D it will be found that the mapping obtained is that of Theorem 1. Here we will develop the details from the beginning for the sake of completeness. Actually in the proof of Theorem 1 one need introduce only linear differentials but in a more general context quadratic differentials would intervene.

Let $\omega(z)$ be the harmonic measure of $\bigcup_{j=1}^{\nu} C_j$ with respect to D and let $\bar{\omega}(z)$ be a conjugate of $\omega(z)$. Then $f(z) = \omega(z) + i\bar{\omega}(z)$ will be analytic but not single-valued in D while $f'(z)$ will be single-valued. Let us join the boundary components $B_1, B_2, \dots, B_{\mu}$ to a point P in D by a finite number of analytic arcs such that just one arc ends on each boundary component and the boundary components $C_1, C_2, \dots, C_{\nu}$ to a point P' in D by a finite number of analytic arcs such that just one arc ends on each boundary component in such a way that on deleting these arcs from D we obtain a single doubly-connected domain D' . Finally we join the point P to the point P' by an analytic arc in D' so that on deleting it from D' we obtain a single simply-connected domain D^* . This is mapped by the function $if(z)$ (which is single-valued on D^*) onto a portion of Riemann surface $\mathfrak{R}$ lying, say, over the w -plane. If we identify boundary points of $\mathfrak{R}'$ arising from the same point of D we obtain a conformal model of D . Under the preceding mapping a boundary component B_j goes into a segment over the real w -axis of length say σ'_j . A boundary component C_j similarly goes into a segment lying above the line $\Im w = 1$ of length say τ'_j . By a well known property of the harmonic measure

$$\sum_{j=1}^{\mu} \sigma'_j = \sum_{j=1}^{\nu} \tau'_j = 2\pi\tau$$

where τ is a certain positive number.

Consider now the function $i(2\pi\tau)^{-1}f(z)$. As before it maps D^* onto a portion of Riemann surface $\mathfrak{R}$ lying, say, over the ζ -plane. Let B_j go into the segment Σ_j

$$s_j \leq \zeta \leq s_j + \sigma_j$$

where $\sigma_j = (2\pi\tau)^{-1}\sigma'_j$ and the end points represent the same point on B_j . Let C_j go into the segment T_j

$$\Im \zeta = (2\pi\tau)^{-1}, \quad t_j \leq \Re \zeta \leq t_j + \tau_j$$

where $\tau_j = (2\pi\tau)^{-1}\tau'_j$ and the end points represent the same point on C_j .

Now we adjoin to $\mathfrak{R}$ along the segment $\Sigma_j, j=1, \dots, \mu$, the half strip F_j

$$\Im \zeta < 0, \quad s_j \leq \Re \zeta \leq s_j + \sigma_j.$$

Also we adjoin to $\mathfrak{R}$ along the segment $T_j, j=1, \dots, \nu$, the half strip G_j

$$\Im \zeta > (2\pi\tau)^{-1}, \quad t_j \leq \Re \zeta \leq t_j + \tau_j.$$

From the covering surface so obtained we produce an abstract Riemann surface $\mathfrak{S}$ by

- (i) performing the identifications arising from the mapping $i(2\pi\tau)^{-1}f(z)$,
- (ii) identifying each two boundary points of F_j for which $\Im\zeta$ has a given value, assigning there as local uniformizing parameter the parameter obtained by covering of the ζ -plane, $j=1, \dots, \mu$,
- (iii) identifying each two boundary points of G_j for which $\Im\zeta$ has a given value, assigning there as local uniformizing parameter the parameter obtained by covering of the ζ -plane, $j=1, \dots, \nu$,
- (iv) assigning a point P_j corresponding to the point at infinity on each of F_j , $j=1, \dots, \mu$, at which we assign the local uniformizing parameter $\exp(-2\pi i\sigma_j^{-1}\zeta)$,
- (v) assigning a point Q_j corresponding to the point at infinity on each of G_j , $j=1, \dots, \nu$, at which we assign the local uniformizing parameter $\exp(2\pi i\tau_j^{-1}\zeta)$.

The surface $\mathfrak{S}$ is seen at once to be simply-connected and of elliptic type. Let it be mapped conformally on the Z -sphere. By this mapping to the points P_j , $j=1, \dots, \mu$, correspond points a_j and to the points Q_j , $j=1, \dots, \nu$, correspond points b_j . We may assume all the a_j , b_j finite (although corresponding considerations would apply if one of them were the point at infinity). As a function of Z , ζ is of course not single-valued but $d\zeta$ is. It is thus a linear differential on the Z -sphere and is regular except at the points a_j , $j=1, \dots, \mu$ and b_j , $j=1, \dots, \nu$. At a_j we have the expansion

$$d\zeta = \left(\frac{i\sigma_j}{2\pi} \frac{1}{Z - a_j} + \text{regular terms} \right) dZ$$

and at b_j the expansion

$$d\zeta = \left(\frac{-i\tau_j}{2\pi} \frac{1}{Z - b_j} + \text{regular terms} \right) dZ.$$

Thus

$$d\zeta = \frac{i}{2\pi} \left\{ \sum_{j=1}^{\mu} \frac{\sigma_j}{Z - a_j} - \sum_{j=1}^{\nu} \frac{\tau_j}{Z - b_j} \right\} dZ.$$

(Note that $d\zeta$ is regular at $Z = \infty$.) Moreover

$$\zeta = \frac{i}{2\pi} \left\{ \sum_{j=1}^{\mu} \sigma_j \log(Z - a_j) - \sum_{j=1}^{\nu} \tau_j \log(Z - b_j) \right\} + K$$

where K is constant. Take now

$$T(Z) = \exp(-2\pi i\zeta)$$

and set $\sigma_j = M_j$, $\tau_j = N_j$, $\exp(-2\pi iK) = A$. Then

$$T(Z) \equiv \frac{A(Z - a_1)^{M_1}(Z - a_2)^{M_2} \cdots (Z - a_\mu)^{M_\mu}}{(Z - b_1)^{N_1}(Z - b_2)^{N_2} \cdots (Z - b_\nu)^{N_\nu}}$$

with $\sum_{j=1}^\mu M_j = \sum_{j=1}^\nu N_j = 1$. On the other hand, on the curves B_j , $j=1, \dots, \mu$, we have $\Im \zeta = 0$; on the curves C_j , $j=1, \dots, \nu$, we have $\Im \zeta = i(2\pi\tau)^{-1}$. Thus on the images of the first set of curves $|T(Z)| = 1$ and on the images of the second set of curves $|T(Z)| = \exp(1/\tau)$. It is seen at once that these are the only points at which these equalities hold and that the points of D go into the points for which

$$1 < |T(Z)| < \exp(1/\tau).$$

This completes the proof of Theorem 1.

3. We give now a proof of the uniqueness result corresponding to Theorem 1 [3, p. 137].

THEOREM 2. *Let D be a domain on the z -sphere defined by the inequalities*

$$1 < |R(z)| < e^{1/\tau}, \quad R(z) \equiv \frac{A(z - a_1)^{m_1} \cdots (z - a_\mu)^{m_\mu}}{(z - b_1)^{n_1} \cdots (z - b_\nu)^{n_\nu}},$$

$$\sum_{j=1}^\mu m_j = \sum_{j=1}^\nu n_j = 1,$$

and whose boundary consists of mutually disjoint Jordan curves $B_1, B_2, \dots, B_\mu, C'_1, C'_2, \dots, C'_\nu$, where B_j separates a_j from D and C_j separates b_j from D . Let Δ be a domain on the Z -sphere defined by the inequalities

$$1 < |R_1(Z)| < e^{1/\tau'}, \quad R_1(Z) \equiv \frac{A'(Z - a'_1)^{m'_1} \cdots (Z - a'_\mu)^{m'_\mu}}{(Z - b'_1)^{n'_1} \cdots (Z - b'_\nu)^{n'_\nu}},$$

$$\sum_{j=1}^\mu m'_j = \sum_{j=1}^\nu n'_j = 1$$

and whose boundary consists of mutually disjoint Jordan curves $B'_1, B'_2, \dots, B'_\mu, C'_1, C'_2, \dots, C'_\nu$, where B'_j separates a'_j from Δ and C'_j separates b'_j from Δ . If there exists a $(1, 1)$ conformal mapping of D onto Δ so that each B_j corresponds to B'_j , $j=1, \dots, \mu$, and each C_j corresponds to C'_j , $j=1, \dots, \nu$, then the mapping is a linear transformation of the complex variable z , defined throughout the z - and Z -spheres. We have $\tau = \tau'$, $m_j = m'_j$, $n_j = n'_j$. If $\arg A'$ is suitably chosen we have $R(z) \equiv R_1(Z)$.

Let $Z = g(z)$ be the conformal mapping of Theorem 2. Define

$$u(z) = \sum_{j=1}^\mu m_j \log |z - a_j| - \sum_{j=1}^\nu n_j \log |z - b_j| + \log |A|,$$

$$U(Z) = \sum_{j=1}^\mu m'_j \log |Z - a'_j| - \sum_{j=1}^\nu n'_j \log |Z - b'_j| + \log |A'|.$$

The function $u(z)$ is single-valued and harmonic in D , takes the value 0 on the curves $B_j, j=1, \dots, \mu$ and the value $1/\tau$ on the curves $C_j, j=1, \dots, \nu$. It is thus equal to $1/\tau$ times the harmonic measure of the set of curves $C_j, j=1, \dots, \nu$ with respect to D . Similarly $U(Z)$ is $1/\tau'$ times the harmonic measure of the set of curves $C'_j, j=1, \dots, \nu$, with respect to Δ . Thus $\tau u(z) = \tau' U(g(z))$. Now $2\pi m_j$ is the period of a conjugate of $u(z)$ about B_j , $-2\pi n_j$ the period of a conjugate of $u(z)$ about C_j while $2\pi m'_j, -2\pi n'_j$ are the corresponding periods for $U(Z)$. Thus

$$\begin{aligned} \tau m_j &= \tau' m'_j, & j &= 1, \dots, \mu, \\ \tau n_j &= \tau' n'_j, & j &= 1, \dots, \nu. \end{aligned}$$

Adding the first set of equations we obtain

$$\tau = \tau'.$$

This means

$$\begin{aligned} m_j &= m'_j, & j &= 1, \dots, \mu, \\ n_j &= n'_j, & j &= 1, \dots, \nu, \end{aligned}$$

while

$$(2) \quad u(z) = U(g(z)).$$

Setting now

$$\begin{aligned} d\zeta &= \left(\sum_{j=1}^{\mu} \frac{m_j}{z - a_j} - \sum_{j=1}^{\nu} \frac{n_j}{z - b_j} \right) dz, \\ d\zeta' &= \left(\sum_{j=1}^{\mu} \frac{m_j}{Z - a'_j} - \sum_{j=1}^{\nu} \frac{n_j}{Z - b'_j} \right) dZ \end{aligned}$$

we have at once that $d\zeta$ and $d\zeta'$ are corresponding differentials under the mapping g . Defining on D and Δ the respective metrics $|d\zeta|$ and $|d\zeta'|$ the mapping g becomes isometric.

Let now Γ and Γ' be a pair of corresponding boundary components of D and Δ ; let c and c' be the distinguished points they separate from D and Δ . Evidently Γ and Γ' are closed trajectories respectively of the quadratic differentials $-d\zeta^2$ and $-(d\zeta')^2$. Let O and Ω be the domains bounded respectively by Γ and Γ' in the complements of D and Δ . Since $O(\Omega)$ contains no pole of $-d\zeta^2(-d\zeta')^2$ other than $c(c')$, by the Basic Structure Theorem [1] $O(\Omega)$ is a subdomain of a circle domain associated with $-d\zeta^2(-d\zeta')^2$. Consider z on Γ and the corresponding point $g(z)$ on Γ' . The portion of orthogonal trajectory of $-d\zeta^2(-d\zeta')^2$ entering $\overline{O}(\overline{\Omega})$ at $z(g(z))$ is a half-open arc $\Lambda_z(\Lambda'_z)$ with limiting end point at $c(c')$. We extend the mapping g to the set O by mapping the point of Λ_z at a given distance from z in the metric $|d\zeta|$ onto the point of Λ'_z at the same distance from $g(z)$ in the metric $|d\zeta'|$ and map c

onto c' . This clearly gives a $(1, 1)$ mapping of O onto Ω . At all points of $\bar{O}$ but c the mapping is isometric for the metrics $|d\zeta|$, $|d\zeta'|$ thus conformal. Finally by the Riemann theorem on removable singularities the mapping is conformal at c . Carrying out this construction for all corresponding pairs of domains complementary to D and Δ we extend g to a conformal mapping of the z -sphere onto the Z -sphere which is thus a linear transformation.

Finally by Equation (2) we have

$$|R(z)| = |R_1(g(z))|.$$

Thus for a suitable choice of $\arg A'$ we have

$$R(z) \equiv R_1(g(z)).$$

This completes the proof of Theorem 2.

3. Walsh gave also an analogue to Theorem 1 when the boundary components B_j , $j=1, \dots, \mu$, are allowed to shrink to points [3, p. 140].

THEOREM 3. *Let D be a domain on the z -sphere with boundary components $C_1, C_2, \dots, C_\nu$, let $\alpha_1, \alpha_2, \dots, \alpha_\mu$ be arbitrary distinct points of D and let $M_1, M_2, \dots, M_\mu$ be arbitrary positive numbers which satisfy $\sum_{j=1}^\mu M_j = 1$. Then there exists a conformal mapping of D onto a domain Δ on the Z -sphere where Δ is defined by*

$$|T(Z)| < 1, \quad T(Z) \equiv \frac{A(Z - a_1)^{M_1}(Z - a_2)^{M_2} \dots (Z - a_\mu)^{M_\mu}}{(Z - b_1)^{N_1}(Z - b_2)^{N_2} \dots (Z - b_\nu)^{N_\nu}},$$

$$N_j > 0, \quad \sum_{j=1}^\nu N_j = 1.$$

The a_j are the respective images of the α_j , the locus $|T(Z)| = 1$ consists of ν mutually disjoint analytic Jordan curves, which are respective images of the C_j and which separate Δ from the b_j .

A remark on boundary correspondence holds as for Theorem 1.

Theorem 3 is proved just as Theorem 1 except the harmonic measure is replaced by the function

$$M_1 g_1(z) + M_2 g_2(z) + \dots + M_\mu g_\mu(z)$$

where $g_j(z)$ is the Green's function of D with pole at α_j , $j=1, \dots, \mu$.

4. The uniqueness result corresponding to Theorem 3 is [3, p. 143]

THEOREM 4. *Let D be a domain on the z -sphere defined by the inequality*

$$|R(z)| < 1, \quad R(z) \equiv \frac{A(z - a_1)^{m_1}(z - a_2)^{m_2} \dots (z - a_\mu)^{m_\mu}}{(z - b_1)^{n_1}(z - b_2)^{n_2} \dots (z - b_\nu)^{n_\nu}},$$

$$m_j > 0, \quad \sum_{j=1}^\mu m_j = \sum_{j=1}^\nu n_j = 1,$$

and whose boundary consists of mutually disjoint Jordan curves $C_1, C_2, \dots, C_\nu$ which separate the b_j respectively from D . Let Δ be a domain on the Z -sphere defined by the inequality

$$|R_1(Z)| < 1, \quad R_1(Z) \equiv \frac{A'(Z - a'_1)^{m_1}(Z - a'_2)^{m_2} \cdots (Z - a'_\mu)^{m_\mu}}{(Z - b'_1)^{n_1'}(Z - b'_2)^{n_2'} \cdots (Z - b'_\nu)^{n_\nu'}},$$

$$\sum_{j=1}^{\nu} n_j' = 1,$$

and whose boundary consists of mutually disjoint Jordan curves $C'_1, C'_2, \dots, C'_\nu$ which separate the b'_j respectively from Δ . If there exists a $(1, 1)$ conformal mapping of D onto Δ so that the a_j correspond respectively to the a'_j , then this correspondence can be continued beyond D and Δ as a linear transformation of the z -sphere onto the Z -sphere. We have $n_j = n_j'$, and if $\arg A'$ is suitably chosen, $R(z) \equiv R_1(Z)$.

There is no essential difference in the proof of Theorem 4 from that of Theorem 2.

BIBLIOGRAPHY

1. James A. Jenkins, *A general coefficient theorem*, Trans. Amer. Math. Soc. vol. 77 (1954) pp. 262–280.
2. James A. Jenkins and D. C. Spencer, *Hyperelliptic trajectories*, Ann. of Math. vol. 53 (1951) pp. 4–35.
3. J. L. Walsh, *On the conformal mapping of multiply connected regions*, Trans. Amer. Math. Soc. vol. 82 (1956) pp. 128–146.

UNIVERSITY OF NOTRE DAME,
NOTRE DAME, IND.