NODAL NONCOMMUTATIVE JORDAN ALGEBRAS AND SIMPLE LIE ALGEBRAS OF CHARACTERISTIC p

BY

R. D. SCHAFFER⁽¹⁾

A noncommutative Jordan algebra A over a field F is a nodal algebra [9]⁽²⁾ in case every element of A may be written in the form $\alpha 1 + z$ where α is in F , 1 is the unity element of A , and z is nilpotent, while the set N of nilpotent elements is not a subalgebra of A . F is necessarily of characteristic $p > 0$.

L. A. Kokoris gave in [5] the first examples of simple nodal noncommutative Jordan algebras. Generalizing these, he has constructed in [7] a class K of nodal noncommutative Jordan algebras A of dimension p^n , and has proved [6; 7] that each simple nodal noncommutative Jordan algebra of characteristic $p \neq 2$ is in K . Although not all of the algebras in K are simple, it turns out that those algebras with which we are principally concerned here are indeed simple.

Any algebra A of dimension p^n in K may be represented as follows: let $B_n = F[x_1, \dots, x_n]$, $x_i^p = 0$, be a truncated polynomial ring with partial differentiation operators $\partial/\partial x_i$. Write $f \cdot g$ for the commutative associative product in B_n . Then A is the same vector space as B_n , but multiplication in A is defined by

$$(1) \quad fg = f \cdot g + \sum_{i,j=1}^n \frac{\partial f}{\partial x_i} \cdot \frac{\partial g}{\partial x_j} \cdot c_{ij}, \quad c_{ij} = -c_{ji},$$

where the c_{ij} ($= -c_{ji}$) are elements in B_n which are arbitrary except for the proviso that at least one of them has an inverse (equivalently, at least one of them is not in the radical N of B_n). Then necessarily $n \geq 2$. Also we note that

$$(2) \quad c_{ij} = \frac{1}{2} [x_i, x_j], \quad i, j = 1, 2, \dots, n.$$

This representation of an algebra A in K is not unique. It is well-known [4, p. 108; or 8, §4.5, Proposition 6] that any representatives in B_n of the elements of any basis of the n -dimensional space $N/N \cdot N$ will serve for the

---

Presented to the Society, January 20, 1959; received by the editors February 13, 1959.

⁽¹⁾ National Science Foundation Senior Postdoctoral Fellow.

⁽²⁾ Numbers in brackets refer to the references cited at the end of the paper.

x_i ($i=1, \dots, n$). By the chain rule $\partial f/\partial x_i = \sum_{k=1}^n (\partial f/\partial y_k) \cdot (\partial y_k/\partial x_i)$, one obtains

$$fg = f \cdot g + \sum_{k,l=1}^n \frac{\partial f}{\partial y_k} \cdot \frac{\partial g}{\partial y_l} \cdot d_{kl}$$

with

$$(3) \quad d_{kl} = \frac{1}{2} [y_k, y_l] = \sum_{i,j=1}^n \frac{\partial y_k}{\partial x_i} \cdot \frac{\partial y_l}{\partial x_j} \cdot c_{ij},$$

where $y_1, \dots, y_n$ are any elements of B_n whose residue classes form a basis for $N/N \cdot N$ over F .

In this paper we study the derivations of the algebras in K . We are led to relationships between some of these algebras and recently discovered simple Lie algebras of characteristic p . In particular, we obtain an intrinsic characterization of the simple Lie algebras V_r of A. A. Albert and M. S. Frank [1]. Also we display each of the simple Lie algebras $L(G, \delta, f)$ of characteristic $\neq 2$ defined by Richard Block [2] as an ideal in the derivation algebra of a suitably chosen simple nodal noncommutative Jordan algebra A . We assume characteristic $p > 2$ throughout.

1. Derivations. Since $c_{ij} = -c_{ji}$ in (1), we have $(fg + gf)/2 = f \cdot g$. That is, the commutative algebra A^+ attached to A is B_n itself.

Clearly any derivation of A is also a derivation of A^+ . But the derivations of B_n are well-known [4, p. 107]:

$$f \rightarrow \sum_{k=1}^n \frac{\partial f}{\partial x_k} \cdot a_k$$

for arbitrary elements a_k in B_n . Hence, if D is a derivation of A , we have

$$(4) \quad fD = \sum_{k=1}^n \frac{\partial f}{\partial x_k} \cdot a_k$$

for all f in A and for certain elements $a_k (= x_k D)$ in A ($k=1, \dots, n$). Occasionally we shall employ the notation $D = (a_1, \dots, a_n)$ used in [3] and [1].

Suppose that D is given by (4). Then D is a derivation of A^+ , and also

$$\begin{aligned} \frac{\partial h}{\partial x_i} D - \frac{\partial(hD)}{\partial x_i} &= \sum_k \left(\frac{\partial^2 h}{\partial x_k \partial x_i} \cdot a_k - \frac{\partial^2 h}{\partial x_i \partial x_k} \cdot a_k - \frac{\partial h}{\partial x_k} \cdot \frac{\partial a_k}{\partial x_i} \right) \\ &= - \sum_k \frac{\partial h}{\partial x_k} \cdot \frac{\partial a_k}{\partial x_i} \end{aligned}$$

for any h in A . Hence

$$\begin{aligned}
(fg)D - (fD)g - f(gD) &= (f \cdot g)D - (fD) \cdot g - f \cdot (gD) \\
&+ \sum_{i,j} \left\{ \left(\frac{\partial f}{\partial x_i} \cdot \frac{\partial g}{\partial x_j} \cdot c_{ij} \right) D - \frac{\partial(fD)}{\partial x_i} \cdot \frac{\partial g}{\partial x_j} \cdot c_{ij} - \frac{\partial f}{\partial x_i} \cdot \frac{\partial(gD)}{\partial x_j} \cdot c_{ij} \right\} \\
&= \sum_{i,j} \left\{ \frac{\partial f}{\partial x_i} D \cdot \frac{\partial g}{\partial x_j} \cdot c_{ij} + \frac{\partial f}{\partial x_i} \cdot \frac{\partial g}{\partial x_j} D \cdot c_{ij} + \frac{\partial f}{\partial x_i} \cdot \frac{\partial g}{\partial x_j} \cdot c_{ij} D \right. \\
&\quad \left. - \frac{\partial(fD)}{\partial x_i} \cdot \frac{\partial g}{\partial x_j} \cdot c_{ij} - \frac{\partial f}{\partial x_i} \cdot \frac{\partial(gD)}{\partial x_j} \cdot c_{ij} \right\} \\
&= \sum_{i,j,k} \left(-\frac{\partial f}{\partial x_k} \cdot \frac{\partial a_k}{\partial x_i} \cdot \frac{\partial g}{\partial x_j} \cdot c_{ij} - \frac{\partial f}{\partial x_i} \cdot \frac{\partial g}{\partial x_k} \cdot \frac{\partial a_k}{\partial x_j} \cdot c_{ij} + \frac{\partial f}{\partial x_i} \cdot \frac{\partial g}{\partial x_j} \cdot \frac{\partial c_{ij}}{\partial x_k} \cdot a_k \right) \\
&= \sum_{i,j} \frac{\partial f}{\partial x_i} \cdot \frac{\partial g}{\partial x_j} \cdot \left\{ \sum_k \left(\frac{\partial c_{ij}}{\partial x_k} \cdot a_k - \frac{\partial a_i}{\partial x_k} \cdot c_{kj} - \frac{\partial a_j}{\partial x_k} \cdot c_{ik} \right) \right\}.
\end{aligned}$$

Hence D is a derivation of A if and only if

$$(5) \quad \sum_{k=1}^n \left(\frac{\partial c_{ij}}{\partial x_k} \cdot a_k + \frac{\partial a_i}{\partial x_k} \cdot c_{jk} + \frac{\partial a_j}{\partial x_k} \cdot c_{ki} \right) = 0$$

for $1 \leq i, j \leq n$. The equations (5) are redundant for $j \geq i$, so we have proved

THEOREM 1. *Let A be a nodal noncommutative Jordan algebra in K , multiplication being defined by (1). Then a mapping D on A is a derivation of A if and only if D has the form (4) for elements $a_1, \dots, a_n$ in A satisfying (5) for $1 \leq i < j \leq n$.*

Solution of the equations (5) in general seems a formidable task. Even a preliminary simplification of the problem by using (3) to normalize the c_{ij} seems exceedingly complex in the general situation. In the next two sections we treat two special cases: (i) n arbitrary, but all of the c_{ij} in $F1$; (ii) $n=2$, but c_{ij} arbitrary in A .

2. Algebras defined by skew-symmetric bilinear forms. Suppose that all of the c_{ij} are in $F1$. That is, $c_{ij} = \phi_{ij}1$, $\phi_{ij} = -\phi_{ji}$ in F , not all ϕ_{ij} zero, where $x_i x_j = x_i \cdot x_j + \phi_{ij}1$. There is a unique skew-symmetric bilinear form ϕ defined on the n -dimensional space $M = \sum_{i=1}^n Fx_i$ (equivalently, defined on $N/N \cdot N$) such that $\phi(x_i, x_j) = \phi_{ij}$. Then

$$fg = f \cdot g + \sum_{i,j} \frac{\partial f}{\partial x_i} \cdot \frac{\partial g}{\partial x_j} \cdot \phi(x_i, x_j)1.$$

If x_i ($i=1, \dots, n$) exist for which $c_{ij} = \phi(x_i, x_j)1$, we shall say that A is defined by the skew-symmetric bilinear form ϕ . (Since $N(N \cdot N) \subseteq N$ by [9, Theorem 5] we know that, if A is defined by different skew-symmetric bilinear forms, the forms are equivalent.) Let $2r$ ($2 \leq 2r \leq n$) be the rank of ϕ .

A change of basis in M gives $\phi(x_i, x_{r+i}) = 1 = -\phi(x_{r+i}, x_i)$ for $i = 1, \dots, r$; $\phi(x_i, x_j) = 0$ otherwise. The form ϕ is nondegenerate if and only if $n = 2r$.

We remark that an algebra A defined by a skew-symmetric bilinear form ϕ is simple if and only if ϕ is nondegenerate. For take the special basis above in M . If $n > 2r$, then $c_{i, 2r+1} = \phi_{i, 2r+1} 1 = 0$ for $i = 1, \dots, n$, so that $fx_{2r+1} = f \cdot x_{2r+1} = x_{2r+1}f$ for every f in A by (1). It follows that $C = x_{2r+1} \cdot A$ is an ideal of A , $C \neq 0$, $C \neq A$. If $n = 2r$, we shall see in §4 that A is a simple nodal noncommutative Jordan algebra associated in a specific way with one of the algebras in a general class of simple Lie algebras of characteristic p . Of course the simplicity of A may also be established directly in this particular case.

If we take the special basis above for M in an algebra A defined by a skew-symmetric form, equations (5) become

$$(6) \quad \frac{\partial a_i}{\partial x_{r+j}} = \frac{\partial a_j}{\partial x_{r+i}}, \quad i, j = 1, \dots, r;$$

$$(7) \quad \frac{\partial a_{r+i}}{\partial x_j} = \frac{\partial a_{r+j}}{\partial x_i}, \quad i, j = 1, \dots, r;$$

$$(8) \quad \frac{\partial a_i}{\partial x_j} + \frac{\partial a_{r+j}}{\partial x_{r+i}} = 0, \quad i, j = 1, \dots, r;$$

and

$$(9) \quad \frac{\partial a_j}{\partial x_i} = 0, \quad 1 \leq i \leq 2r; \quad 2r+1 \leq j \leq n.$$

Equations (6), (7), (8) involve only $a_1, \dots, a_{2r}$. Hence the elements a_{2r+k} ($k = 1, \dots, n - 2r$) are arbitrary in $F[x_{2r+1}, \dots, x_n]$ by (9).

LEMMA 1. *Let f be in $B_n = F[x_1, \dots, x_n]$. For any i ($1 \leq i \leq n$), write $R_i = F[x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n]$ so that $B_n = R_i[x_i]$. Then there exists g in B_n satisfying*

$$(10) \quad \frac{\partial g}{\partial x_i} = f$$

if and only if f is of the form

$$(11) \quad f = \beta_0 + \beta_1 \cdot x_i + \beta_2 \cdot x_i^2 + \dots + \beta_{p-2} \cdot x_i^{p-2}, \quad \beta_j \in R_i.$$

If f satisfies (11), there is always a solution g of (10) having the form

$$(12) \quad g = x_i \cdot h, \quad h \in B_n.$$

If g is one solution of (10), then g^* is a solution if and only if

$$(13) \quad g^* = g + h, \quad h \in R_i.$$

The proof is straightforward.

Using the lemma, we give an inductive proof that the solutions of (6)–(9) are

$$(14) \quad \begin{aligned} a_i &= \frac{\partial g}{\partial x_{r+i}} + \sigma_i \cdot x_{r+i}^{p-1} & \text{for } i = 1, \dots, r, \\ a_{r+i} &= -\frac{\partial g}{\partial x_i} + \sigma_{r+i} \cdot x_i^{p-1} & \text{for } i = 1, \dots, r, \end{aligned}$$

and

$$(15) \quad a_{2r+k} = \sigma_{2r+k} \quad \text{for } k = 1, \dots, n - 2r,$$

for arbitrary g in A and arbitrary $\sigma_i, \sigma_{r+i}, \sigma_{2r+k}$ in $F[x_{2r+1}, \dots, x_n]$. Clearly any elements $a_1, \dots, a_n$ given by (14) and (15) satisfy equations (6)–(9). We are concerned only with establishing the fact that, if $a_1, \dots, a_{2r}$ satisfy (6)–(8), then $a_1, \dots, a_{2r}$ have the form given in (14).

Let $a_1 = \beta_0 + \beta_1 \cdot x_{r+1} + \dots + \beta_{p-2} \cdot x_{r+1}^{p-2} + \beta_{p-1} \cdot x_{r+1}^{p-1}$, $\beta_j \in R_{r+1}$, so that, for any $k \neq r+1$,

$$\frac{\partial a_1}{\partial x_k} = \frac{\partial \beta_0}{\partial x_k} + \dots + \frac{\partial \beta_{p-1}}{\partial x_k} \cdot x_{r+1}^{p-1}.$$

Then, putting $i=1$ in (6) and (8), we have

$$\frac{\partial \beta_{p-1}}{\partial x_k} = 0 \quad \text{for } k = 1, \dots, r, r+2, \dots, 2r$$

by (11). Also $\partial \beta_{p-1} / \partial x_{r+1} = 0$. By the lemma there exist $g \in B_n$ and $\sigma_1 \in F[x_{2r+1}, \dots, x_n]$ such that $a_1 = \partial g / \partial x_{r+1} + \sigma_1 \cdot x_{r+1}^{p-1}$. For any t ($2 \leq t \leq 2r$) we assume that (14) holds for $i=1, \dots, t-1$. There are two cases to be considered: $t \leq r$, and $t \geq r+1$. If $t \leq r$, we replace i in (6) by t , and obtain

$$\frac{\partial a_t}{\partial x_{r+j}} = \frac{\partial^2 g}{\partial x_{r+t} \partial x_{r+j}} \quad \text{for } j = 1, \dots, t-1,$$

and

$$\frac{\partial a_t}{\partial x_{r+j}} = \frac{\partial a_j}{\partial x_{r+t}} \quad \text{for } j = t+1, \dots, r.$$

It follows that $a_t = \partial g / \partial x_{r+t} + f$, where

$$f = \frac{\partial h}{\partial x_{r+t}} + \sigma_t \cdot x_{r+t}^{p-1}$$

for some h satisfying $\partial h / \partial x_{r+j} = 0$ for $j = 1, \dots, t-1$, and some σ_t satisfying

$$(16) \quad \frac{\partial \sigma_t}{\partial x_k} = 0$$

for $k = r+1, \dots, 2r$. Write $g^* = g + h$. Then $\partial g^*/\partial x_{r+i} = \partial g/\partial x_{r+i}$ for $i=1, \dots, t-1$, so that the inductive hypothesis remains valid with g^* replacing g . Also $a_t = \partial g^*/\partial x_{r+t} + \sigma_t \cdot x_{r+t}^{p-1}$ for σ_t satisfying (16) for $k=r+1, \dots, 2r$. Putting $i=t$ in (8), we have

$$\frac{\partial \sigma_t}{\partial x_j} \cdot x_{r+t}^{p-1} = - \frac{\partial}{\partial x_{r+t}} \left(a_{r+j} + \frac{\partial g^*}{\partial x_j} \right),$$

so (16) holds also for $k=1, \dots, r$. That is, $\sigma_t \in F[x_{2r+1}, \dots, x_n]$, as desired. In the second case ($t \geq r+1$), we write $t=r+s$. Then the assumption of the induction is that the first line of (14) holds for $i=1, \dots, r$, and the second line for $i=1, \dots, s-1$. Putting $j=s$ in (8), we have $a_{r+s} = -\partial g/\partial x_s + f$, where

$$(17) \quad \frac{\partial f}{\partial x_k} = 0$$

for $k=r+1, \dots, 2r$. Also $j=s$ in (7) yields (17) for $k=1, \dots, s-1$, and

$$\frac{\partial f}{\partial x_i} = \frac{\partial}{\partial x_s} \left(a_{r+i} + \frac{\partial g}{\partial x_i} \right)$$

for $i=s+1, \dots, r$. Hence $f = -\partial h/\partial x_s + \sigma_{r+s} \cdot x_s^{p-1}$ for some h satisfying $\partial h/\partial x_k = 0$ for $k=1, \dots, s-1, r+1, \dots, 2r$, and some σ_{r+s} satisfying

$$(18) \quad \frac{\partial \sigma_{r+s}}{\partial x_k} = 0 \quad \text{for } k = 1, \dots, 2r.$$

Write $g^* = g + h$. Then $\partial g^*/\partial x_k = \partial g/\partial x_k$ for $k=1, \dots, s-1, r+1, \dots, 2r$, so that the inductive hypothesis remains valid with g^* replacing g . Also $a_{r+s} = -\partial g^*/\partial x_s + \sigma_{r+s} \cdot x_s^{p-1}$ with $\sigma_{r+s} \in F[x_{2r+1}, \dots, x_n]$ by (18). We have established equations (14).

Now any D given by (4), (14), and (15) determines g modulo $F[x_{2r+1}, \dots, x_n]$ and the σ_j ($j=1, \dots, n$) uniquely. Hence we have proved

THEOREM 2. *Let A be a nodal noncommutative Jordan algebra of dimension p^n which is defined by a skew-symmetric bilinear form ϕ of rank $2r$. Then the derivation algebra $D(A)$ of A has dimension*

$$p^n - p^{n-2r} + n p^{n-2r} = p^{n-2r}(p^{2r} - 1 + n).$$

The x_i in A may be chosen so that the derivations D of A have the form (4) with a_k as in (14) and (15).

Let D be given by (4) and E by

$$(19) \quad fE = \sum_{k=1}^n \frac{\partial f}{\partial x_k} \cdot b_k$$

for b_k in A . Then it is well-known [4, p. 107] that $[D, E] = C = (c_1, \dots, c_n)$ where

$$(20) \quad c_i = \sum_{j=1}^n \left(\frac{\partial a_i}{\partial x_j} \cdot b_j - \frac{\partial b_i}{\partial x_j} \cdot a_j \right), \quad i = 1, \dots, n.$$

It is easily seen that, in case $n > 2r$ above, there are many ideals in $D(A)$. If $n = 2r$, however, (that is, if A is defined by a nondegenerate ϕ) we have (14) with $\sigma_k \in F1 (= F)$ for $k = 1, \dots, 2r$. Writing $D(g; \sigma_1, \dots, \sigma_{2r})$ for D in (14), we obtain

$$(21) \quad [D(f; \rho_1, \dots, \rho_{2r}), D(g; \sigma_1, \dots, \sigma_{2r})] = D(h; 0, \dots, 0)$$

from (20), where

$$(22) \quad h = \sum_{j=1}^r \left\{ (\rho_j \sigma_{r+j} - \rho_{r+j} \sigma_j) x_j^{p-1} \cdot x_{r+j}^{p-1} + \left(\frac{\partial f}{\partial x_j} \cdot \frac{\partial g}{\partial x_{r+j}} - \frac{\partial f}{\partial x_{r+j}} \cdot \frac{\partial g}{\partial x_j} \right) \right. \\ \left. + \left(\sigma_{r+j} \frac{\partial f}{\partial x_{r+j}} - \rho_{r+j} \frac{\partial g}{\partial x_{r+j}} \right) \cdot x_j^{p-1} + \left(\sigma_j \frac{\partial f}{\partial x_j} - \rho_j \frac{\partial g}{\partial x_j} \right) \cdot x_{r+j}^{p-1} \right\}.$$

We recognize $D(g) = D(g; 0, \dots, 0)$ as any element of the algebra V_{0r} of Albert and Frank [1, p. 127]. Let $\tilde{A}$ be the subspace of B_n consisting of all elements of B_n for which the coefficient of $x^{p-1} \cdot x_2^{p-1} \cdot \dots \cdot x_n^{p-1}$ is zero. The $(p^{2r} - 2)$ -dimensional simple Lie algebra V_r of Albert and Frank consists of all $D(g)$ with $g \in \tilde{A}$. It is known [1, pp. 127, 128] that V_r is the derived algebra V'_{0r} of V_{0r} and that $\partial f / \partial x_j \cdot \partial g / \partial x_{r+j} - \partial f / \partial x_{r+j} \cdot \partial g / \partial x_j$ is in $\tilde{A}$. It follows from (11) that $\partial f / \partial x_{r+j} \cdot x_j^{p-1} \in \tilde{A}$, etc. Hence only the first term within the braces in (22) could fail to be in $\tilde{A}$. If $r > 1$, h in (22) is in $\tilde{A}$, and $D(h) \in V_r$. Hence $V_r = V'_{0r} \subseteq D(A)' \subseteq V_r$ if $r > 1$. If $r = 1$, there exists h in (22) which is not in $\tilde{A}$, so that $D(A)' \subseteq V_{01}$ but $D(A)' \neq V_1$. However, $V_1 = V'_1 \subseteq D(A)'' \subseteq V'_{01} = V_1$. Of course $D(A)'' = V_r$ for $r > 1$ also.

Thus we obtain the following intrinsic characterization of the $(p^{2r} - 2)$ -dimensional simple Lie algebras V_r of Albert and Frank:

THEOREM 3. *A Lie algebra L is a simple algebra V_r if and only if there is a p^{2r} -dimensional (simple) nodal noncommutative Jordan algebra A defined by a nondegenerate skew-symmetric bilinear form such that $L \cong D(A)'$ in case $r > 1$, and $L \cong D(A)''$ in case $r = 1$ (actually $L \cong D(A)''$ in both cases).*

A second characterization of the algebras V_r results from the following observation concerning Lie-admissible algebras.

In any nonassociative algebra A , $xy = x \cdot y + [x, y]/2$. Therefore a linear transformation D on A is a derivation if and only if D is a derivation of both A^+ and A^- , where A^- is the anticommutative algebra attached to A having the product $[x, y]$. That is,

$$(23) \quad D(A) = D(A^+) \cap D(A^-).$$

Now in any nodal noncommutative Jordan algebra A in K

$$(24) \quad [f, g] = \sum_{i,j} \frac{\partial f}{\partial x_i} \cdot \frac{\partial g}{\partial x_j} \cdot 2c_{ij},$$

and D in (4) is a derivation of A if and only if D is a derivation of A^- .

If A^- is a Lie algebra, then the mappings $\text{ad } g/2$ defined by

$$(25) \quad f \rightarrow \frac{1}{2} [f, g] = \sum_i \frac{\partial f}{\partial x_i} \cdot \left(\sum_j \frac{\partial g}{\partial x_j} \cdot c_{ij} \right)$$

are inner derivations of A^- , and therefore derivations of A since they are of the form (4) with

$$(26) \quad a_i = \sum_{j=1}^n \frac{\partial g}{\partial x_j} \cdot c_{ij}, \quad i = 1, \dots, n.$$

The set $\text{ad } A$ of all inner derivations (25) of A^- , being an ideal of $D(A^-)$, is an ideal of $D(A)$ by (23).

We use the Jacobi identity

$$\begin{aligned} 0 &= [[f, g], h] + [[g, h], f] + [[h, f], g] \\ &= 2 \sum_{i,j} \left\{ \left[\frac{\partial f}{\partial x_i} \cdot \frac{\partial g}{\partial x_j} \cdot c_{ij}, h \right] + \left[\frac{\partial g}{\partial x_i} \cdot \frac{\partial h}{\partial x_j} \cdot c_{ij}, f \right] + \left[\frac{\partial h}{\partial x_i} \cdot \frac{\partial f}{\partial x_j} \cdot c_{ij}, g \right] \right\} \\ &= 4 \sum_{i,j,k,t} \left\{ \frac{\partial}{\partial x_t} \left(\frac{\partial f}{\partial x_i} \cdot \frac{\partial g}{\partial x_j} \cdot c_{ij} \right) \cdot \frac{\partial h}{\partial x_k} \cdot c_{tk} \right. \\ &\quad \left. + \frac{\partial}{\partial x_t} \left(\frac{\partial g}{\partial x_i} \cdot \frac{\partial h}{\partial x_j} \cdot c_{ij} \right) \cdot \frac{\partial f}{\partial x_k} \cdot c_{tk} + \frac{\partial}{\partial x_t} \left(\frac{\partial h}{\partial x_i} \cdot \frac{\partial f}{\partial x_j} \cdot c_{ij} \right) \cdot \frac{\partial g}{\partial x_k} \cdot c_{tk} \right\} \\ &= 4 \sum_{i,j,k,t} \left(\frac{\partial^2 f}{\partial x_i \partial x_t} \cdot \frac{\partial g}{\partial x_j} \cdot \frac{\partial h}{\partial x_k} \cdot c_{ij} \cdot c_{tk} + \frac{\partial f}{\partial x_i} \cdot \frac{\partial^2 g}{\partial x_t \partial x_j} \cdot \frac{\partial h}{\partial x_k} \cdot c_{ij} \cdot c_{tk} \right. \\ &\quad \left. + \frac{\partial f}{\partial x_i} \cdot \frac{\partial g}{\partial x_j} \cdot \frac{\partial h}{\partial x_k} \cdot \frac{\partial c_{ij}}{\partial x_t} \cdot c_{tk} + \frac{\partial^2 g}{\partial x_j \partial x_t} \cdot \frac{\partial h}{\partial x_k} \cdot \frac{\partial f}{\partial x_i} \cdot c_{tk} \cdot c_{ji} + \frac{\partial g}{\partial x_k} \cdot \frac{\partial^2 h}{\partial x_i \partial x_t} \cdot \frac{\partial f}{\partial x_j} \cdot c_{kt} \cdot c_{ij} \right. \\ &\quad \left. + \frac{\partial g}{\partial x_j} \cdot \frac{\partial h}{\partial x_k} \cdot \frac{\partial f}{\partial x_i} \cdot \frac{\partial c_{jk}}{\partial x_t} \cdot c_{ti} + \frac{\partial^2 h}{\partial x_t \partial x_i} \cdot \frac{\partial f}{\partial x_j} \cdot \frac{\partial g}{\partial x_k} \cdot c_{ij} \cdot c_{tk} + \frac{\partial h}{\partial x_k} \cdot \frac{\partial^2 f}{\partial x_i \partial x_t} \cdot \frac{\partial g}{\partial x_j} \cdot c_{kt} \cdot c_{ij} \right. \\ &\quad \left. + \frac{\partial h}{\partial x_k} \cdot \frac{\partial f}{\partial x_i} \cdot \frac{\partial g}{\partial x_j} \cdot \frac{\partial c_{ki}}{\partial x_t} \cdot c_{ij} \right) \\ &= 4 \sum_{i,j,k} \frac{\partial f}{\partial x_i} \cdot \frac{\partial g}{\partial x_j} \cdot \frac{\partial h}{\partial x_k} \cdot \sum_t \left(\frac{\partial c_{ij}}{\partial x_t} \cdot c_{tk} + \frac{\partial c_{jk}}{\partial x_t} \cdot c_{ti} + \frac{\partial c_{ki}}{\partial x_t} \cdot c_{ij} \right) \end{aligned}$$

to see that A^- is a Lie algebra if and only if

$$(27) \quad \sum_{i=1}^n \left(\frac{\partial c_{ij}}{\partial x_i} \cdot c_{ik} + \frac{\partial c_{jk}}{\partial x_i} \cdot c_{ii} + \frac{\partial c_{ki}}{\partial x_i} \cdot c_{ij} \right) = 0$$

for $i, j, k = 1, \dots, n$. The equations (27) are redundant for $i \geq j$ and for $j \geq k$. Hence A^- is a Lie algebra if and only if (27) holds for $1 \leq i < j < k \leq n$. (It follows that A^- is a Lie algebra in case $n = 2$.)

The equations (27) are obviously satisfied in any algebra A defined by a skew-symmetric bilinear form ϕ . Using the basis employed before in $M = \sum_i Fx_i$, we have $\text{ad } g/2$ in the form (4) with a_k given by (14) and (15) with $\sigma_1 = \dots = \sigma_n = 0$. Therefore $\text{ad } A$ is an ideal of dimension $p^{n-2r}(p^{2r}-1)$ in $D(A)$ which is of dimension $p^{n-2r}(p^{2r}-1+n)$. If ϕ is nondegenerate, so that $n = 2r$, then $\text{ad } A$ is the (nonsimple) algebra V_{0r} of all $D(g)$ for $g \in A$.

THEOREM 4. *A Lie algebra L is a simple Lie algebra V_r if and only if there is a p^{2r} -dimensional (simple) nodal noncommutative Jordan algebra A defined by a nondegenerate skew-symmetric bilinear form such that $L \cong (\text{ad } A)'$.*

3. The case $n = 2$. Let A in K have least possible dimension p^2 . Then it is known [5, §3] that A is simple. The vector space of A coincides with $B_2 = F[x_1, x_2]$, $x_1^p = x_2^p = 0$, and multiplication in A is defined by

$$(28) \quad fg = f \cdot g + \left(\frac{\partial f}{\partial x_1} \cdot \frac{\partial g}{\partial x_2} - \frac{\partial f}{\partial x_2} \cdot \frac{\partial g}{\partial x_1} \right) \cdot c$$

where c has an inverse c^{-1} in B_2 . Also (5) reduces to the single equation

$$\frac{\partial c}{\partial x_1} \cdot a_1 - \frac{\partial a_1}{\partial x_1} \cdot c + \frac{\partial c}{\partial x_2} \cdot a_2 - \frac{\partial a_2}{\partial x_2} \cdot c = 0,$$

which is equivalent to

$$(29) \quad \frac{\partial}{\partial x_1} (c^{-1} \cdot a_1) + \frac{\partial}{\partial x_2} (c^{-1} \cdot a_2) = 0.$$

If, given D in (4), we write $b \cdot D$ for the derivation

$$f \rightarrow \sum_{k=1}^n \frac{\partial f}{\partial x_k} \cdot (b \cdot a_k)$$

of B_n (that is, $b \cdot D = (b \cdot a_1, \dots, b \cdot a_n)$), and if we write $\delta(D)$ for the *divergence* [3, p. 715]

$$\delta(D) = \sum_{k=1}^n \frac{\partial a_k}{\partial x_k},$$

then condition (29) becomes

$$(30) \quad \delta(c^{-1} \cdot D) = 0.$$

Hence D is a derivation of A if and only if $c^{-1} \cdot D$ is in the (p^2+1) -dimensional Lie algebra M_2 of derivations of B_2 having divergence zero [3]. But $D \leftrightarrow c^{-1} \cdot D$ is a vector space isomorphism, so the derivation algebra $D(A)$ is a (p^2+1) -dimensional algebra $D(A) = c \cdot M_2$. If $c=1$, so that A is defined by a (non-degenerate) form ϕ , then $D(A) = M_2$ consists of the derivations (4) of B_2 given by (14) with $r=1$, while $D(A)' = V_{01}$, and $D(A)'' = V_1$ is simple. More generally two distinct situations arise, depending upon whether c^{-1} is in $\tilde{A}$ or not.

We digress momentarily to point out that (30) defines Lie algebras which generalize the algebras M_n of Frank [3], not only for $n=2$, but for general n . Let c be any invertible element of B_n , and let D range over the derivations of B_n satisfying (30). The set $c \cdot M_n$ of these derivations of B_n is a Lie algebra of dimension $(n-1)p^n+1$ since g in B_n implies

$$(31) \quad \delta(g \cdot [D, E]) = \delta(\delta(g \cdot D) \cdot E) - \delta(\delta(g \cdot E) \cdot D),$$

generalizing [3, Lemma 2]. For (20) gives

$$\begin{aligned} & \delta(\delta(g \cdot D) \cdot E) - \delta(\delta(g \cdot E) \cdot D) \\ &= \sum_j \frac{\partial}{\partial x_j} \left\{ \sum_i \left(\frac{\partial(g \cdot a_i)}{\partial x_i} \cdot b_j - \frac{\partial(g \cdot b_i)}{\partial x_i} \cdot a_j \right) \right\} \\ &= \sum_{i,j} \left(\frac{\partial g}{\partial x_i} \cdot \frac{\partial a_i}{\partial x_j} \cdot b_j + g \cdot \frac{\partial^2 a_i}{\partial x_j \partial x_i} \cdot b_j - \frac{\partial g}{\partial x_i} \cdot \frac{\partial b_i}{\partial x_j} \cdot a_j - g \cdot \frac{\partial^2 b_i}{\partial x_j \partial x_i} \cdot a_j \right) \\ &= \sum_i \frac{\partial g}{\partial x_i} \cdot c_i + g \cdot \sum_i \frac{\partial c_i}{\partial x_i} \\ &= \delta(g \cdot [D, E]). \end{aligned}$$

Putting $g=c^{-1}$ in (31), we see that $[D, E]$ satisfies (30) in case D and E do, or $c \cdot M_n$ is a Lie algebra. Also

$$(32) \quad (c \cdot M_n)' \subseteq c \cdot M_n'.$$

For suppose that D and E satisfy (30). Writing $g=c^{-1}$, we have

$$\begin{aligned} & \sum_{j=1}^n \frac{\partial}{\partial x_j} (g \cdot a_i \cdot b_j - g \cdot b_i \cdot a_j) \\ &= g \cdot \sum_{j=1}^n \left(\frac{\partial a_i}{\partial x_j} \cdot b_j - \frac{\partial b_i}{\partial x_j} \cdot a_j \right) + a_i \cdot \delta(g \cdot E) - b_i \cdot \delta(g \cdot D) \\ &= g \cdot c_i \end{aligned}$$

for c_i in (20). Hence $g \cdot [D, E] = (h_1, \dots, h_n)$ where

$$h_i = \sum_{j \neq i} \frac{\partial}{\partial x_j} (g \cdot a_i \cdot b_j - g \cdot b_i \cdot a_j).$$

Hence $g \cdot [D, E]$ is in the algebra $T_n = S_n$ defined by Frank [3, Lemma 3]. But $S_n = M'_n$ (because S_n is simple in case $n > 2$ and by our earlier remarks in case $n = 2$). Hence $[D, E] \in c \cdot M'_n$, establishing (32). It follows that $c \cdot M'_n = c \cdot S_n$ is a Lie algebra (of dimension $(n-1)(p^n-1)$, the known dimension of S_n), for $(c \cdot M'_n)' \subseteq (c \cdot M_n)' \subseteq c \cdot M'_n$ by (32).

Returning to the case $n=2$, we shall prove that $D(A)'' = (\text{ad } A)' \cong V_1$ is a simple Lie algebra of dimension p^2-2 if $c^{-1} \in \tilde{A}$, whereas $D(A)' = \text{ad } A$ is a simple Lie algebra of dimension p^2-1 if $c^{-1} \notin \tilde{A}$. We begin with a normalization of c by proper choice of x_i in A .

THEOREM 5. *Let A be a (simple) nodal noncommutative Jordan algebra of dimension p^2 in K so that multiplication in A is defined by (28). Then x_i may be chosen in A so that c is in the form*

$$(33) \quad c = 1 + \alpha x_1^{p-1} \cdot x_2^{p-1}, \quad \alpha \in F.$$

According as c^{-1} is or is not in $\tilde{A}$ (for any choice of x_i), we have $\alpha=0$ or $\alpha \neq 0$ in (33).

Proof. Write $c^{-1} = \beta_0 + \beta_1 \cdot x_2 + \cdots + \beta_{p-2} \cdot x_2^{p-2} + \beta_{p-1} \cdot x_2^{p-1}$, $\beta_j \in F[x_1]$. Then β_0^{-1} exists, and $c^{-1} \cdot (1 - \beta_0^{-1} \cdot \beta_{p-1} \cdot x_2^{p-1}) = \beta_0 + \beta_1 \cdot x_2 + \cdots + \beta_{p-2} \cdot x_2^{p-2}$. But then (11) and (12) imply that there exists $y_2 = x_2 \cdot h$ such that

$$(34) \quad \frac{\partial y_2}{\partial x_2} = c^{-1} \cdot (1 - \beta_0^{-1} \cdot \beta_{p-1} \cdot x_2^{p-1}).$$

Now $\partial y_2 / \partial x_2 = x_2 \cdot \partial h / \partial x_2 + h$ implies that h and β_0 are congruent modulo N . Hence h^{-1} exists, so that $y_2 = \delta x_2 + n$, $n \in N \cdot N$, $\delta \neq 0$. Let $y_1 = x_1$. Then $A = F[y_1, y_2]$, $y_1^p = y_2^p = 0$. Now $y_2^{p-1} = x_2^{p-1} \cdot h^{p-1}$, so that $x_2^{p-1} \in A \cdot y_2^{p-1}$. But every element of $A \cdot y_2^{p-1}$ has the form $\rho \cdot y_2^{p-1}$ for $\rho \in F[y_1]$. Then (3) and (34) imply $[y_1, y_2]/2 = (\partial y_2 / \partial x_2) \cdot c = 1 - \beta_0^{-1} \cdot \beta_{p-1} \cdot x_2^{p-1} = 1 + \sigma \cdot y_2^{p-1}$ where

$$\sigma = -\beta_0^{-1} \cdot \beta_{p-1} \cdot \rho \in F[y_1].$$

That is, we may as well take c in (28) in the form

$$c = 1 + \sigma \cdot x_2^{p-1}, \quad \sigma \in F[x_1]$$

Now $\sigma = \alpha_0 1 + \alpha_1 x_1 + \cdots + \alpha_{p-2} x_1^{p-2} + \alpha_{p-1} x_1^{p-1}$ for $\alpha_j \in F$. Then

$$\begin{aligned} c^{-1} \cdot (1 + \alpha_{p-1} x_1^{p-1} \cdot x_2^{p-1}) &= (1 - \sigma \cdot x_2^{p-1}) \cdot (1 + \alpha_{p-1} x_1^{p-1} \cdot x_2^{p-1}) \\ &= 1 - \alpha_0 x_2^{p-1} - \alpha_1 x_1 \cdot x_2^{p-1} - \cdots - \alpha_{p-2} x_1^{p-2} \cdot x_2^{p-1}. \end{aligned}$$

By (11) and (12) there exists

$$y_1 = x_1 \cdot (1 + \pi \cdot x_2^{p-1}), \quad \pi \in F[x_1],$$

such that

$$(35) \quad \frac{\partial y_1}{\partial x_1} = c^{-1} \cdot (1 + \alpha x_1^{p-1} \cdot x_2^{p-1})$$

where we have written α for $\alpha_{p-1} \in F$. Then $y_1^{p-1} = x_1^{p-1} \cdot (1 - \pi \cdot x_2^{p-1})$, or $x_1^{p-1} = y_1^{p-1} \cdot (1 + \pi \cdot x_2^{p-1})$. Let $y_2 = x_2$. Then $A = F[y_1, y_2]$, while (3) and (35) imply that $[y_1, y_2]/2 = (\partial y_1 / \partial x_1) \cdot c = 1 + \alpha x_1^{p-1} \cdot x_2^{p-1} = 1 + \alpha y_1^{p-1} \cdot (1 + \pi \cdot y_2^{p-1}) \cdot y_2^{p-1} = 1 + \alpha y_1^{p-1} \cdot y_2^{p-1}$. That is, we may take c in the form (33).

The final statement in the theorem could probably be established by a careful analysis of the argument above. Instead we note that, for any choice of x_i , V_{01} consists of all $D(g)$ with $g \in A$, and $V_1 = V'_{01}$ of all $D(g)$ for which $g \in \tilde{A}$, while $[D(f), D(g)] = D(h)$ where

$$(36) \quad h = \frac{\partial f}{\partial x_1} \cdot \frac{\partial g}{\partial x_2} - \frac{\partial f}{\partial x_2} \cdot \frac{\partial g}{\partial x_1}.$$

Since 1 is in $\tilde{A}$, as well as being of the form (36), it follows that $\tilde{A}$ consists of all linear combinations of elements of the form (36). By (24) we have $[f, g]/2 = h \cdot c$ for h in (36), or $[A, A] = \tilde{A} \cdot c$. Hence $c^{-1} \in \tilde{A}$ if and only if $1 \in [A, A]$. But the latter condition is independent of the choice of x_i . Hence, for any choice of x_i , $c^{-1} \in \tilde{A}$ if and only if the $c^{-1} = 1 - \alpha x_1^{p-1} \cdot x_2^{p-1}$ given by (33) is in $\tilde{A}$.

One point in the proof of the next theorem is deferred to the final section where we consider the simple Lie algebras $L(G, \delta, f)$.

THEOREM 6. *Let A be a (simple) nodal noncommutative Jordan algebra of dimension p^2 in K so that multiplication in A is defined by (28). Then*

$$(37) \quad D(A) = c \cdot M_2$$

where M_2 is the (p^2+1) -dimensional algebra consisting of all derivations (4) of B_2 given by (14) with $r=1$, and $D(A)''$ is simple. If $c^{-1} \in \tilde{A}$, then $D(A) \cong M_2$, $D(A)' \cong V_{01}$, and $D(A)'' \cong V_1$ is of dimension p^2-2 . If $c^{-1} \notin \tilde{A}$, then $D(A)' = D(A)'' = c \cdot M'_2$ is of dimension p^2-1 .

Proof. The case $c^{-1} \in \tilde{A}$ has already been established since we may take $c=1$ by Theorem 5. Suppose that $c^{-1} \notin \tilde{A}$ so we may take c in the form (33) with $\alpha \neq 0$. We have seen that (37) holds. Hence (32) implies that

$$(38) \quad \dim D(A)' \leq \dim(c \cdot M'_2) = p^2 - 1.$$

We shall see in the next section that a class of central simple Lie algebras L_0 of dimension p^2-1 is obtained as follows: $L_0 \cong \text{ad } A$ where A has multiplication defined by (28) with $c = \gamma(1+x_1) \cdot (1+x_2)$ for $\gamma \neq 0 \in F$. We trace through

the steps of the proof of Theorem 5 to see that x_i may be chosen in this A so that

$$(39) \quad \alpha = -\gamma^{p-1}$$

in (33): $c^{-1} = \gamma^{-1}(1+x_1)^{-1} \cdot (1+x_2)^{-1} = \gamma^{-1}(1+x_1)^{-1}(1-x_2+x_2^2 - \cdots + x_2^{p-1})$ so that $\beta_0 = \beta_{p-1} = \gamma^{-1}(1+x_1)^{-1}$. Then $y_1 = x_1$,

$$\begin{aligned} y_2 &= \gamma^{-1}(1+x_1)^{-1} \cdot x_2 \cdot \left(1 - \frac{1}{2}x_2 + \cdots - \frac{1}{p-1}x_2^{p-2}\right), \\ y_2^{p-1} &= \gamma^{-(p-1)}(1+x_1)^{-(p-1)} \cdot x_2^{p-1} \cdot \left(1 - \frac{1}{2}x_2 + \cdots - \frac{1}{p-1}x_2^{p-2}\right)^{p-1} \\ &= \gamma^{-(p-1)}(1+x_1)^{-(p-1)} \cdot x_2^{p-1}, \end{aligned}$$

so that $x_2^{p-1} = \gamma^{p-1}(1+x_1)^{p-1} \cdot y_2^{p-1} = \gamma^{p-1}(1+y_1)^{p-1} \cdot y_2^{p-1}$. Then α in (33) is the coefficient of $y_1^{p-1} \cdot y_2^{p-1}$ in $-\beta_0^{-1} \cdot \beta_{p-1} \cdot x_2^{p-1} = -x_2^{p-1}$; that is, we have (39). Let $H = F(\gamma)$ where γ satisfies (39). Then $(\text{ad } A)_H = \text{ad}(A_H) \cong L_0$ (an algebra defined over H) is simple and of dimension p^2-1 over H . Hence $\text{ad } A$ is simple and of dimension p^2-1 over F . But $\text{ad } A \subseteq D(A)$ implies $p^2-1 = \dim(\text{ad } A) = \dim(\text{ad } A)' \leq \dim D(A)' \leq p^2-1$ by (38). Hence $\text{ad } A = D(A)' = D(A)'' = (c \cdot M_2)' = c \cdot M_2'$.

We remark that equality holds in (32) for $n=2$.

4. The simple algebras $L(G, \delta, f)$. The simple Lie algebras L_0 and L_δ of Albert and Frank [1] have been generalized by Block⁽³⁾ in [2] to an extensive class of simple Lie algebras $L(G, \delta, f)$. Block has shown [2, Lemma 3] that each V_r is an algebra $L(G, \delta, f)$. In this section we prove

THEOREM 7. *For any simple Lie algebra $L(G, \delta, f)$ (of characteristic $\neq 2$) there exists a simple nodal noncommutative Jordan algebra A in K such that A^- is a Lie algebra and $L(G, \delta, f) \cong (\text{ad } A)'$, an ideal in $D(A)$. Actually $L_0 \cong \text{ad } A$.*

If $L(G, \delta, f)$ is simple, then $G = G_0 + G_1 + \cdots + G_m$ is an elementary p -group [2, Theorem 2], so that each G_k may be regarded as an n_k -dimensional vector space over the prime field F_p of characteristic p . The order of G is p^n

⁽³⁾ I am indebted to Dr. Block for furnishing me with a copy of his excellent dissertation [2] before its publication. My Theorem 7 was suggested by his Lemma 3. The following remarks about [2] may be of interest: (i) each of the algebras $V_{m,\mu}$ is isomorphic to V_m , for $y_i = \mu_i^{-1}x_i$, $y_{m+i} = \mu_i^{-1}x_{m+i}$ ($i=1, \dots, m$) implies $\mu_i \partial \phi / \partial x_{m+i} = \partial \phi / \partial y_{m+i}$, $-\mu_i \partial \phi / \partial x_i = -\partial \phi / \partial y_i$, and the coefficient of $(x_1 \cdots x_{2m})^{p-1}$ is zero if and only if the coefficient of $(y_1 \cdots y_{2m})^{p-1}$ is; (ii) if $F = F_p$, the prime field of characteristic p , then any simple $L(G, \delta, f)$ for which $G_0=0$ is isomorphic to V_m , for [2, Theorem 4] implies that in each $G_i \{h(\delta), h(\beta_1), \dots, h(\beta_k)\}$ and $\{g(\beta_{k+1}), \dots, g(\beta_r)\}$ are linearly independent sets (of elements in F) over F_p , so that $k=0$ and $r=k+1=1$, requiring that each G_i be 2-dimensional over F_p and that $L(G, \delta, f) \cong V_m$ by [2, Lemma 3] and (i) above.

where $n = n_0 + n_1 + \cdots + n_m$, and we write $q_{-1} = 0$, $q_k = n_0 + n_1 + \cdots + n_k$ ($k = 0, 1, \cdots, m$). Let $\sigma_1, \cdots, \sigma_{n_0}$ be any basis for G_0 over F_p . Since $\delta = \delta_0 + \delta_1 + \cdots + \delta_m$ where $\delta_0 = 0$ and $\delta_k \neq 0$ in G_k for $k = 1, \cdots, m$, we may take a basis $\sigma_{q_{k-1}+1}, \cdots, \sigma_{q_k-1}, \delta_k$ for G_k over F_p ($k = 1, \cdots, m$). But then, defining σ_{q_k} by

$$(40) \quad \delta_k = \sigma_{q_{k-1}+1} + \cdots + \sigma_{q_k-1} + \sigma_{q_k},$$

we also have $\sigma_{q_{k-1}+1}, \cdots, \sigma_{q_k}$ a basis for G_k over F_p ($k = 1, \cdots, m$). Then $\sigma_1, \cdots, \sigma_n$ is a basis for G over F_p , and any $\alpha \in G$ may be written uniquely in the form

$$(41) \quad \alpha = \sum_{i=1}^n s_i \sigma_i, \quad s_i \in F_p.$$

Now $L(G, \delta, f)$ is the derived algebra of a Lie algebra L/Fu_0 of dimension $p^n - 1$ over F , where L has a basis consisting of p^n elements u_α in (1-1) correspondence with the elements α of G . By (41) the u_α are in (1-1) correspondence with the n -tuples $(s_1, \cdots, s_n)$, $s_i \in F_p$, and we shall represent the u_α in this way. The skew-symmetric biadditive function $f(\alpha, \beta)$ on G to F may be taken so that $f(\alpha_k, \beta_l) = 0$ for $k \neq l$, $\alpha_k \in G_k$, $\beta_l \in G_l$ ($k, l = 0, 1, \cdots, m$). Writing $f(\sigma_i, \sigma_j) = \alpha_{ij} \in F$, we see that

$$(42) \quad \alpha_{ij} = 0 \text{ unless } q_{k-1} + 1 \leq i, j \leq q_k \text{ for some } k \quad (0 \leq k \leq m).$$

Now (41) and $\beta = \sum_{j=1}^n t_j \sigma_j$ imply $f(\alpha_k, \beta_k) = \sum_{i, j=q_{k-1}+1}^{q_k} s_i t_j \alpha_{ij}$. Since $\delta = \sum_{k=n_0+1}^n \sigma_k$ by (40), we see that $[2, (4)]$ defines multiplication in L by

$$(43) \quad (s_1, \cdots, s_n)(t_1, \cdots, t_n) = \sum_{i, j=1}^{n_0} s_i t_j \alpha_{ij} (s_1 + t_1, \cdots, s_n + t_n) \\ + \sum_{k=1}^m \left(\sum_{i, j=q_{k-1}+1}^{q_k} s_i t_j \alpha_{ij} (s_1 + t_1, \cdots, s_{q_{k-1}} + t_{q_{k-1}}, s_{q_{k-1}+1} + t_{q_{k-1}+1}, \cdots, s_{q_k} + t_{q_k} - 1, s_{q_{k+1}} + t_{q_{k+1}}, \cdots, s_n + t_n) \right).$$

Instead of the nilpotent generators x_i of $B_n = F[x_1, \cdots, x_n]$ used in previous sections, we use at this point generators $z_i = 1 + x_i$ ($i = 1, \cdots, n$). We have $z_i^p = 1$, and every element of B_n may be written uniquely in the form

$$(44) \quad f = \sum_{s_i \in F_p} \alpha_{s_1 \cdots s_n} z_1^{s_1} \cdots z_n^{s_n}, \quad \alpha_{s_1 \cdots s_n} \in F.$$

Let A in K be of dimension p^n so that $A^+ = B_n$. Then (24) implies that multiplication in A^- is defined by

$$[f, g] = \sum_{i, j} \frac{\partial f}{\partial z_i} \cdot \frac{\partial g}{\partial z_j} \cdot 2c_{ij}$$

since $\partial f / \partial z_i = \partial f / \partial x_i$. Equivalently, multiplication in A^- is defined by

$$(45) \quad [z_1^{s_1} \cdots z_n^{s_n}, z_1^{t_1} \cdots z_n^{t_n}] = \sum_{i,j=1}^n s_i t_j z_1^{s_1+t_1} \cdots z_i^{s_i+t_i-1} \cdots z_j^{s_j+t_j-1} \cdots z_n^{s_n+t_n} \cdot 2c_{ij}.$$

Let

$$(46) \quad \begin{aligned} c_{ij} &= 0 && \text{unless } q_{k-1} + 1 \leq i, j \leq q_k \text{ for some } k \ (0 \leq k \leq m), \\ 2c_{ij} &= \alpha_{ij} z_i \cdot z_j && \text{for } 1 \leq i, j \leq n_0, \\ 2c_{ij} &= \alpha_{ij} z_i \cdot z_j \cdot (z_{q_{k-1}+1} \cdots z_{q_k})^{-1} && \text{for } q_{k-1} + 1 \leq i, j \leq q_k \ (k = 1, \dots, m). \end{aligned}$$

For typographical reasons we write $\{s_1, \dots, s_n\}$ for $z_1^{s_1} \cdots z_n^{s_n}$. Then (45) and (46) imply

$$(47) \quad \begin{aligned} [\{s_1, \dots, s_n\}, \{t_1, \dots, t_n\}] &= \sum_{i,j=1}^{n_0} s_i t_j \alpha_{ij} \{s_1 + t_1, \dots, s_n + t_n\} \\ &+ \sum_{k=1}^n \left(\sum_{i,j=q_{k-1}+1}^{q_k} s_i t_j \alpha_{ij} \{s_1 + t_1, \dots, s_{q_{k-1}} + t_{q_{k-1}}, s_{q_{k-1}+1} + t_{q_{k-1}+1} - 1, \dots, \right. \\ &\quad \left. s_{q_k} + t_{q_k} - 1, s_{q_k+1} + t_{q_k+1}, \dots, s_n + t_n \} \right). \end{aligned}$$

That is, $L \cong A^-$ by (43) and (47), and $L/Fu_0 \cong A^-/F1$.

In order to complete the proof of the theorem, we shall require the following

LEMMA 2. *Let A be a noncommutative Jordan algebra such that A^+ is associative. If $[f, g] = 0$ for every $g \in A$, then $f \cdot A$ ($= fA = Af$) is an ideal of A . Thus, if A is any simple nodal noncommutative Jordan algebra, $[f, g] = 0$ for every $g \in A$ if and only if $f \in F1$.*

Proof. Clearly $fg = gf = f \cdot g$ for every $g \in A$. Then [6, (4)] implies $(f \cdot g)h = -(g \cdot h)f + (gh) \cdot f + (gf) \cdot h = -f \cdot g \cdot h + f \cdot (gh) + f \cdot g \cdot h = f \cdot (gh) \in f \cdot A$. If $A = F1 + N$ is simple, then either $f \cdot A = 0$, implying $f = 0$, or $f \cdot A = A$. In the latter case $f = \alpha 1 + z$, $\alpha \neq 0$, $z \in N$. But then $[z, g] = [f - \alpha 1, g] = 0$ for every g , implying $z = 0$, $f = \alpha 1 \in F1$.

We return to the proof of Theorem 7. A is in K , since at least one of the α_{ij} in (46) is not zero [2, Theorem 2]. If $G = G_0$, then $L_0 = L(G, \delta, f) = L/Fu_0$ has dimension $p^n - 1$. If $G \neq G_0$, then $L(G, \delta, f) = (L/Fu_0)'$ has dimension $p^n - 2$. We shall show in both cases that A is simple since $L(G, \delta, f)$ is. If A is not simple, then A has a nonzero ideal $B \subseteq N$. Since N is not an ideal of A ,

$$(48) \quad 1 \leq \dim B \leq p^n - 2.$$

B^- is an ideal of A^- , and either $F1 \cap B^- = F1$, implying $1 \in B$, $B = A$, a con-

tradiction, or $F1 \cap B^- = 0$. Hence $C^- = F1 \oplus B^-$ is an ideal of A^- . In case $G = G_0$, then $A^-/F1 \cong L(G, \delta, f)$ is simple, so the kernel $C^-/F1$ of the natural homomorphism of $A^-/F1$ onto A^-/C^- is either 0 or all of $A^-/F1$. That is, either $B^- = 0$ or $\dim B^- = p^n - 1$, contradicting (48) in either event. In case $G \neq G_0$, $L/Fu_0 = L(G, \delta, f) + Fv$ contains an ideal M corresponding to $C^-/F1$ in $A^-/F1$. Clearly $M \cong B^-$. Then $M \cap L(G, \delta, f)$ is an ideal of the simple algebra $L(G, \delta, f)$. In view of (48), it follows that either (i) $M = L(G, \delta, f)$, or (ii) $L/Fu_0 = L(G, \delta, f) \oplus M$ where $\dim B = \dim M = 1$. In case (i), $L/Fu_0 = M + Fv$. Correspondingly, $A^-/F1 = C^-/F1 + F\bar{z}$ where z may be taken to be in N . Then $A = C + Fz = F1 + B + Fz$, $N = B + Fz$. Now B an ideal of A implies $NN = (B + Fz)(B + Fz) \subseteq B + Fz^2 \subseteq N$, a contradiction, since A is a nodal algebra. There remains the possibility (ii), $L/Fu_0 = (L/Fu_0)' \oplus M$. Correspondingly, $(A^-/F1) = (A^-/F1)' \oplus F\bar{w}$ where w may be taken in B . Then $B = Fw$. Since B^+ is a 1-dimensional ideal in $A^+ = F[x_1, \dots, x_n]$, we have $w = \sigma x_1^{p-1} \cdot \dots \cdot x_n^{p-1}$ for $\sigma \neq 0 \in F$. Write $c_{ij} = \alpha_{ij}1 + z_{ij}$ in (24), $\alpha_{ij} \in F$, $z_{ij} \in N$. There exist i_0, j_0 such that $\alpha_{i_0 j_0} \neq 0$. Then $[x_{i_0}, x_1^{p-1} \cdot \dots \cdot x_n^{p-1}] \in B$ implies

$$\begin{aligned} 0 &\equiv [x_{i_0}, x_1^{p-1} \cdot \dots \cdot x_n^{p-1}] \equiv (p-1) \sum_i x_1^{p-1} \cdot \dots \cdot x_j^{p-2} \cdot \dots \cdot x_n^{p-1} \cdot 2c_{i_0 j} \\ &\equiv 2(p-1) \sum_i \alpha_{i_0 j} x_1^{p-1} \cdot \dots \cdot x_j^{p-2} \cdot \dots \cdot x_n^{p-1} \pmod{B} = Fx_1^{p-1} \cdot \dots \cdot x_n^{p-1}, \end{aligned}$$

implying $\alpha_{i_0 j_0} = 0$, a contradiction. That is, A must be simple if $L(G, \delta, f)$ is. That $\text{ad } A$ is isomorphic to $A^-/F1$ follows directly from Lemma 2.

In §2 we referred to the proof above for justification of the statement that any A defined by a nondegenerate form ϕ is simple. This follows from the fact that $V_r = L(G, \delta, f)$ where $G_0 = 0$, $m = r$, and G_k is of dimension 2 over F_p for $k = 1, \dots, r$ [2, Lemma 3]. For then the c_{ij} defined by (46) are all in $F1$. In the proof of Theorem 6 we relied on (46) for the case $n = n_0 = 2$. In that instance $2c = 2c_{12} = \alpha_{12}z_1 \cdot z_2 = \alpha_{12}(1+x_1) \cdot (1+x_2)$ with $\alpha_{12} \neq 0$.

We have not computed the derivations of the algebras A in Theorem 7. Instead we conclude with the following result which generalizes (26) in the direction of (14).

THEOREM 8. *Let A be in K so that multiplication is defined by (1). If A^- is a Lie algebra, then the mappings D defined by (4) with*

$$(49) \quad a_i = \sum_{j=1}^n \left(\frac{\partial g}{\partial x_j} + \alpha_j x_j^{p-1} \right) \cdot c_{ij}, \quad i = 1, \dots, n,$$

for any $g \in A$ and any $\alpha_j \in F$ ($j = 1, \dots, n$), are derivations of A .

Since $\text{ad } g/2$ in (25) is a derivation of A , it is sufficient to verify that D in (4) is a derivation in case

$$(50) \quad a_i = \sum_{k=1}^n \alpha_k x_k^{p-1} \cdot c_{ik}, \quad \alpha_k \in F.$$

Now D is a derivation of A in case (5) with k replaced by t is satisfied. But (50) implies

$$\begin{aligned}
 & \sum_{i=1}^n \left(\frac{\partial c_{ij}}{\partial x_t} \cdot a_t + \frac{\partial a_i}{\partial x_t} \cdot c_{jt} + \frac{\partial a_j}{\partial x_t} \cdot c_{ti} \right) \\
 &= \sum_{k,t} \left(\frac{\partial c_{ij}}{\partial x_t} \cdot \alpha_k x_k^{p-1} \cdot c_{ik} + \alpha_k \frac{\partial (x_k^{p-1} \cdot c_{ik})}{\partial x_t} \cdot c_{jt} + \alpha_k \frac{\partial (x_k^{p-1} \cdot c_{jk})}{\partial x_t} \cdot c_{ti} \right) \\
 &= \sum_k \alpha_k x_k^{p-1} \cdot \left\{ \sum_t \left(\frac{\partial c_{ij}}{\partial x_t} \cdot c_{ik} + \frac{\partial c_{ik}}{\partial x_t} \cdot c_{jt} + \frac{\partial c_{jk}}{\partial x_t} \cdot c_{ti} \right) \right\} \\
 &\quad + \sum_k \alpha_k x_k^{p-2} \cdot (c_{ik} \cdot c_{jk} + c_{jk} \cdot c_{ki}) \\
 &= 0
 \end{aligned}$$

by (27).

REFERENCES

1. A. A. Albert and M. S. Frank, *Simple Lie algebras of characteristic p* , Rend. Sem. Mat. Torino vol. 14 (1954–1955) pp. 117–139.
2. Richard Block, *New simple Lie algebras of prime characteristic*, Trans. Amer. Math. Soc. vol. 89 (1958) pp. 421–449.
3. M. S. Frank, *A new class of simple Lie algebras*, Proc. Nat. Acad. Sci. U.S.A. vol. 40 (1954) pp. 713–719.
4. Nathan Jacobson, *Classes of restricted Lie algebras of characteristic p . II*, Duke Math. J. vol. 10 (1943) pp. 107–121.
5. L. A. Kokoris, *Some nodal noncommutative Jordan algebras*, Proc. Amer. Math. Soc. vol. 9 (1958) pp. 164–166.
6. ———, *Simple nodal noncommutative Jordan algebras*, Proc. Amer. Math. Soc. vol. 9 (1958) pp. 652–654.
7. ———, *Nodal noncommutative Jordan algebras*, to appear in Canad. J. Math.
8. D. G. Northcott, *Ideal theory*, Cambridge, 1953.
9. R. D. Schafer, *On noncommutative Jordan algebras*, Proc. Amer. Math. Soc. vol. 9 (1958) pp. 110–117.

INSTITUTE FOR ADVANCED STUDY,
PRINCETON, NEW JERSEY
UNIVERSITY OF CONNECTICUT,
STORRS, CONNECTICUT