METRICAL THEOREMS ON FRACTIONAL PARTS OF SEQUENCES

BY
WOLFGANG M. SCHMIDT⁽¹⁾

1. Introduction. Let C be the additive group of real numbers modulo 1, and let $x \rightarrow \{x\}$ be the natural mapping from the reals onto C . It is clear what we shall mean by an interval I in C and by the length $l(I)$ of I . Denote the distance of the real number α to the closest integer by $\|\alpha\|$. The image in C of the set of reals ξ satisfying $\|\xi - \theta\| \leq \varepsilon$ with given θ and $0 < \varepsilon < 1/2$ is an example of an interval of C of length 2ε .

THEOREM 1. Let $n \geq 1$ and let $P_1(q), \dots, P_n(q)$ be nonconstant polynomials with integral coefficients. For each of the integers $j = 1, \dots, n$ let $I_{j1} \supseteq I_{j2} \supseteq \dots$ be a sequence of nested intervals in C . Put $\psi(q) = l(I_{1q}) \cdots l(I_{nq})$ and

$$(1.1) \quad \Psi(h) = \sum_{q=1}^h \psi(q).$$

Put $N(h; \alpha_1, \dots, \alpha_n)$ for the number of integers q , $1 \leq q \leq h$, with

$$(1.2) \quad \{\alpha_j P_j(q)\} \in I_{jq} \quad (j = 1, \dots, n).$$

Let $\varepsilon > 0$. Then

$$(1.3) \quad N(h; \alpha_1, \dots, \alpha_n) = \Psi(h) + O(\Psi(h)^{1/2+\varepsilon})$$

for almost every n -tuple of real numbers $\alpha_1, \dots, \alpha_n$.

The theorem implies, for example, that the number of solutions of

$$|\alpha q - p - \theta| \leq q^{-1}$$

in integers p and q , $1 \leq q \leq h$, is asymptotically equal to $2 \log h$ for every $\alpha \notin \sigma(\theta)$ where $\sigma(\theta)$ is a set of measure zero. To see this we only have to put $n = 1$, $P(q) = q$ and to define intervals I_q as the images of the sets $\|\xi - \theta\| \leq q^{-1}$.

On the other hand, let $P(q) = a_0 q^d + \dots + a_d$ be a polynomial of degree $d > 0$ with integral coefficients, let μ be real, and let $M(h; \alpha)$ be the number of solutions in integers p, q , $1 \leq q \leq h$, of

$$(1.4) \quad |\alpha - p/P(q)| \leq q^{-\mu}.$$

Received by the editors August 29, 1962 and, in revised form, December 11, 1962.

⁽¹⁾ This paper was written with partial support of the National Science Foundation Grant No. NSF-G19894.

Then $M(h; \alpha)$ is bounded for almost every α if $\mu > d + 1$; $M(h; \alpha) \sim 2|a_0| \log h$ if $\mu = d + 1$; and $M(h; \alpha) \sim 2|a_0|h^{d+1-\mu}(d+1-\mu)^{-1}$ for almost every α if $\mu < d + 1$.

To see this, we remark that for $\mu > d$ and large q , (1.4) is equivalent to $\|\alpha P(q)\| \leq |P(q)|q^{-\mu}$. Thus our interval I_q has length $\psi(q) = 2|P(q)|q^{-\mu} = |2a_0q^{d-\mu} + 2a_1q^{d-\mu-1} + \dots|$, and the theorem gives the result. For $\mu = d$, (1.4) becomes $|\alpha P(q) - p| \leq |a_0 + a_1q^{-1} + \dots|$, and $M(h; \alpha)$ becomes $2|a_0|h$ plus (or minus) the number of solutions of $\|\alpha P(q)\| \leq |a_1q^{-1} + \dots|$ for $1 \leq q \leq h$, whence $M(h; \alpha) \sim 2|a_0|h$ almost everywhere. Finally for $\mu < d$ our formula for $M(h; \alpha)$ is in fact true for every α . The reader should have no difficulty in proving this elementary result.

There can be at most countably many α_j 's such that $\{\alpha_j P_j(q)\}$ is an endpoint of I_{jq} for some q , and hence we may assume I_{jq} to be closed ($j = 1, \dots, n$; $q = 1, 2, \dots$). The intersections $J_j = \bigcap_q I_{jq}$ ($j = 1, \dots, n$) are then nonempty. The case where $0 \in J_j$ for each j is usually called the homogeneous case, the general case the inhomogeneous case.

Our theorem implies in particular that $N(h; \alpha_1, \dots, \alpha_n)$ remains bounded almost everywhere if $\Psi(h)$ is bounded, while it will tend to infinity almost everywhere if $\Psi(h)$ tends to infinity. This had been proved by Khintchine [9] in the homogeneous case under the assumption that $P_j(q) = q$ ($j = 1, \dots, n$) and that $q\psi(q)$ is decreasing. Szűsz [13] generalized Khintchine's result to the inhomogeneous case. Szűsz' method involves continued fractions and therefore applies only to the case $n = 1$. Before Szűsz, Cassels [2] had shown that Khintchine's conclusion is true for "almost every inhomogeneous case," that is, if $(I_{1q}, \dots, I_{nq})$ is replaced by its translation by a vector $(\theta_1, \dots, \theta_n)$ of reals mod 1 ($q = 1, 2, \dots$), then the conclusion is true for almost every $\theta_1, \dots, \theta_n$. Thus Cassel's result was "doubly metrical."

Erdős [5] proved for the homogeneous case with $n = 1$, $P(q) = q$, that $N(h; \alpha) \sim \Psi(h)$ almost everywhere, and the author [12]⁽²⁾ proved (1.3) in this case. Our generalization from the homogeneous to the inhomogeneous case is not trivial. We shall choose $\theta_j \in J_j$ ($j = 1, \dots, n$) and use rational approximations to θ_j . The generalization from linear to general polynomials also causes some difficulty.

Le Veque [10] proved a general theorem where polynomials $P(q)$ are replaced by general sequences $a(q)$ which have to satisfy a certain condition. However, this condition is not satisfied for $a(q) = q$, and it is difficult to decide whether it is satisfied for nonlinear polynomials.

It would be possible to replace (1.2) by $(\{\alpha_1 P_1(q)\}, \dots, \{\alpha_n P_n(q)\}) \in H_q$, thus replacing products of intervals $I_{1q} \times \dots \times I_{nq}$ by somewhat more general sets H_q of $C \times \dots \times C$.

(2) We use this opportunity to mention two errors in [12]: In Theorem 1 of [12] one has to assume that the functions $\psi_j(q)$ are bounded. Everywhere in §6 except in $\beta(Q, \theta)$, θ should be replaced by $\Theta = (\theta_1, \dots, \theta_n)$.

In §10 we shall point out how one could prove a more general theorem where the expressions $\alpha_j P_j(q)$ are replaced by linear forms $\alpha_{j1} P_{j1}(q_1) + \dots + \alpha_{jm} P_{jm}(q_m)$. A special case of such a result is contained in Theorem 2 of [12].

THEOREM 2. *Let a sequence of positive integers $a_i(1) < a_i(2) < \dots$ be given for $i = 1, \dots, n$. Let θ be arbitrary but fixed, and put*

$$\Sigma(h; \alpha_1, \dots, \alpha_n) = \sum_{q_1=1}^h \dots \sum_{q_n=1}^h \left(q_1 \dots q_n \left\| \sum_{i=1}^n \alpha_i a_i(q_i) + \theta \right\| \right)^{-1}.$$

Then one has for $\varepsilon > 0$ and almost every $\alpha_1, \dots, \alpha_n$

$$(1.5) \quad (\log h)^{n+1} \ll \Sigma(h; \alpha_1, \dots, \alpha_n) \ll (\log h)^{n+1+\varepsilon}.$$

Using Theorem 2, together with an n -dimensional generalization of a result of Erdős and Turán [7, Theorem 3], we shall easily deduce

THEOREM 3. *Assume the hypotheses of Theorem 1 to be satisfied, and assume we deal with the special case $P_j(q) = q$ ($j = 1, \dots, n$) and $I_{j1} = I_{j2} = \dots$ ($j = 1, \dots, n$). Write ψ for $\psi(1) = \psi(2) = \dots$, and let $\varepsilon > 0$. Then*

$$N(h; \alpha_1, \dots, \alpha_n) = h\psi + O(\log h)^{n+1+\varepsilon}$$

for almost every $\alpha_1, \dots, \alpha_n$.

Khinchine [8, §3], proved the surprisingly small error-term $O(\log h)^{1+\varepsilon}$ for $n = 1$, and hence our result is not best possible. However, Khinchine's method involves continued fractions and cannot easily be generalized to $n > 1$. It seems that Theorem 1 cannot much be improved for nonlinear polynomials. Behnke [1, Theorem XXV] showed for $n = 1$, $P(q) = q^2$ and $I_1 = I_2 = \dots = I$, say, that the relation $D_\alpha(h) = \text{def } \sup_I |N_I(h; \alpha) - hI(I)| \ll \sqrt{h}$ is wrong for every α .

2. Notation and simplification. Throughout, $[\alpha]$ will be the integral part of the real number α . U will denote the unit interval $0 \leq \xi < 1$.

We shall prove the case $n = 1$ of Theorem 1 in §§2–8. In §9 we shall point out the necessary changes for $n > 1$.

The set of α 's in U where $\{\alpha P(q)\} \in I_q$ has measure $\psi(q)$. Assume now that $\Psi(h)$ is bounded. Given $\varepsilon > 0$ there is a q_0 such that $\sum_{q > q_0} \psi(q) < \varepsilon$, and the set of α 's in U such that $\{\alpha P(q)\} \in I_q$ for some $q > q_0$ has measure $< \varepsilon$. Hence $N(h; \alpha)$ is bounded for almost every α .

From now on, we shall assume that $\Psi(h)$ tends to infinity.

Let $\theta \in J = \bigcap I_q$. Then each I_q is union of θ and of two intervals I_q^l and I_q^r , where I_q^l is of the type $0 < \{\theta - \xi\} \leq \psi^l(q)$, where I_q^r is of the type $0 < \{\xi - \theta\} \leq \psi^r(q)$, and where $\psi^l(q) + \psi^r(q) = \psi(q)$. (I_q^l or I_q^r may be empty.) Now $\Psi^l(h)$, $\Psi^r(h)$, $N^l(h; \alpha)$, $N^r(h; \alpha)$ can be defined in the obvious way. One has $\Psi(h) = \Psi^l(h) + \Psi^r(h)$ and $N(h; \alpha) = N^l(h; \alpha) + N^r(h; \alpha)$ for almost every α . Hence it will suffice to prove the theorem for the case of intervals of type I^l and the case of intervals of type I^r .

Since the mapping $\xi \rightarrow -\xi$, $\theta \rightarrow -\theta$ transforms intervals of type I^l into intervals of type I^r , we may restrict ourselves to intervals of type I^r .

From now on, I_q will denote the interval

$$0 < \{\xi - \theta\} \leq \psi(q).$$

Replacing $P(q)$ by $-P(q)$ and α by $-\alpha$ if necessary, we may assume that $P(q) > 0$, $P'(q) > 0$ for $q > q_0$. Making a translation by q_0 we may even assume $P(q) > 0$, $P'(q) > 0$ for $q > 0$.

The introduction of a parameter k is essential for our proof. Put $\phi(k, x)$ for the number of integers y between 1 and x , $1 \leq y \leq x$, such that $\text{g.c.d.}(x, y) \leq k$. $\phi(1, x)$ is the well-known Euler ϕ -function.

Given $q \geq 1$ there are pairs of relatively prime integers a, b such that

$$(2.1) \quad 1 \leq a \leq q^{1/2} \quad \text{and} \quad |\theta - b/a| < a^{-1} q^{-1/2}.$$

This follows from Dirichlet's principle. For every integer $q \geq 1$ we pick integers $a = a(q)$, $b = b(q)$ with these properties. We define $S(k, q)$ as the set of integers p where

$$(2.2) \quad \text{g.c.d.}(pa(q) + b(q), P(q)) \leq k.$$

The sets $S(k, q)$ have two important properties:

(1) If $p \in S(k, q)$ and $p \equiv p' \pmod{P(q)}$, then $p' \in S(k, q)$.

(2) The number $\phi^*(k, q)$ of integers of $S(k, q)$ in $1 \leq x \leq P(q)$ satisfies $\phi^*(k, q) \geq \phi(k, P(q))$.

To prove (2), put $P(q) = q_1 q_2$ where every prime factor of q_1 divides a and where q_2 and a are relatively prime. Now $\text{g.c.d.}(a, b) = 1$ yields $\text{g.c.d.}(pa + b, P(q)) = \text{g.c.d.}(pa + b, q_2)$ and $\phi^*(k, q) = q_1 \phi(k, q_2) \geq \phi(k, P(q))$.

We now put

$$\beta(q, \alpha) = \begin{cases} 1 & \text{if } \alpha \in U \text{ and } \{\alpha\} \in I_q, \\ 0 & \text{otherwise,} \end{cases}$$

$$\gamma(q, \alpha) = \sum_p \beta(q, \alpha P(q) - p),$$

$$\gamma(k, q, \alpha) = \sum_{p \in S(k, q)} \beta(q, \alpha P(q) - p),$$

$$\Gamma(q) = \int_0^1 \gamma(q, \alpha) d\alpha,$$

$$\Gamma(k, q) = \int_0^1 \gamma(k, q, \alpha) d\alpha,$$

$$\Gamma(k, q, r) = \int_0^1 \gamma(k, q, \alpha) \gamma(k, r, \alpha) d\alpha,$$

$$A(k, q, r) = \Gamma(k, q, r) - \psi(q)\psi(r),$$

and

$$\Psi(u, v) = \sum_{q=u+1}^v \psi(q).$$

It is easy to see that $N(h; \alpha) = \sum_{q=1}^h \gamma(q, \alpha)$, and we define

$$N(k; u, v; \alpha) = \sum_{q=u+1}^v \gamma(k, q, \alpha).$$

One has

$$(2.3) \quad \Gamma(q) = \sum_p \int_0^1 \beta(q, P(q)\alpha - p) d\alpha = P(q) \int_{-\infty}^{\infty} \beta(q, P(q)\alpha) d\alpha = \psi(q),$$

and similarly

$$(2.4) \quad \Gamma(k, q) = \psi(q) \phi^*(k, q) P(q)^{-1}.$$

Summing over q we find

$$(2.5) \quad \int_0^1 N(h; \alpha) d\alpha = \Psi(h)$$

and

$$(2.6) \quad \int_0^1 N(k; u, v; \alpha) d\alpha = \sum_{q=u+1}^v \psi(q) \phi^*(k, q) P(q)^{-1}.$$

3. Deduction of Theorem 1 from two propositions.

PROPOSITION 1. *Let $\delta > 0$. Then*

$$(3.1) \quad \sum_{q=1}^h (P(q) - \phi(k, P(q))) P(q)^{-1} \ll h k^{\delta-1} + h^{\delta} k^{\delta}.$$

PROPOSITION 2. *For every $\delta > 0$*

$$(3.2) \quad \sum_{q=1}^h \sum_{r=1}^h A(k, q, r) \ll \Psi^{1+\delta}(h) + \Psi(h) k^{\delta}.$$

REMARK. Here and later, the estimate $\ll$ holds simultaneously in h and k . That is, the constant implied by $\ll$ depends only on δ .

We are going to show that Theorem 1 is a consequence of these two propositions. The propositions will be proved later.

LEMMA 1. *Let $\omega_1(q)$, $\omega_2(q)$, $\omega_3(q)$ be positive bounded functions of positive integers q , and put*

$$\Omega_i(h) = \sum_{q=1}^h \omega_i(q) \quad (i = 1, 2, 3).$$

Assume that ω_1 and ω_3 are decreasing, and that $\Omega_2(r) \leq \Omega_3(r)$ for every r . Then

$$(3.3) \quad \sum_{q=1}^h \omega_1(q) \omega_2(q) \ll \Omega_3([\Omega_1(h)]).$$

Proof. If Ω_1 is bounded, then so is the sum in (3.3). Hence we assume Ω_1 to be unbounded. Since ω_1 is decreasing, and since $\Omega_2 \leq \Omega_3$, one finds by partial summation that $\sum_{q=1}^h \omega_1(q) \omega_2(q) \leq \sum_{q=1}^h \omega_1(q) \omega_3(q)$.

To estimate the latter sum we may assume $\omega_1(q) \leq 1$. Put $m_0 = 0$ and for integral $a > 0$ put m_a for the largest m with $\Omega_1(m) \leq a$. Then $m_a \geq a$ and $\omega_1(m_a + 1) + \dots + \omega_1(m_{a+1}) \leq 2$. Putting $b = [\Omega_1(h)]$ we obtain

$$\begin{aligned} \sum_{q=1}^h \omega_1(q) \omega_3(q) &\leq \sum_{a=0}^b (\omega_1(m_a + 1) \omega_3(m_a + 1) + \dots + \omega_1(m_{a+1}) \omega_3(m_{a+1})) \\ &\leq 2 \sum_{a=0}^b \omega_3(m_a + 1) \leq 2 \sum_{a=1}^{b+1} \omega_3(a) = 2\Omega_3([\Omega_1(h)] + 1). \end{aligned}$$

Denote by J_r the set of intervals $(u, v]$, $0 \leq u = t \cdot 2^s < v = (t+1)2^s \leq 2^r$ where r, s, t are non-negative integers. Every interval $(0, w]$ where w is integral and $w \leq 2^r$ is union of not more than $\max(1, r)$ intervals of J_r . Given an integer $u > 0$ put n_u for some integer satisfying $[\Psi(n_u)] = u$, and put $n_0 = 0$. Since $\psi(q) \leq 1$ and since $\Psi(h)$ tends to infinity, such an n_u will always exist. Put $h_r = n_{2^r}$.

For the remainder of this section, k and r will be connected by

$$(3.4) \quad k = 2^r.$$

LEMMA 2. Let $\delta > 0$. Then

$$(3.5) \quad 0 \leq \int_0^1 (N(h_r; \alpha) - N(k; 0, h_r; \alpha)) d\alpha \ll 2^{r\delta}$$

and

$$(3.6) \quad \sum_{(u,v] \in J_r} \int_0^1 (N(k; n_u, n_v; \alpha) - \Psi(n_u, n_v))^2 d\alpha \ll 2^{r+r\delta}.$$

Proof. Formulae (2.5) and (2.6) yield

$$\begin{aligned} S_r &= \int_0^1 (N(h_r; \alpha) - N(k; 0, h_r; \alpha)) d\alpha = \sum_{q=1}^{h_r} \psi(q) (P(q) - \phi^*(k, q)) P(q)^{-1} \\ &\leq \sum_{q=1}^{h_r} \psi(q) (P(q) - \phi(k, P(q))) P(q)^{-1}. \end{aligned}$$

We now put

$$\omega_1(q) = \psi(q), \quad \omega_2(q) = (P(q) - \phi(k, P(q))) P(q)^{-1}, \quad \omega_3(q) = c(k^{\delta-1} + q^{\delta-1} k^{\delta}).$$

Proposition 1 shows that Lemma 1 is applicable if $c > 0$ is chosen large enough. Under our conditions we actually obtain the bound $2\Omega_3([\Omega_1(h)] + 1)$. Hence

$$S_r \ll \Psi(h_r)k^{\delta-1} + \Psi(h_r)^{\delta}k^{\delta} \ll 2^{r+r(\delta-1)} + 2^{2r\delta} \ll 2^{2r\delta}.$$

This is true for every $\delta > 0$, and hence (3.5) is proved.

$$N(k; u, v; \alpha) - \Psi(u, v) = \sum_{q=u+1}^v (\gamma(k, q, \alpha) - \psi(q)).$$

Hence by (2.3), (2.4) and the estimate just derived,

$$\begin{aligned} & \int_0^1 (N(k; u, v; \alpha) - \Psi(u, v))^2 d\alpha \\ &= \sum_{q=u+1}^v \sum_{q'=u+1}^v (\Gamma(k, q, q') - \Gamma(k, q)\psi(q') - \Gamma(k, q')\psi(q) + \psi(q)\psi(q')) \\ &= \sum_{q=u+1}^v \sum_{q'=u+1}^v A(k, q, q') + 2 \sum_{q=u+1}^v \sum_{q'=u+1}^v \psi(q)\psi(q')(P(q) - \phi^*(k, q))P(q)^{-1} \\ &\ll \sum_{q=u+1}^v \sum_{q'=u+1}^v A(k, q, q') + \sum_{q'=u+1}^v \psi(q')2^{r\delta}. \end{aligned}$$

We first consider the part of the sum (3.6) where $(u, v]$ are intervals of J_r with fixed s (see the definition of J_r). These intervals cover $(0, 2^r]$ exactly once, and hence the corresponding intervals $(n_u, n_v]$ cover $(0, h_r]$ exactly once. Our part of the sum (3.6) has the upper bound

$$\sum_{q=1}^{h_r} \sum_{q'=1}^{h_r} A(k, q, q') + 2^{r\delta} \sum_{q=1}^{h_r} \psi(q) \ll \Psi(h_r)^{1+\delta} + \Psi(h_r)k^{\delta} + 2^{r\delta}\Psi(h_r) \ll 2^{r+r\delta}.$$

Summing over s from 0 to r we find the bound $\ll r2^{r+r\delta}$ for the sum (3.6). Since $\delta > 0$ is arbitrary, Lemma 2 is proved.

LEMMA 3. *Let δ be positive and fixed. Then there is a sequence of subsets $\sigma_1, \sigma_2, \dots$ of U with measures*

$$\mu_r = \int_{\sigma_r} d\alpha \ll r^{-2}$$

such that

$$N(n_w; \alpha) = \Psi(n_w) + O(r^2 2^{r/2+r\delta})$$

for every $w \leq 2^r$ and every $\alpha \in U$ which is not in σ_r .

Proof. We define σ_r to be the subset of U where not both of the following inequalities hold:

$$(3.7) \quad 0 \leq N(h_r; \alpha) - N(k; 0, h_r; \alpha) \leq r^2 2^{r/2},$$

$$(3.8) \quad \sum_{(u, v] \in J_r} (N(k; n_u, n_v; \alpha) - \Psi(n_u, n_v))^2 \leq r^2 2^{r+r\delta}.$$

Lemma 2 implies $\mu_r \ll r^{-2}$. Every interval $(0, w]$, $w \leq 2^r$, is union of at most $\max(1, r)$ intervals of J_r , hence $(0, n_w]$ is union at most $\max(1, r)$ intervals $(n_u, n_v]$ where $(u, v] \in J_r$. Thus $N(k; 0, n_w; \alpha) - \Psi(n_w) = \sum (N(k; n_u, n_v; \alpha) - \Psi(n_u, n_v))$, where the sum is over at most $r + 1$ pairs $(u, v] \in J_r$. This relation together with (3.8) and Cauchy's inequality gives for $\alpha \in U$, $\alpha \notin \sigma_r$,

$$(3.9) \quad (N(k; 0, n_w; \alpha) - \Psi(n_w))^2 \leq r^2(r+1)2^{r+r\delta}.$$

Lemma 3 is a consequence of (3.7) and (3.9).

Proof of Theorem 1. Since $\sum r^{-2}$ is convergent, there exists for almost every $\alpha \in U$ an $r_0 = r_0(\alpha)$ such that $\alpha \notin \sigma_r$ for $r \geq r_0$. Assume α has such an r_0 , and assume $w > 2^{r_0}$. Choose r such that $2^{r-1} \leq w < 2^r$. Then $r > r_0$, $\alpha \notin \sigma_r$, and Lemma 3 implies

$$\begin{aligned} N(n_w; \alpha) &= \Psi(n_w) + O(r^2 2^{r/2+r\delta}) \\ (3.10) \quad &= \Psi(n_w) + O(w^{1/2+\delta} \log^2 w) \\ &= \Psi(n_w) + O(\Psi^{1/2+\delta}(n_w) \log^2 \Psi(n_w)). \end{aligned}$$

Since $\Psi(n_{w+1}) = \Psi(n_w) + O(1)$, (3.10) is true for arbitrary integers h and not only the n_w 's. And since $\delta > 0$ was arbitrary, we find

$$N(h; \alpha) = \Psi(h) + O(\Psi(h)^{1/2+\epsilon})$$

for almost every $\alpha \in U$. Hence (1.3) is true for almost every α .

4. The number of solutions of $P(x) \equiv 0 \pmod{d}$. Put $D(q)$ for the number of positive divisors of q . As is well known,

$$(4.1) \quad D(q) \ll q^\delta$$

for every $\delta > 0$. Put $z_P(d) = z_P(d)$ for the number of solutions of $P(x) \equiv 0 \pmod{d}$. Here, as always, $P(x)$ is a nonconstant polynomial with integral coefficients. Define the discriminant Δ of $P(x)$ in the usual way if $P(x)$ is nonlinear, and put $\Delta = \alpha_0$ if $P(x) = a_0x + a_1$.

LEMMA 4. Let $P(x)$ be a polynomial of degree f and with discriminant $\Delta \neq 0$. Then $z_P(p^k) \leq f\Delta^2$ for every prime-power p^k .

Proof. For linear $P(x)$ it is well known that $z(m) \leq \text{g.c.d.}(m, \Delta) \leq \Delta \leq f\Delta^2$. The case where $P(x)$ is nonlinear and primitive, that is, where the coefficients of $P(x)$ are relatively prime, is Theorem 54 of [11]. A proof can be found there. In the general nonlinear case one has $P(x) = cQ(x)$ with primitive $Q(x)$, whence $z_P(p^k) \leq cz_Q(p^k) \leq cf\Delta_Q^2 \leq f\Delta_P^2$.

COROLLARY. Let $P(x)$ be a polynomial with no multiple factors. Let $\delta > 0$. Then

$$(4.2) \quad z_P(d) \ll d^\delta.$$

Proof. The set τ of prime-powers p^k such that $p^{k\delta} \geq f\Delta^2$ is finite. For every d ,

$$z(d)d^{-\delta} \leq \prod_{p^k \in \tau} z(p^k)p^{-k\delta} \ll 1.$$

Given an integer $g > 0$ we define a function $^g(d)$ of positive integers d as follows: $^g(d)$ is multiplicative, and $^g(p^{gx+y}) = p^{x+1}$ if p is a prime and $1 \leq y \leq g$. Our function has the property that $d \mid m^g$ implies $^g(d) \mid m$.

LEMMA 5. *Let $P(x)$ be a nonconstant polynomial, g a positive integer and $s > 1$. Then the two sums*

$$(4.3) \quad \sum_{d=1}^{\infty} z_P(d)d^{-s}$$

and

$$(4.4) \quad \sum_{d=1}^{\infty} (^g(d))^{-s}$$

are convergent.

Proof. There is an integer m and a polynomial $Q(x)$ without multiple factors such that $P(x) \mid Q(x)^m$. Now $P(x) \equiv 0 \pmod{d}$ implies $Q(x) \equiv 0 \pmod{d}$, and hence $z_P(d)d^{-1} \leq z_Q(m(d))(m(d))^{-1}$. Thus one has for $d = p^{mx+y}$ where p is prime and $0 < y \leq m$,

$$\begin{aligned} z_P(d)d^{-s} &\leq z_Q(m(d))(m(d))^{-1}d^{1-s} = z_Q(p^{x+1})p^{(1-s)(mx+y)-x-1} \\ &\leq f\Delta_Q^2 p^{-x(m(s-1)+1)-y(s-1)-1} \leq f\Delta_Q^2 p^{-sx-s}. \end{aligned}$$

This implies

$$\sum_{e=1}^{\infty} z_P(p^e)p^{-es} \leq mf\Delta_Q^2 p^{-s} \sum_{x=0}^{\infty} p^{-xs} \leq c_s p^{-s}.$$

Since the product $\prod_p (1 + cp^{-s})$ over all primes p is convergent, the convergence of (4.3) follows.

The convergence of (4.4) is proved similarly.

5. Proof of Proposition 1. The Euler ϕ -function $\phi(x) = \phi(1, x)$ can be expressed $\phi(x) = x \sum_{y \mid x} \mu(y)y^{-1}$, where $\mu(y)$ is the Moebius function. Now

$$\phi(k, P(q)) = \sum_{x \leq k; x \mid P(q)} \phi(P(q)x^{-1}) = \sum_{x \leq k; x \mid P(q)} P(q)x^{-1} \sum_{y \mid P(q)x^{-1}} \mu(y)y^{-1},$$

hence

$$\begin{aligned} T_{k,h} &\equiv \sum_{q=1}^h \phi(k, P(q))P(q)^{-1} = \sum_{q=1}^h \sum_{x \leq k; x \mid P(q)} x^{-1} \sum_{y \mid P(q)x^{-1}} \mu(y)y^{-1}, \\ &= \sum_{x \leq k; x \leq P(h)} x^{-1} \sum_{y \leq P(h)x^{-1}} \mu(y)y^{-1} \sum_{q \leq h: xy \mid P(q)} 1. \end{aligned}$$

The number of $q \leq h$ such that $xy \mid P(q)$ equals $hz(xy)(xy)^{-1} + O(z(xy))$. Therefore

$$T_{k,h} = h \sum_{x \leq k; x \leq P(h)} \sum_{y \leq P(h)x^{-1}} z(xy)(xy)^{-2} \mu(y) + O\left(\sum_{x \leq k} \sum_{y \leq P(h)} z(xy)(xy)^{-1}\right) \\ = hU_{k,h} + O(V_{k,h}),$$

say. Putting $xy = w$ and using (4.1) with $\delta = \varepsilon/2$ and Lemma 5 with $s = 1 + \varepsilon/2$, $\varepsilon > 0$, we find

$$U_{k,h} = \sum_{w \leq k; w \leq P(h)} z(w)w^{-2} \sum_{y \mid w} \mu(y) + O\left(\sum_{w > k} z(w)w^{-2} D(w)\right) \\ = 1 + O\left(k^{\varepsilon-1} \left(\sum_{w=1}^{\infty} z(w)w^{-1-\varepsilon/2} D(w)w^{-\varepsilon/2}\right)\right) = 1 + O(k^{\varepsilon-1}).$$

Similarly,

$$V_{k,h} \leq \sum_{w \leq P(h)k} z(w)w^{-1} D(w) \leq P(h)^{\varepsilon} k^{\varepsilon} \sum_{w=1}^{\infty} (z(w)w^{-1-\varepsilon/2} D(w)w^{-\varepsilon/2}) \ll P(h)^{\varepsilon} k^{\varepsilon}.$$

Combining our formulae and observing that $\varepsilon > 0$ was arbitrary we obtain $T_{k,h} = h + O(hk^{\delta-1} + h^{\delta}k^{\delta})$, thereby proving the proposition.

We use the remainder of this section to prove four related lemmas.

LEMMA 6. Let $P(x)$ be a polynomial of degree $f > 1$, and let $\varepsilon > 0$. Then

$$W_h \equiv \sum_{q=1}^h P(q)^{-1} \sum_{d \mid P(q); d < q^{f-\varepsilon}} d \sum_{r \leq q; d \mid P(r)} 1 \leq h.$$

Proof. Choose $\delta > 0$ so small that $2\delta f \leq \varepsilon(1 - f^{-1})$.

There is an integer $g \geq 1$ and a polynomial $Q(x)$ with no multiple factors such that $P(x) \mid Q(x)^g$. We may choose $g \leq f$. Now $d \mid P(r)$ implies $^g(d) \mid Q(r)$, hence the number of $r \leq q$ with $d \mid P(r)$ is not larger than $(q(^g(d))^{-1} + 1)z_Q(^g(d))$ and therefore by the corollary to Lemma 4 not larger than

$$\ll (q(^g(d))^{-1} + 1) d^{\delta} \leq (qd^{-1/g} + 1) d^{\delta} \leq (qd^{-1/f} + 1) d^{\delta}.$$

Using $D(P(q)) \ll q^{f\delta}$ we obtain

$$W_h \ll \sum_{q=1}^h q^{-f} \sum_{d \mid P(q); d < q^{f-\varepsilon}} (qd^{1-1/f+\delta} + d^{1+\delta}) \\ \ll \sum_{q=1}^h D(P(q)) (q^{-f+1+(1-1/f+\delta)(f-\varepsilon)} + q^{-f+(1+\delta)(f-\varepsilon)}) \\ \ll \sum_{q=1}^h q^{2\delta f - \varepsilon(1-1/f)} \ll h.$$

LEMMA 7. Let $P(x)$ be arbitrary and $\delta > 0$. Then $\sum_{q=1}^h \sum_{d \mid P(q)} d^{-\delta} \ll h$.

Proof. The part of the sum where $d \geq q$ is not larger than

$$\sum_{q=1}^h q^{-\delta} D(P(q)) \ll h.$$

The part of the sum where $d < q$ is estimated by

$$\sum_{d=1}^h d^{-\delta} \sum_{d < q \leq h; d|P(q)} 1 \leq \sum_{d=1}^h d^{-\delta} h d^{-1} z(d) \leq h \sum_{d=1}^{\infty} z(d) d^{-1-\delta} \ll h.$$

LEMMA 8. Write $D_k(x)$ for the number of positive divisors of x which are not larger than k , and let $\delta > 0$. Then

$$\sum_{q=1}^h D_k(P(q)) \ll h k^{\delta}.$$

Proof. We break the sum into two parts, $\sum_{q=1}^{\min(k,h)} + \sum_{k < q \leq h}$, where the second part may be empty. For q contributing to the first part of the sum, $D_k(P(q)) \leq D(P(q)) \ll k^{\delta}$, and we obtain the desired estimate. The second part equals

$$\begin{aligned} \sum_{k < q \leq h} D_k(P(q)) &= \sum_{d \leq k} \sum_{k < q \leq h; d|P(q)} 1 \leq \sum_{d \leq k} h d^{-1} z(d) \\ &\leq h k^{\delta} \sum_{d=1}^{\infty} z(d) d^{-1-\delta} \ll h k^{\delta}. \end{aligned}$$

LEMMA 9. Write $D(x, y)$ for the number of common positive divisors of integers $x, y \neq 0, 0$. Let $P_1(x), P_2(x)$ be polynomials with integral coefficients such that $P_i(x) \neq 0$ for $x > 0$. Then

$$(5.1) \quad X_{h_1, h_2} = \sum_{q_1=1}^{h_1} \sum_{q_2=1}^{h_2} D(P_1(q_1), P_2(q_2)) \ll h_1 h_2.$$

This estimate holds simultaneously in h_1, h_2 .

Proof. It is sufficient to prove (5.1) with $P_1(x), P_2(x)$ both replaced by the product $P_1(x)P_2(x)$. We may therefore assume $P_1(x) = P_2(x) = P(x)$, say. There is an integer $g > 0$ and a polynomial $Q(x)$ without multiple factors such that $P(x) | Q(x)^g$.

Let σ be the set of positive divisors of $P(x)$ where $1 \leq x \leq \min(h_1, h_2)$. The number of elements of σ is $\ll (\min(h_1, h_2))^{1+\delta}$ for every $\delta > 0$.

$$\begin{aligned} X_{h_1, h_2} &\leq \sum_{d \in \sigma} \left(\sum_{q_1 \leq h_1; d|P(q_1)} 1 \right) \left(\sum_{q_2 \leq h_2; d|P(q_2)} 1 \right) \\ &\leq \sum_{d \in \sigma} \left(\sum_{q_1 \leq h_1; g(d)|Q(q_1)} 1 \right) \left(\sum_{q_2 \leq h_2; g(d)|Q(q_2)} 1 \right) \\ &\leq \sum_{d \in \sigma} (h_1(g(d))^{-1} + 1)(h_2(g(d))^{-1} + 1) z_Q^2(g(d)) \\ &\ll \sum_{d \in \sigma} (h_1(g(d))^{-1} + 1)(h_2(g(d))^{-1} + 1)(g(d))^{2\delta}. \end{aligned}$$

Using the distributive law we can break this sum into four parts, and Lemma 5 implies that each part is $\ll h_1 h_2$.

6. Estimates for $A(k, q, r)$. In what follows, $d^* = d^*(q, r)$ will mean g.c.d. $(P(q), P(r))$. Put $B(k, q, r)$ for the number of pairs of integers $p, s, p \in S(k, q), s \in S(k, r), 0 \leq p < P(q)$, such that

$$|P(q)(s + \theta) - P(r)(p + \theta)| < \min(d^*, P(q)\psi(r)).$$

LEMMA 10. For $r \leq q, A(k, q, r) \leq \psi(q) P(q)^{-1} B(k, q, r)$.

Proof. All the expressions $P(q)s - P(r)p$ are multiples of d^* . Write $C(l, k, q, r)$ for the number of pairs $p, s, p \in S(k, q), s \in S(k, r), 0 \leq p < P(q)$ such that $P(q)s - P(r)p = ld^*$. The congruence $P(r)p \equiv ld^* \pmod{P(q)}$ has d^* solutions in p , and therefore

$$C(l, k, q, r) \leq d^*.$$

By definition,

$$\Gamma(k, q, r) = \sum_{p \in S(k, q)} \sum_{s \in S(k, r)} \int_0^1 \beta(q, P(q)\alpha - p) \beta(r, P(r)\alpha - s) d\alpha.$$

We now make the substitution $P(q)\alpha' = P(q)\alpha - p - \theta$. Then $P(r)\alpha - s = P(r)\alpha' + \theta - (P(q)(s + \theta) - P(r)(p + \theta))P(q)^{-1}$ and

$$\begin{aligned} \Gamma(k, q, r) &= \sum_{p \in S(k, q)} \sum_{s \in S(k, r)} \int_{-(p+\theta)P(q)^{-1}}^{1-(p+\theta)P(q)^{-1}} \beta(q, P(q)\alpha' + \theta) \beta(r, P(r)\alpha' + \theta \\ &\quad - (P(q)(s + \theta) - P(r)(p + \theta))P(q)^{-1}) d\alpha' \\ &= \sum_l C(l, k, q, r) \\ &\quad \cdot \int_{-\infty}^{\infty} \beta(q, P(q)\alpha + \theta) \beta(r, P(r)\alpha + \theta - ld^* + (P(q) - P(r))\theta) P(q)^{-1} d\alpha \\ &= \sum_l C(l, k, q, r) D(q, r, ld^* + (P(q) - P(r))\theta), \end{aligned}$$

where $D(q, r, t) = \int_{-\infty}^{\infty} \beta(q, P(q)\alpha + \theta) \beta(r, P(r)\alpha + \theta - tP(q)^{-1}) d\alpha$.

For the following estimates we recall that $\beta(q, \xi + \theta)$ is the characteristic function of $0 < \xi \leq \psi(q)$. We note

$$\int_{-\infty}^{\infty} D(q, r, t) dt = \psi(q) \psi(r)$$

as well as $0 \leq D(q, r, t) \leq \psi(q) P(q)^{-1}$ and the fact that D is zero outside the interval $(-P(q)\psi(r), P(r)\psi(q))$, hence in particular if $|t| \geq P(q)\psi(r)$. Furthermore, $D(q, r, t)$ is decreasing for $t > 0$, increasing for $t < 0$. Hence

$$\begin{aligned}
 \Gamma(k, q, r) &\leq d^* \sum_{l: |ld^* + (P(q) - P(r))\theta| \geq d^*} D(q, r, ld^* + (P(q) - P(r))\theta) \\
 &\quad + \sum_{l: |ld^* + (P(q) - P(r))\theta| < d^*} C(l, k, q, r) D(q, r, ld^* + (P(q) - P(r))\theta) \\
 (6.1) \quad &\leq d^* \int_{-\infty}^{\infty} D(q, r, \lambda d^* + (P(q) - P(r))\theta) d\lambda + \psi(q) P(q)^{-1} B(k, q, r) \\
 &= \psi(q) \psi(r) + \psi(q) P(q)^{-1} B(k, q, r),
 \end{aligned}$$

and the lemma follows.

Put $E_\delta(k, q, r)$ for the number of $p \in S(k, q)$, $s \in S(k, r)$, $0 \leq p < P(q)$ with $|P(q)(s + \theta) - P(r)(p + \theta)| < P(q)q^{-1}d^{*\delta}$.

LEMMA 11. Let $P(q)$ be a polynomial of degree $f > 0$, and let $\varepsilon = 1$ if $f = 1$, $\varepsilon > 0$ if $f > 1$. Let $\delta > 0$. Then

$$\begin{aligned}
 (6.2) \quad &\sum_{q=1}^h \sum_{r=1}^q \psi(q) P(q)^{-1} B(k, q, r) \ll \Psi(h)^{1+\delta} \\
 &\quad + \sum_{q=1}^h \sum_{r \leq q \text{ with } d^* \geq q^{f-\varepsilon}} \psi(q) P(q)^{-1} E_\delta(k, q, r).
 \end{aligned}$$

Proof. Choose $\delta_1 > 0$, $\delta_2 > 0$ such that $\delta_1 + 1/\delta_2 < \delta$. We shall use the easily proved estimate

$$(6.3) \quad B(k, q, r) \leq 2d^*.$$

We consider four parts $\Sigma_1, \Sigma_2, \Sigma_3, \Sigma_4$ of the sum we want to estimate.

Σ_1 : $d^* < q^{f-\varepsilon}$. We may assume $f > 1$, since $d^* < q^{f-\varepsilon}$ is otherwise impossible.

$$\begin{aligned}
 \Sigma_1 &\leq 2 \sum_{q=1}^h \sum_{r \leq q \text{ with } d^* < q^{f-\varepsilon}} \psi(q) P(q)^{-1} d^* \\
 &\leq 2 \sum_{q=1}^h \psi(q) P(q)^{-1} \sum_{d|P(q); d < q^{f-\varepsilon}} d \sum_{r \leq q; d|P(r)} 1.
 \end{aligned}$$

Using Lemma 6 and partial summation we obtain $\Sigma_1 \ll \Psi(h)$.

Σ_2 : $d^* < (q/r)^{1/\delta_1}$.

$$\begin{aligned}
 \sum_{r < qd^{*- \delta_1}} B(k, q, r) &\ll \sum_{d|P(q)} \sum_{r < qd^{-\delta_1}; d|P(r)} d \leq \sum_{d|P(q)} d \sum_{x \leq P(qd^{-\delta_1}); d|x} 1 \\
 &\ll \sum_{d|P(q)} d P(q) d^{-1-f\delta_1} \\
 \Sigma_2 &\ll \sum_{q=1}^h \psi(q) \sum_{d|P(q)} d^{-f\delta_1} \ll \Psi(h)
 \end{aligned}$$

by Lemma 7 and partial summation.

$$\Sigma_3 : d^* < \Psi(q)^{\delta_2}.$$

$$\begin{aligned} \sum_{r < q \text{ with } d^* < \Psi(q)} \delta_2 B(k, q, r) &\ll \sum_{d | P(q); d < \Psi(q)^{\delta_2}} \sum_{r \leq q; d | P(r)} d \\ &\leq \sum_{d | P(q); d < \Psi(q)^{\delta_2}} d \sum_{x \leq P(q); d | x} 1 \\ &\leq \sum_{d | P(q); d < \Psi(q)^{\delta_2}} d P(q) d^{-1} = P(q) \sum_{d | P(q); d < \Psi(q)^{\delta_2}} 1. \end{aligned}$$

Putting $l = \Psi(h)^{\delta_2}$ we obtain

$$\Sigma_3 \ll \sum_{q=1}^h \psi(q) D_l(P(q)).$$

Lemma 8 together with partial summation gives $\Sigma_3 \ll \Psi(h)^{1+\delta}$.

$\Sigma_4 : d^* \geq q^{f-\varepsilon}$, $d^* \geq (q/r)^{1/\delta_1}$, $d^* \geq \Psi(q)^{\delta_2}$. Under these conditions, $P(q)\psi(r) \leq P(q)\psi(qd^{*- \delta_1(3)}) = P(q)q^{-1}d^{*\delta_1}qd^{*- \delta_1}\psi(qd^{*- \delta_1}) \leq P(q)q^{-1}d^{*\delta_1}\Psi(q) \leq P(q)q^{-1}d^{*\delta_1+1/\delta_2} \leq P(q)q^{-1}d^{*\delta}$; therefore $B(k, q, r) \leq E_\delta(k, q, r)$. Obviously, Σ_4 is bounded by the right-hand sum of (6.2).

7. Proof of Proposition 2 for nonlinear polynomials. In the case of polynomials of degree $f > 1$ we may use Lemma 6, which ceases to be true if $f = 1$. On the other hand, much of the preceding discussion could be simplified for $f = 1$.

We assume now $f > 1$.

We define $(x, y; k)$ by

$$(x, y; k) = \begin{cases} \text{g.c.d.}(x, y) & \text{if this divisor is } \geq xk^{-1}, \\ 0 & \text{otherwise.} \end{cases}$$

LEMMA 12. Assume that $\delta > 0$ is so small that $f - 1/2 < (f - 1/4)(1 - \delta)$. Further assume $q > q_0(P, \delta)$, $r \leq q$, $d^* = \text{g.c.d.}(P(q), P(r)) \geq q^{f-1/4}$. Then

$$(7.1) \quad E_\delta(k, q, r) \leq (P(q), P(r); k).$$

Proof. Put $c = |a_0| + \dots + |a_f|$, where the a_i 's are the coefficients of $P(q)$. Choose q_0 so large that the two inequalities

$$q_0^{1/4} > 2c, \quad 2cq_0^{f-1/2} < q_0^{(f-1/4)(1-\delta)}$$

hold, and let $q \geq q_0$.

The numbers $a = a(q)$, $b = b(q)$ satisfy

$$\theta = b/a + R, \quad |R| < a^{-1}q^{-1/2}, \quad a \leq q^{1/2}$$

and

$$(7.2) \quad 2caq^{f-1} \leq 2cq^{f-1/2} < q^{(f-1/4)(1-\delta)} \leq d^{*1-\delta}.$$

(3) For $0 \leq a < 1$ define $\psi(n-a) = \psi(n)$.

$E_\delta(k, q, r)$ is bounded by the number of pairs p, s , $0 \leq p < P(q)$, $p \in S(k, q)$ satisfying

$$|P(q)s - P(r)p + (P(q) - P(r))(b/a + R)| < P(q)q^{-1}d^{*\delta}.$$

This equation together with (7.2) yields

$$\begin{aligned} |P(q)(s + b/a) - P(r)(p + b/a)| &< P(q)q^{-1}d^{*\delta} + |P(q) - P(r)||R| \\ (7.3) \qquad \qquad \qquad &\leq cq^{f-1}d^{*\delta} + cq^fa^{-1}q^{-1/2} \\ &< d^*/2a + q^{f-1/4}/2a \leq d^*/a. \end{aligned}$$

The left-hand side of (7.3) is an integral multiple of d^*/a , hence it must be zero.

$$(7.4) \qquad P(q)(as + b) = P(r)(ap + b).$$

It remains to estimate the number of solutions of (7.4) in pairs p, s , $0 \leq p < P(q)$, $p \in S(k, q)$. Putting $P(q) = d^*P(q)^*$ we find that (7.4) implies

$$ap + b \equiv 0 \pmod{P(q)^*}.$$

Since a and b are relatively prime, this congruence has at most one solution in p modulo $P(q)^*$, hence at most d^* solutions in $0 \leq p < P(q)$. On the other hand, the congruence gives $\text{g.c.d.}(ap + b, P(q)) \geq P(q)^* = P(q)d^{*-1}$, while $p \in S(k, q)$ implies $\text{g.c.d.}(ap + b, P(q)) \leq k$. Thus $E_\delta(k, q, r)$ is zero unless $d^* \geq P(q)k^{-1}$.

Lemma 12 is proved.

Proof of Proposition 2 ($f > 1$). We may assume that $\delta > 0$ is so small that $f - 1/2 < (f - 1/4)(1 - \delta)$. Combining Lemma 10, Lemma 11 with $\varepsilon = 1/4$ and Lemma 12, we obtain

$$\sum_{q=1}^h \sum_{r=1}^q A(k, q, r) \ll \Psi(h)^{1+\delta} + \sum_{q=1}^h \sum_{r=1}^q \psi(q) P(q)^{-1} (P(q), P(r); k).$$

Using Lemma 8 and partial summation we find

$$\begin{aligned} \sum_{q=1}^h \sum_{r=1}^q P(q)^{-1} (P(q), P(r); k) &\leq \sum_{q=1}^h P(q)^{-1} \sum_{d|P(q); d \geq P(q)k^{-1}} d \sum_{r \leq q; d|P(r)} 1 \\ &\leq \sum_{q=1}^h P(q)^{-1} \sum_{d|P(q); d \geq P(q)k^{-1}} d P(q) d^{-1} \\ &= \sum_{q=1}^h \sum_{d|P(q); d \leq k} 1 \\ &= \sum_{q=1}^h D_k(P(q)) \ll hk^\delta \end{aligned}$$

and

$$\sum_{q=1}^h \sum_{r=1}^q \psi(q) P(q)^{-1} (P(q), P(r); k) \ll \Psi(h)k^\delta.$$

8. Proof of Proposition 2 for linear polynomials.

LEMMA 13. Let $P(x)$ be a linear polynomial, let $0 < \delta < 1/4$, $q \geq 1$ and $d^* \mid P(q)$. Then

$$(8.1) \quad \sum_{r \leq q; (P(q), P(r)) = d^*} E_\delta(k, q, r) \ll q d^{*-1/4} + d^* + q^{1/2} + \begin{cases} q & \text{if } d^* \geq P(q)k^{-1}, \\ 0 & \text{otherwise.} \end{cases}$$

REMARK. The constant involved in the symbol $\ll$ depends on $P(x)$ only.

Proof. Put $c = |a_0| + |a_1|$ where $P(x) = a_0x + a_1$. The relation $|P(q)s - P(r)p + (P(q) - P(r))\theta| < P(q)q^{-1}d^{*\delta} \leq cd^{*1/4}$ in the definition of $E_\delta(k, q, r)$ implies $\|(P(q) - P(r))d^{*-1}\theta\| < cd^{*-3/4}$. Given an r such that the last inequality holds, there are at most $2c$ integers l with

$$|(P(q) - P(r))d^{*-1}\theta - l| < cd^{*-3/4}.$$

Given r and l , $P(q)s - P(r)p = ld^*$ has at most d^* solutions in p , $0 \leq p < P(q)$. Putting

$$F(q, r) = \begin{cases} 1 & \text{if } \|(P(q) - P(r))d^{*-1}\theta\| < cd^{*-3/4}, \\ 0 & \text{otherwise,} \end{cases}$$

we thus find

$$(8.2) \quad E_\delta(k, q, r) \ll d^* F(q, r).$$

Assume now that r runs through those values between 1 and q where $d^* \mid P(r)$. Then $P(q) - P(r)$ runs through some or all of the numbers $0, d^*, 2d^*, \dots, [a_0q/d^*]d^* = q^*d^*$. Thus if we put $G(q, d^*)$ for the number of integers x in $0 \leq x \leq q^*$ satisfying

$$(8.3) \quad \|x\theta\| < cd^{*-3/4},$$

then

$$(8.4) \quad \sum_{r \leq q; d^* \mid P(r)} F(q, r) \leq G(q, d^*).$$

We now distinguish three cases: A , B and C .

A. $2ca(q) \geq d^{*1/4}$. How often does (8.3) hold when x runs through an interval $m < x \leq m + a(q)$? Putting $\theta = b/a + R$ and $x = m + y$, the inequality becomes $\|m\theta + yb/a + yR\| < cd^{*-3/4}$ and this implies $\|m\theta + yb/a\| < cd^{*-3/4} + a|R| < cd^{*-3/4} + q^{-1/2}$. The number of solutions of (8.3) for x in an interval of length a is therefore $\ll (d^{*-3/4} + q^{-1/2})a + 1 \ll d^{*-3/4}a + 1$. Hence

$$G(q, d^*) \ll (d^{*-3/4}a + 1)(q^*a^{-1} + 1) \ll q^*d^{*-1/4} + q^{1/2}d^{*-3/4} + 1,$$

and

$$d^*(G(q, d^*)) \ll qd^{*-1/4} + d^*.$$

(8.1) now follows from (8.2), (8.4) and the last inequality.

B. $2ca(q) < d^{*1/4}$, $2aq^*|R| \geq 1$. Putting $\theta = b/a + R$ again, we rewrite (8.3) as $\|xb/a + xR\| < cd^{*-3/4}$. This implies that

$$(8.5) \quad |m/a + xR| < cd^{*-3/4}$$

for some integer m . For fixed m the number of solutions in x of (8.5) is at most $2cd^{*-3/4}|R|^{-1} + 1$. On the other hand, $x \leq q^*$, whence $|m| \leq cad^{*-3/4} + aq^*|R|$. Thus

$$\begin{aligned} G(q, d^*) &\ll (d^{*-3/4}|R|^{-1} + 1)(ad^{*-3/4} + aq^*|R| + 1) \\ &\leq a^2q^*d^{*-3/2} + ad^{*-3/4} + aq^*d^{*-3/4} + q^*q^{-1/2} + 2aq^*d^{*-3/4} + 1 \\ &\ll a^2q^*d^{*-3/4} + q^*q^{-1/2} + 1 \end{aligned}$$

and

$$d^*G(q, d^*) \ll qd^{*-1/4} + q^{1/2} + d^*.$$

C. $2ca(q) < d^{*1/4}$, $2aq^*|R| < 1$. $E_\delta(k, q, r)$ is bounded by the number of solutions in integers p, s , $0 \leq p < P(q)$, $p \in S(k, q)$, of

$$|P(q)(s + b/a) - P(r)(p + b/a) + (P(q) - P(r))R| < cd^{*1/4}.$$

Now for $r \leq q$, $d^*|P(r)|$, one has $|P(q) - P(r)| \leq q^*d^*$, and we obtain the inequality

$$|P(q)(s + b/a) - P(r)(p + b/a)| \leq cd^{*1/4} + q^*d^*|R| < d^*/2a + d^*/2a = d^*/a.$$

Just as in the proof of Lemma 12 we may conclude that (7.4) holds, and we obtain (7.1). The number of $r \leq q$ with $d^*|P(r)|$ is $\ll qd^{*-1}$, and therefore

$$\sum_{r \leq q; d^*|P(r)} E_\delta(k, q, r) \ll \begin{cases} q & \text{if } d^* \geq P(q)k^{-1}, \\ 0 & \text{otherwise.} \end{cases}$$

Proof of Proposition 2 ($f = 1$). We may assume $0 < \delta < 1/4$. By applying Lemma 10 and Lemma 11 with $\varepsilon = 1$ we obtain

$$\sum_{q=1}^h \sum_{r=1}^h A(k, q, r) \ll \Psi(h)^{1+\delta} + \sum_{q=1}^h \sum_{r=1}^q \psi(q)P(q)^{-1}E_\delta(k, q, r).$$

By Lemma 13,

$$\begin{aligned} &\sum_{q=1}^h \sum_{r=1}^h P(q)^{-1}E_\delta(k, q, r) \\ &\ll \sum_{q=1}^h q^{-1} \sum_{d^*|P(q)} (qd^{*-1/4} + d^* + q^{1/2}) + \sum_{q=1}^h \sum_{d^*|P(q); d^* \geq P(q)k} -1 \\ &\ll \sum_{q=1}^h \sum_{d|P(q)} d^{-1/4} + \sum_{q=1}^h q^{-1/2}D(P(q)) + \sum_{q=1}^h D_k(P(q)) \\ &\ll h + h + hk^\delta \ll hk^\delta. \end{aligned}$$

Here we used (4.1) and Lemmas 7 and 8 to estimate the last three sums. Proposition 2 now follows by partial summation.

9. The higher dimensional case. Most of the arguments used for the case $n = 1$ carry over immediately to $n > 1$, but some of them have to be modified.

We may assume that I_{jq} is of the type $0 < \{\xi_j - \theta_j\} \leq \psi_j(q)$. For each of the integers $j = 1, \dots, n$ we can now define $a_j(q)$, $b_j(b)$, $S_j(k, q)$, $\beta_j(q, \alpha_j)$, $\gamma_j(q, \alpha_j), \dots, \Gamma_j(k, q, r)$. For given q, r we write d_j^* for the greatest common divisor of $P_j(g)$ and $P_j(r)$, and we may now define $B_j(k, q, r), \dots, E_{j\delta}(k, q, r)$, $F_j(q, r)$, $G_j(q, d_j^*)$. We put $\beta(q, \alpha_1, \dots, \alpha_n) = \prod_j \beta_j(q, \alpha_j)$, $\gamma(q, \alpha_1, \dots, \alpha_n) = \prod_j \gamma_j(q, \alpha_j), \dots$, $\Gamma(k, q, r) = \prod_j \Gamma_j(k, q, r)$, and we define $A(k, q, r)$ as in paragraph 2.

PROPOSITION 1a. *Let $\delta > 0$. Then*

$$\sum_{q=1}^h (P_1(q) \cdots P_n(q) - \phi(k, P_1(q)) \cdots \phi(k, P_n(q))) (P_1(q) \cdots P_n(q))^{-1} \ll hk^{\delta-1} + h^{\delta} k^{\delta}.$$

PROPOSITION 2a. *Let $\delta > 0$. Then (3.2) holds.*

The argument of paragraph 3 can be used to deduce the general theorem from these propositions.

Proposition 1a follows from Proposition 1 and

$$\begin{aligned} & P_1(q) \cdots P_n(q) - \phi(k, P_1(q)) \cdots \phi(k, P_n(q)) \\ &= \sum_{j=1}^n P_1 \cdots P_{j-1} (P_j - \phi(k, P_j)) \phi(k, P_{j+1}) \cdots \phi(k, P_n). \end{aligned}$$

(6.1) now becomes

$$\Gamma_j(k, q, r) \leq \psi_j(q) \psi_j(r) + \psi_j(q) P_j(q)^{-1} B_j(k, q, r),$$

and therefore for $r \leq q$

$$A(k, q, r) \leq \sum_{m=1}^n \sum_{\Delta_m} H(k, q, r; m, \Delta_m),$$

where Δ_m runs through all divisions of the integers $1, \dots, n$ into two classes $i_1, \dots, i_m$ and $j_1, \dots, j_{n-m}$, and where

$$H(k, q, r; m, \Delta_m) = \psi(q) \prod_{s=1}^m (P_{i_s}^{-1}(q) B_{i_s}(k, q, r)) \prod_{t=1}^{n-m} \psi_{j_t}(r).$$

For reasons of symmetry it will suffice to estimate $H(k, q, r; m, \Delta_m^0)$, where Δ_m^0 is the division with $i_1 = 1, \dots, i_m = m$. We shall use

$$(9.1) \quad B_i(k, q, r) \leq 2d_i^* \quad (i = 1, \dots, n).$$

LEMMA 14. *Let $m > 1$. Then*

$$\sum_{q=1}^h \sum_{r=1}^q H(k, q, r; m, \Delta_m^0) \ll \Psi(h).$$

Proof. We use the estimate

$$H(k, q, r; m, \Delta_m^0) \ll \psi(q) P_1(q)^{-1} P_2(q)^{-1} d_1^* d_2^*.$$

By Schwartz' inequality,

$$\begin{aligned} Y_h &= \sum_{q=1}^h \sum_{r=1}^q P_1(q)^{-1} d_1^* P_2(q)^{-1} d_2^* \\ &\leq \left(\sum_{q=1}^h \sum_{r=1}^q P_1(q)^{-2} d_1^{*2} \right)^{1/2} \left(\sum_{q=1}^h \sum_{r=1}^q P_2(q)^{-2} d_2^{*2} \right)^{1/2}. \end{aligned}$$

Now

$$\begin{aligned} \sum_{q=1}^h \sum_{r=1}^q P(q)^{-2} d^{*2} &\leq \sum_{q=1}^h P(q)^{-2} \sum_{d|P(q)} d^2 \sum_{r \leq q; d|P(r)} 1 \\ &\leq \sum_{q=1}^h P(q)^{-2} \sum_{d|P(q)} d^2 P(q) d^{-1} = \sum_{q=1}^h \sum_{d|P(q)} d^{-1} \ll h \end{aligned}$$

by Lemma 7. Hence $Y_h \ll h$, and Lemma 14 follows by partial summation.

Everything can be completed as in the case $n = 1$ once we have shown

LEMMA 11a. *Let $\varepsilon = 1$ if the degree f_1 of $P_1(x)$ equals 1, $\varepsilon > 0$ if $f_1 > 1$. Let $\delta > 0$. Then*

$$\begin{aligned} \sum_{q=1}^h \sum_{r=1}^q H(k, q, r; 1, \Delta_1^0) \\ \ll \Psi(h)^{1+\delta} + \sum_{q=1}^h \sum_{r \leq q \text{ with } d_1^* \geq q^{f_1-\varepsilon}} \psi(q) P_1(q)^{-1} E_{1\delta}(k, q, r). \end{aligned}$$

Proof. Choose $\delta_1 > 0$, $\delta_2 > 0$ such that $\delta_1 + 1/\delta_2 < \delta$. We write $\chi_1(q) = \psi_2(q) \cdots \psi_n(q)$ and put

$$\begin{aligned} \Psi_1(h) &= \sum_{q=1}^h \psi_1(q), \\ X_1(h) &= \sum_{q=1}^h \chi_1(q). \end{aligned}$$

Since both $\psi_1(q)$ and $\chi_1(q)$ are decreasing, one has

$$(9.2) \quad h\Psi(h) \geq \Psi_1(h) \chi_1(h).$$

$H(k, q, r; 1, \Delta_1^0)$ equals $\psi(q) P_1(q)^{-1} B_1(k, q, r) \chi_1(r)$.

We consider four parts of the sum we want to estimate. Σ_1 consists of terms with $d_1^* < q^{f_1-\varepsilon}$, Σ_2 of terms with $d_1^* < (q/r)^{1/\delta_1}$, Σ_3 of terms where $d_1^* < \Psi_1(q)^{\delta_2}$, and Σ_4 consists of the remaining terms, that is, terms where $d_1^* \geq q^{f_1-\varepsilon}$,

$d_1^* \geq (q/r)^{1/\delta_1}$, $d_1^* \geq \Psi_1(q)^{\delta_2}$. For the parts Σ_1 , Σ_2 , Σ_4 we estimate $H(\dots) \leq \psi(q)P_1(q)^{-1}B_1(k, q, r)$ and proceed as in paragraph 6. The difficulty lies in estimating Σ_3 .

Let $d|P_1(q)$ and denote the numbers r having $r \leq q$ and $d|P_1(r)$ by $r_1 < r_2 < \dots < r_j$. One has $j \leq P_1(q)d^{-1}$ and $r_j \geq c(jd)^{1/f_1} \geq c'djP_1(q)^{-1}q$ for large q . Hence

$$\begin{aligned} \sum_{r \leq q \text{ with } d_1^* < \Psi_1(q)} \delta_2 B_1(k, q, r) \chi_1(r) &\ll \sum_{d|P_1(q); d < \Psi_1(q)} \delta_2 d \sum_{r \leq q; d|P_1(r)} \chi_1(r) \\ &\ll \sum_{d|P_1(q); d < \Psi_1(q)} \delta_2 d \chi_1(q) P_1(q) q^{-1} d^{-1} \\ &= \chi_1(q) P_1(q) q^{-1} D_f(P_1(q)) \end{aligned}$$

with $f = f(q) = \Psi_1(q)^{\delta_2}$. Hence

$$(9.3) \quad \Sigma_3 \ll \sum_{q=1}^h \psi(q) q^{-1} X_1(q) D_f(P_1(q)).$$

We estimate the last sum in three parts, which overlap somewhat.

$T_1: \Psi(q) \geq q^{1/4\delta_2}$. Unless this part is empty, there is a largest $q \leq h$ in T_1 say q_1 . By (4.1),

$$\begin{aligned} T_1 &\leq \sum_{q=1}^{q_1} \psi(q) D(P_1(q)) \ll \sum_{q=1}^{q_1} \psi(q) q^{\delta/4\delta_2} \leq \Psi(q_1) q_1^{\delta/4\delta_2} \\ &\leq \Psi(q_1)^{1+\delta} \leq \Psi(h)^{1+\delta}. \end{aligned}$$

$T_2: \chi_1(q) \leq q^{1-1/4\delta_2}$. Again using (4.1) we obtain

$$T_2 \ll \sum_{q=1}^h \psi(q) q^{-1/4\delta_2} q^{1/4\delta_2} = \Psi(h).$$

$T_3: \Psi(q) < q^{1/4\delta_2}$, $X_1(q) > q^{1-1/4\delta_2}$. We denote the set of q 's involved by σ , by h_1 the largest element of σ , and we write $g(q) = \psi(q) q^{-1} \chi_1(q)$. Observing (9.2) we find $\Psi_1(q) \leq q^{1/2\delta_2}$ and $f(q) \leq q^{1/2}$ for $q \in \sigma$.

$$T_3 = \sum_{q \in \sigma} g(q) D_f(P_1(q)) = \sum_{d \leq f(h_1)} \sum_{q \in \sigma; d|P_1(q); f(q) \geq d} g(q).$$

Let $x_1, \dots, x_{z(d)}$ be the solutions of $P_1(x) \equiv 0 \pmod{d}$. Since $g(q)$ is decreasing, one has

$$\sum_{q \in \sigma; f(q) \geq d; q \equiv x_i \pmod{d}} g(q) \leq g(q_i) + d^{-1} \sum_{f(q) \geq d} g(q),$$

where q_i is the smallest $q \equiv x_i \pmod{d}$ such that $q \in \sigma$ and $f(q) \geq d$. Since $f(q) \leq q^{1/2}$, one finds $q_i \geq d^2$. Therefore

$$\sum_{q \in \sigma; f(q) \leq d; d|P_1(q)} g(q) \leq z(d) g(d^2) + z(d) d^{-1} \sum_{f(q) \geq d} g(q).$$

Observing $z(d)g(d^2) \leq dg(d^2) \leq \sum_{q=(d-1)^2+1}^{d^2} g(q)$ we obtain

$$\begin{aligned} T_3 &\leq \sum_{d \leq f(h_1)} \left(\sum_{q=(d-1)^2+1}^{d^2} g(q) + z(d)d^{-1} \sum_{f(q) \geq d} g(q) \right) \\ &\leq \sum_{q=1}^{f^2(h_1)} \psi(q) + \sum_{q=1}^h g(q) \sum_{d \leq f(q)} z(d)d^{-1} \\ &\ll \Psi(h) + \sum_{q=1}^h g(q) \Psi_1(q)^\delta \end{aligned}$$

by Lemma 5. (9.2) finally yields

$$T_3 \ll \Psi(h) + \sum_{q=1}^h \psi(q)(q^{-1}\chi_1(q))^{1-\delta} \Psi(q)^\delta \ll \Psi(h)^{1+\delta}.$$

Lemma 11a is proved.

10. Linear forms. We restrict ourselves to the case of one form only.

PROPOSITION 3. Let $P_1(q_1), \dots, P_n(q_n)$ be nonconstant polynomials, $n+1$, and let $I_{q_1, \dots, q_n}$ be intervals of C ($q_i = 1, 2, \dots$; $i = 1, \dots, n$). We assume that the length of $I_{q_1, \dots, q_n}$ is $\psi_1(q_1)\psi_2(q_2)\dots\psi_n(q_n)$, where $\psi_i(x)$ are decreasing functions ($i = 1, \dots, n$), and we put

$$\Psi_i(h) = \sum_{q_i=1}^h \psi_i(q_i).$$

We write $M(h_1, \dots, h_n; \alpha_1, \dots, \alpha_n)$ for the number of solutions of $\{\alpha_1 P_1(q_1) + \dots + \alpha_n P_n(q_n)\} \in I_{q_1, \dots, q_n}$, where $1 \leq q_i \leq h_i$ ($i = 1, \dots, n$). Let $\varepsilon > 0$. Then for almost all $\alpha_1, \dots, \alpha_n$,

$$M(h_1, \dots, h_n; \alpha_1, \dots, \alpha_n) = \Psi_1(h_1) \dots \Psi_n(h_n) + O(\Psi_1(h_1) \dots \Psi_n(h_n))^{1/2+\varepsilon}.$$

This estimate holds simultaneously for $h_1, \dots, h_n$.

Proof. We restrict ourselves to a few hints. The reader might compare paragraph 6 of [12]. We assume $n > 1$.

We put $\beta(q_1, \dots, q_n, \xi)$ equal to 1 if $\{\xi\} \in I_{q_1, \dots, q_n}$ and $\xi \in U$, $\beta(\dots) = 0$ otherwise. $\Gamma(q_1, \dots, q_n; r_1, \dots, r_n)$ stands for the integral

$$\int_0^1 \dots \int_0^1 \left(\sum_p \left(\beta(q_1, \dots, q_n, \sum \alpha_i P_i(q_i) - p) \right) \right) \left(\sum_s \left(\beta(r_1, \dots, r_n, \sum \alpha_i P_i(r_i) - s) \right) \right) d\alpha_1 \dots d\alpha_n,$$

and $A(q_1, \dots, q_n; r_1, \dots, r_n)$ for

$$\Gamma(q_1, \dots, r_n) - \psi_1(q_1) \dots \psi_n(q_n) \psi_1(r_1) \dots \psi_n(r_n).$$

PROPOSITION 2b. $\sum_{q_1=1}^{h_1} \dots \sum_{r_n=1}^{h_n} A(q_1, \dots, r_n) \ll \Psi_1(h_1) \dots \Psi_n(h_n).$

To deduce Proposition 3 from Proposition 2b we put

$$M(h_1, \dots, h_n; k_1, \dots, k_n; \alpha_1, \dots, \alpha_n)$$

for the number of $q_1, \dots, q_n$, $h_i < q_i \leq k_i$ ($i = 1, \dots, n$) such that $\{\sum \alpha_i P_i(q_i)\} \in I_{q_1, \dots, q_n}$ and we put $\Psi_i(h, k) = \sum_{h < q \leq k} \psi_i(q)$. We choose integers $m_u^i = m_u^i(r_1, \dots, r_n)$ such that $[2^{r_1 + \dots + r_n - r_i} \Psi_i(m_u^i)] = u$. The following two lemmas are now used.

LEMMA 2b. Let $\delta > 0$. Then one has for $T = T_{r_1 + \dots + r_n}$

$$\begin{aligned} \sum_{(u_1, v_1) \in J} \dots \sum_{(u_n, v_n) \in J} \int_0^1 (M(m_{u_1}^1, \dots, m_{u_n}^n; m_{v_1}^1, \dots, m_{v_n}^n; \alpha_1, \dots, \alpha_n) \\ - \Psi_1(m_{u_1}^1, m_{v_1}^1) \dots \Psi_n(m_{u_n}^n, m_{v_n}^n))^2 d\alpha_1 \dots d\alpha_n \\ \ll 2^{(r_1 + \dots + r_n)(1 + \delta)}. \end{aligned}$$

LEMMA 3b. Let $\delta > 0$. There are subsets $\sigma_{r_1, \dots, r_n}(r_i = 1, 2, \dots; i = 1, \dots, n)$ of $U \times \dots \times U$ with measures

$$\mu_{r_1, \dots, r_n} \ll r_1^{-2} \dots r_n^{-2}$$

such that

$$M(m_{w_1}^1, \dots, m_{w_n}^n; \alpha_1, \dots, \alpha_n) = \Psi_1(m_{w_1}^1) \dots \Psi_n(m_{w_n}^n) + O(r_1^2 \dots r_n^2 2^{(r_1 + \dots + r_n)(1/2 + \delta)})$$

for every $w_1, \dots, w_n$ with $w_i \leq 2^{r_1 + \dots + r_i}$ ($i = 1, \dots, n$) and $(\alpha_1, \dots, \alpha_n)$ in $U \times \dots \times U$ but not in $\sigma_{r_1, \dots, r_n}$.

To prove Proposition 2b we require

LEMMA 10b. A. If the matrix

$$\begin{pmatrix} P_1(q_1), \dots, P_n(q_n) \\ P_1(r_1), \dots, P_n(r_n) \end{pmatrix}$$

has rank 2, then

$$A(q_1, \dots, r_n) = 0.$$

B. If the matrix has rank 1, then

$$A(q_1, \dots, r_n) \leq \psi_1(q_1) \dots \psi_n(q_n) P_1(q_1)^{-1} B_1(q_1, \dots, r_n),$$

where $B_1(q_1, \dots, r_n)$ is the number of solutions of $|P_1(q_1)(s + \theta') - P_1(r_1)(p + \theta)| < d_1^*$ in integers p, s , $0 \leq p < P_1(q_1)$, where θ, θ' are the left endpoints of $I_{q_1, \dots, q_n}, I_{r_1, \dots, r_n}$, respectively, and where $d_1^* = \text{g.c.d.}(P_1(q_1), P_1(r_1))$.

We leave the proof of A to the reader. As for B, we make the substitution $\alpha_2 = \xi_2, \dots, \alpha_n = \xi_n$, $\sum \alpha_i P_i(q_i) = \xi_1 P_1(q_1)$, hence $\sum \alpha_i P_i(r_i) = \xi_1 P_1(r_1)$. When $\xi_2, \dots, \xi_n$ is fixed, ξ_1 ranges in an interval of length 1, and Γ equals

$$\int_0^1 \left(\sum_p \beta(q_1, \dots, q_n, \xi_1 P_1(q_1) - p) \right) \left(\sum_s \beta(r_1, \dots, r_n, \xi_1 P_1(r_1) - s) \right) d\xi_1.$$

This one-dimensional integral can be estimated by the method of paragraph 6.

The proof of Proposition 2b now proceeds as follows. We may restrict ourselves to terms $r_1 \leq q_1$. For fixed $q_1, \dots, q_n$, let $\Delta = \text{g.c.d.}(P_1(q_1), \dots, P_n(q_n))$ and $P_i(q_i) = P_i(q_i) * \Delta$ ($i=1, \dots, n$). In view of Lemma 10b we may restrict ourselves to $r_1 \leq q_1$ where $P_1(r_1)$ is of the type $lP_1(q_1)^*$, whence $d_1^* = P_1(q_1)^*(\Delta, l)$. Since $B(q_1, \dots, r_n) \leq 2d_1^*$, one has

$$\begin{aligned} \sum_{q_1=1}^{h_1} \dots \sum_{r_n=1}^{h_n} A(q_1, \dots, r_n) &\leq 4 \sum_{q_1=1}^{h_1} \dots \sum_{q_n=1}^{h_n} \psi_1(q_1) \dots \psi_n(q_n) \sum_{l=1}^{\Delta} P_1(q_1)^{-1} (P_1(q_1)^*(\Delta, l)) \\ &\leq 4 \sum_{q_1=1}^{h_1} \dots \sum_{q_n=1}^{h_n} \psi_1(q_1) \dots \psi_n(q_n) D(\Delta) \\ &\leq 4 \left(\sum_{q_1=1}^{h_1} \sum_{q_2=1}^{h_2} \psi_1(q_1) \psi_2(q_2) D(P_1(q_1), P_2(q_2)) \right) \\ &\quad \Psi_3(h_3) \dots \Psi_n(h_n). \end{aligned}$$

Using Lemma 9 and partial summation both for the sum over q_1 and over q_2 we obtain

$$\sum_{q_1=1}^{h_1} \sum_{q_2=1}^{h_2} \psi_1(q_1) \psi_2(q_2) D(P_1(q_1), P_2(q_2)) \ll \Psi_1(h_1) \Psi_2(h_2).$$

11. Theorem 2. To prove the lower bound in (1.5) we shall need

PROPOSITION 4. Let $a(1) < a(2) < \dots$ be a sequence of positive integers and put $M_I(h; \alpha)$ for the number of $q \leq h$ such that $\{\alpha a(q)\} \in I$. Then for $\varepsilon > 0$ and almost all α the inequality

$$|M_I(h; \alpha) - hl(I)| < h^{1/2} \log^{5/2+\varepsilon} h$$

holds for all intervals I and all $h > h_1$, where h_1 depends only on α and ε (but not on I).

Proof. This proposition is a special case of Theorem 1 of [3] and of Theorem 1 of [6].

Proof of Theorem 2. We use the abbreviation

$$\|\Sigma\| = \left\| \sum_{i=1}^n \alpha_i a_i(q_i) + \theta \right\|.$$

Put $\delta = \varepsilon/(n+1)$. Using an idea of Littlewood [4, Appendix A], we consider the integral

$$J(q_1, \dots, q_n) = \int_0^1 \dots \int_0^1 (\|\Sigma\| |\log \|\Sigma\||^{1+\delta})^{-1} d\alpha_1 \dots d\alpha_n.$$

This integral has a finite value independent of $q_1, \dots, q_n$. Hence the sum

$$\sum_{q_1=1}^{\infty} \cdots \sum_{q_n=1}^{\infty} (q_1 \log^{1+\delta} q_1 \cdots q_n \log^{1+\delta} q_n)^{-1} J(q_1, \dots, q_n)$$

is convergent and

$$(11.1) \quad \sum_{q_1=1}^{\infty} \cdots \sum_{q_n=1}^{\infty} (q_1 \log^{1+\delta} q_1 \cdots q_n \log^{1+\delta} q_n \|\Sigma\| |\log \|\Sigma\||^{1+\delta})^{-1}$$

is convergent for almost all $\alpha_1, \dots, \alpha_n$.

It is easy to see that the inequality

$$(11.2) \quad \|\Sigma\| \leq (q_1 \cdots q_n)^{-2}$$

has only a finite number of solutions in integers $q_1, \dots, q_n$ for almost every $\alpha_1, \dots, \alpha_n$.

We are going to show that the upper estimate for $\Sigma(h; \alpha_1, \dots, \alpha_n)$ in (1.5) is true for every $\alpha_1, \dots, \alpha_n$ such that (11.1) is convergent and (11.2) has only finitely many solutions. There is a constant $c > 0$ such that $\|\Sigma\| \geq c^{-1}(q_1 \cdots q_n)^{-2}$, whence

$$|\log \|\Sigma\|| \leq 2 \log(q_1 \cdots q_n) + \log c.$$

We obtain

$$\begin{aligned} \Sigma(h; \alpha_1, \dots, \alpha_n) &\leq \left(\max_{q_i \leq h} (\log^{1+\delta} q_1 \cdots \log^{1+\delta} q_n |\log \|\Sigma\||^{1+\delta}) \right) \\ &\cdot \sum_{q_1=1}^{\infty} \cdots \sum_{q_n=1}^{\infty} (q_1 \log^{1+\delta} q_1 \cdots q_n \log^{1+\delta} q_n \|\Sigma\| |\log \|\Sigma\||^{1+\delta})^{-1} \\ &\ll (\log h)^{(1+\delta)n} (\log h)^{1+\delta} \ll (\log h)^{n+1+\varepsilon}. \end{aligned}$$

We now turn to the proof of the lower bound in (1.5). We are going to apply Proposition 4 to the sequence $a(q) = a_1(q)$. For almost all reals α and $h \geq h_1(\alpha)$ one has $|M_I(h; \alpha) - hI(I)| < h^{3/4}$. Let α_1 have this property. Denote the number of $q \leq h$ such that $\|\alpha_1 a_1(q) + \eta\| \leq \gamma$ by $M_{\gamma, \eta}(h; \alpha_1)$. Then

$$|M_{\gamma, \eta}(h; \alpha_1) - 2\gamma h| < h^{3/4} \quad (h \geq h_1(\alpha_1), 0 \leq \gamma \leq 1/2, \eta \text{ arbitrary}).$$

Let $k_0 = k_0(h)$ be the largest integer with $2^{k_0+1} \leq h^{1/4}$. Then $k_0 \geq 0$ for $h \geq h_2(\alpha_1) = \max(h_1(\alpha_1), 2^8)$. The number $N_{k, \eta}(h; \alpha_1)$ of $q \leq h$ such that

$$(11.3) \quad 2^{-k-1} < \|\alpha_1 a_1(q) + \eta\| \leq 2^{-k}$$

satisfies $N_{k, \eta}(h; \alpha_1) \geq 2^{-k} h - 2h^{3/4} \geq 2^{-k-1} h$ for every k in $0 \leq k \leq k_0$.

By considering the parts of the sum where (11.3) is satisfied for $k = 0, \dots, k_0$ we obtain

$$\sum_{q=1}^h \|\alpha_1 a_1(q) + \eta\|^{-1} \geq \sum_{k=0}^{k_0} 2^k 2^{-k-1} h > \frac{1}{2} h k_0(h) \geq c_1(\alpha_1) h \log h.$$

Partial summation yields

$$\sum_{q=1}^h \|\alpha_1 a_1(q) + \eta\|^{-1} q^{-1} \geq c_2(\alpha_1) \log^2 h.$$

This inequality holds for arbitrary η . By writing $\eta = \alpha_2 a_2(q_2) + \cdots + \alpha_n a_n(q_n) + \theta$ and taking the sum over $q_2, \dots, q_n$ one finds

$$\sum(h; \alpha_1, \dots, \alpha_n) \geq c_3(\alpha_1) \log^{n+1} h.$$

REMARK. Our method could be used to show the following: The left inequality of (1.5) is true for arbitrary $\alpha_1, \dots, \alpha_{n-1}$ and $\alpha_n \in \sigma(\alpha_1, \dots, \alpha_{n-1})$, where $\sigma(\dots)$ is a set containing almost all numbers. The other inequality of (1.5) holds for n -tuples such that $\alpha_n \in \tau$ where τ is independent of $\alpha_1, \dots, \alpha_{n-1}$ and contains almost all numbers.

12. **Theorem 3.** We define a function $\pi(k_1, \dots, k_n)$ as follows. $\pi(0, \dots, 0) = 0$, and if $k_{i_1}, \dots, k_{i_m}$ are those k_i 's which are different from zero, then $\pi(k_1, \dots, k_n) = |k_{i_1} \cdots k_{i_m}|^{-1}$. In our applications $k_1, \dots, k_n$ will always be integers. Write $\exp \xi$ for $e^{2\pi i \xi}$.

GENERALIZED THEOREM OF ERDÖS AND TURAN. *There are absolute constants c_n , $n = 1, 2, \dots$ with the following properties.*

Let $n \geq 1$, $h \geq 1$, and let vectors $(\alpha_{1q}, \dots, \alpha_{nq})$ be given ($q = 1, \dots, h$). Put

$$\omega(k_1, \dots, k_n) = \left| \sum_{q=1}^h \exp \left(\sum_{i=1}^n \alpha_{iq} k_i \right) \right|.$$

Let $I_1, \dots, I_n$ be intervals of C of lengths $l(I_j) = \psi_j$ and put $\psi = \prod \psi_j$. Write N for the number of q , $1 \leq q \leq h$, such that simultaneously $\{\alpha_{jq}\} \in I_j$ ($j = 1, \dots, n$). Let m be a positive integer. Then

$$|N - \psi l| \leq c_n \left(hm^{-1} + \sum_{k_1, \dots, k_n: |k_j| \leq m} \pi(k_1, \dots, k_n) \omega(k_1, \dots, k_n) \right).$$

This theorem is a generalization to n dimensions of a result of Erdős and Turan [7, Theorem 3]. We shall not give a proof, since the argument in [7] can easily be extended to our situation.

Proof of Theorem 3. Put $\alpha_{jq} = \alpha_j q$ ($j = 1, \dots, n$; $q = 1, 2, \dots$).

$$\omega_h(k_1, \dots, k_n) = \left| \sum_{q=1}^h \exp \left(\sum_{i=1}^n k_i \alpha_{iq} \right) \right| \ll \|k_1 \alpha_1 + \cdots + k_n \alpha_n\|^{-1}.$$

Theorem 3 is an immediate consequence of the generalized Erdős-Turan Theorem with $m = h$ and the fact that

$$\sum_{k_1, \dots, k_n: |k_i| \leq m} \pi(k_1, \dots, k_n) \|k_1 \alpha_1 + \cdots + k_n \alpha_n\|^{-1} \ll (\log m)^{n+1+\varepsilon}$$

for almost every $\alpha_1, \dots, \alpha_n$. This fact follows from Theorem 2.

REFERENCES

1. H. Behnke, *Zur Theorie der diophantischen Approximationen*, Abh. Math. Sem. Univ. Hamburg **3** (1924), 261–318.
2. J. W. S. Cassels, *Some metrical theorems in diophantine approximation*. I, Proc. Cambridge Philos. Soc. **46** (1950), 209–218.
3. ———, *Some metrical theorems in diophantine approximation*. III, Proc. Cambridge Philos. Soc. **46** (1950), 219–225.
4. J. W. S. Cassels and H. P. F. Swinnerton-Dyer, *On the product of three homogeneous linear forms and indefinite ternary quadratic forms*, Philos. Trans. Roy. Soc. London A **248** (1956), 73–96.
5. P. Erdős, *Some results on diophantine approximation*, Acta Arith. **5** (1959), 359–369.
6. P. Erdős, and J. F. Koksma, *On the uniform distribution modulo 1 of sequences $f(n, \theta)$* , Indag. Math. **11** (1949), 299–302.
7. P. Erdős and P. Turan, *On a problem in the theory of uniform distribution*, Indag. Math. **10** (1948), 370–382 and 406–413.
8. A. Khintchine, *Ein Satz über Kettenbrüche, mit arithmetischen Anwendungen*, Math. Z. **18** (1923), 289–306.
9. ———, *Einige Sätze über Kettenbrüche mit Anwendungen auf die Theorie der diophantischen Approximationen*, Math. Ann. **92** (1924), 115–125.
10. W. J. Le Veque, *On the frequency of small fractional parts in certain real sequences*. III, J. Reine Angew. Math. **202** (1959), 215–220.
11. T. Nagell, *Introduction to number theory*, Wiley, New York, 1951.
12. W. Schmidt, *A metrical theorem in diophantine approximation*, Canad. J. Math. **12** (1960), 619–631.
13. P. Szűsz, *Über die metrische Theorie der diophantischen Approximationen*, Acta Math. Acad. Sci. Hungar. **9** (1958), 177–193.

COLUMBIA UNIVERSITY,
 NEW YORK, NEW YORK
 UNIVERSITY OF COLORADO,
 BOULDER, COLORADO