THE STRUCTURE AND IDEAL THEORY OF THE PREDUAL OF A BANACH LATTICE

BY

E. B. DAVIES

1. Introduction. If V is an ordered Banach space over the real field, then the Banach dual V^* has a natural induced partial ordering. In Theorems 3.1 and 3.2 we present necessary and sufficient conditions for V^* to be a Banach lattice, extending partial results obtained by Choquet [5] and Andô [1]. The theorems include as special cases the characterisations of the predual of an M -space, [8], [14], and the predual of an L -space, [6], called a simplex space in [11]. We show how the theory is a natural generalisation of Choquet simplex theory.

If V^* is a Banach lattice, we study the set of closed ideals of V . For the special case of simplex spaces, the results provide direct proofs of theorems in [11].

2. Basic theorems on ordered Banach spaces. An ordered normed space V over the real field is defined as a normed vector space V with a closed cone C which is *proper* in the sense that $C \cap (-C) = \{0\}$ and with the partial ordering given by saying that $x \leq y$ if and only if $y - x \in C$. V is said to be *positively generated* if $C - C = V$ and to have a *monotone norm* if $0 \leq x \leq y$ implies $\|x\| \leq \|y\|$. V is said to be *regular* if it has the properties

- (i) if $x, y \in V$ and $-x \leq y \leq x$ then $\|y\| \leq \|x\|$;
- (ii) if $x \in V$ and $\varepsilon > 0$ then there is some $y \in V$ with $y \geq x$, $-x$ and $\|y\| < \|x\| + \varepsilon$.

A *normed lattice* is a regular ordered normed space which is a lattice under its partial ordering, and a *Banach lattice* is a complete normed lattice. Banach lattices are discussed in many places, for example [7], [12], and we follow the standard terminology. A partially ordered vector space V is said to have the *Riesz separation property* if when $a, b \leq c, d \in V$ we can find some $x \in V$ with

$$a, b \leq x \leq c, d.$$

For alternative formulations of this condition see [1], [9].

LEMMA 2.1. *Let V be an ordered positively generated Banach space with a monotone norm $\|\cdot\|$, and define*

$$\|x\|_1 = \inf \{\|y\| : x, -x \leq y \in V\}$$

for all $x \in V$. Then $\|\cdot\|_1$ is a regular norm on V which is equivalent to $\|\cdot\|$, and for any $0 \leq x \in V$ we have $\|x\|_1 = \|x\|$.

For a proof of this see [16].

Let V be a positively generated ordered Banach space with Banach dual V^* . The set C^* of continuous positive functionals is a weak*-closed proper cone in V^* . By the Hahn-Banach theorem if $x \in V$ then $x \geq 0$ if and only if $(x, \phi) \geq 0$ for all $\phi \in C^*$. If V is a positively generated ordered Banach space with a monotone norm then, by [16], every positive functional on V is continuous; also if $0 \leq x \in V$ then

$$\|x\| = \max \{(x, \phi) : \phi \in C^*, \|\phi\| \leq 1\}.$$

This follows from the fact that $-x$ is not in the open convex set

$$\{v \in V : \exists c \in C \cdot \|v - c\| < \|x\|\}$$

by the use of the Hahn-Banach theorem. If V is a regular ordered Banach space then for any $0 \leq \phi \in V^*$ we have

$$\|\phi\| = \sup \{(x, \phi) : x \in C, \|x\| \leq 1\}.$$

LEMMA 2.2. *If V is a regular ordered normed space then for any $\phi, \psi \in V^*$ with $-\phi \leq \psi \leq \phi$ we have $\|\phi\| \geq \|\psi\|$.*

For let $x \in V$ and $\|x\| < 1$. If $z \geq x$, 0 then

$$\begin{aligned} (\psi, x) &= (\psi, z) + (\psi, x - z) \\ &\leq (\phi, z) + (\phi, z - x) \\ &= (\phi, 2z - x). \end{aligned}$$

If $y \in V$ satisfies $y \geq x$, $-x$ and $\|y\| < 1$ and we put $z = (x + y)/2$ then we obtain

$$(\psi, x) \leq \|\phi\| \|y\| < \|\phi\|.$$

The result now follows immediately.

Very much more information can be obtained about V^* , but we first need to consider a special case.

LEMMA 2.3. *Let V be a regular ordered normed space with the Riesz separation property. Then V^* is a Banach lattice.*

Let $\phi \in V^*$ and $x \in V$, $x \geq 0$. Then we define

$$(\phi^+, x) = \sup \{(\phi, y) : 0 \leq y \leq x\}.$$

It is easy to see that ϕ^+ can be uniquely extended to a positive linear functional on V , and that ϕ^+ is continuous with $\|\phi^+\| \leq \|\phi\|$. Clearly, V^* is a lattice with $\phi \vee 0 = \phi^+$. Define $|\phi| = 2\phi^+ - \phi$. For any $y \in V$, $y \geq 0$ we can find some $a \in V$ with $0 \leq a \leq y$ and

$$(\phi, a) \leq (\phi^+, y) \leq (\phi, a) + \varepsilon/2,$$

which gives

$$|(\phi, y) - (\phi, 2a - y)| < \varepsilon.$$

As $-y \leq 2a - y \leq y$ so

$$|(\phi, 2a - y)| \leq \|\phi\| \|y\|$$

from which we conclude that $\|\phi\| \leq \|\phi\|$. The result now follows immediately.

Let V be an ordered positively generated Banach space and define:

$$\Delta = C^* \cap \{\phi \in V^* : \|\phi\| \leq 1\}.$$

Δ is a weak*-compact convex set with $0 \in \Delta$ and there is a natural map $\lambda: V \rightarrow A_0(\Delta)$, the space of all continuous affine functionals ϕ on Δ such that $\phi(0)=0$. By our previous remarks λ is an order isomorphism of V into $A_0(\Delta)$, and if $\|\cdot\|_u$ denotes the supremum norm in $A_0(\Delta)$, we have for all $f \in V$

$$\|\lambda f\|_u \leq \|f\|.$$

Identifying V with λV we define S as the cone of functions of the form $f_1 \vee \cdots \vee f_n$ where $f_i \in V$. Then if $L = S - S$, L is a vector lattice of continuous functions on Δ . See [5]. For $\phi \in L$ define

$$\|\phi\| = \inf \{\|\psi\| : \psi \in V, \psi \geq \phi, -\phi\}.$$

Then, provided V is positively generated and has a monotone norm, L becomes a normed lattice and V is canonically embedded as a Banach subspace of L .

LEMMA 2.4. *If V is a regular ordered Banach space then V^* is also regular.*

Let $\phi \in V^*$ and $\|\phi\| \leq 1$. We can, by the Hahn-Banach theorem, extend ϕ to a functional $\xi \in L^*$ with $\|\xi\| \leq 1$. As L^* is a Banach lattice so by defining ψ as the restriction to V of $|\xi|$, we see that $\psi \geq \phi$, $-\phi$ and $\|\psi\| \leq 1$.

LEMMA 2.5. *If V is an ordered Banach space such that V^* is regular then the map $\lambda: V \rightarrow A_0(\Delta)$ is a one-one onto order isomorphism such that for all $f \in V$*

$$\|\lambda f\|_u \leq \|f\| \leq 2\|\lambda f\|_u,$$

and for all $0 \leq f \in V$,

$$\|\lambda f\|_u = \|f\|.$$

That $\|\lambda f\|_u \leq \|f\|$ is clear. Suppose $f \in V$ and $|(\phi, f)| = \|f\|$ where $\|\phi\| = 1$. Then for any $\varepsilon > 0$ we find $\psi \in V^*$ with $\psi \geq \phi$, $-\phi$, $\|\psi\| < 1 + \varepsilon$.

$$|(\phi, f)| = |((\psi + \phi)/2, f) - ((\psi - \phi)/2, f)|$$

so that

$$\max |((\psi \pm \phi)/2, f)| \geq \frac{1}{2}\|f\|.$$

As $(\psi \pm \phi)/2 \geq 0$ and $\|(\psi \pm \phi)/2\| < 1 + \varepsilon$ so

$$\|f\| \leq 2\|\lambda f\|_u.$$

If $f \geq 0$ then $((\psi \pm \phi)/2, f) \geq 0$ so that

$$|(\phi, f)| \leq \max |((\psi \pm \phi)/2, f)|,$$

from which we see that

$$\|f\| \leq \|\lambda f\|_u.$$

We see that V maps onto a uniformly closed subspace of $A_0(\Delta)$ which separates the points of Δ , and so must conclude by Lemma 4.3 of [13] that

$$\lambda V = A_0(\Delta).$$

A partial converse to Lemma 2.4 is contained in the following

LEMMA 2.6. *Let V be an ordered Banach space such that V^* has a proper positive dual cone C^* and is regular. Then V is positively generated and for any $x, y \in V$ with $-x \leq y \leq x$ we have $\|y\| \leq \|x\|$.*

For let $\psi \in V^*$ and $\|\psi\| < 1$. Let $\phi \in V^*$ satisfy $\phi \geq \psi$, $-\psi$ and $\|\phi\| < 1$. Then

$$\begin{aligned} (\psi, y) &= ((\psi + \phi)/2, y) - ((\phi - \psi)/2, y) \\ &\leq ((\psi + \phi)/2, x) + ((\phi - \psi)/2, x) \\ &= (\phi, x) < \|x\|. \end{aligned}$$

Thus $\|y\| \leq \|x\|$. That V is positively generated follows from [1] upon observing that V^* is $\sigma(0)$ complete.

We continue to suppose that V is an ordered Banach space and that V^* is regular. Following [5], we now define a *conical measure* μ as a positive functional in L^* . Such a functional defines by restriction to V a point of C^* called the *barycentre* of μ . The set P of conical measures μ with $\|\mu\| \leq 1$ is compact in the $\sigma(L^*, L)$ topology. We define a closed partial ordering, $<$, on P by

$$\mu < \nu \quad \text{if} \quad (\mu, f) \leq (\nu, f) \quad \text{for all } f \in S.$$

We say $\mu \in P$ is a *representing conical measure* for $x \in \Delta$ if $f(x) \leq (\mu, f)$ for all $f \in S$, and observe that every $x \in \Delta$ has at least one maximal representing conical measure.

If $x_1, \dots, x_n \in \Delta$ and $x_1 + \dots + x_n = x \in \Delta$ then the functional

$$f \leadsto f(x_1) + \dots + f(x_n)$$

is called a *discrete conical measure*, and is obviously a representing conical measure for x . We show that such conical measures are dense in P in a very good sense.

LEMMA 2.7. *Suppose $\mu \in P$ is a representing conical measure for $x \in \Delta$ and that $f_r \in S$ for $r = 1, \dots, n$. Then there is a discrete representing conical measure ν for x such that $\mu(f_r) = \nu(f_r)$ for $r = 1, \dots, n$.*

We can write the weak*-closed cone $C^* \subseteq V^*$ as a union of closed cones $D_1, \dots, D_m$ such that $f_r|_{D_s}$ are equal to the restrictions of functions in $A_0(\Delta)$. If $f \in L$ and ≥ 0 , we define

$$(\mu_{D_s}, f) = \inf \{(\mu, g) : 0 \leq g \in L \cdot g|_{D_s} > f|_{D_s}\}.$$

Then $\mu_{D_s} \in P$ and $\mu \leq \mu_{D_1} + \dots + \mu_{D_m}$. Now we choose $\mu_s \in P$ such that $0 \leq \mu_s \leq \mu_{D_s}$ and $\mu = \mu_1 + \dots + \mu_m$. If $f \in L$ and $f|_{D_s} \geq 0$ then

$$(f, \mu_s) \geq (f \wedge 0, \mu_s) \geq (f \wedge 0, \mu_{D_s}) = 0.$$

Thus μ_s is determined by the value of f on D_s alone. The barycentre x_s of μ_s is in D_s . Finally for $r=1, \dots, n$

$$(f_r, \mu) = \sum_{s=1}^m (f_r, \mu_s) = \sum_{s=1}^m (f_r(x_s)).$$

LEMMA 2.8. Suppose $f_r \in V$ and $f_1 \vee \dots \vee f_n = f \in S$. Then

$$\max \{(f, \mu) : \varepsilon_x < \mu\} = \max \left\{ \sum_{r=1}^n f(x_r) : x_r \in \Delta, \sum x_r = x \right\}$$

and this quantity is called $\hat{f}(x)$. The map $x \rightarrow \hat{f}(x)$ is upper semicontinuous (u.s.c.).

The set

$$\left\{ \{x_r \in \Delta\}_{r=1}^n : \sum_{r=1}^n x_r = x \in \Delta \right\}$$

is a closed subset of Δ^n and so by the continuity of the f_r we see that the supremum of the right-hand side is always attained. The two sides are equal by Lemma 2.7. That $\hat{f}$ is upper semicontinuous follows from the equation

$$\{x \in \Delta : \hat{f}(x) \geq \alpha\} = \left\{ x = \sum_{r=1}^n x_r : \sum_{r=1}^n f_r(x_r) \geq \alpha \right\}.$$

The following extension of the above lemma to semicontinuous functions will be important in our later discussion of closed ideals.

LEMMA 2.9. Let $f_r : \Delta \rightarrow [-\infty, \infty)$ be upper semicontinuous affine functionals with $f_r(0)=0$. Then for $f=f_1 \vee \dots \vee f_n$ and $x \in \Delta$ the supremum

$$\sup \left\{ \sum_{r=1}^n f(x_r) : \sum_{r=1}^n x_r = x, x_r \in \Delta \right\}$$

is always attained and is denoted $\hat{f}(x)$. $\hat{f}$ is an upper semicontinuous function with $\hat{f}(0)=0$ and $\hat{f}(\alpha x) = \alpha \hat{f}(x)$ for all $x \in \Delta$.

The proof is as for Lemma 2.8.

3. The main theorems.

THEOREM 3.1. Let V be an ordered Banach space, such that V^* is regular. Then the following statements are equivalent:

- (i) V is regular and has the Riesz separation property;
- (ii) V^* is a lattice;
- (iii) the map $f \leadsto \hat{f}(x)$ is linear for any $x \in \Delta$;
- (iv) every $x \in \Delta$ has a unique maximal representing conical measure;
- (v) the map $f \leadsto \hat{f}(x)$ is linear for any $f \in S$.

As an immediate corollary we have

THEOREM 3.2. *Let V be an ordered Banach space. Then V^* is a Banach lattice if and only if V is regular and has the Riesz separation property.*

We now prove Theorem 3.1. The implication (i) $\rightarrow$ (ii) was shown in Lemma 2.3. The implications (ii) $\rightarrow$ (iii) $\rightarrow$ (iv) $\rightarrow$ (v) can be found in [5]. The equivalence (i) $\leftrightarrow$ (ii) is closely related to a Theorem in [1], but is more specific. We now show (v) $\rightarrow$ (i).

Suppose $f, g, h \in V$. Then $(f \vee g)^\wedge$ is u.s.c. affine and so we can define $((f \vee g)^\wedge \vee h)^\wedge$ as in Lemma 2.9. It is easy to show that

$$((f \vee g)^\wedge \vee h)^\wedge = (f \vee g \vee h)^\wedge$$

and so is u.s.c. affine.

Next suppose that f, g are u.s.c. affine on Δ and $f, g \leq h$ where h is affine. Then $(f \vee g)^\wedge \leq h$ without any continuity assumptions for h .

For $f \in -S$ we define $f^\sim$ by $f^\sim = -(-f)^\wedge$. Now let $f_1, f_2, g_1, g_2 \in V$ and $f_1 \vee f_2 \leq g_1 \wedge g_2$. Then if $f = (f_1 \vee f_2)^\wedge$ and $g = (g_1 \wedge g_2)^\sim$ we have $f \leq g$. We construct a sequence $h_n \in A_0(\Delta) \equiv V$ such that

$$f - 1/2^n < h_n < g + 1/2^n.$$

Suppose h_n is given. Let $z \in \Delta$ and $z = x + y$ where $x, y \in \Delta$ and

$$(f \vee h_n)^\wedge(z) = f(x) + h_n(y) < (h_n(z) + 1/2^n) \wedge (g(z) + 1/2^n).$$

Now $(f \vee h_n)^\wedge$ is u.s.c. affine and so by simple convexity arguments to be found in [3], we can find $k_n \in A(\Delta)$ with

$$(f \vee h_n)^\wedge - 1/2^{n+1} < k_n < h_n \wedge g + 1/2^{n+1},$$

and also satisfying $k_n(0) < 0$.

Similarly we can find $l_n \in A(\Delta)$ with

$$f \vee h_n - 1/2^{n+1} < l_n < (h_n \wedge g)^\sim + 1/2^{n+1},$$

and also satisfying $h_n(0) > 0$.

Putting $h_{n+1} = \lambda k_n + (1 - \lambda) l_n$ for a suitable $0 < \lambda < 1$ we obtain

$$(f \vee h_n)^\wedge - 1/2^{n+1} < h_{n+1} < (h_n \wedge g)^\sim + 1/2^{n+1},$$

and $h_{n+1}(0) = 0$.

Then

$$\|h_n - h_{n+1}\|_u < 1/2^{n+1},$$

so $h_n \rightarrow h \in A_0(\Delta)$. It is then clear that

$$f_1 \vee f_2 \leq f \leq h \leq g \leq g_1 \wedge g_2,$$

so that V has the Riesz separation property.

We now show that V is regular. Let $f \in A_0(\Delta)$ and $z \in \Delta$. Then we can find $x, y \in \Delta$ with $x+y=z$ and satisfying

$$\begin{aligned}\{f \vee (-f)\}^\wedge(z) &= f(x) - f(y) \\ &= f(x-y).\end{aligned}$$

Now $-z \leq x-y \leq z$ so by the regularity of V^*

$$\{f \vee (-f)\}^\wedge(z) \leq \|f\|.$$

Now suppose $\varepsilon > 0$. We define $h_1 \in A_0(\Delta)$ so that

$$\{f \vee (-f)\}^\wedge - \varepsilon/2 < h_1 < \|f\| + \varepsilon/2.$$

Inductively we construct $h_n \in A_0(\Delta)$ with

$$\{f \vee (-f)\}^\wedge - \varepsilon/2^n < h_n,$$

and more exactly with

$$\{f \vee (-f)\}^\wedge \vee h_n - \varepsilon/2^{n+1} < h_{n+1} < h_n + \varepsilon/2^{n+1}.$$

This can be done by the same procedure as above. As before $h_n \rightarrow h \in A_0(\Delta)$ and $f \vee (-f) \leq h$. Thus $0 \leq h \in V$ and

$$\|h\| = \|h\|_u \leq \sum_{n=1}^{\infty} \|h_{n+1} - h_n\|_u + \|h_1\|_u < \|f\| + \frac{\varepsilon}{2} + \sum_{n=2}^{\infty} \frac{\varepsilon}{2^n} = \|f\| + \varepsilon.$$

Together with Lemma 2.6 this proves the regularity of V .

4. Some special cases. We define an *R-space* V as a regular ordered Banach space with the Riesz separation property. If V is an ordered Banach space we say it is of *type M* if for any $x, y \geq 0$ we can find $z \geq x, y$ with

$$\|z\| \leq \max\{\|x\|, \|y\|\}$$

and of *type L* if for any $x, y \geq 0$, we have

$$\|x+y\| = \|x\| + \|y\|.$$

LEMMA 4.1. *If V is an R-space of type L then it is a Banach lattice.*

See [2].

LEMMA 4.2. *If V is an R-space of type M then V^* is a Banach lattice of type L.*

The proof of this is trivial.

We can now give an immediate proof of a theorem of Dixmier and Kakutani [8], [14].

THEOREM 4.3. *Let V be an ordered Banach space. Then V is an L-type Banach lattice if and only if V^* is an M-type Banach lattice.*

If V is an L -type Banach lattice, it is an easy corollary of Lemma 2.3 that V^* is an M -type Banach lattice. If V^* is an M -type Banach lattice then V is an R -space and V^{**} is an L -type Banach lattice by Lemma 4.2. As V is canonically embedded in V^{**} , so V is of type L and is an L -type Banach lattice.

THEOREM 4.4. *Let V be an ordered Banach space. Then V is an R -space of type M if and only if V^* is a Banach lattice of type L .*

For the proof see [6]. Such spaces V are referred to as *simplex spaces* in [11].

We show how this theory is related to Choquet boundary theory, as expounded in [3], [9] and [17]. Let V be an ordered Banach space with a distinguished order unit e such that for all $x \in V$ we have

$$\|x\| = \inf \{\alpha : -\alpha e \leq x \leq \alpha e\}.$$

Then V is regular and if

$$\Omega = \{\phi \in V^* : \phi \geq 0 \cdot \phi(e) = 1\},$$

then Ω is a weak*-compact base for the cone C^* in the sense of [10]. V is isometrically and order isomorphic with $A_0(\Delta)$ and with $A(\Omega)$, the space of continuous affine functionals on Ω . The conical measures can be identified with the regular Borel measures on Ω . Ω is called a *simplex* [4] if C^* is lattice-ordered, and it is shown in [9], [15] that this occurs if and only if $A(\Omega)$ has the Riesz separation property.

5. The ideals in an R -space. If V is an ordered Banach space, an *order ideal* I in V is a subspace such that if $0 \leq x \leq y \in I$, then $x \in I$. An ideal is defined as a positively generated order ideal. We now investigate the properties of the closed ideals of an R -space. These generalise the results on the closed ideals of a simplex space in [11], and provide direct proofs for those theorems.

LEMMA 5.1. *Let I be a closed ideal in an R -space V . Then with the restriction norm and ordering I is an R -space.*

The only part not immediate is the second half of the regularity condition. Let $x \in I$ and let $z \in I$, $z \geq x$, $-x$. Let $y \in V$ with $y \geq x$, $-x$ and $\|y\| < \|x\| + \varepsilon$. Then by the Riesz separation property there is some $\omega \in V$ with x , $-x \leq \omega \leq y$, z . As $0 \leq \omega \leq z \in I$ so $\omega \in I$ and as $0 \leq \omega \leq y$ so $\|\omega\| \leq \|y\| < \|x\| + \varepsilon$. Thus I is indeed regular.

LEMMA 5.2. *Let I, J be closed ideals in an R -space V . Then $I \cap J$ is a closed ideal.*

We need only verify that $I \cap J$ is positively generated. Let $x \in I \cap J$ and let $y \in I$, $z \in J$ be such that x , $-x \leq y$ and x , $-x \leq z$. Then we can find $\omega \in V$ such that x , $-x \leq \omega \leq y$, z . As $0 \leq \omega \leq y$ so $\omega \in I$ and as $0 \leq \omega \leq z$ so $\omega \in J$. This proves the lemma.

THEOREM 5.3. *Let I, J be closed ideals in an R -space V . Then $I+J$ is a closed ideal.*

We first show that $I+J$ is positively generated. Let $x=i+j$ where $i \in I$ and $j \in J$. Then let $i_1 \in I$ and $j_1 \in J$ be such that $i, -i \leq i_1$ and $j, -j \leq j_1$. If $x_1 = i_1 + j_1$ then $x_1 \geq 0$ and $x, -x \leq x_1$.

If in particular in the above $x \geq 0$ then by the Riesz separation property as $0 \leq x \leq i_1 + j_1$ we can find $0 \leq i_2 \leq i_1 \in I$ and $0 \leq j_2 \leq j_1 \in J$ such that $x = i_2 + j_2$. Thus the positive cone of $I+J$ is the sum of the positive cones of I and J . It is now immediate that $I+J$ is an order ideal. As in Theorem 5.1 we see that it is regular.

We now have only to show that $I+J$ is a closed subspace of V . Suppose $\varepsilon > 0$ is given. Let $z \in I+J$ and let $z, -z \leq i+j$ where $0 \leq i \in I, 0 \leq j \in J$ and $\|i+j\| < \|z\| + \varepsilon$. Now observe that $z-j, -i \leq z, z+j$ so by the Riesz separation property if we let $x \in V$ satisfy

$$z-j, -i \leq x \leq i, z+j$$

and put $y = z - x$ we have

$$-i \leq x \leq i \quad \text{and} \quad -j \leq y \leq j$$

so that $x \in I$ and $y \in J$. We also have

$$\|x\| \leq \|i\| \leq \|i+j\| < \|z\| + \varepsilon \quad \text{and} \quad \|y\| \leq \|j\| \leq \|j+i\| < \|z\| + \varepsilon.$$

Now let $z_n \in I+J$ and $\sum \|z_n\| < \infty$. Let $z_n = x_n + y_n$ where $x_n \in I, \|x_n\| < 2\|z_n\|$ and $y_n \in J, \|y_n\| < 2\|z_n\|$. Then we see that $\sum x_n \rightarrow x \in I$ and $\sum y_n \rightarrow y \in J$, so that $\sum z_n \rightarrow x+y \in I+J$. This proves that $I+J$ is closed.

THEOREM 5.4. *Let I be a closed ideal in an R -space V and let V/I be given the quotient norm and the positive cone which is the image of the positive cone of V . Then V/I is an R -space.*

Let $\tilde{C}$ be the positive cone in V/I so that $\tilde{C} = \pi C$ where $\pi: V \rightarrow V/I$ is the quotient map. Then $\tilde{C} \cap (-\tilde{C}) = \{0\}$. For let $x, y \in C$ and $\pi x = -\pi y$. Then $\pi(x+y) = 0$ so that $x+y \in I$. Then $x, y \in I$ so that $\pi x = \pi y = 0$.

It is clear that V/I is a positively generated partially ordered space, but we do not yet show that its positive cone is closed.

Suppose $0 \leq \tilde{y} \in V/I$. For some $x \in V$ we have $\pi x = \tilde{y}$ and $\|x\| < \|\tilde{y}\| + \varepsilon/2$. There is also some $z \in V$ with $z \geq 0$ and $\pi z = \tilde{y}$. As I is positively generated there is some $v \in V$ such that $v \geq x, z$ and $\pi v = \tilde{y}$. Also as V is regular there is some $w \in V$ such that $w \geq x, 0$ and $\|w\| < \|x\| + \varepsilon/2$. If now $y \in V$ is chosen so that $0, x \leq y \leq w, v$ then $0 \leq y \in V, \pi y = \tilde{y}$ and $\|y\| < \|\tilde{y}\| + \varepsilon$.

Suppose $0 \leq \tilde{y} \leq \tilde{z} \in V/I$ and

$$0 \leq a \in V \text{ has } \pi a = \tilde{y},$$

$$0 \leq b \in V \text{ has } \pi b = \tilde{z} - \tilde{y},$$

$$0 \leq c \in V \text{ has } \pi c = \tilde{z} \quad \text{and} \quad \|c\| < \|\tilde{z}\| + \varepsilon.$$

Then $\pi(a+b)=\pi c=\tilde{z}$. As I is an ideal we can find $z \in V$ such that $0 \leq z \leq a+b, c$ and $\pi z=\tilde{z}$. By the Riesz decomposition property we can find $y, x \in V$ such that $0 \leq y \leq a$ and $0 \leq x \leq b$ and $x+y=z$. Then $0 \leq \pi y \leq \tilde{y}$ and $0 \leq \pi x \leq \tilde{z}-\tilde{y}$ and $\pi y+\pi x=\tilde{z}$. Thus $\pi y=\tilde{y}$. We now have $0 \leq y \leq z, \pi y=\tilde{y}, \pi z=\tilde{z}$ and $\|z\| \leq \|c\| < \|\tilde{z}\| + \varepsilon$. We immediately obtain

$$\|\tilde{y}\| \leq \|y\| \leq \|z\| < \|\tilde{z}\| + \varepsilon$$

and, as $\varepsilon > 0$ is arbitrary, so $\|\tilde{y}\| \leq \|\tilde{z}\|$.

Suppose $-\tilde{z} \leq \tilde{y} \leq \tilde{z} \in V/I$. Then $0 \leq \tilde{y} + \tilde{z} \leq 2\tilde{z}$. We can from the previous paragraph find $0 \leq u \leq v \in V$ with $\pi u = \tilde{y} + \tilde{z}, \pi v = 2\tilde{z}$ and $\|v\| < 2\|\tilde{z}\| + \varepsilon$. Then $-v/2 \leq u - v/2 \leq v/2$ and $\pi(u - v/2) = \tilde{y}$. Thus $\|\tilde{y}\| \leq \|u - v/2\| \leq \|v/2\| < \|\tilde{z}\| + \varepsilon/2$. Thus $\|\tilde{y}\| \leq \|\tilde{z}\|$, and we have shown that V/I is a regular partially ordered Banach space.

We can now show that the positive cone in V/I is closed. Let $\tilde{x}_n \in V/I$ where $\sum \|\tilde{x}_n\| < \infty$ and $\sum_{r=1}^n \tilde{x}_r \geq 0$ for all n . Let $\tilde{x}_n = \tilde{a}_n - \tilde{b}_n$ where $\tilde{a}_n \geq 0, \tilde{b}_n \geq 0$ and $\sum \|\tilde{a}_r\| < \infty, \sum \|\tilde{b}_r\| < \infty$. We have $\sum_{r=1}^n \tilde{a}_r \geq \sum_{r=1}^n \tilde{b}_r$. Now let $0 \leq a_n \in V$ where $\pi a_n = \tilde{a}_n$ and $\sum \|a_n\| < \infty$. $\sum a_n$ converges in V to a limit $a \geq 0$. We now construct $0 \leq b_n \in V$ such that $\sum_{r=1}^n b_r \leq \sum_{r=1}^n a_r$ and $\|b_n\| < 2\|\tilde{b}_n\|$. Suppose $b_1, \dots, b_{n-1}$ have been so constructed. As $\tilde{b}_n \geq 0$ so we can find $0 \leq c_n \in V$ with $\pi c_n = \tilde{b}_n$ and $\|c_n\| < 2\|\tilde{b}_n\|$. As

$$\tilde{b}_{n+1} \leq \sum_{r=1}^n \tilde{a}_r - \sum_{r=1}^{n-1} \tilde{b}_r,$$

so we can find $d_n \in V$ with $\pi d_n = \tilde{b}_n$ and

$$d_n \leq \sum_{r=1}^n a_r - \sum_{r=1}^{n-1} b_r = k_n.$$

We can now find $e_n \in V$ with $e_n \leq c_n, d_n$ and $\pi e_n = \tilde{b}_n$. As $0, e_n \leq c_n, k_n$, so we can find $b_n \in V$ with $0, e_n \leq b_n \leq c_n, k_n$. Then $0 \leq b_n, \|b_n\| \leq \|c_n\| < 2\|\tilde{b}_n\|, \pi b_n = \tilde{b}_n$ and

$$\sum_{r=1}^n b_r \leq \sum_{r=1}^n a_r \leq a.$$

It is now clear that $\sum_{r=1}^n b_r \rightarrow b \leq a$. Then $\sum_{r=1}^n (a_r - b_r) \rightarrow a - b \geq 0$. Thus $\sum_{r=1}^n \tilde{x}_r \rightarrow x = \pi(a - b) \geq 0$, and we have shown V/I has a closed cone.

Finally we show that V/I has the Riesz separation property. Suppose $\tilde{a}, \tilde{b} \leq \tilde{c}, \tilde{d}$ and let $\pi a = \tilde{a}, \pi b = \tilde{b}$. We can find $c, d \geq a, b$ with $\pi c = \tilde{c}$ and $\pi d = \tilde{d}$. By the Riesz separation property for V we can find $e \in V$ with $a, b \leq e \leq c, d$ and if $\tilde{e} = \pi e$ we have $\tilde{a}, \tilde{b} \leq \tilde{e} \leq \tilde{c}, \tilde{d}$. This concludes the proof of the theorem.

Another way of investigating closed ideals, pursued for simplex spaces in [11], is to characterise their annihilators in the dual space.

Returning to the notation of §2, we define a conical face in Δ as a closed convex set F with $0 \in F$ such that if $x, y \in \Delta$ and $\alpha x + \beta y \in \Delta$ for any $\alpha > 0, \beta > 0$ then $x, y \in F$.

THEOREM 5.5. *Let V be an R -space, V^* its dual and $\Delta = \{\phi \in V^* : \phi \geq 0 \text{ and } \|\phi\| \leq 1\}$. Then the maps $I \rightarrow I^0 \rightarrow I^0 \cap \Delta$ determine a one-one correspondence between the set of all closed ideals I in V , the set of all weak*-closed lattice ideals in V^* , and the set of all conical faces in Δ . I may naturally be identified with the space of all continuous affine functionals f on Δ with $f|_{I^0 \cap \Delta} = 0$.*

It is elementary to show that if I is a closed ideal then I^0 is a weak*-closed lattice ideal and $F = I^0 \cap \Delta$ is a conical face. Now let F be a conical face. Let J be the subspace of V^* generated by Δ . Then J is obviously an ideal and it meets the unit ball B in a weak*-closed set. Specifically,

$$J \cap B = (1 + \varepsilon)(\Delta - \Delta) \cap B.$$

By a well-known theorem on Banach spaces [18] it follows that J is a weak*-closed subspace. J is a lattice ideal with $J \cap \Delta = F$. Let I_1 be defined as the subspace of V such that $f \in I$ and $\phi \in F$ implies $(f, \phi) = 0$. Then $I_1 = {}^0J$ is a closed subspace of $V = A^0(\Delta)$. I_1 determines and is determined by Δ , and if Δ is derived from a closed ideal I then $I_1 = I$.

All we need to show is that I_1 is always an ideal. It is clearly an order ideal. Suppose $f \in I_1$. Let g be the function $g: \Delta \rightarrow (-\infty, \infty]$ defined by

$$\begin{aligned} g(x) &= 0 & \text{if } x \in F, \\ &= \infty & \text{if } x \notin F. \end{aligned}$$

Then g is l.s.c. affine and $f, -f \leq g$. If we can find $h \in A_0(\Delta)$ with $f, -f \leq h \leq g$, then $h|_F = 0$, so we see that I_1 is positively generated and so is an ideal. The theorem is completed by the following separation theorem.

THEOREM 5.6. *Let V be an R -space and*

$$\Delta = \{\phi \in V^* : \phi \geq 0, \|\phi\| \leq 1\}.$$

If $-f_1, \dots, -f_n, g_1, \dots, g_m$ are l.s.c. affine functionals on Δ vanishing at the origin such that $f_i \leq g_j$ for all i, j , then we can find $h \in A_0(\Delta)$ with $f_i \leq h \leq g_j$ for all i, j .

First observe that if $k_1, \dots, k_p$ are u.s.c. affine functionals vanishing at the origin then $(k_1 \vee \dots \vee k_p)^\wedge$ defined as in Lemma 2.7 is easily shown to be u.s.c. affine. The result now follows by a simple application of the technique of forcing convergence developed in the first part of the proof of Theorem 3.1.

REFERENCES

1. T. Andô, *On fundamental properties of a Banach space with a cone*, Pacific J. Math. **12** (1962), 1163–1169.
2. H. Bauer, *Geordnete Gruppen mit Zerlegungseigenschaft*, Bayer. S.-B. Akad. Wiss. Math.-Natur. Kl. **21** (1958), 25–35.
3. ———, *Konvexität in topologischen Vektorräumen*, Lecture notes, Univ. of Hamburg, Hamburg, 1963/1964.

4. G. Choquet and P.-A. Meyer, *Existence et unicité des représentations intégrales dans les convexes compacts quelconques*, Ann. Inst. Fourier (Grenoble) **13** (1963), 139–154.
5. G. Choquet, *Séminaire du théorie du potentiel*, Faculté des Sciences de Paris, fasc. 2 (1961/1962), 12.01–12.15.
6. E. B. Davies, *On the Banach duals of certain spaces with the Riesz decomposition property*, Quart. J. Math. Oxford Ser. (2) **18** (1967), 109–111.
7. M. M. Day, *Normed linear spaces*, Springer-Verlag, Berlin, 1958.
8. J. Dixmier, *Sur certains espaces considérés par M. H. Stone*, Summa Brasil. Math. **2** (1951), 151–182.
9. D. A. Edwards, *Séparations des fonctions réelles définies sur un simplexe de Choquet*, C. R. Acad. Sci. Paris **261** (1965), 2798–2800.
10. ———, *On the homeomorphic affine embedding of a locally compact cone into a Banach dual space endowed with the vague topology*, Proc. London Math. Soc. **14** (1964), 399–414.
11. E. G. Effros, *Structure in simplexes*, Lecture notes, Aarhus Univ., Aarhus, Denmark, 1965.
12. A. J. Ellis, *The duality of partially ordered normed linear spaces*, J. London Math. Soc. **39** (1964), 730–744.
13. R. V. Kadison, *Transformations of states in operator theory and dynamics*, Topology **3** (1964), 177–198.
14. S. Kakutani, *Concrete representations of abstract M -spaces*, Ann. of Math. **42** (1941), 994–1024.
15. J. Lindenstrauss, *Extension of compact operators*, Mem. Amer. Math. Soc., No. 48, 1964, 112 pp.
16. I. Namioka, *Partially ordered linear topological spaces*, Mem. Amer. Math. Soc., No. 24, 1957.
17. R. R. Phelps, *Lectures on Choquet's theorem*, Van Nostrand, New York, 1966.
18. R. P. Robertson and W. J. Robertson, *Topological vector spaces*, Cambridge Tracts in Math. and Math. Phys., No. 53, Cambridge Univ. Press., 1964.

OXFORD UNIVERSITY,
OXFORD, ENGLAND