COMMUTATORS MODULO THE CENTER IN A PROPERLY INFINITE VON NEUMANN ALGEBRA⁽¹⁾

BY
HERBERT HALPERN

1. Introduction. An element C in a von Neumann algebra $\mathcal{A}$ is said to be a commutator in $\mathcal{A}$ if there are elements A and B in $\mathcal{A}$ such that $C = AB - BA$. For finite homogeneous discrete algebras and for properly infinite factor algebras the set of commutators has been completely described [1]–[5], [10]. In each of these special cases any element C is a commutator modulo a central element depending on C . In this paper we show that given any element C in a properly infinite von Neumann algebra $\mathcal{A}$ there is an element C_0 in the center of $\mathcal{A}$ depending on C such that $C - C_0$ is a commutator in $\mathcal{A}$. The element C_0 is an arbitrary element in the intersection $\mathcal{K}_C$ of the center with the uniform closure of the convex hull of $\{U^*CU \mid U \text{ unitary in } \mathcal{A}\}$ [6, III, §5]. We then present a few facts about those elements C such that $0 \in \mathcal{K}_C$ or what is the same as far as determining commutators is concerned about those elements C such that $0 \in \mathcal{K}_{S^{-1}CS}$ for some invertible S in $\mathcal{A}$.

2. Commutators. Let $\mathcal{A}$ be a C^* -algebra with identity and let I be a closed two-sided ideal in $\mathcal{A}$. The image of the element $A \in \mathcal{A}$ in the factor algebra $\mathcal{A}(I) = \mathcal{A}/I$ under the canonical homomorphism of $\mathcal{A}$ onto $\mathcal{A}/I$ will be denoted by $A(I)$. If ζ is a maximal ideal of the center of $\mathcal{A}$, the smallest closed two-sided ideal in $\mathcal{A}$ containing ζ is denoted by $[\zeta]$. For simplicity we write $A([\zeta])$ as $A(\zeta)$. The set of maximal (respectively, primitive) ideals of $\mathcal{A}$ with the hull-kernel topology is called the strong structure space (respectively, structure space) of $\mathcal{A}$. If $\mathcal{A}$ is a von Neumann algebra, then the strong structure space $M(\mathcal{A})$ of $\mathcal{A}$ is homeomorphic with the spectrum of the center $\mathcal{Z}$ of $\mathcal{A}$ under the map $M \rightarrow M \cap \mathcal{Z}$ [13]. This means $M(\mathcal{A})$ is extremely disconnected.

PROPOSITION 1. *Let $\mathcal{A}$ be a properly infinite von Neumann algebra and let A be a fixed element of $\mathcal{A}$. The function $M \rightarrow \|A(M)\|$ of the strong structure space $M(\mathcal{A})$ of $\mathcal{A}$ into the real numbers is continuous.*

Proof. For every $\alpha \geq 0$ we know that the set $X = \{M \in M(\mathcal{A}) \mid \|A(M)\| \leq \alpha\}$ is closed. If $I = \bigcap X$, then $\|A(I)\| \leq \alpha$ [8, Lemma 1.9] and so $\|A(M)\| \leq \alpha$ for every $M \in M(\mathcal{A})$ containing I . Thus $X = \{M \in M(\mathcal{A}) \mid I \subset M\}$.

Received by the editors September 27, 1968.

⁽¹⁾ The author was supported by the National Science Foundation.

Copyright © 1970, American Mathematical Society

Conversely, let $\alpha > 0$; we show that

$$Y = \{M \in M(\mathcal{A}) \mid \|A(M)\| \geq \alpha\}$$

is closed in $M(\mathcal{A})$. Let J be the strong radical of $\mathcal{A}$ and let $\mathcal{P}$ be the structure space of $\mathcal{A}(J)$. The set

$$Y' = \{K \in \mathcal{P} \mid \|A(J)(K)\| \geq \alpha\}$$

is compact (but not necessarily closed) in $\mathcal{P}$ [16, 4.9.18]. If $\mathcal{P}'$ is the structure space of $\mathcal{A}$, then $M \rightarrow M(J)$ defines a homeomorphism of $\{M \in \mathcal{P}' \mid M \supset J\} = h(J)$ onto $\mathcal{P}$ [16, 2.6.6]. But if $M \in \mathcal{P}'$, then the intersection of M with the center of $\mathcal{A}$ is a maximal ideal. So $M \in h(J)$ implies M is of the form $J + [\zeta]$ for some maximal ideal ζ of the center. It is then clear that $h(J)$ is the set of maximal ideals of $\mathcal{A}$ [10, Proposition 2.3]. Furthermore, the topology of $h(J)$ and $M(\mathcal{A})$ coincide. This proves that Y is compact in $M(\mathcal{A})$ since it is the inverse image of Y' under the homeomorphism $M \rightarrow M(J)$ of $M(\mathcal{A})$ onto $\mathcal{P}$. Because $M(\mathcal{A})$ is homeomorphic to the spectrum of the center which is Hausdorff, every compact set of $M(\mathcal{A})$ is closed. Thus Y is a closed subset of $M(\mathcal{A})$. Q.E.D.

REMARK. If $\mathcal{A}$ is not properly infinite, Proposition 1 is certainly not true.

Let H be a Hilbert space and let A be a bounded linear operator on H . Let F be a projection on H . Define the numerical gauge $\eta_A(F)$ to be

$$\eta_A(F) = \text{lub} \{\|Ax - (Ax, x)x\| \mid x \text{ is a unit vector in } F(H)\}.$$

Let $\mathcal{W}_A(F)$ be the closure of the convex set

$$\{(Ax, x) \mid x \text{ a unit vector in } F(H)\}.$$

For every $\alpha \in \mathcal{W}_A(F)$ we have that

$$(1) \quad \|(A - \alpha)F\| \leq 65\eta_A(F).$$

This can be obtained by a simple reworking of Lemma 2.3 [2].

Let $\mathcal{A}$ be a properly infinite von Neumann algebra with no σ -finite type III direct summands; then for each projection F in $\mathcal{A}$ and each element A in $\mathcal{A}$ define $\nu_A(F)$ to be

$$(2) \quad \nu_A(F) = \text{lub} \{\|AE - EAE\| \mid E \in (J), E \leq F\}$$

where (J) is the set of projections in the strong radical J of $\mathcal{A}$. For every irreducible representation ϕ of $\mathcal{A}$ on a Hilbert space such that $\phi(J) \neq (0)$ we have that

$$(3) \quad \eta_{\phi(A)}(\phi(F)) \leq \nu_A(F)$$

[10, Proposition 3.1]. Define $\nu(A)$ to be

$$\nu(A) = \text{glb} \{\nu_A(F) \mid 1 - F \in (J)\}.$$

Let $\mathcal{A}$ now be the product of σ -finite type III algebras; let

$$\nu(A) = \text{lub} \{ \|AE - EAE\| \mid E \text{ a projection in } \mathcal{A} \}$$

for each $A \in \mathcal{A}$. If A is in the complement in $\mathcal{A}$ of the set of all elements of $\mathcal{A}$ equal to scalar (zero included) multiples of the identity modulo some maximal ideal of $\mathcal{A}$, then there is a $\nu > 0$ such that $\nu(AP) \geq \nu$ for every nonzero central projection P since there is a projection E in $\mathcal{A}$ with $E \sim 1 - E \sim 1$ such that $EA^*(1 - E)AE \geq \alpha E$ for some scalar $\alpha > 0$ [10, Theorem 3.7]. Also it is easy to see from Proposition 3.1 [10] that

$$(4) \quad \eta_{\phi(A)}(1) \leq \nu(A)$$

for every irreducible representation of $\mathcal{A}$.

LEMMA 2. Let $\mathcal{A}$ be a properly infinite von Neumann algebra such that

- (i) either $\mathcal{A}$ has no σ -finite type III direct summands or
- (ii) $\mathcal{A}$ is a product of σ -finite type III algebras.

Let $\{P_n\}$ be a set of mutually orthogonal central projections of $\mathcal{A}$ of sum P . Then $\nu(AP) = \text{lub}_n \nu(AP_n)$ for every $A \in \mathcal{A}$.

Proof. Suppose $\mathcal{A}$ satisfies condition (i). Let (J) be the set of projections in the strong radical of $\mathcal{A}$. Then given $\varepsilon > 0$ there is a projection F with $1 - F \in (J)$ such that

$$\nu(AP) + \varepsilon \geq \text{lub} \{ \|(1 - E)APE\| \mid E \leq F, E \in (J) \}.$$

But

$$\|(1 - E)AEP_n\| \leq \|(1 - E)APE\|$$

for every P_n . So $\nu(AP_n) \leq \nu(AP) + \varepsilon$ for every n . Since $\varepsilon > 0$ is arbitrary, we have that $\text{lub}_n \nu(AP_n) \leq \nu(AP)$. Conversely, given that $\varepsilon > 0$ there is for each P_n a projection F_n with $1 - F_n \in (J)$ such that

$$\text{lub} \{ \|(1 - E)AP_nE\| \mid E \leq F_n, E \in (J) \} \leq \nu(AP_n) + \varepsilon.$$

Setting $F = \sum F_n P_n$, we have that $P - F \in (J)$ [10, Corollary, Proposition 2.2] and that

$$\|(1 - E)APE\| = \text{lub}_n \|(1 - E)AP_nE\| \leq \text{lub}_n \nu(AP_n) + \varepsilon$$

for every E in (J) with $E \leq F$. Thus $\nu(AP) \leq \text{lub}_n \nu(AP_n) + \varepsilon$. Since $\varepsilon > 0$ is arbitrary, we have that $\nu(AP) \leq \text{lub}_n \nu(AP_n)$. This completes the proof for case (i).

The proof for case (ii) is similar.

LEMMA 3. Let $\mathcal{A}$ be a properly infinite von Neumann algebra with no σ -finite type III direct summands. Let G be a projection in $\mathcal{A}$ such that the orthogonal complement $1 - G$ of G is in the set of projections (J) of the strong radical of $\mathcal{A}$. Let (J') be the strong radical of the reduced algebra $G\mathcal{A}G$. For any element A in $\mathcal{A}$ we have that

$$\nu'(GAG) = \text{glb} \{ \text{lub} \{ \|(1 - E)GAGE\| \mid E \leq F, E \in (J') \} \mid G - F \in (J') \}$$

is equal to $\nu(A)$.

Proof. First we show that $(J') = \{E \in (J) \mid E \leq G\}$. Suppose $E \notin (J')$ for some $E \in (J)$ majorized by G . There is a nonzero projection P in the center of $G\mathcal{A}G$ such that $PE \sim P$ [10, §2]. But $P = QG$ for some nonzero Q in the center of $\mathcal{A}$ [6, I, §2, Corollary, Proposition 2]. Then $QE \sim QG \sim Q$ since $G \sim 1$ [cf. 10, §2]. So $E \notin (J)$. This is a contradiction. Hence $\{E \in (J) \mid E \leq G\} \subset (J')$. Conversely, suppose $E \in (J')$. If Q is a central projection in $\mathcal{A}$ with $EQ \sim Q$, then $EQ \sim Q \sim QG$. This implies $Q = 0$. So $(J') \subset \{E \in (J) \mid E \leq G\}$.

Now let $\varepsilon > 0$ be given. There is by relation (2) a projection F in $\mathcal{A}$ with $1 - F \in (J)$ such that $\nu_A(F) \leq \nu(A) + \varepsilon$. Let G' be the least upper bound of $1 - G$ and $1 - F$. Then $G' \in (J)$ and $1 - G' \leq F$ and $1 - G' \leq G$. We see that

$$G - (1 - G') = G' - (1 - G) \in (J')$$

by the first paragraph. So

$$\begin{aligned} \nu'(GAG) &\leq \text{lub} \{ \|(1 - E)GAGE\| \mid E \in (J'), E \leq 1 - G' \} \\ &= \text{lub} \{ \|G(1 - E)AE\| \mid E \in (J), E \leq 1 - G' \} \\ &\leq \text{lub} \{ \|(1 - E)AE\| \mid E \in (J), E \leq F \} \leq \nu(A) + \varepsilon. \end{aligned}$$

Since $\varepsilon > 0$ is arbitrary, we have that $\nu'(GAG) \leq \nu(A)$.

Conversely, let $\varepsilon > 0$ be given; there is a projection $F \in G\mathcal{A}G$ with $G - F \in (J')$ and

$$\text{lub} \{ \|(G - E)AE\| \mid E \leq F, E \in (J') \} \leq \nu'(GAG) + \varepsilon.$$

The domain support G' of $(1 - G)A$ is a projection in (J) since $G' < 1 - G$ [6, III, §1, Proposition 2]; similarly, the domain support G'' of $G'F$ is a projection in (J) majorized by F . So $1 - (F - G'') = (1 - G) + (G - F) + G''$ is a projection in (J) . Then

$$\nu(A) \leq \text{lub} \{ \|(1 - E)AE\| \mid E \leq F - G'', E \in (J) \}.$$

But

$$GAGE = AE - (1 - G)AE = AE - (1 - G)AG'FE = AE - (1 - G)AG'FG''E = AE.$$

So we see that

$$\begin{aligned} \nu(A) &\leq \text{lub} \{ \|(1 - E)GAGE\| \mid E \leq F - G'', E \in (J') \} \\ &\leq \text{lub} \{ \|(1 - E)GAGE\| \mid E \leq F, E \in (J') \} \leq \nu'(GAG) + \varepsilon. \end{aligned}$$

Since $\varepsilon > 0$ is arbitrary, we have that $\nu(A) \leq \nu'(GAG)$. Q.E.D.

THEOREM 4. *Let C be an element in a properly infinite von Neumann algebra $\mathcal{A}$ and let C_0 be an element in the intersection $\mathcal{K}_C$ of the center of $\mathcal{A}$ with the uniform closure of the convex hull of $\{U^*CU \mid U \text{ unitary in } \mathcal{A}\}$. Then $C - C_0$ is a commutator in $\mathcal{A}$.*

Proof. We first make a preliminary reduction. There is a maximal central projection P in $\mathcal{A}$ such that $(C - C_0)P$ is in the strong radical J of $\mathcal{A}$ [10, Corollary,

Proposition 2.2]. However, the structure of the strong radical of $\mathcal{A}$ allows us to conclude that $(C - C_0)P$ is in the strong radical of $\mathcal{A}P$ [10, §2]. Since $(C - C_0)P$ is a commutator in $\mathcal{A}P$, it is necessary to prove that $(C - C_0)(1 - P)$ is a commutator in $\mathcal{A}(1 - P)$. It is easy to see that the uniform closure of the convex hull of $\{U^*(C - C_0)U \mid U \text{ unitary in } \mathcal{A}(1 - P)\}$ contains 0. So without loss of generality we may assume that C is an element in $\mathcal{A}$ such that $0 \in \mathcal{K}_C$ and such that $CP \in J$ for some central projection P in $\mathcal{A}$ implies that $P = 0$.

Now there is a central projection Q in $\mathcal{A}$ such that

- (i) $\mathcal{A}Q$ has no σ -finite type III direct summands and
- (ii) $\mathcal{A}(1 - Q)$ is the product of σ -finite type III algebras.

It is clearly necessary to prove only that CQ and $C(1 - Q)$ are commutators in $\mathcal{A}Q$ and $\mathcal{A}(1 - Q)$ respectively. Here $0 \in \mathcal{K}_{CQ}$ and $0 \in \mathcal{K}_{C(1 - Q)}$ is also true when these sets are formed relative to $\mathcal{A}Q$ and $\mathcal{A}(1 - Q)$ respectively. In the ensuing paragraphs we shall assume that either $\mathcal{A}$ satisfies condition (i) or condition (ii).

Let $\|C\| = \alpha$. We construct by induction a sequence $\{P_n\}$ of mutually orthogonal central projections of sum 1 such that for each nonzero central projection P majorized by P_n the number $\nu(CP)$ lies in the real interval $[2^{-n}\alpha, 2^{-n+1}\alpha]$. The induction hypothesis may be stated as follows: let $P_0 = 0$; then P_n is the largest central projection majorized by $1 - \sum \{P_k \mid 0 \leq k \leq n-1\}$ such that $\nu(CP) \in I_n$ for every nonzero central projection P majorized by P_n . Suppose we have constructed $P_1, P_2, \dots, P_n$. We find P_{n+1} . We may assume $R = 1 - \sum \{P_k \mid 0 \leq k \leq n\}$ is nonzero. There is a maximal ideal M in the strong structure space $M(\mathcal{A})$ of $\mathcal{A}$ such that $CR(M)$ is not in the center of $\mathcal{A}(M)$. Indeed, suppose $CR(M)$ is in the center of $\mathcal{A}(M)$ for every $M \in M(\mathcal{A})$. Given $\varepsilon > 0$ there are unitary operators $U_1, U_2, \dots, U_m$ in $\mathcal{A}$ and positive scalars $\alpha_1, \alpha_2, \dots, \alpha_m$ of sum 1 such that $\|\sum \alpha_i U_i^* C U_i\| < \varepsilon$ since $0 \in \mathcal{K}_C$. Thus

$$\|CR(M)\| = \left\| \left(\sum \alpha_i U_i^* C U_i \right) (M) \right\| < \varepsilon$$

for every $M \in M(\mathcal{A})$. Because $\varepsilon > 0$ is arbitrary, we have that $CR(M) = 0$ for every $M \in M(\mathcal{A})$. This means that CR is in the strong radical of $\mathcal{A}$. This is contrary to the choice of C . Hence, we must conclude that $CR(M)$ is not in the center of $\mathcal{A}(M)$ for at least one M in $M(\mathcal{A})$. Then there is a projection E in $\mathcal{A}$ such that $\|(1 - E)CRE(M)\| \neq 0$. By the continuity of $M' \rightarrow \|(1 - E)CRE(M')\|$ on $M(\mathcal{A})$ (Proposition 1), there is an open and closed neighborhood X of M in $M(\mathcal{A})$ such that for every $M' \in X$ the element $CR(M')$ is not in the center of $\mathcal{A}(M')$. Let Q be the nonzero central projection of $\mathcal{A}$ which determines X by the relation $X = \{M' \in M(\mathcal{A}) \mid Q \notin M'\}$ [13]. Then Q is majorized by R and $CQ(M')$ is not in the center of $\mathcal{A}Q(M')$ for every M' in the strong structure space $M(\mathcal{A}Q)$ of the algebra $\mathcal{A}Q$. The latter is true because $M' \rightarrow M'Q$ defines a homeomorphism of

$$\{M' \in M(\mathcal{A}) \mid Q \notin M'\} \text{ onto } M(\mathcal{A}Q) \text{ [16, Theorem 2.6.6].}$$

Then by 3.1 and 3.7 [10] there is a $\nu > 0$ such that $\nu(CQ') \geq \nu$ for every nonzero central projection Q' majorized by Q . It is clearly immaterial whether $\nu(CQ')$ is evaluated in $\mathcal{A}Q$ or in $\mathcal{A}$. Let m be the smallest integer for which there is a nonzero central projection Q majorized by R such that $\nu(CQ') \geq 2^{-m}\alpha$ for every nonzero central projection Q' majorized by Q . We then have that $\nu(CQ') \in I_m$ for every such Q' . In fact by the choice of m the projection Q is easily seen to be the least upper bound of a set $\{Q_i\}$ of nonzero mutually orthogonal central projections which satisfy $2^{-m}\alpha \leq \nu(CQ_i) < 2^{-m+1}\alpha$. By Lemma 2 we may conclude that $\nu(CQ) \leq 2^{-m+1}\alpha$. So for every nonzero central projection Q' majorized by Q we have that $\nu(CQ') \leq \nu(CQ) \leq 2^{-m+1}\alpha$. Now if $n+1 < m$ set P_{n+1} equal to zero. If $m \leq n+1$, then $m=n+1$. Indeed suppose that $m < n+1$; the projection $P_m + Q \neq P_m$, and for any nonzero central projection Q' majorized by $P_m + Q$ we have that

$$\nu(CQ') = \text{lub} \{\nu(CP_m Q'), \nu(CQ Q')\} \in I_m.$$

This contradicts the definition of P_m . Therefore $m=n+1$. Now we argue as follows. Let $\{Q_n\}$ be a maximal set of nonzero mutually orthogonal central projections majorized by R such that $\nu(CQ') \in I_{n+1}$ for every nonzero central projection Q' majorized by some Q_i . Let $P_{n+1} = \sum Q_i$. It is clear that $P_{n+1} \leq R$. Because $\nu(CQ) = \text{lub}_i \nu(CQ Q_i)$ for any nonzero central projection Q majorized by P_{n+1} (Lemma 2) and since at least one projection $Q Q_i$ is nonzero, we have that $\nu(CQ) \geq 2^{-(n+1)}\alpha$. On the other hand $Q \leq R$ and so by the induction hypothesis there is a set $\{R_i\}$ of nonzero mutually orthogonal central projections of sum Q such that $\nu(CR_i) < 2^{-n}\alpha$ for each R_i . Thus $\nu(CQ) \leq 2^{-n}\alpha$ (Lemma 2). This proves that $\nu(CQ) \in I_{n+1}$. It is clear that P_{n+1} is the largest central projection majorized by R such that $\nu(CQ) \in I_{n+1}$ for every nonzero central projection majorized by P_{n+1} .

Suppose that the sequence $\{P_n\}$ with the required properties has been constructed by induction. We show that $\sum P_n = 1$. If $R = 1 - \sum P_n$, then for each $n=1, 2, \dots$ we may conclude that $\nu(CR) \leq 2^{-n}\alpha$ by performing the construction of the previous paragraph. This means that $\nu(CR) = 0$. The results of the previous paragraph show that $R=0$ by our choice of C . Hence $\sum P_n = 1$.

Let Z be the spectrum of the center of $\mathcal{A}$ and let $X_n = \{\zeta \in Z \mid P_n \notin \zeta\}$ for $n=1, 2, \dots$. Now suppose that $\mathcal{A}$ has no σ -finite type III direct summands. For each $n=1, 2, \dots$ there is a projection F_n in $\mathcal{A}P_n$ with $P_n - F_n \in J$ such that $\nu_{CP_n}(F_n) \leq 2\nu(CP_n)$ by definition. For each $\zeta \in Z$ there is an irreducible representation $\psi_\zeta = \psi$ of $\mathcal{A}$ on a Hilbert space whose kernel is the smallest closed two-sided ideal $[\zeta]$ of $\mathcal{A}$ which contains ζ [11, Theorem 4.7]. Then ψ does not annihilate J . Indeed J contains a projection E of central support 1, cf. [10, §2]. The two-valued continuous function $\zeta' \rightarrow \|E(\zeta')\|$ [9, Lemma 10] on Z assumes the value 0 on an open and closed set given by $\{\zeta' \in Z \mid R \notin \zeta'\}$ where R is a central projection. Then $\|(E(1-R)-E)(\zeta')\| = 0$ for every $\zeta' \in Z$. Since $\bigcap \{[\zeta'] \mid \zeta' \in Z\} = (0)$ cf. [9, §4, remarks preceding Lemma 9], we have that $E(1-R) = E$. But this means that $R=0$ since the central support of E is 1. This proves that $E(\zeta') \neq 0$ for every $\zeta' \in Z$.

and in particular $E(\zeta) \neq 0$. So ψ does not annihilate J . Thus relations (1) and (3) imply that

$$(5) \quad \|(\psi(CP_n) - \beta)\psi(F_n)\| \leq 65\eta_{\psi(CP_n)}(\psi(F_n)) \leq 65\nu_{CP_n}(F_n) \leq 130\nu(CP_n)$$

whenever $\beta \in \mathcal{W}_{\psi(CP_n)}(\psi(F_n)) = \mathcal{W}$ and $\zeta \in X_n$. We show that $0 \in \mathcal{W}$. Indeed, given $\varepsilon > 0$, there is a set $U_1, U_2, \dots, U_m$ of unitary elements in $\mathcal{A}$ and positive scalars $\alpha_1, \alpha_2, \dots, \alpha_m$ of sum 1 such that $\|\sum \alpha_i U_i^* C U_i\| < \varepsilon$. If G_i is the range projection of $U_i^*(P_n - F_n)$ for $i = 1, 2, \dots, m$, then G_i is a projection in J and $G = \text{lub } G_i$ is a projection in J . Thus the projection $P_n - G$ is equivalent to P_n . Let x be a unit vector in the subspace determined by $\psi(P_n - G)$. For each U_i we have that $\psi(U_i)x = y_i$ is in the orthogonal complement of the subspace determined by $\psi(P_n - F_n)$ and thus $\psi(F_n)y_i = y_i$. Therefore

$$\sum \alpha_i (\psi(CP_n)y_i, y_i) \in \mathcal{W}$$

since $\mathcal{W}$ is convex. But we have that

$$\left| \sum (\alpha_i \psi(CP_n)y_i, y_i) \right| \leq \|\psi\| \left\| \sum \alpha_i U_i^* C U_i \right\| \|x\|^2 < \varepsilon.$$

Because $\mathcal{W}$ is closed and because $\varepsilon > 0$ is arbitrary, we see that $0 \in \mathcal{W}$. The relation (5) now becomes

$$\|CP_n F_n(\zeta)\| = \|CF_n(\zeta)\| = \|\psi_\zeta(CP_n)\psi_\zeta(F_n)\| \leq 130\nu(CP_n),$$

for every $\zeta \in X_n$. The orthogonal complement of the projection $F = \sum F_n$ is in J since $P_n(1 - F) \in J$ for every $n = 1, 2, \dots$ [10, Proposition 2.2]. For every nonzero central projection P majorized by P_n we have that

$$\|CFP\| = \text{lub} \{ \|CF(\zeta)\| \mid P \notin \zeta, \zeta \in Z \} \leq 130\nu(CP_n) \leq 260\nu(CP).$$

So for any central projection P we have that

$$(6) \quad \begin{aligned} \|CFP\| &= \text{lub} \{ \|CFPP_n\| \mid n = 1, 2, \dots \} \\ &\leq \text{lub} \{ 260\nu(CPP_n) \mid n = 1, 2, \dots \} \\ &\leq 260\nu(CP) \end{aligned}$$

by Lemma 2.

Now for an algebra which is the product of σ -finite type III algebras, we may show that $\|CP\| \leq 260\nu(CP)$ for any central projection P . The proof is entirely similar to that just given except that relation (4) replaces relations (1) and (3).

Now let us suppose that $\mathcal{A}$ has no σ -finite type III direct summands. Let $D = FCF$ and let $\mathcal{B}$ be the von Neumann algebra $F\mathcal{A}F$. By setting $Q_n = P_n F$ we obtain a sequence $\{Q_n\}$ of mutually orthogonal central projections in $\mathcal{B}$ of sum F such that $\nu(DQ) \in I_n$ for any nonzero central projection Q in $\mathcal{B}$ majorized by Q_n . Here $\nu(DQ)$ is evaluated in $\mathcal{B}$ and Lemma 3 is employed. By relation (6) we see that

$\{\|2^n DQ_n\|\}$ is a bounded sequence and hence $B = \sum 2^n DQ_n$ defines an element of $\mathcal{B}$ such that $\nu(BQ) \geq \alpha$ for every nonzero central projection Q in $\mathcal{B}$. Indeed,

$$\begin{aligned}\nu(BQ) &= \text{lub } \{\nu(BQ_n) \mid n = 1, 2, \dots\} \\ &= \text{lub } \{2^n \nu(DQ_n) \mid n = 1, 2, \dots\} \geq \alpha\end{aligned}$$

since at least one projection Q_n is nonzero. There is an invertible element S in $\mathcal{B}$ and a projection G in $\mathcal{B}$ with $F \sim G \sim F - G$ such that $U^*S^{-1}BSU = 0$ and $V^*S^{-1}BSV$ is a commutator in $\mathcal{B}$. Here U and V are partial isometries in B such that $U^*U = V^*V = F$, $UU^* = G$ and $VV^* = F - G$ [10, Theorem 3.6]. By multiplying both $U^*S^{-1}BSU$ and $V^*S^{-1}BSV$ by the central element $\sum 2^{-n}Q_n$ we see that $U^*S^{-1}DSU = 0$ and $V^*S^{-1}DSV$ is a commutator in $\mathcal{B}$. Now let $T = S + (1 - F)$ in $\mathcal{A}$; the element T is invertible with inverse $T^{-1} = S^{-1} + (1 - F)$ where S^{-1} still denotes the inverse of S in $\mathcal{B}$. Let W be a partial isometry in $\mathcal{A}$ with domain support 1 and range support F . Then $V_1 = VW$ is a partial isometric operator of domain support 1 and range support $F - G$. Then it is easy to see that $V_1^*T^{-1}CTV_1 = W^*V^*S^{-1}DSVW$ is a commutator in $\mathcal{A}$. We have that $1 \sim F \sim G \leq G + (1 - F)$. Thus there is a partial isometry U_1 in $\mathcal{A}$ with domain support 1 and range support $G + (1 - F)$. Then

$$U_1^*T^{-1}CTU_1 = U_1^*T^{-1}CT(1 - F)U_1 + U_1^*(1 - F)T^{-1}CTGU_1$$

is an element of the strong radical of $\mathcal{A}$ and therefore, is a commutator in $\mathcal{A}$ [10, Theorem 2.5]. We have proved that there is an isomorphism of $\mathcal{A}$ onto the algebra $\mathcal{A}_2$ of 2×2 matrices over $\mathcal{A}$ which carries $T^{-1}CT$ into the matrix (B_{ij}) where B_{11} and B_{22} are commutators in $\mathcal{A}$. But this matrix is a commutator in the algebra $\mathcal{A}_2$. Indeed, let $B_{11} = S_{11}T_{11} - T_{11}S_{11}$ and $B_{22} = S_{22}T_{22} - T_{22}S_{22}$ for S_{11} , S_{22} , T_{11} , T_{22} in $\mathcal{A}$. We may assume that S_{11} and S_{22} have disjoint spectra since $B_{11} = (S_{11} + \beta)T_{11} - T_{11}(S_{11} + \beta)$ for a scalar β . There is an operator T_{12} and an operator T_{21} in $\mathcal{A}$ such that $S_{11}T_{12} - T_{12}S_{22} = B_{12}$ and $S_{22}T_{21} - T_{21}S_{11} = B_{21}$ [12]. Setting $S_{21} = S_{12} = 0$, we find by direct calculation that $(S_{ij})(T_{ij}) - (T_{ij})(S_{ij}) = (B_{ij})$ in $\mathcal{A}_2$. This proves (B_{ij}) is a commutator in $\mathcal{A}_2$ and $T^{-1}CT$ is a commutator in $\mathcal{A}$.

Now if $\mathcal{A}$ is the product of σ -finite type III algebras, the preceding paragraph allows us to conclude that there is an invertible S in $\mathcal{A}$ such that $S^{-1}CS$ may be identified with the 2×2 matrix (B_{ij}) over $\mathcal{A}$ with $B_{11} = 0$ and B_{22} a commutator in $\mathcal{A}$. So (B_{ij}) is a commutator in the 2×2 matrices over $\mathcal{A}$ and $S^{-1}CS$ is a commutator in $\mathcal{A}$. Q.E.D.

3. Elements C with $0 \in \mathcal{K}_C$. The construction of Theorem 4 actually depended upon choosing a central element C_0 corresponding to a given element C in a properly infinite von Neumann algebra $\mathcal{A}$ such that $ES^{-1}(C - C_0)SE = 0$ for some invertible S in $\mathcal{A}$ and some projection E in $\mathcal{A}$ equivalent to 1. The next proposition clarifies this choice.

PROPOSITION 5. *If C is an element in a properly infinite von Neumann algebra $\mathcal{A}$ such that $ECE=0$ for some projection E in $\mathcal{A}$ equivalent to 1, then 0 is an element of the intersection $\mathcal{K}_C$ of the center of $\mathcal{A}$ with the uniform closure of the convex hull of the set $\{U^*CU \mid U \text{ unitary in } \mathcal{A}\}$.*

Proof. There are projections E' and E'' such that $E' \sim E'' \sim E$ and $E' + E'' = E$ [6, III, §8, Theorem 1, Corollary 2]. Then $1 \sim E' \leq E' + (1-E) \leq 1$ implies that $E' + (1-E) \sim 1$. So there is no loss of generality in supposing that $E \sim 1-E \sim 1$. The operator $U = E - (1-E)$ is unitary in $\mathcal{A}$ and

$$2^{-1}(C + U^*CU) = (1-E)C(1-E).$$

Now let $E_1, E_2, \dots, E_n$ be orthogonal projections of sum E such that $E_1 \sim \dots \sim E_n \sim E$. There are unitary operators $U_1, U_2, \dots, U_n$ in $\mathcal{A}$ such that $(1-E)U_j$ has domain support E_j for $j=1, 2, \dots, n$ since $E_j \sim 1-E$ and $1-E_j \sim E$ for $j=1, 2, \dots, n$. So we have that

$$\begin{aligned} & \left\| \sum \{n^{-1}U_j^*(1-E)C(1-E)U_j x \mid j=1, 2, \dots, n\} \right\|^2 \\ & \leq n^{-2} \|C\|^2 \sum \|E_j x\|^2 \leq n^{-2} \|C\|^2 \|x\|^2 \end{aligned}$$

for every x in the Hilbert space. Thus

$$\left\| \sum n^{-1}U_j^*(1-E)C(1-E)U_j \right\| \leq n^{-1} \|C\|.$$

This means that 0 is an element of the uniform closure of the convex hull of $\{V^*CV \mid V \text{ unitary in } \mathcal{A}\}$ because n is arbitrary. Hence $0 \in \mathcal{K}_C$. Q.E.D.

Let $\mathcal{A}$ be a properly infinite von Neumann algebra. If we could prove that $0 \in \bigcup \{\mathcal{K}_{S^{-1}CS} \mid S \text{ invertible in } \mathcal{A}\}$ for every commutator C in $\mathcal{A}$, then we would have a complete characterization of the set commutators. This characterization is certainly valid for factor algebras. Indeed C is a commutator in the properly infinite factor algebra $\mathcal{A}$ if and only if $\mathcal{A}$ is not a nonzero scalar multiple of the identity modulo the unique maximal ideal M of $\mathcal{A}$. If $C \in M$, then $0 \in \mathcal{K}_C$ [10, Proposition 2.4]. If C is not a scalar multiple of the identity modulo M , then the canonical form of Brown and Pearcy [2] in conjunction with the preceding proposition shows $0 \in \mathcal{K}_{S^{-1}CS}$ for some invertible S in $\mathcal{A}$. The characterization though is at odds with a conjecture that the set of commutators in $\mathcal{A}$ is the complement (F') in $\mathcal{A}$ of the set of all elements equal to a nonzero scalar multiple of the identity modulo some maximal ideal of $\mathcal{A}$ [4]. In fact let $\{P_n\}$ be a sequence of nonzero mutually orthogonal central projections of sum 1. (This presupposes that $\mathcal{A}$ has a sufficiently large center.) Then let E_n be a projection in $\mathcal{A}P_n$ such that $E_n \sim P_n \sim P_n - E_n$ ($n=1, 2, \dots$). Let $C = \sum (n^{-1}P_n + n^{-2}E_n)$. We have that

$$\bigcup_n \{M \in M(\mathcal{A}) \mid P_n \notin M\}$$

is dense in the strong structure space $M(\mathcal{A})$ of $\mathcal{A}$ by the remarks at the beginning of §2. Then $C(M) = n^{-1} \cdot 1(M) + n^{-2}E_n(M)$ for every M with $P_n \notin M$ and clearly

$C(M)$ is not a scalar (zero included) multiple of the identity. But $S^{-1}CS(M)$ is not a scalar multiple of the identity for every M with $P_n \notin M$. If for example $0 \in \mathcal{K}_S^{-1}CS$, then by the proof of Theorem 4 we would be able to find an invertible T and a projection E equivalent to 1 with $ET^{-1}CTE=0$. Thus

$$\|n^{-1}EP_n\| = n^{-2}\|ET^{-1}E_nTE\| \leq n^{-2}\|T^{-1}\| \|T\|$$

for each $n=1, 2, \dots$. This is obviously impossible.

It might be well to remark that there is no canonical matrix form with 0 on the diagonal in the sense of Brown and Pearcy [2] for operators of class (F') .

LEMMA 6. *Let C be an element in a von Neumann algebra $\mathcal{A}$. Let D_1 and D_2 be elements in $\mathcal{K}_C$ and let A be a central element of $\mathcal{A}$ with $0 \leq A \leq 1$. Then $AD_1 + (1-A)D_2 \in \mathcal{K}_C$.*

Proof. First let A be a projection in the center of $\mathcal{A}$. There are unitary operators $U_1, U_2, \dots, U_n$ (respectively $V_1, V_2, \dots, V_m$) and positive scalars $\alpha_1, \alpha_2, \dots, \alpha_n$ (respectively $\beta_1, \beta_2, \dots, \beta_m$) of sum 1 such that

$$\left\| \sum \alpha_i U_i^* C U_i - D_1 \right\| < \varepsilon \quad (\text{respectively, } \left\| \sum \beta_j V_j^* C V_j - D_2 \right\| < \varepsilon).$$

Here $\varepsilon > 0$ is a preassigned constant. Then $U'_i = U_i A + (1-A)$ ($1 \leq i \leq n$) and $V'_j = V_j (1-A) + A$ ($1 \leq j \leq m$) are unitary in $\mathcal{A}$ with the property

$$\left\| \sum \alpha_i \beta_j V_j'^* U_i'^* C U_i' V_j' - (D_1 A + D_2 (1-A)) \right\| < \varepsilon.$$

Since $\mathcal{K}_C$ is closed, we have that $AD_1 + (1-A)D_2 \in \mathcal{K}_C$.

Suppose that the restriction that A is a projection is removed. Let Z be the spectrum of the center of $\mathcal{A}$ and let $D^\wedge(\zeta)$ denote the Gelfand transform of the central element D evaluated at $\zeta \in Z$. Since $\mathcal{K}_C$ is convex, the set

$$\mathcal{K}_\zeta = \{D^\wedge(\zeta) \mid D \in \mathcal{K}_C\}$$

is convex and so

$$A^\wedge(\zeta) D_1^\wedge(\zeta) + (1-A)^\wedge(\zeta) D_2^\wedge(\zeta) \in \mathcal{K}_\zeta$$

for every $\zeta \in Z$. Thus there is for each $\varepsilon > 0$ a finite set $P_1, P_2, \dots, P_n$ of orthogonal central projections of sum 1 and corresponding elements $B_1, B_2, \dots, B_n$ in $\mathcal{K}_C$ such that

$$\|(AD_1 + (1-A)D_2 - B_j)P_j\| < \varepsilon$$

for $j=1, 2, \dots, n$. Since $\sum B_j P_j \in \mathcal{K}_C$ by the first paragraph and since $\varepsilon > 0$ is arbitrary, we see that $AD_1 + (1-A)D_2$ is in the closed set $\mathcal{K}_C$. Q.E.D.

The next proposition corresponds to a result of C. R. Putnam [14]. We use many of his calculations cf. [15, 1.5.1].

PROPOSITION 7. *Let A, B, C be elements in a properly infinite von Neumann algebra $\mathcal{A}$. If A is seminormal (i.e. if either $\pm(AA^* - A^*A)$ is positive) and if $C = AB - BA$, then $0 \in \mathcal{K}_S^{-1}CS$ for some invertible S in $\mathcal{A}$.*

Proof. There is a projection P in the center of $\mathcal{A}$ such that CP is in the strong radical of $\mathcal{A}P$ and

$$\{M \in M(\mathcal{A}) \mid C(1-P)(M) \neq 0\}$$

is dense in the subset $\{M \in M(\mathcal{A}) \mid 1-P \notin M\}$ of the strong structure space $M(\mathcal{A})$ of $\mathcal{A}$. Since $\mathcal{K}_C = \mathcal{K}_{CP} + \mathcal{K}_{C(1-P)} = \mathcal{K}_{C(1-P)}$ due to the fact that $\mathcal{K}_{CP} = \{0\}$ [10, Proposition 2.4] and since $\{M \in M(\mathcal{A}) \mid 1-P \notin M\}$ is identified with the strong structure space of $\mathcal{A}(1-P)$, we may assume that $\{M \in M(\mathcal{A}) \mid C(M) \neq 0\}$ is dense in $M(\mathcal{A})$.

As we have previously argued we may assume that either $\mathcal{A}$ has no σ -finite type III direct summands or that $\mathcal{A}$ is the product of σ -finite type III algebras.

Now for each nonzero real scalar α and each unitary element U in the center $\mathcal{Z}$ of $\mathcal{A}$ let

$$F(\alpha, U) = |\alpha|^{1/2}UA + |\alpha|^{-1/2}B^*.$$

Then

$$\operatorname{sgn}(\alpha)(FF^* - F^*F) = D(\alpha) + \operatorname{sgn}(\alpha)S(U)$$

where

$$D(\alpha) = \alpha(AA^* - A^*A) - \alpha^{-1}(BB^* - B^*B)$$

and $S(U) = UC + U^*C^*$. The set $\{|D| = (D^*D)^{1/2} \mid D \in \mathcal{K}_C\}$ is monotonely decreasing in $\mathcal{Z}$. Indeed, let D_1 and D_2 be elements of $\mathcal{K}_C$; there is a central projection P such that $|D_1|P \leq |D_2|P$ and $|D_2|(1-P) \leq |D_1|(1-P)$. But $D_1P + D_2(1-P) \in \mathcal{K}_C$ and $|D_1P + D_2(1-P)| = |D_1|P + |D_2|(1-P)$ is majorized by both $|D_1|$ and $|D_2|$. This proves that $\{|D| \mid D \in \mathcal{K}_C\}$ is monotonely decreasing. Let D_0 be the greatest lower bound of this set [6, Appendix II]. Suppose D_1 is a positive central element which majorizes $D(\alpha)$ for some α . We show that $D_0 \leq D_1$. If not, there is an $\varepsilon > 0$ and a nonzero central projection P such that $D_0P \geq (D_1 + \varepsilon)P$. By reducing to $\mathcal{A}P$ we may assume that $P=1$. Let $U_1, U_2, \dots, U_n$ be unitary elements in $\mathcal{Z}$ and let $\alpha_1, \alpha_2, \dots, \alpha_n$ be positive scalars of sum 1; then for any central element R in the sphere of radius $2^{-1}\varepsilon$ about 0 we have that

$$\left| \left(\sum \alpha_i U_i \right) D_1 + R \right|^2 \leq \left(\sum \alpha_i D_1 + |R| \right)^2 = (D_1 + |R|)^2 \leq (D_1 + 2^{-1}\varepsilon)^2.$$

Hence $(D^*Dx, x) \leq ((D_1 + 2^{-1}\varepsilon)^2x, x)$ for any D in the strong closure $\mathcal{K}''$ of the convex hull of the set

$$\{UD_1 + R \mid U, R \text{ in } \mathcal{Z}, U \text{ unitary}, \|R\| \leq 2^{-1}\varepsilon\}$$

and for any x in the Hilbert space H of $\mathcal{A}$. On the other hand we see that $(D^*Dx, x) \geq ((D_1 + \varepsilon)^2x, x)$ for any D in the strong closure $\mathcal{K}'$ of $\mathcal{K}_C$ and any x in H because $R^*R \geq (D_1 + \varepsilon)^2$ for every R in $\mathcal{K}_C$. By the standard separation theorem there is a nonzero strongly continuous functional f on $\mathcal{Z}$ such that

$$(7) \quad \operatorname{lub} \{\operatorname{Re} f(R) \mid R \in \mathcal{K}'\} \leq \operatorname{glb} \{\operatorname{Re} f(R) \mid R \in \mathcal{K}''\}.$$

Here $\operatorname{Re} \beta$ denotes the real part of the complex number β . Indeed the element 0 is not in the strong closure of $\mathcal{K}' - \mathcal{K}''$. Since f is also weakly continuous on $\mathcal{L}$ [6, I, §3, Theorem 1 (i)], there is a unitary U in $\mathcal{L}$ and a nonzero vector x in H such that $f(R) = (RUx, x)$ for every $R \in \mathcal{L}$ [17] and [6, III, §1, Corollary, Theorem 4]. Now let $F = F(\alpha, \operatorname{sgn}(\alpha)U)$. We have that

$$(8) \quad \operatorname{sgn}(\alpha)(FF^* - F^*F) = D(\alpha) + S(U) \leq D_1 + S(U).$$

Let D be an arbitrary element in $\mathcal{K}_{\operatorname{sgn}(\alpha)(FF^* - F^*F)} = \mathcal{K}$. There is an element S in $\mathcal{K}_{S(U)}$ such that $D \leq D_1 + S$ [6, III, §5, Problem 2a]. We may find a T in $\mathcal{K}_{U^*C - UC}$ such that

$$2^{-1}(S+T) \in \mathcal{K}_{UC} \quad \text{and} \quad 2^{-1}(S-T) \in \mathcal{K}_{U^*C}.$$

[6, III, §5, Problem 2a]. We then have that $2^{-1}U^*(S+T)$ and $2^{-1}U^*(S-T)^*$ are elements of $\mathcal{K}_C$. The latter is true because $\mathcal{K}_C = \{R^* \mid R \in \mathcal{K}_C\}$. From relation (7) we obtain that both $\operatorname{Re}(2^{-1}(S+T)x, x)$ and $\operatorname{Re}(2^{-1}(S-T)x, x)$ are majorized by $\operatorname{glb}\{\operatorname{Re} f(R) \mid R \in \mathcal{K}''\}$. Thus

$$(Sx, x) = \operatorname{Re}(Sx, x) \leq 2 \operatorname{glb}\{\operatorname{Re} f(R) \mid R \in \mathcal{K}''\}$$

since S is clearly selfadjoint. But $-U^*D_1 - 2^{-1}\varepsilon U^*$ is an element of $\mathcal{K}''$. So

$$(9) \quad 2^{-1}(Sx, x) \leq \operatorname{Re}(U(-U^*D_1 - 2^{-1}\varepsilon U^*)x, x) = -((D_1 + 2^{-1}\varepsilon)x, x).$$

Therefore,

$$(Dx, x) \leq (D_1x, x) - \varepsilon(x, x) - 2(D_1x, x) \leq -\varepsilon(x, x)$$

by relation (8). Using reasoning similar to that which we used to prove that $\{|R| \mid R \in \mathcal{K}_C\}$ is monotonely decreasing, we may prove that $\mathcal{K}$ is monotonely increasing. Setting $R_0 = \operatorname{lub} \mathcal{K}$, we see that $(R_0x, x) \leq -\varepsilon(x, x)$. We show that this is impossible by showing $(R_0x, x) \geq 0$. Indeed, in proving this then we may certainly assume that F is invertible and that $\alpha > 0$. Because F is invertible, there is a unitary operator V in $\mathcal{A}$ obtained from the polar decomposition of F [6, Appendix III] such that $V^*FF^*V = F^*F$. If $R \in \mathcal{K}_{F^*F}$, then there are unitary operators $U_1, U_2, \dots, U_n$ in $\mathcal{A}$ and positive scalars $\alpha_1, \alpha_2, \dots, \alpha_n$ of sum 1 such that

$$\left\| \sum \alpha_i U_i^* F^* F U_i - R \right\| < \varepsilon'$$

for any preassigned constant $\varepsilon' > 0$. But this means that

$$\left\| \sum \alpha_i (VU_i)^* FF^* (VU_i) - R \right\| < \varepsilon'.$$

Because $\varepsilon' > 0$ is arbitrary we have that $R \in \mathcal{K}_{FF^*}$. By symmetry it is then clear that $\mathcal{K}_{FF^*} = \mathcal{K}_{F^*F}$. Now for any $\varepsilon' > 0$ there is an element R_1 in $\mathcal{K}_{FF^*}$ such that

$$(R_1x, x) \geq (R_2x, x) - \varepsilon'(x, x)$$

where R_2 is the least upper bound of the monotonely increasing set $\mathcal{K}_{FF^*}$. But

there is an element $R \in \mathcal{K}_{F^*F}$ such that $R_1 - R \in \mathcal{K}$ [6, III, §5, Problem 2a]. However we have that

$$(R_0x, x) \geq ((R_1 - R)x, x) \geq (R_2x, x) - (Rx, x) - \varepsilon'(x, x) \geq -\varepsilon'(x, x)$$

since $R_2 \geq R$. Because $\varepsilon' > 0$ is arbitrary, we see that $(R_0x, x) \geq 0$. This is a contradiction. We must conclude that $D_0 \leq D_1$.

We now show that $D_0 = 0$. We may assume that $AA^* - A^*A \leq 0$. Then for $\alpha > 0$ we have that

$$D(\alpha) \leq -\alpha^{-1}(BB^* - B^*B) \leq 2\alpha^{-1}\|B\|^2.$$

Thus we see that $D_0 \leq 2\alpha^{-1}\|B\|^2$ for every $\alpha > 0$. Therefore $D_0 = 0$. If Q is a nonzero central projection in $\mathcal{A}$ and if $\varepsilon > 0$ there is a net $\{Q_n\}$ of mutually orthogonal central projections of sum Q such that each set $\mathcal{K}_{CQ_n}$ contains an element D_n of norm not exceeding ε . Indeed, if $\{Q_n\}$ is a maximal set of mutually orthogonal nonzero central projections majorized by Q with this property, then the assumption that $Q' = Q - \sum Q_n \neq 0$ gives a contradiction. Since $\text{glb } \{|D| \mid D \in \mathcal{K}_{CQ'}\} = 0$, there is a $D \in \mathcal{K}_{CQ'}$ such that $|D| \geq 2^{-1}\varepsilon Q'$ is not true. This means that there is a nonzero central projection Q'' majorized by Q' such that $|D|Q'' \leq \varepsilon Q''$. This contradicts the maximality of $\{Q_n\}$. Hence we have that $\sum Q_n = Q$.

Now suppose $\mathcal{A}$ has no σ -finite type III direct summands. In Theorem 4 we constructed a sequence $\{P_n\}$ of mutually orthogonal central projections of sum 1 and a projection F whose orthogonal complement $1 - F$ was in the strong radical such that $\|(C - R)FP\| \leq 260\nu(CP)$ whenever P is a central projection majorized by P_n and whenever $R \in \mathcal{K}_{CP}$. Also either $P_n = 0$ or $\nu(CP) \in [2^{-n}\|C\|, 2^{-n+1}\|C\|]$ for every nonzero central projection P majorized by P_n . By the preceding paragraph there is a set $\{P_{nj}\}$ of mutually orthogonal central projections of sum P_n such that each set $\mathcal{K}_{CP_{nj}}$ contains an element D_{nj} of norm not exceeding $\nu(CP_n)$. Then for each nonzero central projection P majorized by P_n we have that

$$\begin{aligned} \|CFP\| &= \text{lub}_j \|CFPP_{nj}\| \leq \text{lub } \|(C - D_{nj})PP_{nj}\| + \text{lub } \|D_{nj}PP_{nj}\| \\ &\leq 260\nu(CP) + \nu(CP_n) \leq 262\nu(CP) \end{aligned}$$

by relation (6). By the same reasoning as found in Theorem 4, we may find an invertible W in $\mathcal{A}$ such that $EW^{-1}CWE = 0$ for some projection E in $\mathcal{A}$ which is equivalent to 1. However, this means that $0 \in \mathcal{K}_{S^{-1}(F^*F - FF^*)S}$ by Proposition 5.

If A is the product of σ -finite type III algebras a similar proof holds. Q.E.D.

COROLLARY. *If F is an element in a properly infinite von Neumann algebra $\mathcal{A}$, then there is an invertible S in $\mathcal{A}$ such that $\mathcal{K}_{S^{-1}(F^*F - FF^*)S}$ contains 0.*

Proof. If $A = 2^{-1}i(F - F^*)$ and $B = 2^{-1}(F + F^*)$, then $2^{-1}i(FF^* - FF^*) = AB - BA$. Now Proposition 7 applies.

Added in proof (April 25, 1970). I have improved Proposition 7 by showing that $0 \in \mathcal{K}_C$.

BIBLIOGRAPHY

1. A. Brown, P. Halmos and C. Pearcy, *Commutators of operators on Hilbert space*, Canad. J. Math. **17** (1965), 695–708. MR **34** #3311.
2. A. Brown and C. Pearcy, *Structure of commutators of operators*, Ann. of Math. (2) **82** (1965), 112–127. MR **31** #2612.
3. ———, *Commutators in factors of type III*, Canad. J. Math. **18** (1966), 1152–1160. MR **34** #1864.
4. A. Brown, C. Pearcy and D. Topping, *Commutators and the strong radical*, Duke Math. J. **35** (1968), 853–859. MR **38** #1534.
5. D. Deckard and C. Pearcy, *On continuous matrix-valued functions on a Stonian space*, Pacific J. Math. **14** (1964), 857–869. MR **30** #2356.
6. J. Dixmier, *Les algèbres d'opérateurs dans l'espace hilbertien*, Cahiers Scientifiques, fasc. 25, Gauthier-Villars, Paris, 1957. MR **20** #1234.
7. ———, *Les C^* -algèbres et leurs représentations*, Cahiers Scientifiques, fasc. 29, Gauthier-Villars, Paris, 1964. MR **30** #1404.
8. J. M. G. Fell, *The structure of algebras of operator fields*, Acta Math. **106** (1961), 233–280. MR **29** #1547.
9. J. Glimm, *A Stone-Weierstrass theorem for C^* -algebras*, Ann. of Math. (2) **72** (1960), 216–244. MR **22** #7005.
10. H. Halpern, *Commutators in properly infinite von Neumann algebras*, Trans. Amer. Math. Soc. **139** (1969), 55–73.
11. ———, *Irreducible homomorphisms of a von Neumann algebra into its center*, Trans. Amer. Math. Soc. **140** (1969), 195–221. MR **39** #3322.
12. G. Lumer and M. Rosenblum, *Linear operator equations*, Proc. Amer. Math. Soc. **10** (1959), 32–41. MR **21** #2927.
13. Y. Misonou, *On a weakly central operator algebra*, Tôhoku Math. J. (2) **4** (1952), 194–202. MR **14**, 566.
14. C. R. Putnam, *On commutators of bounded matrices*, Amer. J. Math. **73** (1951), 127–131. MR **12**, 836.
15. ———, *Commutation properties of Hilbert space operators and related topics*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 36, Springer-Verlag, New York, 1967. MR **36** #707.
16. C. Rickart, *General theory of Banach algebras*, The University Series in Higher Math., Van Nostrand, Princeton, N. J., 1960. MR **22** #5903.
17. S. Sakai, *On linear functionals of W^* -algebras*, Proc. Japan Acad. **34** (1958), 571–574. MR **21** #5915.

ILLINOIS INSTITUTE OF TECHNOLOGY,
CHICAGO, ILLINOIS 60616