ON THE COMPLEX BORDISM OF EILENBERG-MAC LANE SPACES AND CONNECTIVE COVERINGS OF BU

BY

PETER S. LANDWEBER⁽¹⁾

Abstract. Explicit computations show that the universal coefficient spectral sequence from complex bordism to integral homology collapses for the spectra $K(\mathbf{Z})$ and bu , and also for their mod p reductions. Moreover the complex bordism modules of these spectra have infinite projective dimension.

1. Introduction. The aim of this paper is to continue the study of the complex bordism modules $\Omega_*^U(K(\mathbf{Z}, n))$ and $\Omega_*^U(BU(2n, \dots, \infty))$ which was begun in the earlier note [5]. Since our interest is in the stable ranges, it is convenient to introduce the Eilenberg-Mac Lane spectrum $K(\mathbf{Z}) = \{K(\mathbf{Z}, n)\}$ and the connective BU -spectrum (see [1] or [3, §10])

$$bu = \{\dots, BU(2n, \dots, \infty), U(2n+1, \dots, \infty), BU(2n+2, \dots, \infty), \dots\}.$$

Thus the objects of study are the bordism modules $\Omega_*^U(K(\mathbf{Z}))$ and $\Omega_*^U(bu)$; in general if $M = \{M_n\}$ is a spectrum, we define

$$\Omega_k^U(M) = \lim_{n \rightarrow \infty} \tilde{\Omega}_{k+n}^U(M_n).$$

In [5] we determined the images of the Thom homomorphisms

$$\Omega_*^U(K(\mathbf{Z})) \xrightarrow{\mu} H_*(K(\mathbf{Z})), \quad \Omega_*^U(bu) \xrightarrow{\mu} H_*(bu).$$

We are now able to obtain a more complete understanding of the relation between the complex bordism and homology of $K(\mathbf{Z})$ and bu . In [3, §4] P. Conner and L. Smith introduce a natural spectral sequence for finite (CW) complexes

$$(1.1) \quad E^r\langle X \rangle \Rightarrow H_*(X)$$

with

$$(1.2) \quad E_{p,q}^2\langle X \rangle = \text{Tor}_{p,q}^{\Omega_*^U}(\Omega_*^U(X), \mathbf{Z}),$$

Received by the editors July 16, 1970.

AMS 1969 subject classifications. Primary 5530.

Key words and phrases. Complex bordism, Eilenberg-Mac Lane spectra, connective BU -spectrum, Koszul resolution, Hurewicz homomorphisms for MU .

⁽¹⁾ Research supported in part by NSF grant GP-7993.

Copyright © 1971, American Mathematical Society

where Z is made a module over Ω_*^U by the augmentation. By taking limits there are also spectral sequences (1.1) for the spectra $K(Z)$ and bu , as well as for their mod p reductions $K(Z) \wedge Z_p = K(Z_p)$ and $bu \wedge Z_p$. We shall compute the E^2 -terms and prove the following:

THEOREM. *For each of the spectra $K(Z)$, $K(Z_p)$, bu and $bu \wedge Z_p$ (p a prime) the spectral sequence (1.1) of [3, §4] collapses.*

Thus in each case $H_*(X)$ has a filtration by graded subgroups $0 \subset F_0 \subset F_1 \subset \dots$, $\bigcup F_n = H_*(X)$, such that F_p/F_{p-1} is isomorphic to $\text{Tor}_{p,*}^{\Omega_*^U}(\Omega_*^U(X), Z)$. The edge homomorphism of (1.1) is the reduced Thom homomorphism

$$\bar{\mu}: \Omega_*^U(X) \otimes_{\Omega_*^U} Z \rightarrow H_*(X),$$

hence $\bar{\mu}$ is an isomorphism onto $F_0 \subset H_*(X)$ in each case of the theorem.

The steps involved in the proof of the theorem are outlined in §2. The analysis for $K(Z)$ and $K(Z_p)$ is made in §3, and in §4 we carry out the study of bu and $bu \wedge Z_p$.

From the computation of the E^2 -terms (see (3.3) and (4.6)) we conclude the following:

COROLLARY. *The complex bordism of each spectrum $K(Z)$, $K(Z_p)$, bu and $bu \wedge Z_p$ (p a prime) is an Ω_*^U -module of infinite projective dimension.*

In fact L. Smith has pointed out that from [3, §5] one can obtain the more precise result that $\Omega_*^U(K(Z_p, n))$ has projective dimension $\geq n$.

ACKNOWLEDGEMENT. It is a pleasure to thank Larry Smith for sending me preliminary versions of the paper [3], and for an illuminating correspondence that has been very helpful in the development of this paper.

2. Sketch of the argument. The first step is to note that, by switching the roles of the spectra involved, one obtains as in [5] the following isomorphisms:

$$\begin{aligned}\Omega_*^U(K(Z)) &\cong H_*(MU), \\ \Omega_*^U(K(Z_p)) &\cong H_*(MU; Z_p) = H_*(MU) \otimes Z_p, \\ \Omega_*^U(bu) &\cong k_*(MU)\end{aligned}$$

and

$$\Omega_*^U(bu \wedge Z_p) \cong k_*(MU; Z_p) = k_*(MU) \otimes Z_p.$$

These are isomorphisms of Ω_*^U -modules if the graded rings on the right are made Ω_*^U -modules via the following diagram of Hurewicz homomorphisms and reductions mod p .

$$\begin{array}{ccccc}\Omega_*^U = \pi_*(MU) & \begin{array}{c} \nearrow \mathcal{H} \\ \searrow \mathcal{H} \end{array} & \begin{array}{c} k_*(MU) \\ \downarrow \\ H_*(MU) \end{array} & \begin{array}{c} \xrightarrow{\rho_p} \\ \xrightarrow{\rho_p} \end{array} & \begin{array}{c} k_*(MU; Z_p) \\ \downarrow \\ H_*(MU; Z_p) \end{array}\end{array}$$

Next we describe convenient choices of polynomial generators for Ω_*^U , $H_*(MU)$ and $k_*(MU)$ following the detailed study of the complex bordism ring made by R. Stong in [8, Chapter 7] (see (3.1) and (4.3)). However it is rather difficult to deal directly with the Ω_*^U -modules $H_*(MU)$ and $k_*(MU)$ for the computations we have in mind. Thus we first compute the bigraded groups (see (3.3) and (4.6))

$$\mathrm{Tor}_{*,*}^{\Omega_*^U}(H_*(MU; \mathbb{Z}_p), \mathbb{Z}), \quad \mathrm{Tor}_{*,*}^{\Omega_*^U}(k_*(MU; \mathbb{Z}_p), \mathbb{Z})$$

and by a comparison of their totalizations with $H_*(K(\mathbb{Z}_p))$ and $H_*(bu \wedge \mathbb{Z}_p)$ we conclude that the spectral sequences (1.1) for $K(\mathbb{Z}_p)$ and $bu \wedge \mathbb{Z}_p$ must collapse.

Recall from [5, p. 526] that for each prime p the homology groups $H_*(K(\mathbb{Z}))$ and $H_*(bu)$ have no elements of order p^2 , and that this property implies that a homology class vanishes if all its reduction mod p vanish. In order to show that the spectral sequences (1.1) for $K(\mathbb{Z})$ and bu collapse, it suffices to show that also

$$\mathrm{Tor}_{*,*}^{\Omega_*^U}(H_*(MU), \mathbb{Z}), \quad \mathrm{Tor}_{*,*}^{\Omega_*^U}(k_*(MU), \mathbb{Z})$$

have no elements of order p^2 . This we do by applying the elementary result:

LEMMA 2.1. *Let $\dots \rightarrow C_n \xrightarrow{d} C_{n-1} \rightarrow \dots$ be a complex of free abelian groups, and let ∂_p denote the Bockstein homomorphism*

$$H(C; \mathbb{Z}_p) \xrightarrow{\partial} H(C) \xrightarrow{\rho} H(C; \mathbb{Z}_p)$$

where ∂ is the boundary associated to the coefficient sequence $0 \rightarrow \mathbb{Z} \xrightarrow{p} \mathbb{Z} \rightarrow \mathbb{Z}_p \rightarrow 0$ and ρ is reduction mod p . If ∂_p has zero homology in dimension $n+1$ then $H_n(C)$ has no elements of order p^2 .

Proof. Let $x \in H_n(C)$ with $p^2x=0$ but $px \neq 0$; we shall reach a contradiction. Since $p(px)=0$ we have $px = \partial y$ for a class $y \in H_{n+1}(C; \mathbb{Z}_p)$. Then $\partial_p y = \rho \partial y = \rho px = 0$, hence $y \in \mathrm{Ker}(\partial_p)$. By assumption $y = \rho \partial(z)$ for some class $z \in H_{n+2}(C; \mathbb{Z}_p)$. But then $px = \partial y = \partial \rho \partial(z) = 0$ since $\partial \rho = 0$, violating the assumption that $px \neq 0$. Q.E.D.

We use this result as follows: Let

$$\dots \rightarrow F_n \xrightarrow{d} F_{n-1} \rightarrow \dots \rightarrow F_0 \xrightarrow{e} \mathbb{Z} \rightarrow 0$$

be a free Ω_*^U -resolution of $\mathbb{Z}$, and let $\mathcal{M}_*$ denote either of the Ω_*^U -modules $H_*(MU)$, $k_*(MU)$ (both are free abelian). Then $\mathrm{Tor}_{*,*}^{\Omega_*^U}(\mathcal{M}_*, \mathbb{Z})$ is the homology of the complex of free abelian groups

$$(2.2) \quad \dots \rightarrow \mathcal{M}_* \otimes_{\Omega_*^U} F_n \xrightarrow{1 \otimes d} \mathcal{M}_* \otimes_{\Omega_*^U} F_{n-1} \rightarrow \dots \rightarrow \mathcal{M}_* \otimes_{\Omega_*^U} F_0 \rightarrow 0$$

and hence $\mathrm{Tor}_{*,*}^{\Omega_*^U}(\mathcal{M}_* \otimes \mathbb{Z}_p, \mathbb{Z})$ is the mod p homology of the complex (2.2) (for clarity we indicate only the homological degree). We shall compute the Bockstein

homomorphism ∂_p in $\text{Tor}_{**}^{\Omega_*^U}(\mathcal{M}_* \otimes \mathbf{Z}_p, \mathbf{Z})$ and show that its homology is zero in positive dimensions (see (3.6) and (4.8)), hence by the lemma $\text{Tor}_{**}^{\Omega_*^U}(\mathcal{M}_*, \mathbf{Z})$ has no elements of order p^2 .

Finally we exploit the commutative diagrams of differentials

$$\begin{array}{ccc} E_r\langle X \rangle & \xrightarrow{d^r} & E_r\langle X \rangle \\ \downarrow \rho_p & & \downarrow \rho_p \\ E_r\langle X \wedge \mathbf{Z}_p \rangle & \xrightarrow{d^r} & E_r\langle X \wedge \mathbf{Z}_p \rangle \end{array}$$

to conclude, by an induction on $r \geq 2$, that the spectral sequences (1.1) for $K(\mathbf{Z})$ and $\mathbf{bu}$ must collapse. That is, if we suppose that $d^2 = \dots = d^{r-1} = 0$ for $X = K(\mathbf{Z})$ or $\mathbf{bu}$ then $E_r\langle X \rangle = E_2\langle X \rangle$ and we find a commutative diagram

$$\begin{array}{ccc} E_r\langle X \rangle & \xrightarrow{d^r} & E_r\langle X \rangle \\ \downarrow \prod \rho_p & & \downarrow \prod \rho_p \\ \prod_p E_r\langle X \wedge \mathbf{Z}_p \rangle & \xrightarrow{\prod d^r} & \prod_p E_r\langle X \wedge \mathbf{Z}_p \rangle \end{array}$$

in which $\prod \rho_p$ is a monomorphism and $\prod d^r = 0$, hence also $d^r = 0$ for the spectrum X .

3. $H_*(\mathbf{MU})$ and $H_*(\mathbf{MU}; \mathbf{Z}_p)$ as Ω_*^U -modules. We begin by recalling the relation of the complex bordism ring Ω_*^U to the ring $H_*(\mathbf{MU}) \cong H_*(\mathbf{BU})$ (multiplication is provided by the Whitney sum). The Hurewicz homomorphism

$$\Omega_*^U = \pi_*(\mathbf{MU}) \xrightarrow{\mathcal{H}} H_*(\mathbf{MU})$$

is a monomorphism (Ω_*^U has no torsion) which records the Chern numbers of stably complex manifolds. The following result of J. Cohen, taken from [8, p. 130], provides all the information we need.

PROPOSITION 3.1. *There exist polynomial generators x_i ($i \geq 1$) of Ω_*^U and z_i ($i \geq 1$) of $H_*(\mathbf{MU})$, $\dim x_i = \dim z_i = 2i$ such that $\mathcal{H}x_i = m_i z_i$ where $m_i = p$ if $i+1 = p^s$ for some prime p and $m_i = 1$ otherwise.*

Recall that we regard $\mathbf{Z}$ as a module over $\Omega_*^U = \mathbf{Z}[x_1, x_2, \dots]$ by means of the augmentation. The Koszul resolution (see [6, p. 204]) of the Ω_*^U -module $\mathbf{Z}$ consists of the bigraded exterior algebra

$$E_{*,*} = E_{\Omega^U}[y_1, y_2, \dots]$$

where y_i has bidegree $(1, 2i)$ and elements of Ω_{2i}^U have bidegree $(0, 2i)$, and homomorphisms of Ω_*^U -modules

$$\dots \longrightarrow E_{n,*} \xrightarrow{d} E_{n-1,*} \longrightarrow \dots \longrightarrow E_{0,*} \xrightarrow{\varepsilon} \mathbf{Z} \longrightarrow 0$$

such that d is a derivation satisfying $d(y_i) = x_i \cdot 1$ on the generators and ε is the augmentation $E_{0,*} = \Omega_*^U \rightarrow \mathbf{Z}$. This is a free resolution.

We now fix a prime p and continue to denote by z_i the images of the polynomial generators in $H_*(MU; \mathbf{Z}_p)$, hence $H_*(MU; \mathbf{Z}_p) = \mathbf{Z}_p[z_1, z_2, \dots]$. Then the bigraded algebra

$$(3.2) \quad \text{Tor}_{*,*}^{\Omega_*^U}(H_*(MU; \mathbf{Z}_p), \mathbf{Z})$$

is the homology of

$$H_*(MU; \mathbf{Z}_p) \otimes E_{\mathbf{Z}}[y_1, y_2, \dots]$$

under a derivation d' of $H_*(MU; \mathbf{Z}_p)$ -modules which satisfies $d'(1 \otimes y_i) = \rho \mathcal{H} x_i \otimes 1$ if $i+1 \neq p^s$ (note that $\rho \mathcal{H} x_i$ is then a generator of $H_*(MU; \mathbf{Z}_p)$ in dimension $2i$) and $d'(1 \otimes y_i) = 0$ if $i+1 = p^s$ for some $s > 0$. Thus (3.2) is the homology of the tensor product of the complexes

$$\begin{aligned} & \mathbf{Z}_p[\rho \mathcal{H} x_i; i+1 \neq p^s] \otimes E_{\mathbf{Z}_p}[y_i; i+1 \neq p^s], \\ & \mathbf{Z}_p[z_{p^s-1}; s > 0] \otimes E_{\mathbf{Z}_p}[y_{p^s-1}; s > 0] \end{aligned}$$

under a derivation d' which satisfies $d'(y_i) = \rho \mathcal{H} x_i$ if $i+1 \neq p^s$ and which annihilates z_{p^s-1} and y_{p^s-1} for $s > 0$. This is simply the tensor product of a Koszul resolution for $\mathbf{Z}_p$ and a complex with zero differential, hence the Künneth formula yields

PROPOSITION 3.3. *There is an algebra isomorphism*

$$\text{Tor}_{*,*}^{\Omega_*^U}(H_*(MU; \mathbf{Z}_p), \mathbf{Z}) \cong \mathbf{Z}_p[z_{p^s-1}; s > 0] \otimes E_{\mathbf{Z}_p}[y_{p^s-1}; s > 0]$$

where y_i has bidegree $(1, 2i)$ and z_i has bidegree $(0, 2i)$ for $i = p^s - 1$.

Thus we have computed the E^2 -term of the spectral sequence

$$(3.4) \quad E^r\langle K(\mathbf{Z}_p) \rangle \Rightarrow H_*(K(\mathbf{Z}_p)).$$

By interchanging the roles of the Eilenberg-Mac Lane spectra $K(\mathbf{Z})$ and $K(\mathbf{Z}_p)$ as in [5] we find that $H_*(K(\mathbf{Z}_p)) \cong H_*(K(\mathbf{Z}); \mathbf{Z}_p)$. It is well known that

$$H_*(K(\mathbf{Z}); \mathbf{Z}_p) \rightarrow H_*(K(\mathbf{Z}_p); \mathbf{Z}_p)$$

is a monomorphism whose image we now describe. Recall from [7] that $H_*(K(\mathbf{Z}_p); \mathbf{Z}_p)$ is a Hopf algebra dual to the mod p Steenrod algebra and that there is an algebra isomorphism

$$(3.5) \quad H_*(K(\mathbf{Z}_p); \mathbf{Z}_p) = E_{\mathbf{Z}_p}[\eta_i; i \geq 0] \otimes \mathbf{Z}_p[\zeta_i; i > 0]$$

where $\deg \eta_i = 2p^i - 1$ and $\deg \zeta_i = 2(p^i - 1)$. Then the image of $H_*(K(\mathbf{Z}); \mathbf{Z}_p)$ is the subalgebra generated by the η_i and ζ_i for $i > 0$, and so we conclude by a comparison of (3.3) and (3.5) that $H_*(K(\mathbf{Z}_p))$ is (algebra) isomorphic to the totalization of $E_{*,*}^2\langle K(\mathbf{Z}_p) \rangle$. Since both are $\mathbf{Z}_p$ -vector spaces of finite type, the spectral sequence (3.4), i.e. the spectral sequence (1.1) for $K(\mathbf{Z}_p)$, must collapse.

It remains to compute the Bockstein ∂_p in $\text{Tor}_{*,*}^{\Omega_*^U}(H_*(MU; \mathbb{Z}_p), \mathbb{Z})$, as in §2, in order to show that also the spectral sequence (1.1) for $K(\mathbb{Z})$ collapses. We shall prove

PROPOSITION 3.6. *The Bockstein ∂_p is a derivation of the algebra*

$$\text{Tor}_{*,*}^{\Omega_*^U}(H_*(MU; \mathbb{Z}_p), \mathbb{Z})$$

which satisfies $\partial_p(y_{p^s-1}) = z_{p^s-1}$ and $\partial_p(z_{p^s-1}) = 0$ on the generators.

It then follows that the homology of ∂_p is isomorphic to $\mathbb{Z}_p$ (in bidegree $(0, 0)$) since we have simply the Koszul resolution for $\mathbb{Z}_p$ over the polynomial algebra $\mathbb{Z}_p[z_{p^s-1}; s > 0]$. We then argue as in §2 that the spectral sequence (1.1) for $K(\mathbb{Z})$ must also collapse.

Proof of (3.6). We omit the standard argument that ∂_p is a derivation and concentrate on identifying $\partial_p(y_i)$, where the y_i are the exterior algebra generators of (3.3). Thus we consider the projection

$$H_*(MU) \otimes Ez[y_1, y_2, \dots] \rightarrow H_*(MU; \mathbb{Z}_p) \otimes Ez[y_1, y_2, \dots]$$

of complexes, lift $1 \otimes y_i$ ($i = p^s - 1$) back to $1 \otimes y_i$ in $H_*(MU) \otimes Ez[y_1, y_2, \dots]$, apply the differential to obtain $\mathcal{H}x_i \otimes 1 = pz_i \otimes 1$ (see (3.1)), divide by p to obtain $z_i \otimes 1$ and finally apply reduction mod p which yields z_i as desired. Q.E.D.

4. $k_*(MU)$ and $k_*(MU; \mathbb{Z}_p)$ as Ω_*^U -modules. First recall that $k_*()$ is the multiplicative homology theory represented by the connective BU -spectrum bu , and that the coefficient ring $k_* = \mathbb{Z}[\beta]$, $\dim \beta = 2$ (for example see [3, §10]). For a spectrum $M = \{M_n\}$ we define

$$k_m(M) = \lim_{n \rightarrow \infty} \tilde{k}_{m+n}(M_n).$$

Maps which result from the Whitney sum make $k_*(MU)$ an algebra over k_* . Since $H_*(MU)$ is a polynomial algebra over $\mathbb{Z}$, hence has no torsion, it follows easily that

$$(4.1) \quad k_*(MU) = k_*[t_1, t_2, \dots], \quad \dim t_i = 2i.$$

It is possible to choose the generators so that the Hurewicz homomorphism

$$(4.2) \quad \Omega_*^U = \pi_*(MU) \xrightarrow{\mathcal{H}} k_*(MU),$$

induced by the map of the sphere spectrum $S = \{S^n\}$ into bu which consists of maps $S^{2n} \rightarrow BU(2n, \dots, \infty)$ and $S^{2n+1} \rightarrow U(2n+1, \dots, \infty)$ that lift generators of $\pi_{2n}(BU)$ and $\pi_{2n+1}(U)$, takes a convenient form. Namely, the generators can be chosen so that we have

$$\mathcal{H}[M^{2n}] = \sum [M^{2n}]_{i_1, \dots, i_r, t_{i_1} \dots t_{i_r} \beta^{n-i_1-\dots-i_r}}$$

where the coefficient is the tangential K -theory characteristic number (see [2] or [8])

$$[M^{2n}]_{i_1, \dots, i_r} = s_{(i_1, \dots, i_r)}(\gamma(\tau)) [M^{2n}]$$

of the U -manifold M^{2n} determined by the partition $(i_1, \dots, i_r)$. (We choose tangential rather than normal K -theory numbers in order to agree with [8].)

It follows directly from the celebrated theorem of A. Hattori [4] and R. Stong [8, p. 129] that the Hurewicz homomorphism (4.2) is (additively) a split monomorphism. We shall deduce the following further information from [8, Chapter 7].

PROPOSITION 4.3. *There exist polynomial generators x_i ($i \geq 1$) of Ω_*^U and z_i ($i \geq 1$) of $k_*(MU)$ over $k_* = \mathbb{Z}[\beta]$, $\dim x_i = \dim z_i = 2i$, such that $\mathcal{H}x_{p-1} = pz_{p-1} + \beta^{p-1}$, $\mathcal{H}x_i = pz_i \bmod \beta^{p-1}$ if $i+1 = p^s$ ($s > 1$) and $\mathcal{H}x_i = z_i$ if $i+1$ is not a prime power.*

We remark that (3.1) is an immediate consequence of this result, in view of the natural transformation $\lambda: k_*() \rightarrow H_*()$ of [5, §2]. For the generators z_i of $k_*(MU)$ over k_* are carried under λ to generators of $H_*(MU)$, and on the coefficient rings λ is the augmentation $k_* = \mathbb{Z}[\beta] \rightarrow \mathbb{Z}$ (so $\lambda(\beta) = 0$).

When $i+1 = p^s$ and $s > 1$ we have an equality $\mathcal{H}x_i = pz_i + \beta^{p-1}w^{(s)}$ where $w^{(s)} \in k_{2(p^s-p)}(MU)$. Although this knowledge is sufficient for our purposes, it would be interesting to have a better grasp on the elements $w^{(s)}$. We conjecture that it is possible to choose the generators so that

$$\mathcal{H}x_{p^s-1} = pz_{p^s-1} + (\mathcal{H}x_{p-1})(z_{p^s-1-1})^p$$

for $s > 1$. The best supporting evidence is the lemma on p. 121 of [8].

Proof of (4.3). We assume familiarity with the relevant portion of [8, Chapter 7]. If $i+1$ is not a prime power let $x_i = [M^{2i}]$ be any generator of Ω_*^U in dimension $2i$; then by [8, p. 128] we have $s_{(i)}(c(\tau))[M^{2i}] = \pm 1$ and therefore $\mathcal{H}x_i$ is a generator of $k_*(MU)$ over k_* in dimension $2i$. We put $z_i = \mathcal{H}x_i$.

If $i+1 = p$ then put $x_i = [CP(p-1)]$. Since $s_{(p-1)}(c(\tau))[CP(p-1)] = p$ it follows from [8, p. 128] that $[CP(p-1)]$ is a generator of Ω_*^U in dimension $2(p-1)$. From [2, (14.1)] we see that $s_\omega(c(\tau))[CP(p-1)]$ is zero mod p unless ω is the empty partition, and then as is well known we obtain the Todd genus of $CP(p-1)$ which is 1. The equation $\mathcal{H}x_{p-1} = pz_{p-1} + \beta^{p-1}$ now defines the generator z_{p-1} .

Finally suppose $i+1 = p^{s+1}$ and $s > 0$. Let $x_i = [M^{2i}]$ be a generator of Ω_*^U in dimension $2i = 2(p^{s+1} - 1)$ which is congruent to a multiple of $[H_{p^s, \dots, p^s}] \bmod p$ (see [8, p. 121]). Then the key lemma on p. 121 of [8] implies that $\mathcal{H}x_i$ is divisible by $\beta^{p-1} \bmod p$; for the mod p K -theory characteristic numbers of M^{2i} are a multiple of those of $H_{p^s, \dots, p^s}$, and therefore $s_{(i)}(c(\tau))[M^{2i}] = 0 \bmod p$ if $i_1 + \dots + i_r > p^{s+1} - p = i - (p-1)$. From [8, p. 128] we have $s_{(i)}(c(\tau))[M^{2i}] = \pm p$, hence any solution of the congruence $\mathcal{H}x_i = pz_i \bmod \beta^{p-1}$ is a generator of $k_*(MU)$ in dimension $2i$. Q.E.D.

We are now ready to compute the bigraded algebra

$$(4.4) \quad \text{Tor}_{*,*}^{\Omega_*^U}(k_*(MU; \mathbb{Z}_p), \mathbb{Z})$$

for a fixed prime p . We continue to denote by z_i the reductions of the generators for $k_*(MU)$, so that $k_*(MU; \mathbb{Z}_p) = \mathbb{Z}_p[\beta; z_1, z_2, \dots]$. Then, as in §3, (4.4) is the homology of

$$(4.5) \quad k_*(MU; \mathbb{Z}_p) \otimes_{\mathbb{Z}} E_{\mathbb{Z}}[y_1, y_2, \dots]$$

under a derivation d' of $k_*(MU; \mathbb{Z}_p)$ -modules which satisfies $d'(1 \otimes y_i) = \rho \mathcal{H} x_i \otimes 1$ if $i+1 \neq p^s$ (note that $\rho \mathcal{H} x_i$ is then a generator of $k_*(MU; \mathbb{Z}_p)$ in dimension $2i$) and $d'(1 \otimes y_{p^s-1}) = \rho \mathcal{H} x_{p^s-1} \otimes 1 = \beta^{p-1} w^{(s)} \otimes 1$ for some $w^{(s)} \in k_{2(p^s-p)}(MU; \mathbb{Z}_p)$. In particular we have $d'(1 \otimes y_{p-1}) = \beta^{p-1} \otimes 1$.

For $s > 1$ we shall replace the exterior algebra generator $1 \otimes y_{p^s-1}$ by the cycle $y'_{p^s-1} = 1 \otimes y_{p^s-1} - w^{(s)} \otimes y_{p-1}$. One checks easily that these cycles generate an exterior algebra (each y_i has odd total degree). Hence the complex (4.5) is the tensor product of the complexes

$$\mathbb{Z}_p[\rho \mathcal{H} x_i; i+1 \neq p^s] \otimes E_{\mathbb{Z}_p}[y_i; i+1 \neq p^s], \quad \mathbb{Z}_p[\beta] \otimes E_{\mathbb{Z}_p}[y_{p-1}]$$

and

$$\mathbb{Z}_p[z_{p^s-1}; s > 0] \otimes E_{\mathbb{Z}_p}[y'_{p^s-1}; s > 1]$$

under a derivation d' which satisfies $d'(y_i) = \rho \mathcal{H} x_i$ if $i+1 \neq p^s$, $d'(\beta) = 0$ and $d'(y_{p-1}) = \beta^{p-1}$, and which annihilates z_{p^s-1} and y'_{p^s-1} . In view of the Koszul resolution and the Künneth formula we now find

PROPOSITION 4.6. *There is an isomorphism*

$$\mathrm{Tor}_{*,*}^{\Omega_U}(k_*(MU; \mathbb{Z}_p), \mathbb{Z}) = B_* \otimes P_* \otimes E_*$$

with the tensor product of the truncated polynomial ring $B_* = \mathbb{Z}_p[\beta]/(\beta^{p-1})$, the polynomial ring $P_* = \mathbb{Z}_p[z_{p^s-1}; s > 0]$ and the exterior algebra $E_* = E_{\mathbb{Z}_p}[y'_{p^s-1}; s > 1]$.

Notice that this result immediately implies that the spectral sequence

$$(4.7) \quad E^r\langle bu \wedge \mathbb{Z}_p \rangle \Rightarrow H_*(bu \wedge \mathbb{Z}_p)$$

collapses. For J. F. Adams showed in [1] that $H^*(bu; \mathbb{Z}_p)$ is isomorphic to a direct sum of cyclic modules $A_p/A_p Q_0 + A_p Q_1$ over the mod p Steenrod algebra on generators in $H^{2i}(bu; \mathbb{Z}_p)$ for $i=0, 1, \dots, p-2$ (recall that $Q_0 \in A_p^1$ and $Q_1 \in A_p^{2p-1}$). Hence $H_*(bu; \mathbb{Z}_p)$ and the totalization of $E^2\langle bu \wedge \mathbb{Z}_p \rangle \cong \mathrm{Tor}_{*,*}^{\Omega_U}(k_*(MU; \mathbb{Z}_p), \mathbb{Z})$ are graded $\mathbb{Z}_p$ -modules of finite type which have the same dimension in each degree, so they are isomorphic. Therefore the spectral sequence (4.7), i.e. the spectral sequence (1.1) for $bu \wedge \mathbb{Z}_p$, must collapse.

It only remains to compute the Bockstein ∂_p in $\mathrm{Tor}_{*,*}^{\Omega_U}(k_*(MU; \mathbb{Z}_p), \mathbb{Z})$, as in §2, in order to show that also the spectral sequence (1.1) for bu collapses. We shall prove, in the notation of (4.6),

PROPOSITION 4.8. *The Bockstein ∂_p is a derivation of the algebra*

$$\mathrm{Tor}_{*,*}^{\Omega_U}(k_*(MU; \mathbf{Z}_p), \mathbf{Z})$$

which satisfies $\partial_p(y_p'^{s-1}) = z_p^{s-1}$ for $s > 1$, $\partial_p(\beta) = 0$ and $\partial_p(z_p^{s-1}) = 0$ for $s > 0$.

Proof. By a standard argument ∂_p is a derivation sending $\mathrm{Tor}_{r,s}$ to $\mathrm{Tor}_{r-1,s}$, and so we concentrate on identifying $\partial_p(y_p'^{s-1})$. Thus we consider the projection

$$k_*(MU) \otimes Ez[y_1, y_2, \dots] \rightarrow k_*(MU; \mathbf{Z}_p) \otimes Ez[y_1, y_2, \dots]$$

of complexes, lift the cycle $y_p'^{s-1}$ back to $1 \otimes y_p^{s-1} - w^{(s)} \otimes y_{p-1}$ (where $w^{(s)} \in k_{2(p^s-p)}(MU)$ satisfies $\mathcal{H}x_{p^s-1} = pz_{p^s-1} + \beta^{p-1}w^{(s)}$), apply the differential to obtain $(\mathcal{H}x_{p^s-1} - \beta^{p-1}w^{(s)}) \otimes 1 = pz_{p^s-1} \otimes 1$, divide by p to obtain $z_{p^s-1} \otimes 1$, and finally apply reduction mod p which yields z_{p^s-1} as desired. Q.E.D.

We now obtain immediately

COROLLARY 4.9. *The homology of the Bockstein ∂_p in $\mathrm{Tor}_{*,*}^{\Omega_U}(k_*(MU; \mathbf{Z}_p), \mathbf{Z})$ is algebra isomorphic to $\mathbf{Z}_p[\beta, z_{p-1}]/(\beta^{p-1})$, where β has bidegree $(0, 2)$ and z_{p-1} has bidegree $(0, 2p-2)$.*

Thus the homology of ∂_p is concentrated in bidegrees $(0, *)$, and then (2.1) implies that $\mathrm{Tor}_{*,*}^{\Omega_U}(k_*(MU), \mathbf{Z})$ has no elements of order p^2 . We now may conclude as in §2 that the spectral sequence (1.1) for *bu* must also collapse.

REFERENCES

1. J. F. Adams, *On Chern characters and the structure of the unitary group*, Proc. Cambridge Philos. Soc. **57** (1961), 189–199. MR **22** #12525.
2. P. E. Conner and E. E. Floyd, *The relation of cobordism to K-theories*, Lecture Notes in Math., no. 28, Springer-Verlag, Berlin and New York, 1966. MR **35** #7344.
3. P. E. Conner and L. Smith, *On the complex bordism of finite complexes*, Inst. Hautes Études Sci. Publ. Math. **37** (1970).
4. A. Hattori, *Integral characteristic numbers for weakly almost complex manifolds*, Topology **5** (1966), 259–280. MR **33** #742.
5. P. S. Landweber, *Steenrod representability of stable homology*, Proc. Amer. Math. Soc. **18** (1967), 523–529. MR **37** #4815.
6. S. MacLane, *Homology*, Die Grundlehren der math. Wissenschaften, Band 114, Academic Press, New York; Springer-Verlag, Berlin, 1963. MR **28** #122.
7. J. W. Milnor, *The Steenrod algebra and its dual*, Ann. of Math. (2) **67** (1958), 150–171. MR **20** #6092.
8. R. E. Stong, *Notes on cobordism theory*, Mathematical Notes, Princeton Univ. Press, Princeton, N. J.; Univ. of Tokyo Press, Tokyo, 1968. MR **40** #2108.

YALE UNIVERSITY,

NEW HAVEN, CONNECTICUT 06520

RUTGERS UNIVERSITY,

NEW BRUNSWICK, NEW JERSEY 08903