SLICING THEOREMS FOR n -SPHERES IN EUCLIDEAN $(n+1)$ -SPACE

BY

ROBERT J. DAVERMAN⁽¹⁾

Abstract. This paper describes conditions on the intersection of an n -sphere Σ in Euclidean $(n+1)$ -space E^{n+1} with the horizontal hyperplanes of E^{n+1} sufficient to determine that the sphere be nicely embedded. The results generally are pointed towards showing that the complement of Σ is 1-ULC (uniformly locally 1-connected) rather than towards establishing the stronger property that Σ is locally flat. For instance, the main theorem indicates that $E^{n+1} - \Sigma$ is 1-ULC provided each non-degenerate intersection of Σ and a horizontal hyperplane be an $(n-1)$ -sphere bicollared both in that hyperplane and in Σ itself ($n \neq 4$).

1. Introduction. Much of the literature that treats the problem of determining properties of the embedding of an object Σ in E^n from information about the intersection of Σ with the horizontal hyperplanes of E^n focuses on the case $n=3$. Such a problem first arose in this dimension when J. W. Alexander [1] suggested that a 2-sphere in E^3 might be embedded just as a round sphere if each of its intersections with the horizontal planes were either a point or a simple closed curve. Recently Eaton [11] and Hosay [12] showed this to be true. After generalizations by Loveland [15] and Jensen [13], Cannon proved that the same property is held by any 2-sphere Σ in E^3 such that no intersection of Σ with a horizontal plane has a degenerate component [7].

For higher dimensions Bryant has proved that a k -dimensional compact set X in E^n , where $n-k \geq 3$, has a 1-ULC complement if the complement of X with respect to each member of some dense subset of the horizontal hyperplanes of E^n is 1-ULC [5].

The main results of this paper are found in §5, where it is shown that a closed n -manifold Σ topologically embedded in E^{n+1} ($n \neq 4$) is nice (meaning, $E^{n+1} - \Sigma$ is 1-ULC) if each nondegenerate intersection Σ_i of Σ and a horizontal, n -dimensional hyperplane in E^{n+1} is a PL $(n-1)$ -manifold bicollared in that hyperplane and nice in Σ ($\Sigma - \Sigma_i$ is 1-ULC). Most of the techniques required are consigned to §4. In §6 other generalizations to the results mentioned in the first paragraph are given, the methods for which are taken from [7] and [12]. In addition, we describe

Received by the editors March 15, 1971.

AMS 1970 subject classifications. Primary 57A45, 57A50; Secondary 57A15.

Key words and phrases. Horizontal hyperplane of Euclidean space, bicollared submanifold, locally flat submanifold, locally simply connected, homeomorphic approximation, topological embedding.

⁽¹⁾ Research was supported in part by NSF Grant GP-19966.

Copyright © 1972, American Mathematical Society

in §3 some methods, similar to but weaker than those of the three-dimensional case (see [3]), for improving mappings of a disk into the closure of a complementary domain of an n -manifold in an $(n+1)$ -manifold.

2. Definitions and notation. An n -manifold is a separable metric space which is locally homeomorphic to E^n ; thus, the term manifold is reserved for manifolds without boundary. For simplicity we shall assume all manifolds to be connected, but they need not be compact or triangulated. A manifold that is compact (and without boundary) is said to be *closed*.

A subset S of a metric space is called an ε -subset if and only if the diameter of S , written $\text{diam } S$, is less than ε .

Suppose f and g are maps of a space X into a space Y that has a metric ρ . The symbol $\rho(f, g) < \varepsilon$ means that $\rho(f(x), g(x)) < \varepsilon$ for each x in X . The maps f and g are said to be ε -homotopic (ε -isotopic) if and only if there exists a homotopy (isotopy) h_t sending X into Y such that $h_0 = f$, $h_1 = g$ and $\rho(h_s, h_t) < \varepsilon$ for all s, t in $[0, 1]$.

A map f of the metric space Y into a subset A is an ε -map if and only if $\rho(y, f(y)) < \varepsilon$ for each $y \in Y$.

The symbol Δ^2 denotes a 2-simplex fixed throughout this paper and $\partial\Delta^2$ denotes its boundary. Given a triangulation R of Δ^2 , we use $R^{(i)}$ to denote the i -skeleton of R ($i=0, 1$).

For any point p in a metric space S and any positive number δ , $N_\delta(p)$ denotes the set of points in S whose distance from p is less than δ .

Let A denote a subset of a metric space X and p a limit point of A . We say that A is *locally simply connected at p* , written 1-LC at p , if and only if for each $\varepsilon > 0$ there is a $\delta > 0$ such that each map of $\partial\Delta^2$ into $A \cap N_\delta(p)$ can be extended to a map of Δ^2 into $A \cap N_\varepsilon(p)$. Furthermore, we say that A is *uniformly locally simply connected*, written 1-ULC, if for each $\varepsilon > 0$ there is a $\delta > 0$ such that each map of $\partial\Delta^2$ into a δ -subset of A can be extended to a map of Δ^2 into an ε -subset of A .

In the same context we say that A is *locally arcwise connected at p* , or 0-LC at p , if and only if for each $\varepsilon > 0$ there exists a $\delta > 0$ such that any map of ∂I (where $I = [0, 1]$) into $A \cap N_\delta(p)$ extends to a map of I into $A \cap N_\varepsilon(p)$. We define analogously the uniform condition 0-ULC.

For a subset U of a space S we use the symbol $\text{Cl } U$ to denote the closure of U and $\text{Bd } U$ to denote the (topological) boundary of U in S .

Let Σ be an n -manifold embedded in the interior of an $(n+1)$ -manifold M as a closed subset, and let U be an open subset of $M - \Sigma$. Then Σ is *collared from U* if and only if there exists a homeomorphism g of $\Sigma \times I$ into $\text{Cl } U$ such that $g(s, 0) = s$ for each s in Σ . Similarly, Σ is *bicollared in M* if and only if there exists a homeomorphism h of $\Sigma \times [-1, 1]$ into M such that $h(s, 0) = s$ for each s in Σ . In addition, Σ is *locally flat in M* if and only if each point s of Σ has a neighborhood N (relative to M) such that $N \cap \Sigma$ is bicollared in N .

Let n be a positive integer. For each real number t define E_t , the hyperplane of E^{n+1} at the t -level, as $\{(x_1, \dots, x_{n+1}) \in E^{n+1} \mid x_{n+1} = t\}$. For any subset Σ of E^{n+1} we define Σ_t as $\Sigma \cap E_t$, and for any set C of real numbers we define $\Sigma(C)$ as $\bigcup \{\Sigma_t \mid t \in C\}$; however, for an interval (a, b) we simplify $\Sigma((a, b))$ to $\Sigma(a, b)$.

3. Altering maps of a disk.

LEMMA 3.1. *Suppose Σ is an n -manifold embedded in the interior of an $(n+1)$ -manifold M as a closed separating subset, U a component of $M - \Sigma$, X a closed subset of Σ such that $\text{Cl } U - X$ is 1-LC at each point of X , R a triangulation of Δ^2 , T a subdivision of R , and F a map of Δ^2 into $\text{Cl } U$ such that $F(|R^{(1)}|) \subset U$.*

Then for each $\varepsilon > 0$ there exists a map G of Δ^2 into $\text{Cl } U$ such that

- (a) $G \mid |R^{(1)}| = F \mid |R^{(1)}|$,
- (b) $\rho(F, G) < \varepsilon$,
- (c) $G(|T^{(1)}|) \subset U$,
- (d) $G(\Delta^2) \cap X = \emptyset$.

Proof. Note that, since $F(\Delta^2)$ is compact, there exist a neighborhood V of $F(\Delta^2)$ (relative to $\text{Cl } U$) and a positive number δ such that any map of $\partial\Delta^2$ into a δ -subset of $V - X$ can be extended so as to send Δ^2 into an $(\varepsilon/3)$ -subset of $\text{Cl } U - X$.

Now we simply modify F , beginning with the 0-skeleton of T and working up. In case $v \in T^{(0)}$ and $F(v) \in \Sigma$, define $G(v)$ as a point of $U \cap V$ very close to $F(v)$; when $F(v) \notin \Sigma$, define $G(v) = F(v)$. Since U is 0-LC at each point of Σ [17, Theorem II.5.35], then, for each 1-simplex σ of T , G can be extended along σ in such a way that $G(\sigma) \subset U \cap V$ and $\rho(G|_{\sigma}, F|_{\sigma}) < \varepsilon/3$; in case $\sigma \subset |R^{(1)}|$, define $G|_{\sigma} = F|_{\sigma}$. Because first we could have subdivided T (if necessary), we can assume that, for each 2-simplex τ of T , $\text{Diam } F(\tau) < \varepsilon/3$ and $\text{Diam } G(\partial\tau) < \delta$. According to the previous paragraph, G can be extended over τ into an $\varepsilon/3$ -subset of $\text{Cl } U - X$. It follows easily that $\rho(F, G) < \varepsilon$.

THEOREM 3.2. *Suppose Σ is an n -manifold embedded in the interior of an $(n+1)$ -manifold M as a closed separating subset, U a component of $M - \Sigma$, and f a map of Δ^2 into $\text{Cl } U$ with $f(\partial\Delta^2) \subset U$. Suppose $\{X^i\}$ is a countable collection of closed subsets of Σ such that $\text{Cl } U - X^i$ is 1-LC at each point of X^i ($i=1, 2, \dots$). Then for each $\varepsilon > 0$ there exists a map g of Δ^2 into $\text{Cl } U$ such that*

- (1) $g|_{\partial\Delta^2} = f|_{\partial\Delta^2}$,
- (2) $\rho(f, g) < \varepsilon$,
- (3) $g^{-1}(\Sigma \cap g(\Delta^2))$ is 0-dimensional,
- (4) $g(\Delta^2) \cap X^i = \emptyset$ for $i=1, 2, \dots$

The proof follows from routine applications of Lemma 3.1.

This result yields the following corollary, which can be regarded as a very weak version of [3, Theorem 4.2].

COROLLARY 3.3. *Suppose Σ is an n -manifold embedded in the interior of an $(n+1)$ -manifold M as a closed separating subset, U a component of $M - \Sigma$, and f a map of Δ^2 into $\text{Cl } U$ such that $f(\partial\Delta^2) \subset U$. Then for each $\varepsilon > 0$ there exists a map g of Δ^2 into $\text{Cl } U$ such that (i) $g|_{\partial\Delta^2} = f|_{\partial\Delta^2}$, (ii) $\rho(f, g) < \varepsilon$, and (iii) $g^{-1}(\Sigma \cap g(\Delta^2))$ is 0-dimensional.*

4. Embedding Cartesian products in E^{n+1} . Let n denote a fixed positive integer. Obviously there exists a countable collection $\mathcal{M}$ of closed, PL $(n-1)$ -manifolds such that any closed, PL $(n-1)$ -manifold is homeomorphic to some member of $\mathcal{M}$. (In fact, this holds even without the PL hypothesis [8].)

In this section Σ will denote an n -manifold embedded in E^{n+1} as a closed subset, U a component of $E^{n+1} - \Sigma$, and (a, b) an interval of real numbers such that, for each $t \in (a, b)$, Σ_t is homeomorphic to some member of $\mathcal{M}$ and is collared from U_t . Let $(a, b)_M$ be the set of all t in (a, b) such that Σ_t is homeomorphic to M , where $M \in \mathcal{M}$. For each such M and each $t \in (a, b)_M$ define an embedding of $M \times [-1, 1]$ into $\text{Cl } U_t$ such that $\lambda_t(M \times \{-1\}) = \Sigma_t$. Topologize the set $\{\lambda_t \mid t \in (a, b)_M\}$ by the sup-norm metric in E^{n+1} , producing a separable metric space.

As suggested by Bryant [4] (and in the unpublished work of Bing [2]), one can easily establish the following lemma.

LEMMA 4.1. *There exists a countable subset $\mathcal{D}$ of (a, b) such that to each t in $(a, b) - \mathcal{D}$ there correspond two sequences $\{s(i)\}$ and $\{u(i)\}$ of real numbers, with $a < s(i) < t < u(i) < b$, such that each of the associated sequences $\{\lambda_{s(i)}\}$ and $\{\lambda_{u(i)}\}$ converges (homeomorphically) to λ_t .*

Throughout the rest of §§4 and 5, $\mathcal{D}$ will denote the subset of (a, b) described in Lemma 4.1, and n will denote a fixed integer other than 4.

The following result can be established by adding simple epsilons to the proof of Borsuk's theorem (see [10, Theorem 10.2]).

LEMMA 4.2. *Let A be an ANR embedded as a closed subset of a metric space X , $\varepsilon > 0$, and $f: X \rightarrow A$ an ε -map such that $f|_A$ is ε -homotopic (in A) to the identity map. Then there exists a 2ε -retraction of X onto A .*

LEMMA 4.3. A. *If $t \in (a, b)_M - \mathcal{D}$, then there exists a homeomorphism h of $M \times [-1, 1]$ onto a subset A_t of $\text{Cl } U$ such that*

$$\Sigma \cap A_t = h(M \times \{-1\}) \cup h(M \times \{1\}) = \Sigma_s \cup \Sigma_u,$$

where $a < s < t < u < b$ and $A_t \cap E_z = \emptyset$ for z not in $[s, u]$.

B. *For any such A_t , let X_t denote the closure of the component of $\text{Cl } U - A_t$ containing Σ_t . Then, for each $\varepsilon > 0$, A_t can be obtained so that there exists an ε -retraction of X_t onto A_t .*

Proof. Fix a point t of $(a, b)_M - \mathcal{D}$. By restricting $[-1, 1]$ to a subinterval containing -1 , if necessary, we may assume that $\text{diam } \lambda_t(\{p\} \times [-1, 1])$ is less than

$\varepsilon/18$ for each p in M . Since $\lambda_t(M \times \{0\})$ is an ANR, the obvious retraction of $\lambda_t(M \times [-1, 1])$ onto $\lambda_t(M \times \{0\})$ can be extended over a neighborhood N of $\lambda_t(M \times [-1, 1])$ to an $\varepsilon/18$ -retraction R of N onto $\lambda_t(M \times \{0\})$. Furthermore, N can be chosen as a product $N' \times (s', u') \subset E^n \times E^1$ where N' is a bounded open subset of E^n and $N \cap \Sigma = \Sigma(s', u')$.

It is sufficient to describe the homeomorphism h of $M \times [-1, 1]$ onto some (as yet undefined) A_t subject to the following conditions:

- (1) $X_t \subset N$,
- (2) $\text{diam } h(\{p\} \times [-1, 1]) < \varepsilon/6$ for each p in M ,
- (3) $h(p, 0) = \lambda_t(p, 0)$ for each p in M .

Let g denote the map of A_t to $h(M \times \{0\})$ sending each $h(\{p\} \times [-1, 1])$ to $h(p, 0)$. The product structure on A_t will provide the natural guide for defining an $\varepsilon/6$ -homotopy $G_s: A_t \rightarrow A_t$ between g and the identity map. Note that condition (2) above and the definition of R imply that

$$\text{diam } Rh(\{p\} \times [-1, 1]) < \varepsilon/3$$

for each p in M . Thus, RG_s will be an $\varepsilon/3$ -homotopy between $Rg = g$ and R . This means that $R|_{A_t}$ will be $\varepsilon/2$ -homotopic in A_t to the identity map, and part B of this lemma will be a consequence of Lemma 4.2.

Let δ_t denote the distance from $\lambda_t(M \times \{0\})$ to $\Sigma \cap (E^{n+1} - N)$. It follows from [18] (see also [16, Lemma 5]) in case $n > 4$ and from [14, Lemma 4] or [9, Theorem 8.2] in case $n = 3$ that there exists a $d_t > 0$ such that any locally flat n -manifold in E_t homeomorphically within d_t of Σ_t is δ_t -isotopic to Σ_t in E_t . According to Lemma 4.1 there exist real numbers s and u , with $s' < s < t < u < u'$, such that λ_s and λ_u are homeomorphically within d_t of λ_t . If p_t denotes the map projecting $E^n \times E^1$ onto $E^n \times \{t\}$, then $p_t \lambda_s(M \times \{0\})$ and $p_t \lambda_u(M \times \{0\})$ are each δ_t -isotopic to $\lambda_t(M \times \{0\})$ in E_t . By lifting this isotopy through the levels E_r ($s \leq r \leq u$), we construct an embedding h of $M \times [-\frac{1}{2}, \frac{1}{2}]$ into $U \cap N$ such that condition (3), as well as the following conditions, holds:

- (4) $h(M \times \{-\frac{1}{2}\}) = \lambda_s(M \times \{0\})$,
- (5) $h(M \times \{\frac{1}{2}\}) = \lambda_u(M \times \{0\})$,
- (6) for each $w \in [-\frac{1}{2}, \frac{1}{2}]$, there exists a distinct $z \in [s, u]$ such that $h(M \times \{w\}) \subset E_z$,
- (7) $\text{diam } h(\{p\} \times [-\frac{1}{2}, \frac{1}{2}]) < \varepsilon/18$ for each p in M .

Since both λ_s and λ_u are homeomorphically close to λ_t , we may assume s and u were chosen so that

- (8) $\text{diam } \lambda_s(\{p\} \times [-1, 0]) < \varepsilon/18$ for each p in M ,
- (9) $\text{diam } \lambda_u(\{p\} \times [-1, 0]) < \varepsilon/18$ for each p in M .

Now $h|M \times [-\frac{1}{2}, \frac{1}{2}]$ can be extended to a homeomorphism h of $M \times [-1, 1]$ onto

$$A_t = \lambda_s(M \times [-1, 0]) \cup h(M \times [-\frac{1}{2}, \frac{1}{2}]) \cup \lambda_u(M \times [-1, 0]).$$

To verify that $X_t \subset N$, observe that N has connected boundary, as does X_t (where $\text{Bd } X_t$ is taken in E^{n+1} , not in $\text{Cl } U$). Since $\text{Bd } X_t \subset N$ by construction and since both X_t and N are bounded, X_t must be a subset of N . The only unverified requirement on the construction, condition (2), follows easily from (7)–(9).

ADDENDUM. Suppose $\varepsilon > 0$, X_t and A_t satisfy the conclusions of Lemma 4.3B, and Z is a compact subset of X_t . Then there exists an ε -map of Z into A_t such that $f(Z) \cap \text{Bd } A_t \subset Z$ and $f|Z \cap A_t = \text{identity}$.

Proof. Follow the ε -retraction of X_t onto A_t by a small homeomorphism g of A_t into $(Z \cap \text{Bd } A_t) \cup \text{Int } A_t$ such that $g|Z \cap A_t = \text{identity}$.

5. Submanifolds of E^{n+1} whose levels are twice bicollared. The basic result in this section is the following application of Lemma 4.3.

THEOREM 5.1. *Let Σ denote an n -manifold embedded in E^{n+1} ($n \neq 4$) as a closed subset, U a component of $E^{n+1} - \Sigma$, and (a, b) an interval such that for each $t \in (a, b)$ (i) Σ_t is a closed, PL $(n-1)$ -manifold that is collared from U_t and (ii) $\text{Cl } U - \Sigma_t$ is 1-LC at each point of Σ_t . Then U is 1-LC at each point of $\Sigma(a, b)$.*

Proof. Suppose $f: \Delta^2 \rightarrow \text{Cl } U$ is a map such that $f(\partial\Delta^2) \subset U$ and $f(\Delta^2) \cap \Sigma \subset \Sigma(a, b)$. Let ε be a positive number less than both $\rho(f(\Delta^2), \Sigma_a \cup \Sigma_b)$ and $\rho(f(\partial\Delta^2), \Sigma)$, and let $\mathcal{D}$ denote the countable subset of (a, b) described in Lemma 4.1. Then by Theorem 3.2 there exists a map $f_0: \Delta^2 \rightarrow \text{Cl } U$ such that

- (1) $f_0|_{\partial\Delta^2} = f|_{\partial\Delta^2}$,
- (2) $\rho(f, f_0) < \varepsilon/3$,
- (3) $f_0(\Delta^2) \cap \Sigma_d = \emptyset$ for each d in $\mathcal{D}$.

For each $t \in (a, b) - \mathcal{D}$, application of Lemma 4.3 yields an $\varepsilon/3$ -retraction r_t of the X_t onto the A_t associated (by Lemma 4.3B) with t and $\varepsilon/3$, where $\Sigma \cap A_t = \Sigma_{s(t)} \cup \Sigma_{u(t)}$. Let π denote the projection of $E^n \times E^1$ onto the second factor. Then $\pi(\Sigma \cap f_0(\Delta^2))$, a subset of $(a, b) - \mathcal{D}$, is covered by the collection $\mathcal{C}$ of open intervals $\mathcal{C} = \{(s(t), u(t)) \mid t \in \pi(\Sigma \cap f_0(\Delta^2))\}$, from which we can extract a finite subcovering $\mathcal{F}$ of $\pi(\Sigma \cap f_0(\Delta^2))$. After eliminating unnecessary elements of $\mathcal{F}$, we can regard $\mathcal{F}$ as the union of two (finite) collections $\mathcal{F}_1$ and $\mathcal{F}_2$ such that for $i = 1, 2$ no two elements of $\mathcal{F}_i$ intersect. Let F_1 and F_2 denote the underlying point sets of $\mathcal{F}_1$ and $\mathcal{F}_2$ respectively.

In case $(s(t), u(t)) \in \mathcal{F}_1$, the addendum to Lemma 4.3 implies the existence of an $\varepsilon/3$ map R_t of $X_t \cap f_0(\Delta^2)$ into A_t such that

- (4) $R_t(X_t \cap f_0(\Delta^2)) \cap \text{Bd } A_t \subset X_t \cap f_0(\Delta^2)$,
- (5) $R_t|_{f_0(\Delta^2) \cap A_t} = \text{identity}$.

Define a map $f_1: \Delta^2 \rightarrow \text{Cl } U$ by the rule

$$\begin{aligned} f_1(x) &= R_t f_0(x), & \text{if } f_0(x) \in X_t \text{ and } (s(t), u(t)) \in \mathcal{F}_1, \\ &= f_0(x), & \text{otherwise.} \end{aligned}$$

It follows from (5) and the fact that the X_t 's considered are pairwise disjoint that f_1 is well defined and continuous. Note that

$$(6) f_1|_{\partial\Delta^2} = f_0|_{\partial\Delta^2},$$

$$(7) \rho(f_0, f_1) < \varepsilon/3,$$

$$(8) \pi(\Sigma \cap f_1(\Delta^2)) \subset F_2.$$

Similarly, in case $(s(t), u(t)) \in \mathcal{F}_2$, the addendum to Lemma 4.3 implies the existence of an $\varepsilon/3$ -map R_t of $X_t \cap f_1(\Delta^2)$ into A_t such that

$$(9) R_t(X_t \cap f_1(\Delta^2)) \cap \text{Bd } A_t = \emptyset,$$

$$(10) R_t|_{f_1(\Delta^2)} \cap A_t = \text{identity}.$$

Define $f_2: \Delta^2 \rightarrow \text{Cl } U$ by the rule

$$\begin{aligned} f_2(x) &= R_t f_1(x), & \text{if } f_1(x) \in X_t \text{ and } (s(t), u(t)) \in \mathcal{F}_2, \\ &= f_1(x), & \text{otherwise.} \end{aligned}$$

As before, f_2 is a continuous function satisfying

$$(11) f_2|_{\partial\Delta^2} = f_1|_{\partial\Delta^2},$$

$$(12) \rho(f_1, f_2) < \varepsilon/3,$$

$$(13) f_2(\Delta^2) \cap \Sigma = \emptyset.$$

Since $f_2|_{\partial\Delta^2} = f_1|_{\partial\Delta^2}$ and $\rho(f_1, f_2) < \varepsilon$, it follows immediately that U is 1-LC at each point of $\Sigma(a, b)$.

REMARKS. Although condition (ii) clearly is a necessary hypothesis for Theorem 5.1, one questions whether it might be superfluous. Even without this condition it follows from [4], by means of the trick emerging here in Lemma 4.1, that $\text{Cl } U - \Sigma_t$ is 1-LC at each point of Σ_t for all but at most countably many points t in (a, b) , but this fact, obviously, is no help. By attacking the problem differently in the next section, we shall prove, under the hypotheses of Theorem 5.1 without condition (ii), that U is 1-LC at many points of Σ (see Corollary 6.2).

Without much extra effort one can prove the following slightly stronger version of Theorem 5.1. Statements of the other results in this section can be altered in a similar manner.

THEOREM 5.2. *Let Σ denote an n -manifold embedded in E^{n+1} ($n \neq 4$) as a closed subset, U a component of $E^{n+1} - \Sigma$, and (a, b) an interval such that (i) for each $t \in (a, b)$, $\text{Cl } U - \Sigma_t$ is 1-LC at each point of Σ_t and (ii) for all but countably many points t in (a, b) , Σ_t is a closed, PL $(n-1)$ -manifold that is collared from U_t . Then U is 1-LC at each point of $\Sigma(a, b)$.*

Proof. Simply incorporate those countably many t 's of (a, b) that fail to satisfy condition (ii) into the set $\mathcal{D}$ and reapply the proof of Theorem 5.1.

THEOREM 5.3. *Let Σ denote an n -manifold embedded in E^{n+1} ($n \neq 4$) as a closed subset, U a component of $E^{n+1} - \Sigma$, and (a, b) an interval such that for each $t \in (a, b)$ (i) Σ_t is a closed, PL $(n-1)$ -manifold that is collared from U_t and (ii) $\Sigma - \Sigma_t$ is 1-LC at each point of Σ_t . Then U is 1-LC at each point of $\Sigma(a, b)$.*

Proof. Let p be a point of Σ_t and $\varepsilon > 0$. There exists a $\delta > 0$ such that any loop in $N_\delta(p) \cap (\Sigma - \Sigma_t)$ is contractible in $N_\varepsilon(p) \cap (\Sigma - \Sigma_t)$. Since U is locally 1-connected at p in the homology sense [17, Theorem II, 5.35], it follows from [6, Proposition 3.3] that there exists an $\alpha > 0$ such that any loop in $N_\alpha(p) \cap U$ is contractible in $N_\delta(p) \cap (E^{n+1} - \Sigma_t)$. By cutting off such a contraction in $\Sigma - \Sigma_t$ one can show that each loop in $N_\alpha(p) \cap U$ is contractible in $N_\varepsilon(p) \cap (\text{Cl } U - \Sigma_t)$. Using this property one easily can prove that $\text{Cl } U - \Sigma_t$ is 1-LC at p . Hence, Theorem 5.1 gives the desired result.

THEOREM 5.4. *Let Σ denote a closed n -manifold in E^{n+1} ($n \neq 4$), U a component of $E^{n+1} - \Sigma$, and $[a, b]$ the interval such that $\Sigma = \Sigma([a, b])$. Suppose that for each t in (a, b) , Σ_t is a closed PL $(n-1)$ -manifold that is collared from U_t and that for each t in $[a, b]$, $\text{Cl } U - \Sigma_t$ is 1-ULC. Then U is 1-ULC.*

Proof. Obviously U is 1-LC at points of $\Sigma(a, b)$. Furthermore, by hypothesis, any small loop in U near a point of Σ_a is contractible in a small subset of $\text{Cl } U - \Sigma_a$. According to Theorem 3.2 such a contraction can be modified slightly so that the range of the resulting map is a small subset of U . Thus, U is 1-LC at points of Σ_a . The same argument applies to points of Σ_b . Consequently, U is 1-ULC.

Using Theorem 5.4 one can extend results of [11] and [12] to higher dimensions in the following ways.

COROLLARY 5.5. *Suppose Σ is a closed n -manifold in E^{n+1} ($n \neq 4$) such that (i) $\Sigma = \Sigma([-1, 1])$, (ii) $\Sigma - (\Sigma_{-1} \cup \Sigma_1)$ is 1-ULC, and (iii) for each $t \in (-1, 1)$, Σ_t is a closed, PL $(n-1)$ -manifold bicollared in E_t and $\Sigma - \Sigma_t$ is 1-ULC. Then $E^{n+1} - \Sigma$ is 1-ULC.*

Proof. For either component U of $E^{n+1} - \Sigma$ and each t in $[-1, 1]$, the proof of Theorem 5.3 indicates that $\text{Cl } U - \Sigma_t$ is 1-ULC. Although the hypotheses of Proposition 3.3 of [6] do not apply when $t = \pm 1$, the argument there can be used to establish the property employed in proving Theorem 5.3, namely, for each $\delta > 0$ there exists an $\alpha > 0$ such that each α -loop in U is contractible in a δ -subset of $E^{n+1} - \Sigma_t$ ($t \neq \pm 1$).

COROLLARY 5.6. *Suppose Σ is an n -sphere in E^{n+1} ($n \neq 4$) such that both Σ_1 and Σ_{-1} are points and, for each t in $(-1, 1)$, Σ_t is an $(n-1)$ -sphere bicollared in E_t and $\Sigma - \Sigma_t$ is 1-ULC. Then $E^{n+1} - \Sigma$ is 1-ULC.*

From Theorem 9 of [16] we obtain the following flatness conditions.

THEOREM 5.7. *Suppose Σ is a closed PL n -manifold in E^{n+1} ($n \geq 5$) satisfying the hypothesis of Corollary 5.5. Then Σ is locally flat if and only if Σ can be homeomorphically approximated by locally flat manifolds.*

THEOREM 5.8. *Let Σ denote the boundary of an $(n+1)$ -cell B in E^{n+1} ($n \geq 5$) and U the complement of B . Suppose (i) $\Sigma = \Sigma([a, b])$, (ii) for each $t \in (a, b)$, Σ_t is a PL $(n-1)$ -manifold that is collared from U_t , and (iii) for each $t \in [a, b]$, $\text{Cl } U - \Sigma_t$ is 1-ULC. Then Σ is locally flat.*

6. Submanifolds of E^{n+1} whose levels satisfy 1-ULC conditions. In this section we give sufficient conditions, in the spirit of Cannon's work in E^3 [7], for a complementary domain of a closed n -manifold in E^{n+1} to be 1-ULC. In place of the hypothesis typical of the results found in §5 that the Σ_i 's be collared PL manifolds stands the weakened hypothesis that the U_i 's be 1-ULC, together with strong restrictions on the embeddings of the Σ_i 's in Σ . As one advantage of this approach, the case $n=4$ need not be excluded.

THEOREM 6.1. *Suppose that Σ is a closed n -manifold in E^{n+1} , Z a component of $E^{n+1} - \Sigma$, and (a, b) an interval such that (i) $\Sigma_t = \text{Bd } Z_t$ and (ii) Z_t is both 0-ULC and 1-ULC for each t in (a, b) . Then (a, b) contains a dense G_δ -subset G such that Z is 1-LC at each point of $\Sigma(G)$.*

Proof. Equivalently we shall show that the set F of levels t in (a, b) at which Σ_t contains a point q such that Z fails to be 1-LC at q is a 0-dimensional F_σ -set.

For each positive integer n let X_n denote the set of points x in Σ such that for no neighborhood V of x is every loop of $V \cap Z$ contractible in a $(1/n)$ -subset of Z . Then X_n is a compact subset of Σ , and therefore $\pi(X_n)$ is a closed subset of E^1 , where π denotes the map projecting $E^n \times E^1$ onto the second factor. Define F_n as $(a, b) \cap \pi(X_n)$. Obviously $F = \bigcup F_n$. Hence, we only need show F_n to be 0-dimensional.

Suppose to the contrary that some F_n contains a subinterval $[a', b']$ of (a, b) . By the Baire Category Theorem $[a', b']$ then contains a subinterval $[c, d]$ such that corresponding to some dense subset Y of $[c, d]$ there exists a positive number δ with the property that each δ -loop in Z_t is null homotopic in a $(1/3n)$ -subset of Z_t ($t \in Y$). To reach the required contradiction we shall apply Hosay's argument [12] to prove that each point q of $\Sigma(c, d)$ has a neighborhood V such that every loop in $V \cap Z$ is null-homotopic in a $(1/n)$ -subset of Z .

Given any such point q let U be a round open ball in E^{n+1} containing q of diameter less than $\min\{\delta, 1/3n\}$. Assume further that U misses E_c and E_d . Let V be a neighborhood of q such that $V \subset U$ and $V \cap \Sigma$ lies in an n -cell in $U \cap (\Sigma - (\Sigma_c \cup \Sigma_d))$. We must show that any map f of the boundary of a 2-cell D into $V \cap Z$ has an extension g sending D into a $(1/n)$ -subset of Z .

Using the notation developed in [12] we trace the argument given on pp. 371–373 there with certain modifications. First, replace part A by the following observation: the hypotheses that $\Sigma_t = \text{Bd } Z_t$ and Z_t is 0-ULC imply that there exists an arc in $U \cap Z_t$ connecting each pair of points of $h(A_t^i) \cap f(\partial D)$. Let K_t^i denote the union of the (finitely many) arcs obtained in this way. Second, in part B observe that the special levels (the t 's) can be chosen from Y , since Y is dense in $[c, d]$. The last two paragraphs of B may be ignored, noting instead that any map of a simple closed curve into

$$(K^i \cup K^{i+1} \cup h(\partial D)) \cap \bigcup \{E_t | t_i \leq t \leq t_{i+1}\}$$

is homotopic, in the intersection of Z and the $(1/3n)$ -neighborhood of U , to a constant map. Making use of this fact at the end of part D, we can construct the required extension g sending D into Z .

COROLLARY 6.2. *Suppose that Σ is a closed n -manifold in E^{n+1} , U a component of $E^{n+1} - \Sigma$, and (a, b) an interval such that for each $t \in (a, b)$, Σ_t is a closed $(n-1)$ -manifold that is collared from U_t . Then (a, b) contains a dense G_δ -subset G such that U is 1-LC at each point of $\Sigma(G)$.*

THEOREM 6.3. *Suppose that Σ is a closed n -manifold in E^{n+1} , Z a component of $E^{n+1} - \Sigma$, and (a, b) an interval such that (i) $\text{Cl}(Z) - \Sigma_t$ is 1-ULC and (ii) Z_t is 1-ULC for each t in (a, b) . Then (a, b) contains a dense G_δ -subset G such that Z is 1-LC at each point of $\Sigma(G)$.*

Proof. We begin by repeating the first three paragraphs of the proof of Theorem 6.1. Then, with minor modifications similar to those given in the preceding proposition, Cannon's argument [7, Theorem 1] can be applied to complete the proof.

It would be interesting to know whether the hypotheses of either Theorem 6.1 or 6.3 actually imply that Z is 1-LC at each point of $\Sigma(a, b)$. One should note that some restrictions on the set Z are necessary, for it is quite simple to describe a connected open subset Z of E^{n+1} ($n \geq 2$) such that each Z_t is 1-ULC but Z fails to be 1-LC at certain points of $\text{Bd } Z$. In particular, a bounded open subset Z of E^{n+1} ($n \geq 2$) need not be 1-ULC even if each Z_t is 1-ULC.

THEOREM 6.4. *Suppose Σ is a closed n -manifold in E^{n+1} , U a component of $E^{n+1} - \Sigma$, and (a, b) an interval such that (1) for each $t \in (a, b)$, U_t is 1-ULC and (2) for each compact 0-dimensional subset C of (a, b) , $\text{Cl } U - \Sigma(C)$ is 1-ULC. Then U is 1-LC at each point of $\Sigma(a, b)$.*

Proof. Let f be a map of the disk Δ^2 into $\text{Cl } U$ such that $f(\partial\Delta^2) \subset U$ and $f(\Delta^2) \cap \Sigma \subset \Sigma(a, b)$. If G denotes the dense G_δ -subset of (a, b) promised by Theorem 6.3, then from hypothesis (2) above and Theorem 3.2 we find that f can be adjusted slightly, not changing the map on $\partial\Delta^2$, such that $f(\Delta^2) \cap \Sigma \subset \Sigma(G)$ and $f^{-1}(\Sigma \cap f(\Delta^2))$ is 0-dimensional. As a result, U is 1-LC at each point of $\Sigma \cap f(\Delta^2)$, and it is then a simple matter to alter f further so that $f(\Delta^2)$ misses Σ entirely. Hence, U is 1-LC at each point of $\Sigma(a, b)$.

COROLLARY 6.5. *Suppose Σ is a closed n -manifold in E^{n+1} such that each component U of $E^{n+1} - \Sigma$ satisfies (1) for each $t \in E^1$, U_t is 1-ULC and (2) for each compact, 0-dimensional subset C of E^1 , $\text{Cl } U - \Sigma(C)$ is 1-ULC. Then $E^{n+1} - \Sigma$ is 1-ULC.*

Corollary 6.5 can be interpreted as a generalization of [7, Corollary 3].

REMARK. Variations on Theorem 6.4 and Corollary 6.5 can be obtained by exchanging condition (2) in each for the following condition:

(2*) for each compact 0-dimensional subset C of (a, b) , or E^1 , as the context requires, $\Sigma - \Sigma(C)$ is 1-ULC.

The proof given for Theorem 2 of [7] establishes that to each $\delta > 0$ there corresponds an $\alpha > 0$ such that each α -loop in U is contractible in an ε -subset of $E^{n+1} - \Sigma(C)$. Using this we can prove, as in Theorem 5.3, that $\text{Cl } U - \Sigma(C)$ is 1-ULC.

Finally, Corollary 6.5 and [16, Theorem 9] can be combined, as in §5, to produce a local flatness criterion.

THEOREM 6.6. *Suppose Σ is a closed PL n -manifold in E^{n+1} ($n \geq 4$) satisfying the hypotheses of Corollary 6.5. Then Σ is locally flat if and only if Σ can be homeomorphically approximated by locally flat manifolds.*

REFERENCES

1. J. W. Alexander, *On the subdivision of 3-space by a polyhedron*, Proc. Nat. Acad. Sci. U.S.A. **10** (1924), 6–8.
2. R. H. Bing, *E^3 does not contain uncountably many mutually exclusive surfaces*, Bull. Amer. Math. Soc. **63** (1957), 404. Abstract #60-801t.
3. ———, *Pushing a 2-sphere into its complement*, Michigan Math. J. **11** (1964), 33–45. MR **28** #3408.
4. J. L. Bryant, *Concerning uncountable families of n -cells in E^n* , Michigan Math. J. **15** (1968), 477–479. MR **38** #6561.
5. ———, *On embeddings with locally nice cross-sections*, Trans. Amer. Math. Soc. **155** (1971), 327–332.
6. ———, *Euclidean n -space modulo an $(n-1)$ -cell* (to appear).
7. J. W. Cannon, **-taming sets for crumpled cubes. II. Horizontal sections in closed sets*, Trans. Amer. Math. Soc. **161** (1971), 441–446.
8. J. Cheegar and J. M. Kister, *Counting topological manifolds*, Topology **9** (1970), 149–151. MR **41** #1055.
9. Robert Craggs, *Small ambient isotopies of a 3-manifold which transform one embedding of a polyhedron into another*, Fund. Math. **68** (1970), 225–256.
10. C. H. Dowker, *Mapping theorem for non-compact spaces*, Amer. J. Math. **69** (1947), 200–242. MR **8**, 594.
11. W. T. Eaton, *Cross sectionally simple spheres*, Bull. Amer. Math. Soc. **75** (1969), 375–378. MR **39** #957.
12. Norman Hosay, *A proof of the slicing theorem for 2-spheres*, Bull. Amer. Math. Soc. **75** (1969), 370–374. MR **39** #956.
13. R. A. Jensen, *Cross sectionally connected 2-spheres are tame*, Bull. Amer. Math. Soc. **76** (1970), 1036–1038.
14. J. M. Kister, *Isotopies in 3-manifolds*, Trans. Amer. Math. Soc. **97** (1960), 213–224. MR **22** #11378.
15. L. D. Loveland, *Cross sectionally continuous spheres in E^3* , Bull. Amer. Math. Soc. **75** (1969), 396–397. MR **39** #958.
16. C. L. Seebeck, III, *Collaring an $(n-1)$ -manifold in an n -manifold*, Trans. Amer. Math. Soc. **148** (1970), 63–68. MR **41** #2692.
17. R. L. Wilder, *Topology of manifolds*, Amer. Math. Soc. Colloq. Publ., vol. 32, Amer. Math. Soc., Providence, R. I., 1963. MR **32** #440.
18. Perrin Wright, *A uniform generalized Schoenflies theorem*, Bull. Amer. Math. Soc. **74** (1968), 718–721. MR **37** #4799.