ΒY

### HARRIS S. SHULTZ

ABSTRACT. Let (a, b) be any open subinterval of the reals which contains the origin and let  $\mathfrak{V}$  denote the family of all distributions on (a, b) which are regular in some interval  $(\epsilon, 0)$ , where  $\epsilon < 0$ . Then  $\mathfrak{V}$  is a commutative algebra: Multiplication is defined so that, when restricted to those distributions on (a, b) whose supports are contained in [0, b), it is ordinary convolution. Also,  $\mathfrak{V}$  can be injected into an algebra of operators; this family of operators is a sequentially complete locally convex space. Since it preserves multiplication, this injection serves as a generalization (there are no growth restrictions) of the two-sided Laplace transformation.

In [6] there is introduced a new algebra  $\mathfrak{B}$  of distributions on  $(-\infty, \infty)$ , closed under convolution and containing the space of distributions having support in  $[0, \infty)$  as well as all locally integrable functions. No growth or support restrictions are placed on the elements of  $\mathfrak{B}$ . There is also defined a one-to-one transformation of  $\mathfrak{B}$  into a commutative algebra of operators (somewhat analogous to the Fourier transformation). In the present article we generalize these results in obtaining a space  $\mathfrak{B}$  of distributions on  $\Omega$ , where  $\Omega$  is any open subinterval of the reals which contains the origin. A distribution F on  $\Omega$  belongs to  $\mathfrak{B}$  if and only if F is regular in some interval ( $\epsilon$ , 0), where  $\epsilon < 0$ . Convolution is defined and  $\mathfrak{B}$  is shown to be closed under this operation. It is also shown that the algebra  $\mathfrak{C}$  into which  $\mathfrak{B}$  can be injected is a sequentially complete locally convex space in which convergence is defined simply in terms of the ordinary pointwise convergence of functions.

0. Preliminaries. Throughout we assume  $-\infty \le a < 0 < b \le \infty$  and set  $\Omega = (a, b)$ . We define L to be the space of all the complex-valued functions which are Lebesgue integrable on each compact subinterval of  $\Omega$ . We denote by  $L_+$  (respectively,  $L_-$ ) the subspace consisting of those elements of L which vanish on (a, 0) (respectively, (0, b)). If f and g belong to L then the function  $f \land g$  defined by the equation

(0.01) 
$$f \wedge g(t) = \int_0^t f(t-u)g(u) \, du \qquad (t \in \Omega)$$

Copyright © 1972, American Mathematical Society

Presented to the Society, January 19, 1972; received by the editors June 10, 1971.

AMS 1970 subject classifications. Primary 44A40, 44A35, 42A76; Secondary 46F99. Key words and phrases. Generalized functions, operational calculus, Schwartz distri-

butions, two-sided Laplace transformation, Fourier transformation.

also belongs to L, moreover, if we identify functions which are equal almost everywhere on  $\Omega$  then

$$(0.02) f \wedge g = g \wedge f$$

(see [4]). For any f in L we define

$$f_{+}(t) = \begin{cases} f(t), & 0 \le t < b, \\ 0, & t < 0, \end{cases} \text{ and } f_{-}(t) = \begin{cases} 0, & t \ge 0, \\ f(t), & a < t < 0. \end{cases}$$

If  $\Omega_0$  is an open subinterval of the reals we denote by  $\mathfrak{D}(\Omega_0)$  the space of complex-valued infinitely differentiable functions defined on the reals which vanish outside of a compact subset of  $\Omega_0$ . If  $\phi \in \mathfrak{D}(\Omega_0)$  we define the support of  $\phi$ , denoted supp  $\phi$ , to be the closure of the set  $\{t: \phi(t) \neq 0\}$ . Then supp  $\phi \in \Omega_0$  for all  $\phi$  in  $\mathfrak{D}(\Omega_0)$ .

As usual, the dual of  $\mathfrak{D}(\Omega_0)$ , that is, the space of distributions on  $\Omega_0$ , is denoted by  $\mathfrak{D}'(\Omega_0)$ . If R belongs to  $\mathfrak{D}'(\Omega_0)$  and  $\phi$  belongs to  $\mathfrak{D}(\Omega_0)$  the scalar which R assigns to  $\phi$  will be written  $(R(x), \phi(x))$ . If f belongs to the family of locally integrable functions on  $\Omega_0$  and m is a nonnegative integer we shall write  $\partial^m f$  for the element of  $\mathfrak{D}'(\Omega_0)$  defined by

$$\langle \partial^m f(x), \phi(x) \rangle = (-1)^m \int_{\mathbf{\Omega}_0} f(x) \phi^{(m)}(x) dx \qquad (\phi \in \mathfrak{D}(\mathbf{\Omega}_0)).$$

In particular,  $\partial^0 f$  is the *regular* distribution corresponding to the function f. The support of a distribution R on  $\Omega_0$  (denoted supp R) is defined to be the complement with respect to  $\Omega_0$  of the largest open set on which R vanishes.

1. The algebra  $\mathfrak{B}$ . We denote by  $\mathfrak{D}'_b$  the space of elements in  $\mathfrak{D}'((-\infty, b))$  having support in [0, b). We denote by  $\mathfrak{D}'_a$  the space of elements in  $\mathfrak{D}'((a, \infty))$  having support in (a, 0].

1.01. Definition. Suppose  $\{b_n\}$  and  $\{a_n\}$  are sequences of real numbers, that  $\{J_n\}$  and  $\{K_n\}$  are sequences of nonnegative integers and that  $\{F_n\}$  and  $\{G_n\}$  are sequences in L. If the ordered pair (R, S) belongs to the cartesian product  $\mathfrak{D}'_b \times \mathfrak{D}'_a$  we say that the sequence  $\{(F_n, b_n, J_n, G_n, a_n, K_n)\}$  belongs to  $\Sigma_{R,S}$  if  $(1.01.1) \ a \leftarrow \cdots < a_2 < a_1 < a_0 = 0 = b_0 < b_1 < b_2 < \cdots \rightarrow b;$ 

(1.01.2)  $F_n$  vanishes on  $(-\infty, b_n)$  and  $G_n$  vanishes on  $(a_n, \infty)$ ;

(1.01.3) 
$$R = \sum_{n=0}^{\infty} \partial^{J_n} F_n$$
 and  $S = \sum_{n=0}^{\infty} \partial^{K_n} G_n$ .

1.02. Theorem. Given any (R, S) in  $\mathfrak{D}'_b \times \mathfrak{D}'_a$  and any sequences  $\{b_n\}$  and  $\{a_n\}$  satisfying (1.01.1) there exists an element  $\{(F_n, b_n, J_n, G_n, a_n, K_n)\}$  of  $\Sigma_{R,S}$ .

164

**Proof.** By [8, 2.17] there exists a sequence  $\{F_n\}$  in L such that the equation

$$R = \sum_{n=0}^{\infty} \partial^{J_n} F_n$$

holds for some sequence  $\{J_n\}$  of nonnegative integers. We define an element T of  $\mathfrak{D}'((-\infty, -a))$  as follows:

(1) 
$$\langle T(x), \phi(x) \rangle = \langle S(x), \phi(-x) \rangle \quad (\phi \in \mathcal{D}((-b, -a))).$$

Then, since supp  $S \subset (a, 0]$ , the distribution T has support contained in [0, -a). Since  $0 = -a_0 < -a_1 < \cdots < -a$  we may infer from [8, 2.17] the existence of a sequence  $\{H_n\}$  in  $L^{loc}((-\infty, -a))$  such that  $H_n$  vanishes on  $(-\infty, -a_n)$  and such that the equation

(2) 
$$T = \sum_{n=0}^{\infty} \partial^{K_n} H_n$$

holds for some sequence  $\{K_n\}$  of nonnegative integers. If we define

$$G_n(x) = (-1)^{K_n} H(-x)$$

then  $G_n$  vanishes on  $(a_n, \infty)$  and we may combine (1) and (2) to obtain

$$S = \sum_{n=0}^{\infty} \partial^{K_n} G_n$$

Therefore,  $\{(F_n, b_n, J_n, G_n, a_n, K_n)\} \in \Sigma_{R, S}$ .

1.03. Definition. For each  $\phi$  in  $\mathfrak{D}((-\infty, b))$  we define  $[\phi]^+$  to be the family of infinitely differentiable functions  $\lambda$  on the reals such that  $\lambda$  is equal to 1 on a neighborhood of  $[0, \infty)$  and vanishes on some interval  $(-\infty, a')$ , where  $\operatorname{supp} \phi \subset$  $(-\infty, a' + b)$ . For each  $\phi$  in  $\mathfrak{D}((a, \infty))$  we define  $[\phi]^-$  to be the family of infinitely differentiable functions  $\mu$  on the reals such that  $\mu$  is equal to 1 on a neighborhood of  $(-\infty, 0]$  and vanishes on some interval  $(b', \infty)$ , where  $\operatorname{supp} \phi \subset (a + b', \infty)$ .

1.04. Theorem. Suppose (r, s) and (R, S) belong to  $\mathfrak{D}'_b \times \mathfrak{D}'_a$ . If  $\{(f_n, b_n, j_n, g_n, a_n, k_n)\}$  belongs to  $\Sigma_{r,s}$  and  $\{(F_n, b_n, J_n, G_n, a_n, K_n)\}$  belongs to  $\Sigma_{R,S}$ , then for any  $\phi$  in  $\mathfrak{D}((-\infty, b))$  the equation

(1.04.1) 
$$\langle r(y), \langle R(x), \lambda(y)\phi(x+y)\rangle\rangle = \lim_{N\to\infty}\sum_{m=0}^{N}\sum_{n=0}^{N}\langle \partial^{j_m+J_n}(f_m\wedge F_n)(x), \phi(x)\rangle\rangle$$

holds for all  $\lambda$  in  $[\phi]^+$ , and for any  $\phi$  in  $\mathfrak{D}((a, \infty))$  the equation

(1.04.2)  
$$-\langle (s(y), \langle S(x), \mu(y)\phi(x+y)\rangle \rangle$$
$$= \lim_{N \to \infty} \sum_{m=0}^{N} \sum_{n=0}^{N} \langle \partial^{k_{m}+K_{n}}(g_{m} \wedge G_{n})(x), \phi(x) \rangle$$

holds for all  $\mu$  in  $[\phi]^-$ .

**Proof.** Suppose  $\phi \in \mathfrak{D}((-\infty, b))$  and  $\lambda \in [\phi]^+$ . There exist numbers  $\beta$  and a' such that

(1) 
$$\operatorname{supp} \phi \subset (-\infty, \beta] \subset (-\infty, a' + b)$$

and such that  $\lambda$  vanishes on  $(-\infty, a')$ . For any y the function  $x \mapsto \lambda(y)\phi(x + y)$ is infinitely differentiable. From (1) it follows that its support is contained in  $(-\infty, \beta - y]$ . Thus, for  $y \ge a'$ , its support is contained in  $(-\infty, \beta - a']$  and therefore in  $(-\infty, b)$ . And, for  $y \le a'$ , it vanishes identically (since  $\lambda(y) = 0$ ). Consequently, the function  $x \mapsto \lambda(y)\phi(x + y)$  belongs to  $\mathfrak{D}((-\infty, b))$  and has support in  $(-\infty, \beta - a']$  for all y. There exists N such that  $b_{N+1} \ge \beta - a'$  and therefore

(2)  

$$\langle R(x), \lambda(y)\phi(x+y) \rangle = \sum_{n=0}^{\infty} (-1)^{J_n} \int_0^b F_n(x)\lambda(y)\phi^{(J_n)}(x+y) \, dx$$

$$= \sum_{n=0}^N (-1)^{J_n} \int_0^b F_n(x)\lambda(y)\phi^{(J_n)}(x+y) \, dx$$

for all y (recall that  $F_n$  vanishes on  $(-\infty, b_n)$ ). From (2) and [2, 250] it follows that the function

(3) 
$$y \mapsto \langle R(x), \lambda(y)\phi(x+y) \rangle$$

is infinitely differentiable. From (1) comes the equality

(4) 
$$(R(x), \lambda(y)\phi(x+y)) = 0$$
 (all  $y > \beta$ )

(recall supp  $R \subset [0, b)$ ). And

$$\langle R(x), \lambda(y)\phi(x+y)\rangle = 0$$
 (all  $y < a'$ )

since  $\lambda$  vanishes on  $(-\infty, a')$ . Consequently, the function (3) belongs to  $\mathfrak{D}((-\infty, b))$ . Moreover, we may combine (4) and the inequality  $b_{N+1} > \beta$  to obtain

$$\int_0^b f_m(y) \langle R(x), \lambda(y)\phi(x+y) \rangle \, dy = 0 \quad (all \ m > N).$$

Therefore,

166

1972]

....

$$\langle r(y), \langle R(x), \lambda(y)\phi(x+y) \rangle \rangle$$

$$= \sum_{m=0}^{N} (-1)^{j_m} \int_0^b f_m(y) \frac{d^{j_m}}{dy^{j_m}} \langle R(x), \lambda(y)\phi(x+y) \rangle \, dy$$

$$= \sum_{m=0}^{N} \sum_{n=0}^{N} (-1)^{j_m+J_n} \int_0^b f_m(y) \left( \int_0^b F_n(x)\phi^{(j_m+J_n)}(x+y) \, dx \right) dy;$$

the last equality is from (2), [2, 250] and the fact that  $\lambda = 1$  on  $[0, \infty)$ . We may now use the change of variable t = x + y and [2, 283] to obtain

We need only observe now that

$$\int_0^t f_m(y) F_n(t-y) \, dy = 0 \qquad (0 \le t \le \beta)$$

for m > N and n > N to obtain (1.04.1). Suppose now that  $\phi \in \mathfrak{D}((a, \infty))$  and  $\mu \in \mathfrak{D}((a, \infty))$  $[\phi]^-$ . There exist numbers a and b' such that

(5) 
$$\operatorname{supp} \phi \subset [\alpha, \infty) \subset (a + b', \infty)$$

and such that  $\mu$  vanishes on  $(b', \infty)$ . For any y the function  $x \mapsto \mu(y) \phi(x + y)$  is infinitely differentiable. From (5) it follows that its support is contained in  $[a-y,\infty)$ . Thus, for  $y \leq b'$ , its support is contained in  $[a-b',\infty)$  and therefore in  $(a, \infty)$ . And, for y > b', it vanishes identically (since  $\mu(y) = 0$ ). Consequently, the function  $x \mapsto \mu(y) \phi(x + y)$  belongs to  $\mathfrak{D}((a, \infty))$  and has support in  $[a-b',\infty)$  for all y. There exists N such that  $a_{N+1} < a-b'$  and therefore

(6)  
$$\langle S(x), \, \mu(y)\phi(x+y)\rangle = \sum_{n=0}^{\infty} (-1)^{K_n} \int_a^0 G_n(x)\mu(y)\phi^{(K_n)}(x+y) \, dx$$
$$= \sum_{n=0}^{N} (-1)^{K_n} \int_a^0 G_n(x)\mu(y)\phi^{(K_n)}(x+y) \, dx$$

for all y (recall that  $G_n$  vanishes on  $(a_n, \infty)$ ). From (6) and [2, 250] it follows that the function

(7) 
$$y \mapsto \langle S(x), \mu(y)\phi(x+y) \rangle$$

is infinitely differentiable. From (5) comes the equality

(8) 
$$\langle S(x), \mu(y)\phi(x+y) \rangle = 0$$
 (all  $y < \alpha$ )

(recall supp  $S \subset (a, 0]$ ). And

$$\langle S(x), \mu(y)\phi(x+y)\rangle = 0$$
 (all  $y > b'$ )

since  $\mu$  vanishes on  $(b', \infty)$ . Consequently, the function (7) belongs to  $\mathfrak{D}((a, \infty))$ . Moreover, we may combine (8) and the inequality  $a_{N+1} < \alpha$  to obtain

$$\int_a^0 g_m(y) \langle S(x), \mu(y)\phi(x+y) \rangle \, dy = 0 \quad (all \ m > N).$$

Therefore,

$$\langle s(y), \langle S(x), \mu(y)\phi(x+y) \rangle \rangle$$

$$= \sum_{m=0}^{N} (-1)^{k_{m}} \int_{a}^{0} g_{m}(y) \frac{d^{k_{m}}}{dy^{i_{m}}} \langle S(x), \mu(y)\phi(x+y) \rangle dy$$

$$= \sum_{m=0}^{N} \sum_{n=0}^{N} (-1)^{k_{m}+K_{n}} \int_{a}^{0} g_{m}(y) \left( \int_{a}^{0} G_{n}(x)\phi^{(k_{m}+K_{n})}(x+y) dx \right) dy;$$

the last equality is from (5), [2, 250] and the fact that  $\mu = 1$  on  $(-\infty, 0]$ . We may now use the change of variable t = x + y and [2, 283] to obtain

$$\langle s(y), \langle S(x), \mu(y)\phi(x+y)\rangle$$

$$= \sum_{m=0}^{N} \sum_{n=0}^{N} (-1)^{k_{m}+K_{n}} \int_{a}^{0} \left( \int_{a}^{y} g_{m}(y) G_{n}(t-y)\phi^{(k_{m}+K_{n})}(t) dt \right) dy$$

$$= \sum_{m=0}^{N} \sum_{n=0}^{N} (-1)^{k_{m}+K_{n}} \int_{a}^{0} \left( \int_{t}^{0} g_{m}(y) G_{n}(t-y) dy \right) \phi^{(k_{m}+K_{n})}(t) dt.$$

We need only observe now that

$$\int_{t}^{0} g_{m}(y) G_{n}(t-y) \, dy = 0 \qquad (\alpha \leq t \leq 0)$$

for m > N and n > N and that

$$-\int_{t}^{0}g_{m}(y)G_{n}(t-y)\,dy=g_{n}\wedge G_{n}(t)$$

to obtain (1.04.2).

168

1.05. Corollary. Suppose that (r, s) and (R, S) belong to  $\mathfrak{D}'_b \times \mathfrak{D}'_a$ . For any  $\phi$  in  $\mathfrak{D}((-\infty, b))$  the family

$$\{\langle r(y), \langle R(x), \lambda(y)\phi(x+y)\rangle\}: \lambda \in [\phi]^+\}$$

contains a unique element, which will be denoted by  $(r * R(x), \phi(x))$ . For any  $\phi$  in  $\mathfrak{D}((a, \infty))$  the family

$$\{\langle s(y), \langle S(x), \mu(y)\phi(x+y)\rangle\}: \mu \in [\phi]^{-}\}$$

contains a unique element, which will be denoted by  $\langle s * S(x), \phi(x) \rangle$ .

1.06. Definition. Let (r, s) and (R, S) belong to  $\mathfrak{D}'_b \times \mathfrak{D}'_a$ . We denote by r \* R the functional that assigns to any  $\phi$  in  $\mathfrak{D}((-\infty, b))$  the number  $(r * R(x), \phi(x))$ ; we denote by s \* S the functional that assigns to any  $\phi$  in  $\mathfrak{D}((a, \infty))$  the number  $(s * S(x), \phi(x))$ .

1.07. Remark. It follows from 1.04 and (0.02) that r \* R = R \* r and s \* S = S \* s.

1.08. Corollary. If (r, s) and (R, S) belong to  $\mathfrak{D}'_b \times \mathfrak{D}'_a$  then (r \* R, s \* S) belongs to  $\mathfrak{D}'_b \times \mathfrak{D}'_a$ .

**Proof.** It follows from (1.04.1) and the sequential completeness of  $\mathfrak{D}'((-\infty, b))$ (see [1, Proposition 2, p. 315]) that r \* R belongs to  $\mathfrak{D}'((-\infty, b))$ . Moreover, since r \* R is the limit of a sequence of distributions on  $(-\infty, b)$  all having support in [0, b), it too must have support in [0, b), i.e.  $r * R \in \mathfrak{D}'_b$ . Similarly,  $s * S \in \mathfrak{D}'_a$ .

1.09. Remark. If f belongs to L then  $\partial^0 f_+ \in \mathfrak{D}'_b$  and  $\partial^0 f_- \in \mathfrak{D}'_a$ . We may deduce from 1.01 and 1.04 that the equations  $\partial^0 f_+ * \partial^0 g_+ = \partial^0 (f_+ \wedge g_+)$  and  $-\partial^0 f_- * \partial^0 g_- = \partial^0 (f_- \wedge g_-)$  hold for all f and g in L.

1.10. Definition. We denote by  $\mathfrak{B}_{-}$  the linear subspace consisting of those elements of  $\mathfrak{D}'_{a}$  which are regular in a neighborhood of the origin.

1.11. Remark. Thus  $S \in \mathfrak{B}_{-}$  if and only if  $S = \partial^{0}f + T$  where  $f \in L_{-}$  and  $T \in \mathfrak{D}_{a}'$  with supp  $T \subset (a, 0)$ . In particular,  $\partial^{0}f_{-} \in \mathfrak{B}_{-}$  for all  $f \in L$ .

1.12. Lemma. Suppose (R, S) and  $(R_1, S_1)$  belong to  $\mathfrak{D}'_b \times \mathfrak{B}_-$ . If the elements R + S and  $R_1 + S_1$  of  $\mathfrak{D}'(\Omega)$  are equal then  $R = R_1$  and  $S = S_1$ .

**Proof.** Since S and  $S_1$  both vanish on (0, b) we have

(1) 
$$S = S_1$$
 on  $(0, b)$ .

Since R and  $R_1$  both vanish on (a, 0) it follows from  $R + S = R_1 + S_1$  that

(2) 
$$S = S_1$$
 on  $(a, 0)$ .

Now, there exists  $\epsilon > 0$  and elements f and  $f_1$  of  $L_-$  such that  $S = \partial^0 f$  and  $S_1 = \partial^0 f_1$  on  $(-\epsilon, \epsilon)$ . From (1) and (2) it follows that  $\partial^0 f = \partial^0 f_1$  on  $(0, \epsilon)$  and on  $(-\epsilon, 0)$ . Thus, by [9, p. 224] we have  $f = f_1$  almost everywhere on  $(-\epsilon, \epsilon)$ , from which it follows that  $\partial^0 f = \partial^0 f_1$  on  $(-\epsilon, \epsilon)$ . Therefore,

(3) 
$$S = \partial^0 f = \partial^0 f_1 = S_1 \quad \text{on } (-\epsilon, \epsilon).$$

We may now combine (1), (2) and (3) to conclude that  $S = S_1$  on (a, b) (see [9, Theorem 24.1]).

1.13. Definition. We denote by  $\mathfrak{B}$  the linear subspace consisting of those elements of  $\mathfrak{D}'(\Omega)$  of the form R + S where  $R \in \mathfrak{D}'_h$  and  $S \in \mathfrak{B}_-$ .

1.14. **Remark.** Thus,  $F \in \mathfrak{B}$  if and only if  $F \in \mathfrak{D}'(\Omega)$  and is regular in some neighborhood ( $\epsilon$ , 0), where  $a \leq \epsilon \leq 0$ . In particular,  $\partial^0 f \in \mathfrak{B}$  for all  $f \in L$ .

1.15. Theorem. If F belongs to  $\mathfrak{B}$  there exists a unique element of  $\mathfrak{D}'_b \times \mathfrak{B}_-$ , denoted  $(F_+, F_-)$ , such that  $F = F_+ + F_-$ .

Proof. Immediate from 1.12.

1.16. Corollary. The mapping  $F \mapsto (F_+, F_-)$  is an isomorphism of  $\mathfrak{B}$  into  $\mathfrak{D}'_b \times \mathfrak{D}'_a$ .

**Proof.** One may easily verify that the mapping  $F \mapsto (F_+, F_-)$  is linear. The corollary then follows from 1.15.

1.17. Lemma. If V and V<sub>1</sub> belong to  $\mathfrak{D}'_a$  with supp  $V_1 \subset (a, \epsilon]$  for some  $\epsilon < 0$  then supp  $V * V_1 \subset (a, \epsilon]$ .

**Proof.** Let  $\phi \in \mathfrak{D}((a, \infty))$  and have support in  $[\epsilon', \infty)$ , where  $\epsilon < \epsilon' < 0$ . Then, for  $y < \epsilon' - \epsilon$  the function  $x \mapsto \phi(x + y)$  has support contained in  $(\epsilon, \infty)$ . Therefore,

$$\langle V_1(x), \mu(y)\phi(x+y)\rangle = 0$$
 (all  $\mu \in [\phi]$ )

for all  $y < \epsilon' - \epsilon$ . Thus, the function  $y \mapsto \langle V_1(x), \mu(y)\phi(x+y) \rangle$  has support contained in  $(0, \infty)$  for all  $\mu \in [\phi]^-$ . Since V vanishes on this interval it follows that

$$\langle V * V_1(x), \phi(x) \rangle = \langle V(y), \langle V_1(x), \mu(y)\phi(x+y) \rangle = 0$$

for all  $\mu$  in  $[\phi]^-$ . Therefore,  $V * V_1$  vanishes on  $(\epsilon, \infty)$ .

1.18. Theorem. If F and G belong to  $\mathfrak{B}$  then  $F_+ * G_+ - F_- * G_-$  belongs to  $\mathfrak{B}$  with  $(F_+ * G_+ - F_- * G_-)_= - F_- * G_-$ .

**Proof.** It suffices to show that  $F_* \in \mathcal{B}_-$ . By 1.11 there exist f and g in  $L_-$  and T and U in  $\mathfrak{D}'_{\sigma}$  such that  $F_- = \partial^0 f + T$  and  $G_- = \partial^0 g + U$  with

supp  $T \subset (a, \epsilon]$  and supp  $U \subset (a, \epsilon]$  for some  $\epsilon < 0$ . Therefore,

$$F_{-} * G_{-} = (\partial^{0} f + T) * (\partial^{0} g + U) = (\partial^{0} f) * (\partial^{0} g) + (\partial^{0} f) * U + T * (\partial^{0} g) + T * U.$$

From 1.09 and 1.07 it follows that

$$F_* * G_{-} = \partial^0 (f \wedge g) + (\partial^0 f) * U + (\partial^0 g) * T + T * U.$$

If we set  $S = (\partial^0 f) * U + (\partial^0 g) * T + T * U$  we may infer from 1.17 that supp  $S \subset (a, \epsilon]$  and therefore  $F_- * G_- \in \mathfrak{B}_-$ .

1.19. Definition. If F and G belong to  $\mathcal{B}$  we denote the element  $F_+ * G_+ - F_- * G_-$  of  $\mathcal{B}$  by  $F \wedge G$ .

1.20. Remark. As a consequence of 2.23, the space  $\mathfrak{B}$ , with multiplication defined by 1.19, is a commutative algebra.

1.21. Theorem. The equation  $\partial^0(f \wedge g) = (\partial^0 f) \wedge (\partial^0 g)$  holds for all f and g in L.

**Proof.** Since  $f \wedge g = f_+ \wedge g_+ + f_- \wedge g_-$  we may use 1.09 to obtain

$$\partial^{0}(f \wedge g) = \partial^{0}(f_{+} \wedge g_{+}) + \partial^{0}(f_{-} \wedge g_{-}) = \partial^{0}f_{+} * \partial^{0}g_{+} - \partial^{0}f_{-} * \partial^{0}g_{-}$$
$$= (\partial^{0}f)_{+} * (\partial^{0}g)_{+} - (\partial^{0}f)_{-} * (\partial^{0}g)_{-} = (\partial^{0}f) \wedge (\partial^{0}g).$$

2. The algebra of operators. Let W be the space of all the complex-valued infinitely differentiable functions w on  $\Omega$  such that  $w^{(k)}(0) = 0$  for  $k \ge 0$ . In [4] it is shown that  $f \land w$  belongs to W with

$$(2.01) (f \wedge w)' = f \wedge w'$$

whenever f belongs to L and w belongs to W. We denote by  $\langle f \rangle$  the operator which assigns to each w in W the function  $f \wedge w$  in W. Thus,  $\langle f \rangle w = f \wedge w$  (all w in W). Let  $\mathcal{C}$  be the set of all the operators A mapping W into itself such that

for all  $w_1$  and  $w_2$  in W. We make  $\mathfrak{A}$  into a vector space by defining addition and scalar multiplication in the usual way. We define the product of two operators to be the composition of the operators. Then  $\mathfrak{A}$  is a commutative algebra which contains the identity operator I and the differentiation operator D; moreover, the mapping  $f \mapsto \langle f \rangle$  is a linear injection of L into  $\mathfrak{A}$  and

(2.03) 
$$\langle f \wedge g \rangle = \langle f \rangle \langle g \rangle$$

for all f and g in L (see [4]).

2.04. Theorem. If  $\{(F_n, b_n, J_n, G_n, a_n, K_n)\}$  belongs to  $\Sigma_{R,S}$  for some (R, S) in  $\mathfrak{D}'_b \times \mathfrak{D}'_a$  then the equation

$$Aw(t) = \sum_{n=0}^{\infty} \left( \left(F_n \wedge w\right)^{(J_n)}(t) + \left(G_n \wedge w\right)^{(K_n)}(t) \right) \quad (t \in \Omega, w \in W)$$

defines an element of  $\mathfrak{A}$ .

**Proof.** Let  $w \in W$  and  $a < \alpha < \beta < b$ . Then there exists a positive integer N such that  $a_{N+1} < \alpha < \beta < b_{N+1}$ . Since  $F_n$  vanishes on  $(a, b_n)$  we may infer that  $F_n \wedge w$  vanishes on  $(a, \beta)$  for all n > N; since  $G_n$  vanishes on  $(a_n, b)$  we may infer that  $G_n \wedge w$  vanishes on  $(\alpha, b)$  for all n > N. Consequently,

(1) 
$$Aw(t) = \sum_{n=0}^{N} ((F_n \wedge w)^{(J_n)}(t) + (G_n \wedge w)^{(K_n)}(t)) \quad (\alpha < t < \beta).$$

Since each  $F_n \wedge w$  and each  $G_n \wedge w$  is infinitely differentiable on  $(\alpha, \beta)$  it follows from (1) that Aw is infinitely differentiable on  $(\alpha, \beta)$ ; and, clearly, every derivative of Aw vanishes at the origin since the same is true of each term on the right-hand side of (1). Since  $(\alpha, \beta)$  was an arbitrary open subinterval of  $\Omega$ we may conclude that  $Aw \in W$ . There remains to show that the equation  $A(w_1 \wedge w_2) =$  $(Aw_1) \wedge w_2$  holds for all  $w_1$  and  $w_2$  in W. But, using (2.01) and the fact that  $\langle F_n \rangle$  and  $\langle G_n \rangle$  belong to  $\mathcal{A}$  we may deduce that

$$\begin{split} A(w_1 \wedge w_2) &= \sum_{n=0}^{\infty} \left( (F_n \wedge (w_1 \wedge w_2))^{(J_n)} + (G_n \wedge (w_1 \wedge w_2))^{(K_n)} \right) \\ &= \sum_{n=0}^{\infty} \left( ((F_n \wedge w_1) \wedge w_2)^{(J_n)} + ((G_n \wedge w_1)^{(\Lambda w_2)})^{(K_n)} \right) \\ &= \sum_{n=0}^{\infty} \left( (F_n \wedge w_1)^{(J_n)} \wedge w_2 + (G_n \wedge w_1)^{(K_n)} \wedge w_2 \right) \\ &= \left( \sum_{n=0}^{\infty} \left( (F_n \wedge w_1)^{(J_n)} + (G_n \wedge w_1)^{(K_n)} \right) \right) \wedge w_2 = (Aw_1) \wedge w_2 \end{split}$$

For each w in W and each t in  $\Omega$  the equation  $\rho_{w,t}(A) = |Aw(t)|$  defines a seminorm on the space  $\Omega$ . Let  $\Omega$  be endowed with the locally convex topology defined by the family of seminorms  $\{\rho_{w,t}: t \in \Omega, w \in W\}$ .

2.05. Remark. If  $\{A_n\}$  is a sequence in  $\mathfrak{A}$  then  $A_0 = \lim A_n$  if and only if  $A_0 w(t) = \lim A_n w(t)$  for all w in W and all t in  $\Omega$ .

172

2.06. Definition. If  $0 \le t < b$  we denote by [t] the set of all infinitely differentiable functions p which assume the value 1 on a neighborhood of  $[0, \infty)$ and which vanish on some interval  $(-\infty, \alpha_p)$ , where  $t - b < \alpha_p$ . If a < t < 0 we let [t] denote the set of all infinitely differentiable functions q which assume the value 1 on a neighborhood of  $(-\infty, 0]$  and which vanish on some interval  $(\beta_q, \infty)$ , where  $\beta_q < t - a$ .

2.07. Definition. If w belongs to W and  $0 \le t < b$  we define

$$w_t(x) = \begin{cases} w(t-x), & t-b < x < t, \\ 0, & \text{otherwise}; \end{cases}$$

if w belongs to W and a < t < 0 we define

$$w_t(x) = \begin{cases} w(t-x), & t < x < t-a, \\ 0, & \text{otherwise.} \end{cases}$$

2.08. Remark. If  $w \in W$  and  $0 \le t < b$ , the function  $w_t$  is infinitely differentiable on  $(t - b, \infty)$  and vanishes on  $(t, \infty)$ ; thus, if  $p \in [t]$ , the function  $pw_t$  (the pointwise product of the functions p and  $w_t$ ) belongs to  $\mathfrak{D}((-\infty, b))$  with  $\operatorname{supp} pw_t \subset (-\infty, t]$ . If  $w \in W$  and a < t < 0, the function  $w_t$  is infinitely differentiable on  $(-\infty, t-a)$  and vanishes on  $(-\infty, t)$ ; thus, if  $q \in [t]$ , the function  $qw_t$  belongs to  $\mathfrak{D}((a, \infty))$  with  $\operatorname{supp} qw_t \subset [t, \infty)$ .

2.09. Lemma. If f belongs to L and m is a nonnegative integer, the equation

$$\langle \partial^m f_+(x), p(x)w_+(x) \rangle = (f \wedge w)^{(m)}(t) \qquad (0 \le t < b)$$

holds for any p in [t] and any w in W and the equation

$$- \left\langle \partial^m f_{-}(x), q(x)w_t(x) \right\rangle = (f \wedge w)^{(m)}(t) \qquad (a < t < 0)$$

holds for any q in [t] and any w in W.

**Proof.** Let  $0 \le t \le b$  and  $p \in [t]$ . For any  $w \in W$ ,

$$\langle \partial^m f_+(x), p(x)w_t(x) \rangle = (-1)^m \int_0^b f(x) [pw_t]^{(m)}(x) dx.$$

Since p = 1 on [0, b) and since  $w_t$  vanishes on  $(t, \infty)$  it follows that

$$\langle \partial^m f_+(x), p(x)w_t(x) \rangle = (-1)^m \int_0^t f(x) [w_t]^{(m)}(x) dx$$

By observing that  $(-1)^m [w_t]^{(m)}(x) = w^{(m)}(t-x)$  for x > t-b, we may use (2.01) to obtain

$$\langle \partial^m f_+(x), p(x)w_t(x) \rangle = \int_0^t f(x)w^{(m)}(t-x) dx = (f \wedge w)^{(m)}(t).$$

Now, let  $a \le t \le 0$  and  $q \in [t]$ . For any  $w \in W$ ,

$$\begin{aligned} \langle \partial^m f_{-}(x), \ q(x)w_t(x) \rangle &= (-1)^m \int_a^0 f(x) [qw_t]^{(m)}(x) \, dx \\ &= (-1)^m \int_t^0 f(x) [w_t]^{(m)}(x) \, dx = \int_t^0 f(x) w^{(m)}(t-x) \, dx \\ &= -\int_0^t f(x) w^{(m)}(t-x) \, dx = -(f \wedge w)^{(m)}(t). \end{aligned}$$

2.10. Theorem. If (R, S) belongs to  $\mathfrak{D}'_b \times \mathfrak{D}'_a$  there exists an element A of  $\mathfrak{A}$  such that the equation

(2.10.1)  $\langle R(x), p(x)w_t(x) \rangle = Aw(t) \quad (0 \le t < b)$ 

holds for any p in [t] and any w in W and such that the equation

(2.10.2) 
$$-\langle S(x), q(x)w_t(x) \rangle = Aw(t) \quad (a < t < 0)$$

bolds for any q in [t] and any w in W. If  $\{(F_n, b_n, J_n, G_n, a_n, K_n)\}$  belongs to  $\Sigma_{R,S}$  then

(2.10.3) 
$$A = \sum_{n=0}^{\infty} \left( D^{J_n} \langle f_n \rangle + D^{K_n} \langle g_n \rangle \right).$$

**Proof.** Let  $\{(F_n, b_n, J_n, G_n, a_n, K_n)\} \in \Sigma_{R,S}$ . Then, by 2.09,

(1) 
$$\langle R(x), p(x)w_t(x) \rangle = \sum_{n=0}^{\infty} \langle \partial^J {}^n F_n(x), p(x)w_t(x) \rangle = \sum_{n=0}^{\infty} (F_n \wedge w)^{(J_n)}(t)$$

for  $0 \le t \le b$ , any  $p \in [t]$  and any w in W. Similarly,

(2) 
$$-\langle S(x), q(x)w_t(x)\rangle = \sum_{n=0}^{\infty} -\langle \partial^{K_n}G_n(x), q(x)w_t(x)\rangle = \sum_{n=0}^{\infty} (G_n \wedge w)^{(K_n)}(t)$$

for a < t < 0, any  $q \in [t]$  and any w in W. If we define

$$Aw(t) = \sum_{n=0}^{\infty} ((F_n \wedge w)^{(J_n)}(t) + (G_n \wedge w)^{(K_n)}(t)) \quad (t \in \Omega, w \in W)$$

[July

174

then  $A \in \mathfrak{A}$  by 2.04. Moreover, since each  $F_n \in L_+$  and each  $G_n \in L_-$  we have

$$Aw(t) = \begin{cases} \sum_{n=0}^{\infty} (F_n \wedge w)^{(J_n)}(t) & \text{for } 0 \le t < b, \\ \sum_{n=0}^{\infty} (G_n \wedge w)^{(K_n)}(t) & \text{for } a < t < 0, \end{cases}$$

from which follows the theorem.

2.11. Corollary. Suppose that (R,S) belongs to  $\mathbb{D}'_b \times \mathbb{D}'_a$  and  $w \in W$ . If  $0 \leq t < b$ , the family  $\{\langle R(x), p(x)w_t(x) \rangle : p \in [t]\}$  contains a unique element. If a < t < 0, the family  $\{\langle S(x), q(x)w_t(x) \rangle : q \in [t]\}$  contains a unique element. If we define

$$\langle (R, S) \rangle w(t) = \begin{cases} \langle R(x), p(x)w_t(x) \rangle & (p \in [t], 0 \le t < b) \\ - \langle S(x), q(x)w_t(x) \rangle & (q \in [t], a < t < 0) \end{cases}$$

and denote by  $\langle (R, S) \rangle$  the mapping  $w \mapsto \langle (R, S) \rangle w$ , then  $\langle (R, S) \rangle \in \mathbb{C}$ .

2.12. Corollary. If  $(R, S) \in \mathfrak{D}'_b \times \mathfrak{D}'_a$  and  $\{(F_n, b_n, J_n, G_n, a_n, K_n)\} \in \Sigma_{R,S}$ , then

(2.12.1) 
$$\langle (R, S) \rangle = \sum_{n=0}^{\infty} (D^{J_n} \langle F_n \rangle + D^{K_n} \langle G_n \rangle).$$

Proof. Immediate from (2.10.3).

2.13. Corollary. The equation  $\langle (\partial^m f_+, \partial^n f_-) \rangle = D^m \langle f_+ \rangle + D^n \langle f_- \rangle$  holds for all f in L and all nonnegative integers m and n.

Proof. Immediate from 2.09.

2.14. Lemma. There exists a sequence  $\{w_n\}$  in W such that

$$(2.14.1) A = \lim_{n \to \infty} \langle Aw_n \rangle$$

for all A in C.

**Proof.** Choose  $w_n$  in W satisfying

(1) 
$$w_n \ge 0 \quad \text{on } (0, b),$$

$$w_n \leq 0 \quad \text{on } (a, 0),$$

(3) 
$$w_n(t) = 0 \text{ for } |t| \ge 1/n,$$

(4) 
$$\int_{0}^{b} w_{n}(x) dx = 1 = -\int_{a}^{0} w_{n}(x) dx$$

(cf. [1, p. 166]). Let  $A \in \mathbb{C}$  and  $w \in W$ . Then, for  $0 \le t \le b$  and n sufficiently large so that t - 1/n > 0,

$$(Aw) \wedge w_n(t) = \int_{t-1/n}^t Aw(x)w_n(t-x)\,dx.$$

And, by (3)-(4), we may write

$$Aw(t) = \int_{t-1/n}^{t} Aw(t)w_n(t-x)\,dx.$$

Consequently,

$$|(Aw) \wedge w_{n}(t) - Aw(t)| = \left| \int_{t-1/n}^{t} [Aw(x) - Aw(t)]w_{n}(t-x) dx \right|$$
  
$$\leq \sup_{|x-t| \leq 1/n} |Aw(x) - Aw(t)| \int_{t-1/n}^{t} w_{n}(t-x) dx$$
  
$$= \sup_{|x-t| \leq 1/n} |Aw(x) - Aw(t)|.$$

Using (5) and the continuity of Aw at t we obtain

(6) 
$$Aw(t) = \lim_{n \to \infty} (Aw) \wedge w_n(t) \quad (0 < t < b).$$

For a < t < 0 and n sufficiently large so that t + 1/n > 0,

$$(Aw) \wedge w_n(t) = -\int_t^{t+1/n} Aw(x)w_n(t-x)\,dx.$$

And, by (3)-(4), we may write

$$Aw(t) = -\int_t^{t+1/n} Aw(t)w_n(t-x)\,dx.$$

Consequently,

$$|(Aw) \wedge w_n(t) - Aw(t)| = \left| -\int_t^{t+1/n} [Aw(x) - Aw(t)]w_n(t-x) dx \right|$$

(7) 
$$\leq \sup_{|x-t|\leq 1/n} |Aw(x) - Aw(t)| \left(-\int_{t}^{t+1/n} w_n(t-x) dx\right)$$

$$= \sup_{|x-t| \leq 1/n} |Aw(x) - Aw(t)|.$$

Using (7) and the continuity of Aw at t we obtain

(8) 
$$Aw(t) = \lim_{n \to \infty} (Aw) \wedge w_n(t) \quad (a < t < 0).$$

Observing that  $(Aw) \wedge w_n = (Aw_n) \wedge w$  and the fact that  $(Aw_n) \wedge w(0) = 0 = Aw(0)$ we infer from (6) and (8) that  $Aw(t) = \lim_{n \to \infty} (Aw_n) \wedge w(t)$  (all  $t \in \Omega$ ) and therefore that  $A = \lim \langle Aw_n \rangle$ .

2.15. Remark. It follows from 2.14 that each A in  $\mathfrak{A}$  is linear.

2.16. Lemma. If  $\{(R_n, S_n)\}$  is a sequence in  $\mathfrak{D}'_b \times \mathfrak{D}'_a$  and if  $R_0 = \lim R_n$ and  $S_0 = \lim S_n$  then  $\langle (R_0, S_0) \rangle = \lim \langle (R_n, S_n) \rangle$ .

**Proof.** Let  $w \in W$ . If  $R_0 = \lim R_n$  and  $S_0 = \lim S_n$  then

$$\langle R_0(x), \phi(x) \rangle = \lim_{n \to \infty} \langle R_n(x), \phi(x) \rangle \quad (\text{all } \phi \in \mathfrak{D}((-\infty, b))),$$
  
 
$$\langle S_0(x), \phi(x) \rangle = \lim_{n \to \infty} \langle S_n(x), \phi(x) \rangle \quad (\text{all } \phi \in \mathfrak{D}((a, \infty))).$$

Therefore, for 0 < t < b,

$$\langle (R_0, S_0) \rangle w(t) = \langle R_0(x), p(x)w_t(x) \rangle$$
  
=  $\lim_{n \to \infty} \langle R_n(x), p(x)w_t(x) \rangle = \lim_{n \to \infty} \langle (R_n, S_n) \rangle w(t)$  (all  $p \in [t]$ ),

and, for a < t < 0,

$$\langle (R_0, S_0) \rangle w(t) = -\langle S_0(x), q(x)w_t(x) \rangle$$

$$= \lim_{n \to \infty} -\langle S_n(x), q(x)w_t(x) \rangle = \lim_{n \to \infty} \langle (R_n, S_n) \rangle w(t) \quad (all \ q \in [t]).$$

2.17. Definition. For any  $\phi$  in  $\mathfrak{D}((-\infty, \infty))$  and any real t we define  $\phi_t(x) = \phi(t-x)$  for all x.

2.18. Theorem. The mapping  $(R, S) \mapsto \langle\!\!\langle R, S \rangle\!\!\rangle$  is a linear bijection of  $\mathfrak{D}'_b \times \mathfrak{D}'_a$  onto  $\mathfrak{A}$ .

**Proof.** It is easily seen that the mapping is linear. We show first that it is "onto." Let  $A \in \mathcal{C}$  and define

(see 2.14). Then  $\partial^0 f_n \in \mathfrak{D}'_b$  and  $\partial^0 g_n \in \mathfrak{D}'_a$ . For any  $\phi$  in  $\mathfrak{D}((-\infty, b))$  there exists  $t \in (0, b)$  such that  $\operatorname{supp} \phi \subset (-\infty, t]$  and therefore  $\phi_t \in W$ . Thus,

177

(1) 
$$\langle \partial^0 f_n(x), \phi(x) \rangle = \langle \partial^0 f_n(x), (\phi_t)_t(x) \rangle = \langle f_n \rangle \phi_t(t)$$

Combining (1) and 2.14 we have  $\lim_{n\to\infty} \langle \partial^0 f_n(x), \phi(x) \rangle = A(\phi_t)(t)$ . Thus the sequence  $\{ \langle \partial^0 f_n(x), \phi(x) \rangle \}$  converges for all  $\phi$  in  $\mathfrak{D}((-\infty, b))$ . By [1, Proposition 2, p. 315] there exists R in  $\mathfrak{D}'((-\infty, b))$  such that  $R = \lim \partial^0 f_n$ ; it is easily seen that  $R \in \mathfrak{D}'_b$ . For any  $\phi$  in  $\mathfrak{D}((a, \infty))$  there exists  $t \in (a, 0)$  such that supp  $\phi \subset [t, \infty)$  and therefore  $\phi_t \in W$ . Thus,

(2) 
$$- \langle \partial^0 g_n(x), \phi(x) \rangle = - \langle \partial^0 g_n(x), (\phi_t)_t(x) \rangle = \langle g_n \rangle \phi_t(t).$$

Combining (2) and 2.14 we have  $\lim_{n\to\infty} -\langle \partial^0 g_n(x), \phi(x) \rangle = A(\phi_t)(t)$ . Thus the sequence  $\{\langle \partial^0 g_n(x), \phi(x) \rangle\}$  converges for all  $\phi$  in  $\mathfrak{D}((a, \infty))$ . We may similarly infer the existence of S in  $\mathfrak{D}'_a$  such that  $S = \lim \partial^0 g_n$ . We may now use 2.16, 2.13 and 2.14 to obtain

$$\langle (R, S) \rangle = \left\langle \left( \lim_{n \to \infty} \partial^0 f_n, \lim_{n \to \infty} \partial^0 g_n \right) \right\rangle$$
$$= \lim_{n \to \infty} \left\langle (\partial^0 f_n, \partial^0 g_n) \right\rangle = \lim_{n \to \infty} \langle A w_n \rangle = A;$$

whence the mapping  $(R, S) \mapsto \langle (R, S) \rangle$  is "onto." If A = 0 then each  $f_n$  and each  $g_n$  equal 0, from which it follows that  $R = \lim_{n \to \infty} \partial^0 f_n = 0$  and  $S = \lim_{n \to \infty} \partial^0 g_n = 0$ . The mapping  $(R, S) \mapsto \langle (R, S) \rangle$  is therefore one-to-one.

2.19. Theorem. The space  $\mathfrak{A}$  is sequentially complete.

**Proof.** Suppose  $\{A_n\}$  is a Cauchy sequence in  $\mathfrak{A}$ . By 2.17 there exists a unique  $(R_n, S_n)$  in  $\mathfrak{D}'_b \times \mathfrak{D}'_a$  such that  $\langle (R_n, S_n) \rangle = A_n$  and, by assumption, the sequence  $\{\langle (R_n, S_n) \rangle w(t) \}$  converges for all w in W and all  $t \in \Omega$ . For any  $\phi$  in  $\mathfrak{D}((-\infty, b))$  there exists  $t \in (0, b)$  such that  $\operatorname{supp} \phi \subset (-\infty, t]$  and therefore  $\phi_t \in W$ . Since  $\langle R_n(x), \phi(x) \rangle = \langle R_n(x), p(x)(\phi_t)_t(x) \rangle = \langle (R_n, S_n) \rangle \phi_t(t)$  for all  $p \in [t]$ , the sequence  $\{\langle R_n(x), \phi(x) \rangle\}$  converges for all  $\phi$  in  $\mathfrak{D}((-\infty, b))$ . For any  $\phi$  in  $\mathfrak{D}((a, \infty))$  there exists  $t \in (a, 0)$  such that  $\operatorname{supp} \phi \subset [t, \infty)$  and therefore  $\phi_t \in W$ . Since  $-\langle S_n(x), \phi(x) \rangle = -\langle S_n(x), q(x)(\phi_t)_t(x) \rangle = \langle (R_n, S_n) \rangle \phi_t(t)$  the sequence  $\{\langle S_n(x), \phi(x) \rangle\}$  converges for all  $\phi$  in  $\mathfrak{D}((a, \infty))$ . We may again use [1, Proposition 2, p. 315] to infer the existence of (R, S) in  $\mathfrak{D}'_b \times \mathfrak{D}'_a$  such that  $R = \lim R_n$  and  $S = \lim S_n$ . By 2.16 we then have

$$\langle (R, S) \rangle = \lim_{n \to \infty} \langle (R_n, S_n) \rangle = \lim_{n \to \infty} A_n.$$

2.20. Lemma. If (r, s) and (R, S) belong to  $\mathfrak{D}'_b \times \mathfrak{D}'_a$  then  $\langle (r * R, -s * S) \rangle = \langle (r, s) \rangle \langle (R, S) \rangle$ .

178

**Proof.** Let  $\{(F_n, b_n, J_n, G_n, a_n, K_n)\} \in \Sigma_{R,S}$  and  $\{(f_n, b_n, j_n, g_n, a_n, k_n)\} \in \Sigma_{r,S}$ . By 1.05 and 1.04,

$$r * R = \lim_{N \to \infty} \sum_{m=0}^{N} \sum_{n=0}^{N} \partial^{j_m + J_n} (f_m \wedge F_n),$$
  
$$-s * S = \lim_{N \to \infty} \sum_{m=0}^{N} \sum_{n=0}^{N} \partial^{k_m + K_n} (g_m \wedge G_n).$$

Therefore, by 2.16,

$$\langle (r * R, -s * S) \rangle = \lim_{N \to \infty} \sum_{m=0}^{N} \sum_{n=0}^{N} \langle (\partial^{j_m + J_n} (f_m \wedge F_n), \partial^{k_m + K_n} (G_m \wedge G_n)) \rangle$$

$$= \lim_{N \to \infty} \sum_{m=0}^{N} \sum_{n=0}^{N} \langle D^{j_m + J_n} \langle f_m \wedge F_n \rangle + D^{k_m + K_n} \langle g_m \wedge G_n \rangle )$$

$$= \lim_{N \to \infty} \sum_{m=0}^{N} \sum_{n=0}^{N} \langle D^{j_m} \langle f_m \rangle D^{J_n} \langle F_n \rangle + D^{k_m} \langle g_m \rangle D^{K_n} \langle G_n \rangle );$$

the second equality is from 2.13 and the third equality is from (2.03). Let  $w \in W$ and  $t \in \Omega$ . Choose N sufficiently large so that  $a_{N+1} < t < b_{N+1}$ . Suppose first that  $0 \le t < b$ . Then

$$\langle (r * R, -s * S) \rangle w(t) = \sum_{m=0}^{N} \sum_{n=0}^{N} D^{j_m} \langle f_m \rangle D^{J_n} \langle F_n \rangle w(t)$$

$$= \sum_{m=0}^{N} D^{j_m} \langle f_m \rangle \left( \sum_{n=0}^{N} D^{J_n} \langle F_n \rangle w \right) (t)$$

$$= \sum_{m=0}^{\infty} D^{j_m} \langle f_m \rangle \left( \sum_{n=0}^{N} D^{J_n} \langle F_n \rangle w \right) (t)$$

$$= \langle (r, s) \rangle \left( \sum_{n=0}^{N} D^{J_n} \langle F_n \rangle w \right) (t);$$

the last equality is from 2.12. Therefore, by 2.15 and the fact that  ${\mathfrak C}$  is a commutative algebra, we have

$$\langle (r * R, -s * S) \rangle w(t) = \sum_{n=0}^{N} D^{J_n} \langle F_n \rangle (\langle (r, s) \rangle w)(t)$$
$$= \sum_{n=0}^{\infty} D^{J_n} \langle F_n \rangle (\langle (r, s) \rangle w)(t) = \langle (R, S) \rangle (\langle (r, s) \rangle w)(t)$$

 $= \langle (R, S) \rangle \langle (r, s) \rangle w(t) = \langle (r, s) \rangle \langle (R, S) \rangle w(t).$ 

And, if 
$$a < t < 0$$
, then  

$$\langle \langle r * R, -s * S \rangle \rangle w(t) = \sum_{m=0}^{N} \sum_{n=0}^{N} D^{k_m} \langle g_m \rangle D^{K_n} \langle G_n \rangle w(t)$$

$$= \sum_{m=0}^{N} D^{k_m} \langle g_m \rangle \left( \sum_{n=0}^{N} D^{K_n} \langle G_n \rangle w \right)(t) = \sum_{m=0}^{\infty} D^{k_m} \langle g_m \rangle \left( \sum_{n=0}^{N} D^{K_n} \langle G_n \rangle w \right)(t)$$

$$= \langle \langle (r, s) \rangle \left( \sum_{n=0}^{N} D^{K_n} \langle G_n \rangle w \right)(t) = \sum_{n=0}^{N} D^{K_n} \langle G_n \rangle \langle \langle (r, s) \rangle w \rangle(t)$$

$$= \sum_{n=0}^{\infty} D^{K_n} \langle G_n \rangle \langle \langle (r, s) \rangle w \rangle(t) = \langle (R, S) \rangle \langle \langle (r, s) \rangle w \rangle(t) = \langle (r, s) \rangle \langle (R, S) \rangle w(t).$$

2.21. Definition. For any F in  $\mathfrak{B}$  we denote the element  $\langle F_+, F_- \rangle$  of  $\mathfrak{A}$  by  $\langle F \rangle$ .

2.22. Theorem. The equation  $\langle \partial^0 f \rangle = \langle f \rangle$  holds for all f in L.

**Proof.** Observing that  $(\partial^0 f)_+ = \partial^0 f_+$  and  $(\partial^0 f)_- = \partial^0 f_-$  we may combine 2.21 with 2.13 to obtain the theorem.

2.23. Theorem. The mapping  $F \mapsto \langle F \rangle$  is an isomorphism of  $\mathfrak{B}$  into  $\mathfrak{A}$  and the equation  $\langle F \wedge G \rangle = \langle F \rangle \langle G \rangle$  holds for all F and G in  $\mathfrak{B}$ .

**Proof.** The first assertion comes from combining 1.16 and 2.18. As for the second, since  $\langle F \wedge G \rangle = \langle F_+ * G_+ - F_- * G_- \rangle = \langle (F_+ * G_+, -F_- * G_-) \rangle$  (see 1.18 and 1.19), we may use 2.20 to obtain  $\langle F \wedge G \rangle = \langle (F_+, F_-) \rangle \langle (G_+, G_-) \rangle = \langle F \rangle \langle G \rangle$ .

### BIBLIOGRAPHY

1. J. Horvath, Topological vector spaces and distributions. Vol. 1, Addison-Wesley, Reading, Mass., 1966. MR 34 #4863.

180

2. H. Kestelman, Modern theories of integration, 2nd ed., Dover, New York, 1960. MR 23 #A282.

3. G. Krabbe, Operational calculus, Springer-Verlag, New York, 1970.

4. ——, An algebra of generalized functions on an open interval; two-sided operational calculus, Bull. Amer. Math. Soc. 77 (1971), 78-84; Correction, ibid., 633. MR 42 #2262; MR 43 #833.

5. \_\_\_\_\_, Initial-value problems involving generalized functions; two-sided operational calculus, Arch. Math. (Basel) (to appear).

6. — , A new algebra of distributions; initial-value problems involving Schwartz distributions (to appear).

7. -----, Linear operators and operational calculus. I, Studia Math. 40 (1971), 199-223.

8. H. Shultz, Linear operators and operational calculus. II, Studia Math. 41 (to appear).

9. F. Treves, Topological vector spaces, distributions and kernels, Academic Press, New York, 1967. MR 37 #726.

10. G. Krabbe, An algebra of generalized functions on an open interval; two-sided operational calculus, Bull. Amer. Math. Soc. 77 (1971), 78-84. MR 42 #2262.

DEPARTMENT OF MATHEMATICS, CALIFORNIA STATE COLLEGE, FULLERTON, CALIFORNIA 92631