TOPOLOGICAL PROPERTIES OF PARANORMAL OPERATORS ON HILBERT SPACE

BY

GLENN R. LUECKE

ABSTRACT. Let $B(H)$ be the set of all bounded endomorphisms (operators) on the complex Hilbert space H . $T \in B(H)$ is paranormal if $\|(T - zI)^{-1}\| = 1/d(z, \sigma(T))$ for all $z \notin \sigma(T)$ where $d(z, \sigma(T))$ is the distance from z to $\sigma(T)$, the spectrum of T . If $\mathcal{P}$ is the set of all paranormal operators on H , then $\mathcal{P}$ contains the normal operators, $\mathcal{N}$, and the hyponormal operators; and $\mathcal{P}$ is contained in $\mathcal{L}$, the set of all $T \in B(H)$ such that the convex hull of $\sigma(T)$ equals the closure of the numerical range of T . Thus, $\mathcal{N} \subseteq \mathcal{P} \subseteq \mathcal{L} \subseteq B(H)$. Give $B(H)$ the norm topology. The main results in this paper are (1) $\mathcal{N}$, $\mathcal{P}$, and $\mathcal{L}$ are nowhere dense subsets of $B(H)$ when $\dim H \geq 2$, (2) $\mathcal{N}$, $\mathcal{P}$, and $\mathcal{L}$ are arcwise connected and closed, and (3) $\mathcal{N}$ is a nowhere dense subset of $\mathcal{P}$ when $\dim H = \infty$.

Paranormal operators have received considerable attention in the current literature ([16], [17], [19], [20], [21], [23], [24], [25]). However, only the various spectral properties of paranormal operators have been discussed. In this paper, the topological properties of the set of all paranormal operators $\mathcal{P}$ on a Hilbert space H are investigated.

We begin by giving the notation to be used and by defining some of the more specialized terminology. The point spectrum and approximate point spectrum of an operator T are denoted by $\sigma_p(T)$ and $\sigma_{\pi}(T)$, respectively. $z \in \sigma_p(T)$ is a normal eigenvalue if $\{x \in H: (T - zI)x = 0\} = \{x \in H: (T - zI)^*x = 0\}$ where I denotes the identity operator on H . $z \in \sigma_{\pi}(T)$ is a normal approximate eigenvalue of T when (1) $\|x_n\| = 1$ and $\|(T - zI)x_n\| \rightarrow 0$ as $n \rightarrow \infty$ imply $\|(T - zI)^*x_n\| \rightarrow 0$ as $n \rightarrow \infty$, and (2) $\|y_n\| = 1$ and $\|(T - zI)^*y_n\| \rightarrow 0$ as $n \rightarrow \infty$ imply $\|(T - zI)y_n\| \rightarrow 0$ as $n \rightarrow \infty$. If S is a set of complex numbers, then ∂S denotes the boundary of S and $\text{co}(S)$ denotes the convex hull of S . Let $W(T) = \text{closure}\{(Tx, x): x \in H, \|x\| = 1\}$ denote the closure of the numerical range of T . Let $R(T, z) = (T - zI)^{-1}$ for each $z \in \rho(T)$, the resolvent set of T . The spectral radius of T is denoted by $R_{\text{sp}}(T)$. $T \in B(H)$ is hyponormal if $T^*T - TT^* \geq 0$.

1. Elementary properties of paranormal operators. Stampfli [19] has shown that every hyponormal operator is paranormal. Therefore, since every normal

Presented to the Society, August 28, 1970; received by the editors April 16, 1970.

AMS (MOS) subject classifications (1969). Primary 4740.

Key words and phrases. Hilbert space, resolvent, normal operator, convexoid operators, properties and topological properties of G_1 operators.

Copyright © 1973, American Mathematical Society

operator is hyponormal, $\mathcal{N} \subseteq \mathcal{P}$. G. Orland [13] proved that $T \in \mathcal{L}$ if and only if $\|R(T, z)\| \leq 1/d(z, \text{co } \sigma(T))$ for all $z \notin \text{co } \sigma(T)$. From this it follows immediately that if T is a paranormal operator, then $\text{co } \sigma(T) = W(T)$, i.e. $\mathcal{P} \subseteq \mathcal{L}$.

Theorem 1.1. *If T is a paranormal operator and if α and β are complex numbers, then $\alpha T + \beta I$ and T^* are paranormal.*

The proof is a simple computation that depends on $\sigma(\alpha T + \beta I) = \alpha \sigma(T) + \beta$ and $\sigma(T^*) = \sigma(T)^*$.

The following theorem is very useful in constructing examples.

Theorem 1.2. *If A is any operator on H , then there exists a Hilbert space K and a normal operator N on K such that $T = A \oplus N \in B(H \oplus K)$ is paranormal.*

Proof. Since $W(A)$ is a compact set of complex numbers, there exists a Hilbert space K and a normal operator N on K such that $\sigma(N) = W(A)$ [3, p. 581]. Let $T = A \oplus N$. Then $\sigma(T) = \sigma(N) = W(A)$. For $z \notin \sigma(T) = W(A)$, it is well known that $\|R(A, z)\| \leq 1/d(z, W(A)) = 1/d(z, \sigma(T))$. Since N is normal and $z \in \rho(N)$, $\|R(N, z)\| = 1/d(z, \sigma(T))$. Therefore

$$\|R(T, z)\| = \text{Max}\{\|R(A, z)\|, \|R(N, z)\|\} = 1/d(z, \sigma(T)).$$

Thus T is paranormal.

As we shall see, the class of all hyponormal operators on H is distinct, in general, from the class $\mathcal{P}$ of paranormal operators. We know [19] that if T is hyponormal, then $\|T\| = R_{\text{sp}}(T)$ and if T is also invertible, then T^{-1} is hyponormal. These properties do not generalize to paranormal operators.

Theorem 1.3. *There exists an invertible paranormal operator T such that*

1. T is not hyponormal,
2. T^2 is not paranormal,
3. $\|T\| > R_{\text{sp}}(T)$, and
4. T^{-1} is not paranormal.

Proof. Let $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$. Let N be a normal operator such that $\sigma(N) = W(A)$, and let $T = A \oplus N$. Then by Theorem 1.2, T is paranormal. We know from [19] that a hyponormal operator is hyponormal on invariant subspaces. Therefore, since A is not hyponormal, T is not hyponormal. By [1], $W(A)$ is the closed disc of radius $1/2$ about $z = 1$, and $W(A^2)$ is the closed disc of radius 1 about $z = 1$. Therefore

$$0 \in W(A^2) \subseteq W(T^2) \quad \text{and} \quad 0 \notin \text{co } (\sigma(T)^2) = \text{co } \sigma(T^2).$$

Therefore, $\text{co } \sigma(T^2) \neq W(T^2)$ and so T^2 is not paranormal. Let $x = [\sqrt{1/2}, \sqrt{1/2}]$, then $\|x\| = 1$ and $\|Ax\| = \sqrt{10}/2$. Then

$$\|T\| \geq \|Ax\| = \sqrt{10}/2 > 3/2 = R_{\text{sp}}(T).$$

Therefore $\|T\| > R_{\text{sp}}(T)$. If T^{-1} were paranormal, then

$$\|T\| = \|R(T^{-1}, 0)\| = 1/d(0, \sigma(T^{-1})) = R_{\text{sp}}(T).$$

Contradiction. Hence T^{-1} is not paranormal.

II. Topological properties of $\mathcal{P}$. Recall that $\mathcal{N}$ is the set of all normal operators on H , $\mathcal{P}$ is the set of all paranormal operators on H , $\mathcal{Q}$ is the set of all $T \in B(H)$ such that $\text{co } \sigma(T) = W(T)$, and $\mathcal{N} \subseteq \mathcal{P} \subseteq \mathcal{Q}$. It will always be assumed that $B(H)$ has the uniform operator (norm) topology.

The following notation will be used in this section: If S is a compact subset of the complex numbers $\mathbb{C}$ and if $\epsilon > 0$, then let $S + (\epsilon) = \{z: d(z, S) < \epsilon\}$. If S and S_n , $n = 1, 2, 3, \dots$, are compact sets in $\mathbb{C}$, then the sequence $\{S_n\}$ approaches S , written $S_n \rightarrow S$, if for every $\epsilon > 0$ there exists a positive integer N such that, for $n \geq N$, $S_n \subseteq S + (\epsilon)$ and $S \subseteq S_n + (\epsilon)$.

In general $\sigma(T)$ is not a continuous function of T in $B(H)$ (see [7, problem 85]), but $\sigma(T)$ is continuous if we restrict T to $\mathcal{P}$.

Theorem 2.1. *If $\{T_n\}$ is a sequence of paranormal operators approaching the operator T in norm, then $\sigma(T_n) \rightarrow \sigma(T)$ as $n \rightarrow \infty$.*

To prove this theorem we need the following lemma from [7, problem 86].

Lemma. *If $T \in B(H)$ and $\epsilon > 0$, then there exists $\delta > 0$ such that if $S \in B(H)$ and $\|T - S\| < \delta$, then $\sigma(S) \subseteq \sigma(T) + (\epsilon)$.*

Proof of Theorem 2.1. We know by the lemma that for each $\epsilon > 0$ there exists a positive integer N such that, for $n \geq N$, $\sigma(T_n) \subseteq \sigma(T) + (\epsilon)$. Therefore, to show $\sigma(T_n) \rightarrow \sigma(T)$, it suffices to show that for each $\epsilon > 0$ there exists a positive integer N such that $\sigma(T) \subseteq \sigma(T_n) + (\epsilon)$ for all $n \geq N$. If this does not hold, then without loss of generality we may assume that there exists $\epsilon > 0$ and a sequence $\{z_n\} \subseteq \sigma(T)$ such that $d(z_n, \sigma(T_n)) \geq \epsilon$ for all n . Since $\sigma(T)$ is compact, we may assume $z_n \rightarrow z \in \sigma(T)$. If $|z_n - z| < \epsilon/2$, then

$$d(z, \sigma(T_n)) \geq d(z_n, \sigma(T_n)) - |z - z_n| \geq \epsilon - \epsilon/2 = \epsilon/2.$$

Hence

$$\|R(T_n, z)\| = 1/d(z, \sigma(T_n)) \leq 2/\epsilon.$$

Now choose m so that $\|(T_m - T)R(T_m, z)\| < 1$, then $I - (T_m - T)R(T_m, z)$ is invertible [7, problem 173]. Let

$$A = R(T_m, z)(I - (T_m - T)R(T_m, z))^{-1}.$$

Then $A(T - zI) = (T - zI)A = I$ so that $z \in \rho(T)$. Contradiction.

Remark. Note that the full strength of the paranormality assumption was not

used in this proof, and that one can easily devise various larger classes of operators on which $\sigma(T)$ is continuous.

Theorem 2.2. $\mathcal{P}$ is an arcwise connected, closed subset of $B(H)$.

Proof. Since $T \in \mathcal{P}$ implies $aT \in \mathcal{P}$ for every complex number a , we see that the ray in $B(H)$ through T is contained in $\mathcal{P}$. Therefore $\mathcal{P}$ is arcwise connected.

Suppose $T_n \rightarrow T$, $\{T_n\}$ a sequence of operators in $\mathcal{P}$, and $T \in B(H)$. Let $z \in \rho(T)$. By the lemma to Theorem 2.1,

$$\limsup_{n \rightarrow \infty} \frac{1}{d(z, \sigma(T_n))} \leq \frac{1}{d(z, \sigma(T))}.$$

Therefore, since $\|R(T_n, z)\| = 1/d(z, \sigma(T_n))$ whenever $z \in \rho(T_n)$, there exists a positive integer N such that the sequence $\{\|R(T_n, z)\|: n \geq N\}$ is bounded. Then, since $R(T, z) - R(T_n, z) = R(T, z)(T - T_n)R(T_n, z)$, $\|R(T_n, z)\| \rightarrow \|R(T, z)\|$ as $n \rightarrow \infty$. Consequently,

$$\|R(T, z)\| = \lim_{n \rightarrow \infty} \|R(T_n, z)\| = \lim_{n \rightarrow \infty} \frac{1}{d(z, \sigma(T_n))} \leq \frac{1}{d(z, \sigma(T))}.$$

Since the general $\|R(T, z)\| \geq 1/d(z, \sigma(T))$, T is paranormal.

Theorem 2.3. $\mathcal{L}$ is an arcwise connected, closed subset of $B(H)$.

Proof. Since $T \in \mathcal{L}$ implies that $aT \in \mathcal{L}$ for every complex number a , $\mathcal{L}$ is arcwise connected.

Let $T_n \rightarrow T$, $\{T_n\} \subseteq \mathcal{L}$ and $T \in B(H)$. Since $|(T_n x, x) - (Tx, x)| \leq \|T_n - T\|$ for $\|x\| = 1$, $W(T_n) \subseteq W(T) + (2\|T - T_n\|)$ and $W(T) \subseteq W(T_n) + (2\|T - T_n\|)$. Consequently, $W(T_n) \rightarrow W(T)$. Let $\epsilon > 0$, then by the lemma to Theorem 2.1 there exists a positive integer N such that $\sigma(T_n) \subseteq \sigma(T) + (\epsilon)$ for all $n \geq N$. Therefore, for $n \geq N$, $\text{co } \sigma(T_n) \subseteq \text{co } \sigma(T) + (\epsilon)$ and hence

$$W(T) = \lim_{n \rightarrow \infty} W(T_n) = \lim_{n \rightarrow \infty} \text{co } \sigma(T_n) \subseteq \text{co } \sigma(T) + (\epsilon).$$

Since $\epsilon > 0$ is arbitrary, $W(T) \subseteq \text{co } \sigma(T)$. Since in general $\text{co } \sigma(T) \subseteq W(T)$, $T \in \mathcal{L}$.

Let $\mathcal{N}$ be the set of all normal operators on H . Since $\|T_n - T\| \rightarrow 0$ implies $\|T_n^* - T^*\| \rightarrow 0$, $\mathcal{N}$ is closed in the uniform operator topology on $B(H)$. Since $T \in \mathcal{N}$ implies $aT \in \mathcal{N}$ for any complex a , $\mathcal{N}$ is arcwise connected.

We already know that $\mathcal{N} \subseteq \mathcal{P} \subseteq \mathcal{L} \subseteq B(H)$. We will now investigate the relative topological properties of these four sets. Stampfli [20, Theorem C] has shown that $\mathcal{N} = \mathcal{P}$ when $\dim H < \infty$. However, when $\dim H = \infty$, then $\mathcal{N}$ is a very "thin" subset of $\mathcal{P}$.

Theorem 2.4. $\mathcal{N}$ is a nowhere dense subset of $\mathcal{P}$ when $\dim H = \infty$.

Proof. Since $\mathcal{N}$ is closed, to show that $\mathcal{N}$ is a nowhere dense subset of $\mathcal{P}$,

it suffices to show that $\mathcal{N}$ has empty interior in $\mathcal{P}$. Let $T \in \mathcal{N}$ and let $\epsilon > 0$.

First suppose that T has an eigenvalue of infinite multiplicity. We may assume that the eigenvalue is zero. Let M be the eigenspace of zero. Then $\dim M = \infty$, M reduces T , and we can write $T = B \oplus Z$ where Z is the zero operator on M . Let $S = \begin{bmatrix} 0 & \epsilon \\ 0 & 0 \end{bmatrix} \oplus N$ be a nonnormal paranormal operator [see Theorem 1.2] on M with N a normal operator such that $\sigma(N) = W\begin{bmatrix} 0 & \epsilon \\ 0 & 0 \end{bmatrix}$. Then $B \oplus S$ is a nonnormal paranormal operator such that $\|T - B \oplus S\| = \|B \oplus Z - B \oplus S\| = \|S\| = \epsilon$. The last equality holds since $\|N\| = R_{sp}(N) = \epsilon/2$ and $\|\begin{bmatrix} 0 & \epsilon \\ 0 & 0 \end{bmatrix}\| = \epsilon$. Therefore, since $\epsilon > 0$ is arbitrary, T is not contained in the interior of $\mathcal{N}$ in $\mathcal{P}$.

If $\sigma(T)$ is finite and $T \in \mathcal{N}$, then $\sigma(T) = \sigma_p(T)$ and T has an eigenvalue of infinite multiplicity. We therefore assume that $\sigma(T)$ is infinite and that zero is an accumulation point of $\sigma(T)$. Let D be the open disc about zero of radius $\epsilon/2$. Let E be the resolution of the identity for T so that $T = \int_{\sigma(T)} z dE_z$. Let $M = (E(D))(H)$, $P = \sigma(T) - D$, and let $A = \int_P z dE_z$. Then M reduces T , $\dim M = \infty$, and A is a normal operator. Let Z be the zero operator on M . Then $A \oplus Z$ is a normal operator with zero an eigenvalue of infinite multiplicity, and

$$\|T - A \oplus Z\| = \left\| \int_D z dE_z \right\| \leq \epsilon/2.$$

By the first part of this proof, there exists a nonnormal paranormal operator S such that $\|A \oplus Z - S\| < \epsilon/2$. Then

$$\|T - S\| \leq \|T - A \oplus Z\| + \|A \oplus Z - S\| < \epsilon.$$

Therefore, since $\epsilon > 0$ is arbitrary, T is not contained in the interior of $\mathcal{N}$ in $\mathcal{P}$. Hence the interior of $\mathcal{N}$ in $\mathcal{P}$ is empty.

Define C_2 to be the set of all operators $T \in \mathcal{L}$ with $W(T)$ a closed line segment or a point. For $k = 3, 4, 5, \dots$, let C_k be the set of all operators $T \in \mathcal{L}$ such that $W(T)$ is the convex hull of a nondegenerate polygon with k sides. If $T \in C_k$, then each vertex of $W(T)$ must be in the spectrum of T [1]. S. Hildebrandt [10, Theorem 2] has shown that, if $z \in \sigma_p(T) \cap \partial W(T)$, then z is a normal eigenvalue of T . Thus for $T \in C_k$, the vertices of $W(T)$ are normal eigenvalues of T , when $\dim H < \infty$. Hence, all the operators in $C_n \cup C_{n-1}$ are normal operators when $\dim H = n < \infty$.

In [10, Theorem 9] S. Hildebrandt showed that $\mathcal{N} = \mathcal{P} = \mathcal{L}$ when $\dim H \leq 4$, and that $\mathcal{N} \neq \mathcal{L}$ for $5 \leq \dim H < \infty$. The following theorem gives more information on how $\mathcal{P}$ and $\mathcal{L}$ are related when $5 \leq \dim H < \infty$. Recall that $\mathcal{N} = \mathcal{P}$ for $\dim H < \infty$.

Theorem 2.5. *If $5 \leq \dim H = n < \infty$, then the interior of $\mathcal{P}$ in $\mathcal{L}$ equals $C_n \cup C_{n-1}$.*

Proof. Suppose $T \in C_n \cup C_{n-1}$. Since $C_n \cup C_{n-1} \subseteq \mathcal{N}$, T is normal. First, we show that there exists $\epsilon > 0$ such that whenever $S \in \mathcal{L}$ and $\|T - S\| < \epsilon$, then

$S \in C_n \cup C_{n-1}$. To show this, suppose the statement were false. Then there would exist $\{S_n\} \subseteq \mathcal{L}$ such that $\|T - S_n\| \rightarrow 0$ and $S_n \in \bigcup_{i=2}^{n-2} C_i$. Then, since $W(S_n) \rightarrow W(T)$, $T \in \bigcup_{i=2}^{n-2} C_i$. Contradiction. Hence, T is an interior point of $\mathcal{P}$ in $\mathcal{L}$.

Let T be contained in the interior of $\mathcal{P} = \mathcal{N}$ in $\mathcal{L}$. Suppose $T \notin C_n \cup C_{n-1}$. Let $\epsilon > 0$. Since $\text{co } \sigma(T) = W(T)$, $T \in C_k$, for some $k \leq n-2$. Since $\dim H \geq 5$ and since T is a normal operator in C_k , there exists a normal operator N such that

1. $\|T - N\| < \epsilon/2$,
2. $W(N)$ is a polygon with at least three sides, and
3. N has at least two eigenvalues z, w contained in the interior of $W(N)$.

Write $N = A \oplus B$ where B can be written as $B = \begin{bmatrix} z & 0 \\ 0 & w \end{bmatrix}$. Let $a > 0$ and let $C = \begin{bmatrix} z & a \\ 0 & w \end{bmatrix}$; then $\|B - C\| = \|\begin{bmatrix} 0 & a \\ 0 & 0 \end{bmatrix}\| = a$. Choose $a > 0$ small enough so that $W(C) \subseteq W(N)$ and so that $a < \epsilon/2$. Then since $W(A) = W(N)$ and $\sigma(A \oplus C) = \sigma(N)$, $\text{co } \sigma(A \oplus C) = W(A \oplus C)$. Hence $A \oplus C \in \mathcal{L}$. Since $A \oplus C$ is not normal, and since

$$\|T - A \oplus C\| \leq \|T - N\| + \|N - A \oplus C\| \leq \epsilon/2 + \epsilon/2 = \epsilon,$$

T is not an interior point of $\mathcal{P}$ in $\mathcal{L}$. Contradiction. Hence $T \in C_n \cup C_{n-1}$.

It is an open question as to what the interior of $\mathcal{P}$ in $\mathcal{L}$ is when $\dim H = \infty$. However, it can be shown that $\mathcal{P} \neq \mathcal{L}$ when $\dim H = \infty$.

Theorem 2.6. $\mathcal{P} \neq \mathcal{L}$ when $\dim H = \infty$.

Proof. Write $H = M \oplus M^\perp$ where $\dim M = 5$. Let

$$A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \quad \text{and} \quad N = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}$$

where a, b, c are three complex numbers that form a triangle with $W(A)$ contained in the interior of the triangle. Consider $A \oplus N$ as an operator on M and observe that $\text{co } \sigma(A \oplus N) = W(N) = W(A \oplus N)$. Since $A \oplus N$ is not normal and since $\dim M < \infty$, $A \oplus N$ is not paranormal. Hence there exists $z \in \rho(A \oplus N)$ such that

$$\|R(A \oplus N, z)\| > 1/d(z, \sigma(A \oplus N)).$$

Let I be the identity operator on $M^\perp$ and let $T = A \oplus N \oplus aI$. Then, $\sigma(T) = \sigma(A \oplus N)$ and $W(T) = W(A \oplus N)$. Therefore $T \in \mathcal{L}$. Since $d(z, \sigma(T)) \leq |z - a|$,

$$\begin{aligned} \|R(T, z)\| &= \text{Max} \{ \|R(A \oplus N, z)\|, |z - a|^{-1} \} \\ &= \|R(A \oplus N, z)\| > 1/d(z, \sigma(T)). \end{aligned}$$

Therefore T is not paranormal.

We now show that if $\dim H \geq 2$, then $\mathcal{L}$ is a nowhere dense ("thin") subset of $B(H)$. Once this is shown, it follows immediately that $\mathcal{N}$ and $\mathcal{P}$ are also nowhere dense subsets of $B(H)$.

Theorem 2.7. $\mathcal{L}$ is a nowhere dense subset of $B(H)$ when $\dim H \geq 2$.

To prove this theorem we need the following two technical lemmas.

Lemma 1. *If z_1 and z_2 are distinct, normal approximate eigenvalues of $T \in B(H)$, then there exist sequences $\{x_n\}$ and $\{y_n\}$ of unit vectors in H such that*

1. $(x_n, y_n) = 0$ for all n ,
2. $\|(T - z_1 I)x_n\| \rightarrow 0$ as $n \rightarrow \infty$, and
3. $\|(T - z_2 I)y_n\| \rightarrow 0$ as $n \rightarrow \infty$.

Lemma 2. *If $T \in \mathfrak{L}$ such that there exists distinct $a, b \in \partial W(T) \cap \sigma_p(T)$, then T is not contained in the interior of $\mathfrak{L}$.*

Proof of Theorem 2.7. Since $\mathfrak{L}$ is closed (Theorem 2.3), to show that $\mathfrak{L}$ is nowhere dense it suffices to show that $\mathfrak{L}$ has empty interior.

We first show that if T is in the interior of $\mathfrak{L}$, then $\sigma(T)$ must contain at least two points. Suppose $T \in \mathfrak{L}$ and $\sigma(T) = \{a\}$. Then $((T - aI)x, x) = 0$ for all $x \in H$ so that $T = aI$. Since $\dim H \geq 2$, write $H = M \oplus M^\perp$ where $\dim M = 2$. Let $b > 0$ and define $A \in B(H)$ as

$$A = \begin{bmatrix} 0 & b \\ 0 & 0 \end{bmatrix} \text{ on } M, \quad \text{and } A = 0 \text{ on } M^\perp.$$

Then $\sigma(T + A) = \{a\}$ and, since $b \neq 0$, $\{a\} \neq W(T + A)$. Therefore $T + A \notin \mathfrak{L}$. Since $\|A\| = b > 0$ is arbitrary, $T \notin \text{interior } \mathfrak{L}$.

With the above remark completed, we can now finish the proof of Theorem 2.7. Suppose the theorem were false and there exists $T \in \text{interior } \mathfrak{L}$. Then there exists $\epsilon > 0$ such that whenever $V \in B(H)$ and $\|T - V\| < \epsilon$, then $V \in \mathfrak{L}$. From the above remark $\sigma(T)$ must contain at least two points. There must be at least two extreme points of $W(T)$, since extreme points of $W(T)$ for $T \in \mathfrak{L}$ are extreme points of $\sigma(T)$. Hence, after a rotation, if necessary, we may assume there exists $z_1, z_2 \in \sigma_\pi(T) \cap \partial W(T)$ such that

1. $\operatorname{Re} z_1 = \inf \operatorname{Re} W(T)$,
2. $\operatorname{Re} z_2 = \sup \operatorname{Re} W(T)$, and
3. $\operatorname{Re} z_1 < \operatorname{Re} z_2$.

Since $z_1, z_2 \in \partial W(T) \cap \sigma_\pi(T)$, z_1 and z_2 are normal approximate eigenvalues of T [10, Theorem 2, p. 233]. By Lemma 1 there exist unit vectors $x, y \in H$ such that $(x, y) = 0$, $\|(T - z_1 I)x\| < \epsilon/8$, and $\|(T - z_2 I)y\| < \epsilon/8$. Let M be the closed subspace spanned by $\{x, y\}$. Define $C \in B(H)$ as

$$\begin{aligned} Cx &= -(\epsilon/4)x, \\ Cy &= +(\epsilon/4)y, \text{ and} \\ Cz &= 0 \text{ for all } z \in M^\perp. \end{aligned}$$

Since $\|C\| \leq \epsilon/2$, $T + C \in \mathfrak{L}$. Since

$$\begin{aligned} ((T + C)x, x) &= (Tx, x) - \epsilon/4 \quad \text{and} \\ |(Tx, x) - z_1| &\leq \|(T - z_1 I)x\| < \epsilon/8, \end{aligned}$$

we obtain $\inf \operatorname{Re} W(T + C) < \operatorname{Re} z_1$. Since $T + C \in \mathfrak{L}$, there exists $a \in \sigma_\pi(T + C) \cap \partial W(T + C)$ such that $\operatorname{Re} a = \inf \operatorname{Re} W(T + C)$. Since C is a compact operator, Weyl's spectral inclusion theorem [7, problem 143] yields $\sigma(T + C) - \sigma_p(T + C) \subseteq \sigma(T)$. Therefore, $a \in \sigma_p(T + C)$ so that $a \in \sigma_p(T + C) \cap \partial W(T + C)$. Similarly one shows there exists

$$b \in \sigma_p(T + C) \cap \partial W(T + C)$$

such that $\operatorname{Re} b = \sup W(T + C) > \operatorname{Re} z_2$, and hence $a \neq b$. By Lemma 2, there exists $S \in B(H)$ such that $\|S\| < \epsilon/2$ and $T + C + S \notin \mathfrak{L}$. However, $\|T - (T + C + S)\| \leq \|C\| + \|S\| < \epsilon$ and so by assumption $T + C + S \in \mathfrak{L}$. Contradiction.

Proof of Lemma 1. There exist sequences $\{w_n\}$ and $\{y_n\}$ of unit vectors in H such that $\|(T - z_1 I)w_n\| \rightarrow 0$ and $\|(T - z_2 I)y_n\| \rightarrow 0$ as $n \rightarrow \infty$. Then

$$\begin{aligned} |(z_1 - z_2)(w_n, y_n)| &= |(z_1 w_n, y_n) - (w_n, z_2^* y_n)| \\ &\leq |((T - z_1 I)w_n, y_n)| + |(w_n, (T - z_2 I)^* y_n)| \leq \|(T - z_1 I)w_n\| + \|(T - z_2 I)^* y_n\|. \end{aligned}$$

Therefore, $|(z_1 - z_2)(w_n, y_n)| \rightarrow 0$ as $n \rightarrow \infty$. Since $z_1 \neq z_2$, $(w_n, y_n) \rightarrow 0$ as $n \rightarrow \infty$.

There exist complex numbers a_n and b_n and unit vectors x_n in H such that $w_n = a_n y_n + b_n x_n$, $|a_n|^2 + |b_n|^2 = 1$, and $(x_n, y_n) = 0$. From the above paragraph we have that $a_n \rightarrow 0$, so $|b_n| \rightarrow 1$. Therefore, $\|(T - z_1 I)x_n\| \rightarrow 0$ as $n \rightarrow \infty$.

Proof of Lemma 2. Let $\epsilon > 0$. Since $a, b \in \sigma_p(T) \cap \partial W(T)$, a and b are normal eigenvalues of T [10, Theorem 2, p. 233]. Let $u, v \in H$ be unit vectors such that $Tu = au$ and $Tv = bv$. Then $(u, v) = 0$ and the closed subspace N spanned by $\{u, v\}$ reduces T . Define $S \in B(H)$ as

$$Su = \epsilon v, \quad Sv = 0, \quad Sz = 0 \quad \text{for all } z \in N^\perp.$$

Then we may write $T + S = A \oplus B$ corresponding to $H = N \oplus N^\perp$. Then the matrix representation of A relative to $\{u, v\}$ is $A = \begin{bmatrix} a & \epsilon \\ 0 & b \end{bmatrix}$. Hence $\sigma(A) = \{a, b\} \subseteq \sigma(T)$. Clearly $\sigma(B) \subseteq \sigma(T)$. Therefore,

$$\operatorname{co} \sigma(T + S) \subseteq \operatorname{co} \sigma(T) = W(T).$$

Since A is not a normal operator, $W(A)$ is the convex hull of a nondegenerate ellipse (i.e., not a straight line) with foci at a and b (see [1]). Since $W(A) \subseteq W(T + S)$, we must have $\operatorname{co} \sigma(T + S) \neq W(T + S)$. Therefore, $T + S \notin \mathfrak{L}$. Thus, since $\|S\| = \epsilon > 0$ is arbitrary, $T \notin \text{interior } \mathfrak{L}$.

BIBLIOGRAPHY

1. William F. Donoghue, Jr., *On the numerical range of a bounded operator*, Michigan Math. J. 4 (1957), 261–263. MR 20 #2622.
2. ———, *On a problem of Nieminen*, Inst. Hautes Études Sci. Publ. Math. No. 16 (1963), 31–33. MR 27 #2864.
3. N. Dunford and J. T. Schwartz, *Linear operators. I: General theory*, Pure and Appl. Math., vol. 7, Interscience, New York, 1958. MR 22 #8302; *II: Spectral theory. Selfadjoint operators in Hilbert space*, Interscience, New York, 1963. MR 32 #6181.

4. N. Dunford, *A survey of the theory of spectral operators*, Bull. Amer. Math. Soc. 64 (1958), 217–274. MR 21 #3616.
5. P. R. Halmos, *Introduction to Hilbert space and the theory of spectral multiplicity*, Chelsea, New York, 1957.
6. ———, *Numerical ranges and normal dilations*, Acta. Sci. Math. 25 (1964), 1–5. MR 30 #1399.
7. ———, *A Hilbert space problem book*, Van Nostrand, Princeton, N. J., 1967. MR 34 #8178.
8. S. Hildebrandt, *The closure of the numerical range of an operator as spectral set*, Comm. Pure Appl. Math. 17 (1964), 415–421. MR 29 #3882.
9. ———, *Numerischer Wertebereich und normale Dilatationen*, Acta. Sci. Math. (Szeged) 26 (1965), 187–190. MR 32 #2906.
10. ———, *Über den numerischen Wertebereich eines Operators*, Math. Ann. 163 (1966), 230–247. MR 34 #613.
11. C. Meng, *A condition that a normal operator have a closed numerical range*, Proc. Amer. Math. Soc. 8 (1957), 85–88. MR 20 #1223.
12. ———, *On the numerical range of an operator*, Proc. Amer. Math. Soc. 14 (1963), 167–171. MR 26 #601.
13. G. H. Orland, *On a class of operators*, Proc. Amer. Math. Soc. 15 (1964), 75–79. MR 28 #480.
14. C. R. Putnam, *On the spectra of semi-normal operators*, Trans. Amer. Math. Soc. 119 (1965), 509–523. MR 32 #2913.
15. ———, *Commutation properties of Hilbert space operators and related topics*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 36, Springer-Verlag, New York, 1967. MR 36 #707.
16. ———, *Eigenvalues and boundary spectra*, Illinois J. Math. 12 (1968), 278–282. MR 37 #2030.
17. ———, *The spectra of operators having resolvents of first-order growth*, Trans. Amer. Math. Soc. 133 (1968), 505–510. MR 37 #4651.
18. F. Riesz and B. Sz. Nagy, *Leçons d'analyse fonctionnelle*, Akad. Kiadó, Budapest, 1952; English transl., Ungar, New York, 1955. MR 14, 286; MR 17, 175.
19. J. G. Stampfli, *Hyponormal operators*, Pacific J. Math. 12 (1962), 1453–1458. MR 26 #6772.
20. ———, *Hyponormal operators and spectral density*, Trans. Amer. Math. Soc. 117 (1965), 469–476. MR 30 #3375.
21. ———, *Analytic extensions and spectral localization*, J. Math. Mech. 16 (1966), 287–296. MR 33 #4687.
22. ———, *Extreme points of the numerical range of a hyponormal operator*, Michigan Math. J. 13 (1966), 87–89. MR 32 #4551.
23. ———, *Minimal range theorems for operators with thin spectra*, Pacific J. Math. 23 (1967), 601–612. MR 37 #4655.
24. ———, *A local spectral theory for operators. II*, Bull. Amer. Math. Soc. 75 (1969), 803–806. MR 39 #6108.
25. ———, *A local spectral theory for operators*, J. Functional Anal. 4 (1969), 1–10. MR 39 #4698.
26. A. Wintner, *Zur Theorie der beschränkten Bilinearformen*, Math. Z. 30 (1929), 228–282.