CURVATURE TENSORS IN KAEHLER MANIFOLDS⁽¹⁾

BY

MALLADI SITARAMAYYA

ABSTRACT. Curvature tensors of Kaehler type (or type K) are defined on a hermitian vector space and it has been proved that the real vector space $\mathfrak{L}_K(V)$ of curvature tensors of type K on V is isomorphic with the vector space of symmetric endomorphisms of the symmetric product of V^+ , where $V^C = V^+ \oplus V^-$ (Theorem 3.6). Then it is shown that $\mathfrak{L}_K(V)$ admits a natural orthogonal decomposition (Theorem 5.1) and hence every $L \in \mathfrak{L}_K(V)$ is expressed as $L = L_1 + L_W + L_2$. These components are explicitly determined and then it is observed that L_W is a certain formal tensor introduced by Bochner. We call L_W the *Bochner-Weyl* part of L and the space of all these L_W is called the *Weyl subspace* of $\mathfrak{L}_K(V)$.

Introduction. Singer and Thorpe [6] established a natural decomposition of curvature tensors on an n -dimensional vector space with an inner product and then studied the curvature tensor of a 4-dimensional Einstein space and its canonical form. Nomizu [5] using this decomposition discussed generalised curvature tensor fields satisfying the second Bianchi identity (which are called *proper*) on a Riemannian manifold.

The purpose of this paper is to give a similar decomposition of curvature tensors of Kaehler type on a $2n$ -dimensional hermitian vector space and then study curvature tensors on a Kaehler manifold. In this paper the study is purely local in nature. The contents are as follows:

In §1 some basic facts about hermitian vector spaces $(V, J, \langle, \rangle)$ are given. Kaehler metric on a complex manifold is defined and the properties of its curvature tensor are stated in §2. In §3 a curvature tensor of type K on a hermitian vector space $(V, J, \langle, \rangle)$ is defined and then proved that the real vector space $\mathfrak{L}_K(V)$ of

Received by the editors August 1, 1972.

AMS (MOS) subject classifications (1970). Primary 53B35, 53C55; Secondary 32C10.

Key words and phrases. Hermitian and Kaehler metrics, curvature tensor, complex manifold, hermitian vector space, fundamental 2-form, Bochner-Weyl curvature tensor, Chern class, Einstein manifold, tangent bundle.

(1) This formed part of the author's doctoral dissertation submitted to the University of Notre Dame under the direction of Professor Yozo Matsushima. The author is grateful to his adviser for the help and encouragement in this work.

Copyright © 1973, American Mathematical Society

all curvature tensors of type K on V is isomorphic to the real vector space $s(V^+ \cdot V^+)$ of all symmetric endomorphisms of the symmetric product of V^+ (Theorem 3.6). §4 deals with Ricci curvature $K(L)$ and holomorphic sectional curvature k_L of L in $\mathcal{L}_K(V)$. In §5 we have proved that $\mathcal{L}_K(V)$ is an orthogonal direct sum of three subspaces (Theorem 5.1) and hence every $L \in \mathcal{L}_K(V)$ can be uniquely written as $L = L_1 + L_W + L_2$, these components being determined explicitly (Theorem 5.4); L_W is observed to be a certain formal tensor introduced by Bochner [7] and hence we call L_W the Bochner-Weyl part of L . The subspace of these L_W 's is denoted by $\mathcal{L}_K^W(V)$ and called the Weyl subspace. In a forthcoming paper we study curvature tensors in $\mathcal{L}_K^W(V)$.

1. **Algebraic preliminaries.** Let V be a $2n$ -dimensional real vector space with a complex structure J . Let $\langle, \rangle$ be an inner product of hermitian type on V . Denote by V^C the complexification of V and $-$ denotes the conjugation in V^C with respect to (for short, w.r.t.) V . Further, let L be a real endomorphism of V . Then we can extend L to a complex linear map of V^C to itself; in particular, J extends to V^C . Since $J^2 = -\text{identity}$, J is a nonsingular semisimple endomorphism of V^C with eigenvalues $+i$ and $-i$, where $i = \sqrt{-1}$ (i is $\sqrt{-1}$ always unless i is a subscript or superscript). Hence $V^C = V^+ \oplus V^-$ where

$$V^+ = \{u \in V^C \mid Ju = iu\} \quad \text{and} \quad V^- = \{u \in V^C \mid Ju = -iu\}.$$

Moreover, $\overline{V^+} = V^-$ and hence complex dimension of $V^+ = n = \text{complex dimension of } V^-$.

The hermitian inner product $\langle, \rangle$ on V can be extended to V^C as a symmetric complex bilinear form (also denoted by $\langle, \rangle$) and satisfies $\langle V^\pm, V^\pm \rangle = 0$. If we define $(u, v) = \langle u, \bar{v} \rangle$ for $u, v \in V^+$, then $(,)$ is a positive definite hermitian form on V^+ .

2. **Curvature tensor of a Kaehler metric.** Let M be a differentiable (i.e. c^∞ -) manifold of dimension n . $T_p(M)$ denotes the tangent space of M at p which is an n -dimensional vector space. If M is a complex manifold of complex dimension n then the tangent space $T_Z(M)$ at Z of M as a c^∞ -manifold is a real vector space of real dimension $2n$ with a complex structure J . A Riemannian metric g on M is called a *hermitian metric* if the inner product g_Z on $T_Z(M)$ is of hermitian type for each $Z \in M$.

Let g be a hermitian metric on M . Define $\Omega(u, v) = g(Ju, v)$ for $u, v \in T_Z(M)$. Then Ω is a real differential 2-form on M of type $(1, 1)$ and is of maximal rank. We call Ω the fundamental 2-form of the hermitian manifold (M, g) . A hermitian manifold (M, g) is *Kaehlerian* if the fundamental 2-form Ω is closed, i.e. $d\Omega = 0$.

We give two important examples of Kaehler manifolds.

(a) The complex number space $\mathbb{C}^n$ with the usual inner product is a Kaehler manifold and $\Omega = \sqrt{-1} \sum dZ^i \wedge \overline{dZ^i}$ where (Z^i) are the canonical coordinates of $\mathbb{C}^n$.

(b) The complex projective space $\mathbb{P}^n(\mathbb{C})$ with the Fubini-Study metric is a Kaehler manifold and so is every complex submanifold of $\mathbb{P}^n(\mathbb{C})$ (cf. [1, p. 170]).

Let ∇ denote the connection defined by the metric g . Then the curvature tensor R of ∇ is a $(1, 3)$ tensor. That is, $R(X, Y, Z)$ is a vector field. If we write $R(X, Y, *) = R(X, Y)*$ then $R(X, Y)$, for fixed X, Y , is a $(1, 1)$ tensor. Moreover for fixed X, Y , $R(X, Y) = \nabla_X \nabla_Y - \nabla_Y \nabla_X - \nabla_{[X, Y]}$ is a $(1, 1)$ tensor (cf. [2, p. 133]).

The following gives the properties of the curvature tensor R of the Kaehler metric g on a manifold M .

Proposition 2.1 ([3], [4]). *Let (M, g) be a Kaehler manifold and R be the curvature tensor of g . Then (1) $R(X, Y)J = JR(X, Y)$ and (2) $R(JX, JY) = R(X, Y)$ for any two vector fields X, Y on M .*

3. Generalized curvature tensors. In this section, we want to study curvature tensors having properties given in Proposition 2.1 in a more general algebraic setup.

Let V be a $2n$ -dimensional real vector space with an inner product $\langle, \rangle$. Denote by $\mathfrak{o}(V)$ the vector space of skew symmetric endomorphisms of V . We define a curvature tensor L on V as an $\mathfrak{o}(V)$ -valued 2-form on V . Then for each $x, y \in V$, $L(x, y) \in \text{End}(V)$ such that

- (1) $L(x, y)$ is bilinear in x and y ;
- (2) $L(x, y) = -L(y, x)$;
- (3) $\langle L(x, y)z, w \rangle + \langle z, L(x, y)w \rangle = 0$.

Now assume V admits a complex structure J and a hermitian inner product $\langle, \rangle$.

A curvature tensor L on V is said to be of type K on V if L satisfies the following:

- (1) $L(Jx, Jy) = L(x, y)$ for all $x, y \in V$;
- (2) $L(x, y)J = JL(x, y)$;
- (3) $L(x, y)z + L(y, z)x + L(z, x)y = 0$ for all $x, y, z \in V$ (Bianchi identity).

Denote by $\mathfrak{L}_K(V)$ the real vector space of all curvature tensors of type K on V . Our goal is to understand $\mathfrak{L}_K(V)$.

Let $L \in \mathfrak{L}_K(V)$. Extend L to $V^{\mathbb{C}}$. Then L is a skew symmetric bilinear map on $V^{\mathbb{C}}$ with values in $\text{End}(V^{\mathbb{C}})$. Extend J and $\langle, \rangle$ to $V^{\mathbb{C}}$ as in §1. Then

- (1) $L(Jx, Jy) = L(x, y)$ for all $x, y \in V$ is equivalent to $L(V^{\perp}, V^{\perp}) = 0$.
- (2) $L(x, y)J = JL(x, y)$ implies $L(x, y)$ leaves $V^{\perp}$ invariant.

(3) Bianchi identity holds in the complex case.

Since $L(V^\pm, V^\pm) = 0$, the Bianchi identity gives

(a) $L(x, \bar{y})\bar{z} = L(x, \bar{z})\bar{y}$ and hence $L(x, \bar{y})\bar{z}$ is symmetric in y and z , and

(a') $L(x, \bar{w})y = L(y, \bar{w})x$ and hence $L(x, \bar{w})y$ is symmetric in x and y .

Note that since (a') is obtained from (a) by conjugation a curvature tensor L satisfies Bianchi identity if and only if L satisfies (a). Now define the map $F_L: V^+ \times V^+ \times V^+ \times V^+ \rightarrow \mathbb{C}$ by

$$(3.1) \quad F_L(y, z, x, w) = L(y, \bar{z}, x, \bar{w}) = \langle L(x, \bar{w})\bar{z}, y \rangle \quad \text{for all } x, y, z, w \in V^+.$$

Then clearly this map has the properties

(a) F_L is bilinear in x and y and anti-bilinear in w and z ;

(b) F_L is symmetric in x and y and also in z and w .

Since given any $F: V^+ \times V^+ \times V^+ \times V^+ \rightarrow \mathbb{C}$ satisfying (a) and (b) there exists a unique $L \in \mathcal{L}_K(V)$ such that $F = F_L$ (as given in (3.1)); we call such an F also a curvature tensor of type K on V . Note that $F_L(y, z, x, w) = F_L(x, w, y, z)$ (from (b)).

Define hermitian inner product in V^+ by $(x, y) = \langle x, \bar{y} \rangle$. Next we extend $(,)$ on V^+ to $V^+ \otimes V^+$ by defining $(x \otimes y, w \otimes z) = (x, w)(y, z)$.

Now we have

Proposition 3.2. *If $f(w, z)$ is an anti-bilinear function on V^+ then there exists $u \in V^+ \otimes V^+$ such that $f(w, z) = (u, w \otimes z)$.*

Proof. Obvious.

Since, for every fixed $x, y \in V^+$, $F_L(y, *, x, *)$ is anti-bilinear on V^+ , by Proposition 3.2, there exists $u(x, y) \in V^+ \otimes V^+$ such that $F_L(y, z, x, w) = (u(x, y), w \otimes z)$. Since F_L is bilinear and symmetric in x and y the map $u: V^+ \times V^+ \rightarrow V^+ \otimes V^+$ sending (x, y) to $u(x, y)$ is bilinear and symmetric. Hence there exists a linear endomorphism $U': V^+ \otimes V^+ \rightarrow V^+ \otimes V^+$ such that $U'(x \otimes y) = u(x, y)$, and hence we have

$$(3.3) \quad (U'(x \otimes y), w \otimes z) = F_L(y, z, x, w).$$

Moreover, using (3.3) and (3.1) we see easily that U' is a symmetric endomorphism of $V^+ \otimes V^+$ w.r.t. $(,)$. Now we carry this U' one more step forward.

Consider the symmetric product $V^+ \cdot V^+$ of V^+ . $V^+ \cdot V^+$ is, by definition, the quotient of $V^+ \otimes V^+$ by the subspace A consisting of all skew symmetric tensors. Denote by π the canonical projection of $V^+ \otimes V^+$ onto $V^+ \cdot V^+$. Then $t \in V^+ \otimes V^+$ implies $t = s + a$ where s is a symmetric tensor and $a \in A$. Define an inner product in $V^+ \cdot V^+$ by $(\pi(t), \pi(t')) = (s, s')$ where s and s' are respectively the symmetric parts of t and t' and (s, s') is the inner product of s

and s' in $V^+ \otimes V^+$. Since symmetric and skew symmetric tensors are perpendicular, the inner product in $V^+ \cdot V^+$ satisfies the condition $(t, t') = (\pi(t), \pi(t')) + (a, a')$. Moreover, from our definition it follows that $(x \cdot y, w \cdot z) = \frac{1}{2}\{(x, w)(y, z) + (x, z)(y, w)\}$ where we have put $x \cdot y = \pi(x \otimes y)$ for all $x, y \in V^+$.

Now we claim that there exists a symmetric linear map $U: V^+ \cdot V^+ \rightarrow V^+ \cdot V^+$ such that $F_L(y, z, x, w) = (U(x \cdot y), w \cdot z)$. In fact, since $U': V^+ \otimes V^+ \rightarrow V^+ \otimes V^+$ is defined by $U'(x \otimes y) = u(x, y)$ and, since $u(x, y)$ is symmetric in x and y , U' maps kernel π into zero and hence U' induces a unique linear map $U: V^+ \cdot V^+ \rightarrow V^+ \cdot V^+$ such that $U(x, y) = \pi(U'(x \otimes y))$. It remains to show

$$(3.4) \quad (U(x \cdot y), w \cdot z) = (U'(x \otimes y), w \otimes z).$$

But we have $(U'(x \otimes y), w \otimes z) = (U(x \cdot y), w \cdot z) + (a, a')$ where a and a' denote the skew symmetric parts of $U'(x \otimes y)$ and $w \otimes z$ respectively. Since $F_L(y, v, x, u) = (U'(x \otimes y), u \otimes v)$ and since F_L is symmetric in u and v , $U'(x \otimes y)$ is orthogonal to the subspace A of skew symmetric tensors and hence $U'(x \otimes y)$ is a symmetric tensor and hence $a = 0$. This completes the proof of our claim. Moreover, we can prove that this U is a symmetric endomorphism of $V^+ \cdot V^+$.

Denote by $s(V^+ \cdot V^+)$ the vector space of all symmetric endomorphisms of $V^+ \cdot V^+$. In summary, starting with a curvature tensor $L \in \mathcal{L}_K(V)$ we obtained a $U_L \in s(V^+ \cdot V^+)$ such that

$$(3.5) \quad (U_L(x \cdot y), w \cdot z) = F_L(y, z, x, w) = \langle L(y, \bar{z})\bar{w}, x \rangle.$$

Denote this map from $\mathcal{L}_K(V)$ into $s(V^+ \cdot V^+)$ by ϕ . Now we have the following:

Theorem 3.6. *The map ϕ defined above (by (3.5)) is an isomorphism of $\mathcal{L}_K(V)$ onto $s(V^+ \cdot V^+)$.*

Before proceeding with the proof of this theorem we give some facts about bases of V^+ and $V^+ \cdot V^+$.

Let $e_1, e_2, \dots, e_n$ be an orthonormal basis of V^+ , an n -dimensional hermitian vector space, w.r.t. $(\cdot, \cdot)$. Then $\{e_1, e_2, \dots, e_n, \bar{e}_1, \bar{e}_2, \dots, \bar{e}_n\}$ forms a basis of V^C and we have

- (1) $\{e_i \cdot e_j, i \leq j, i, j = 1, 2, \dots, n\}$ is a basis of $V^+ \cdot V^+$.
- (2) $\{e_i \cdot e_j, \sqrt{2} e_i \cdot e_j, i < j, j = 1, 2, \dots, n\}$ is an orthonormal basis of $V^+ \cdot V^+$ w.r.t. $(\cdot, \cdot)$.
- (3) Putting $f_i = (e_i + \bar{e}_i)/\sqrt{2}$, $Jf_i = \sqrt{-1} (e_i - \bar{e}_i)/\sqrt{2}$, $i = 1, 2, \dots, n$, $\{f_i, Jf_i\}$ is a basis for V as a $2n$ -dimensional real vector space.
- (4) $L(\bar{e}_i, e_j) = L(e_i, \bar{e}_j)$.

Proof of Theorem. That ϕ is an injective real linear map is clear from (3.5). We have to show ϕ is onto. That is, given $U \in \mathcal{S}(V^+ \cdot V^+)$, we have to define for each $x, y \in V$ a real endomorphism $L(x, y)$ of V such that $L \in \mathcal{L}_K(V)$ and L when extended to a complex linear map of V^+ should satisfy (3.5).

Taking $x = e_k, y = e_l$ ($k \leq l$), and $w = e_m, z = e_p$ in (3.5) we get

$$(3.7) \quad L_{l\bar{p}k\bar{m}} = \begin{cases} U_{mm,kl} & \text{if } p = m, \\ U_{mp,kl} & \text{if } p \neq m \ (k \leq l, m \leq p). \end{cases}$$

Because of symmetry in k and l and in p and m (3.7) defines L for all suffixes.

Now we want to define the corresponding real endomorphism L of V so that $L \in \mathcal{L}_K(V)$. Let $\{f_i, Jf_i, i = 1, 2, \dots, n\}$ be real basis of V such that $f_k = e_k + \bar{e}_k$ and $Jf_k = i(e_k - \bar{e}_k)$. Then we compute L in terms of components of U (by (3.7)) as in

$$(1) \quad \langle L(f_k, f_m)f_p, f_l \rangle = L_{l\bar{p}k\bar{m}} - L_{p\bar{l}k\bar{m}} + L_{p\bar{l}m\bar{k}} - L_{l\bar{p}m\bar{k}}.$$

Using (3.7) and the fact that U is hermitian we can show the right-hand side of (1) is real.

$$(2) \quad \langle L(f_k, Jf_m)f_p, f_l \rangle = i\{L_{p\bar{l}m\bar{k}} - L_{l\bar{p}k\bar{m}} + L_{p\bar{l}k\bar{m}} - L_{l\bar{p}m\bar{k}}\} \text{ which is also real.}$$

$$(3) \quad \langle L(f_k, f_m)Jf_p, f_l \rangle = i\{L_{p\bar{l}k\bar{m}} - L_{l\bar{p}k\bar{m}} + L_{p\bar{l}m\bar{k}} - L_{l\bar{p}m\bar{k}}\}. \text{ This is also real from (2).}$$

$$(4) \quad \langle L(f_k, Jf_m)Jf_p, f_l \rangle = -\{L_{p\bar{l}k\bar{m}} + L_{l\bar{p}m\bar{k}} - L_{l\bar{p}k\bar{m}} - L_{p\bar{l}m\bar{k}}\}. \text{ This is also real.}$$

From (2) and (3) we get

$$(5) \quad \langle L(f_k, Jf_p)f_m, f_l \rangle = \langle L(f_k, f_m)Jf_p, f_l \rangle.$$

Now using these relations (1)–(5) we define L on the basis elements $\{f_k, Jf_k\}$ of V as follows: Define $L(f_k, f_m)$ by (1) and (3); $L(f_k, Jf_m)$ by (2) and (4); $L(Jf_k, Jf_m) = L(f_k, f_m)$ and $L(f_k, f_m)Jf_p = JL(f_k, f_m)f_p$. Then this L is defined as a real endomorphism of V in terms of the components of given U and L has the properties

- (a) $L(f'_i, f'_j) = -L(f'_j, f'_i)$,
- (b) $L(Jf'_i, Jf'_j) = L(f'_i, f'_j)$, and
- (c) $L(f'_i, f'_j)J = JL(f'_i, f'_j)$, where $\{f'_1, f'_2, \dots, f'_n, f'_{n+1}, \dots, f'_{2n}\}$ denotes $\{f_p, Jf_i\}, i = 1, 2, \dots, n$.

Hence L belongs to $\mathcal{L}_K(V)$ provided L satisfies the Bianchi identity, i.e.

$$(3.8) \quad L(x, y)z + L(y, z)x + L(z, x)y = 0.$$

We verify this on basis elements of V . Let Σ_L denote the left-hand side of (3.8).

Case (a). Let $x = f_k$, $y = f_m$, $z = f_p$ and $w = f_l$. Then from relations (1)–(5) we get

$$\begin{aligned} \langle \Sigma_L, f_l \rangle &= L_{l\bar{p}k\bar{m}} + L_{p\bar{l}m\bar{k}} - L_{p\bar{l}k\bar{m}} - L_{l\bar{p}m\bar{k}} \\ &\quad + L_{l\bar{k}m\bar{p}} + L_{k\bar{l}p\bar{m}} - L_{k\bar{l}m\bar{p}} - L_{l\bar{k}p\bar{m}} \\ &\quad + L_{l\bar{m}p\bar{k}} + L_{m\bar{l}k\bar{p}} - L_{m\bar{l}p\bar{k}} - L_{l\bar{m}k\bar{p}} \end{aligned}$$

which can be seen to be zero using (3.7) and U is hermitian.

Similarly in the other cases

(b) $x = f_k$, $y = f_m$, $z = f_p$ and $w = Jf_l$,

(c) $x = f_k$, $y = Jf_m$, $z = f_p$ and $w = f_l$ or Jf_p ,

(d) $x = f_k$, $y = Jf_m$, $z = Jf_k$;

(e) $x = Jf_k$, $y = Jf_m$, $z = Jf_p$,

etc., we can show that $\langle \Sigma_L, w \rangle = 0$ using $L(Jx, Jy) = L(x, y)$ and $L(x, y)J = JL(x, y)$.

Thus L as defined above belongs to $\mathcal{Q}_K(V)$ and in view of (3.5) we have $\phi(L) = U$, hence ϕ is surjective.

This completes the proof of the theorem.

Remarks. (1) Because of this isomorphism ϕ between $\mathcal{Q}_K(V)$ and $s(V^+ \cdot V^+)$ we call elements of $s(V^+ \cdot V^+)$ also curvature tensors of type K and we identify $\mathcal{Q}_K(V)$ and $s(V^+ \cdot V^+)$. Sometimes for $L \in \mathcal{Q}_K(V)$ we denote its image under ϕ simply by L' instead of U_L or $\phi(L)$.

(2) If $\mathcal{Q}_R(V)$ denotes the vector space of curvature tensors on V satisfying the Bianchi identity (here V is a real inner product space) then $\mathcal{Q}_R(V)$ is isomorphic with a proper subspace of $s(V \wedge V)$ (cf. [6]). But in the Kaehler case we have $\mathcal{Q}_K(V)$ is isomorphic with $s(V^+ \cdot V^+)$.

4. Ricci and holomorphic sectional curvatures of L . We define a linear map $K: \mathcal{Q}_K(V) \rightarrow \text{End}(V^+)$, called the Ricci map, as follows.

For every $L \in \mathcal{Q}_K(V)$, $K(L)$ is an endomorphism of V^+ such that $K(L)x = \sum_{s=1}^n L(x, \bar{e}_s)e_s$ where $x \in V^+$ and $\{e_i\}$ ($i = 1, 2, \dots, n$) is an orthonormal basis of V^+ w.r.t. $(\cdot, \cdot)$. We call $K(L)$ the Ricci tensor of L . It is easily seen that $K(L)$ is a symmetric endomorphism of V^+ w.r.t. $(\cdot, \cdot)$. If we define $K(x, y) = (K(L)x, y)$ then $K(x, y)$ is a symmetric bilinear form on V^+ , called the Ricci form of L in $\mathcal{Q}_K(V)$.

Let $\mathbf{P}(V^+)$ denote the complex projective space of complex lines in V^+ . For any $L \in \mathcal{Q}_K(V)$ we define a function $k_L: \mathbf{P}(V^+) \rightarrow \mathbb{C}$ as follows.

Let v be any vector in a complex line P of V^+ such that $\|v\| = 1$ w.r.t. $(\cdot, \cdot)$. Then

$$k_L(P) = \langle L(\bar{v}, v)\bar{v}, v \rangle = -(L'(v \cdot v), v \cdot v).$$

Then $k_L(P)$ is well defined for any P in $\mathbf{P}(V^+)$.

Remark. $k_L(P)$ is, in fact, a real number because L' is hermitian. Moreover, if we write $v \in V^+$ as $v = (X - iJX)/\sqrt{2}$, $X, JX \in V$ (the corresponding real 2-plane in V is spanned by $X/\sqrt{2}, JX/\sqrt{2}$) then $\langle L(\bar{v}, v)\bar{v}, v \rangle = \langle L(X, JX)JX, X \rangle \in \mathbf{R}$. So our definition of k_L is the usual definition of holomorphic sectional curvature function.

Definition of symmetric (wedge) product of two endomorphisms. Let V be a vector space and $A, B \in \text{End}(V)$. Let $V \cdot V = V \otimes V/A'$ (resp. $V \wedge V = V \otimes V/N$) where A' is the subspace of $V \otimes V$ generated by $x \otimes y - y \otimes x$ for all $x, y \in V$ (resp. N is the subspace of $V \otimes V$ generated by $x \otimes x$ for all $x \in V$). Then $A \otimes B \in \text{End}(V \otimes V)$ is defined by $(A \otimes B)(x \otimes y) = Ax \otimes By$ and $A \otimes B + B \otimes A$ leaves A' (resp. N) invariant and hence induces an endomorphism of $V \cdot V$ (resp. $V \wedge V$). We denote this endomorphism by $A \cdot B$ (resp. $A \wedge B$). Then we have $(A \cdot B)(x \cdot y) = (Ax \cdot By) + (Ay \cdot Bx)$.

Now let $A, B \in s(V^+)$. Then we can show that $A \cdot B \in s(V^+ \cdot V^+)$. In fact

$$\begin{aligned} ((A \cdot B)(x \cdot y), w \cdot z) &= (Ax \cdot By, w \cdot z) + (Ay \cdot Bx, w \cdot z) \\ &= (x \cdot y, Aw \cdot Bz) + (y \cdot x, Aw \cdot Bz) = (x \cdot y, (A \cdot B)(w \cdot z)). \end{aligned}$$

Therefore, by Theorem 3.6, $A \cdot B$ corresponds to a curvature tensor of type K on V and we denote this by $L_{A \cdot B} \in \mathfrak{L}_K(V)$.

Proposition 4.1. Let $L_1, L_2 \in \mathfrak{L}_K(V)$ such that

$$\langle L_1(\bar{v}, v)\bar{v}, v \rangle = \langle L_2(\bar{v}, v)\bar{v}, v \rangle \quad \text{for all } v \in V^+.$$

Then $L_1 = L_2$.

This is the same as Proposition 7.1 in [3, p. 166].

Proposition 4.2. Let $L \in \mathfrak{L}_K(V)$. Then k_L is constant (say c) if and only if $L = -cL_{A \cdot B}$ with $A = B = I/\sqrt{2} \in s(V^+)$, where I denotes the identity endomorphism of V^+ .

Proof. ($\Rightarrow$) Let $P \in \mathbf{P}(V^+)$. Then $k_L(P) = -(L'(v \cdot v), v \cdot v) = c$.

On the other hand, $I \cdot I/2 \in s(V^+ \cdot V^+)$ and

$$(4.3) \quad \langle L_{I \cdot I/2}(\bar{v}, v)\bar{v}, v \rangle = -\langle (I \cdot I/2)(v \cdot v), v \cdot v \rangle = -1 \quad \text{for all } v.$$

Hence $\langle L(\bar{v}, v)\bar{v}, v \rangle = c = -c \langle L_{I \cdot I/2}(\bar{v}, v)\bar{v}, v \rangle$ for all $v \in V^+$ and hence $L = -cL_{I \cdot I/2}$.

Conversely, given $L = -c L_{I, I/2}$ we can see easily from (4.3) that $k_L(P) = c$ for all $P \in \mathbf{P}(V^+)$.

Let $\{e_1, e_2, \dots, e_n\}$ be an orthonormal basis of V^+ w.r.t. $(,)$. $K_{A \cdot B}$ denotes the Ricci tensor of $L_{A \cdot B}$. Then

$$K_{A \cdot B} e_j = \sum_{s=1}^n L_{A \cdot B}(e_j, \bar{e}_s) e_s, \quad K_{A \cdot B} \in s(V^+).$$

Then

$$\begin{aligned} (K_{A \cdot B})_{ij} &= (K_{A \cdot B} e_j, e_i) = - \sum_{s=1}^n ((A \cdot B)(e_j \cdot e_s), e_s \cdot e_i) \\ &= -\frac{1}{2} \{A_{ij} \operatorname{Tr} B + B_{ij} \operatorname{Tr} A + (AB)_{ij} + (BA)_{ij}\}. \end{aligned}$$

So we get

$$(4.4) \quad -2 K_{A \cdot B} = (\operatorname{Tr} A)B + (\operatorname{Tr} B)A + AB + BA$$

and hence

$$(4.5) \quad -\operatorname{Tr} K_{A \cdot B} = \operatorname{Tr} A \operatorname{Tr} B + \operatorname{Tr} AB.$$

Recalling L' is the image of L under ϕ in Theorem 3.6, we can prove that

$$(4.6) \quad \operatorname{Tr} K(L) = 2 \operatorname{Tr} L' = 2 \operatorname{Tr} U_L.$$

We have inner product in $\mathfrak{L}_K(V)$ defined from that of V^+ as

$$(L_1, L_2) = \sum_{i,j,k,l} (L_1)_{ijkl} (L_2)_{ijkl}.$$

The inner product $(,)$ of V^+ also induces an inner product $(,)$ on $s(V^+ \cdot V^+)$ (observe that, in general, if W is a complex vector space with an inner product and $s(W)$ is the real vector space of all symmetric endomorphisms of W then we can define an inner product $(,)$ in $s(W)$ by $(S, T) = \operatorname{Tr}(ST)$). Then we have

$$(4.7) \quad (L_1, L_2) = \sum_{i,j,k,l} (L_1)_{ijkl} (L_2)_{ijkl} = \operatorname{Tr}(L'_1, L'_2).$$

Let $\mathfrak{L}_K^1(V) = \{L \in \mathfrak{L}_K(V) \mid L' = \lambda I \cdot I, \lambda \in \mathbf{R}\}$. Then $\mathfrak{L}_K^1(V)$ is a real 1-dimensional subspace of $\mathfrak{L}_K(V)$. Let $\operatorname{Orth}(\mathfrak{L}_K^1(V))$ denote the orthogonal complement of $\mathfrak{L}_K^1(V)$ in $\mathfrak{L}_K(V)$ w.r.t. inner product defined above. Then in view of (4.6) and (4.7) we have

$$\operatorname{Orth}(\mathfrak{L}_K^1(V)) = \{L \in \mathfrak{L}_K(V) \mid \operatorname{Tr} K(L) = 0\}.$$

Hence we have the decomposition

$$\mathfrak{L}_K(V) = \mathfrak{L}_K^1(V) \oplus \text{Orth}(\mathfrak{L}_K^1(V)).$$

5. Decomposition of the space $\mathfrak{L}_K(V)$. Recall $\mathfrak{L}_K^1(V) = \{L \in \mathfrak{L}_K(V) \mid L' = \lambda I, \lambda \in \mathbb{R}\}$. Let $\mathfrak{L}_K^W(V) = \{L \in \mathfrak{L}_K(V) \mid K(L) = 0\}$ where K is the Ricci map. Then clearly $\mathfrak{L}_K^W(V) \subset \text{Orth}(\mathfrak{L}_K^1(V))$. Now let $\mathfrak{L}_K^2(V)$ be the orthogonal complement of $\mathfrak{L}_K^W(V)$ in $\text{Orth}(\mathfrak{L}_K^1(V))$. Therefore we have the natural decomposition of $\mathfrak{L}_K(V)$ as $\mathfrak{L}_K(V) = \mathfrak{L}_K^1(V) \oplus \mathfrak{L}_K^W(V) \oplus \mathfrak{L}_K^2(V)$. Now we have the following:

Theorem 5.1. *Let V be a real $2n$ -dimensional hermitian vector space. Let $\mathfrak{L}_K(V)$ be the real vector space of all curvature tensors of type K on V . Then $\mathfrak{L}_K(V)$ has the decomposition*

$$\mathfrak{L}_K(V) = \mathfrak{L}_K^1(V) \oplus \mathfrak{L}_K^W(V) \oplus \mathfrak{L}_K^2(V) \quad (\text{orthogonal})$$

where

- (1) $\mathfrak{L}_K^1(V) = \{L \in \mathfrak{L}_K(V) \mid L \text{ has constant holomorphic sectional curvature}\};$
- (2) $\mathfrak{L}_K^W(V) = \{L \in \mathfrak{L}_K(V) \mid L \text{ has Ricci curvature zero}\};$
- (3) $\mathfrak{L}_K^W(V) \oplus \mathfrak{L}_K^2(V) = \{L \in \mathfrak{L}_K(V) \mid \text{Tr } K(L) = 0\};$
- (4) $\mathfrak{L}_K^1(V) \oplus \mathfrak{L}_K^W(V) = \{L \in \mathfrak{L}_K(V) \mid K(L) = \lambda I, \lambda \in \mathbb{R}\}.$

Proof. All except (4) are proved above.

Suppose $L \in \mathfrak{L}_K^1(V) \oplus \mathfrak{L}_K^W(V)$. Then $L = L_1 + L_W$ and hence $K(L) = K(L_1)$ because K is linear. But since $L_1' = I \cdot I$ and using (4.4) we get $K(L_1) = -(n+1)I = K(L)$. Conversely, let $K(L) = \lambda I, \lambda \in \mathbb{R}$. Let L_1 be the curvature tensor corresponding to $L_1' = \mu I \in s(V^+ \cdot V^+)$ with $\mu = -\lambda/(n+1)$. Then $L_1 \in \mathfrak{L}_K^1(V)$ and $K(L - L_1) = 0$. Hence $L - L_1 \in \mathfrak{L}_K^W(V)$. So $L = L_1 + L_W$.

Recall that a Kaehler metric g on a manifold M is called Einstein if R , the Ricci tensor of g , is a scalar multiple of g at each point p of M .

Corollary 5.2. *Suppose (M, g) is a Kaehler-Einstein manifold. Then the curvature tensor L (of g) belongs to*

$$\mathfrak{L}_K^1(T_p(M)) \oplus \mathfrak{L}_K^W(T_p(M)), \quad \forall p \in M \quad (\forall \text{ means "for every"})$$

This is immediate from (4) of Theorem 5.1.

Corollary 5.3. *If M is a connected Kaehler-Einstein manifold of complex dimension ≥ 2 (i.e. $K(L)_p = \lambda(p)I$ for every $p \in M$) then λ is a constant function.*

Proof. By (5.2), $L_p \in \mathfrak{L}_K^1(T_p(M)) \oplus \mathfrak{L}_K^W(T_p(M))$ and hence $R_p = K(L)_p = \lambda(p)I$, i.e. $R_{i\bar{j}} = \lambda(p)g_{i\bar{j}}$ (in local coordinates). Then the first Chern class of M is represented by $\psi = -(1/2\pi)\lambda(p)\Omega$ where $\Omega = ig_{i\bar{j}}d\bar{z}^i \wedge dz^j$ is the fundamental 2-form of g and $\psi = (1/2\pi i)R_{i\bar{j}}dz^i \wedge d\bar{z}^j$ is the Ricci form of g . Since ψ is a

closed 2-form we have $0 = d\psi = -(1/2\pi)d\lambda \wedge \Omega$. Then $d\lambda = 0$ because wedge product with Ω is an isomorphism of complex vector spaces for complex dimension of $M \geq 2$. Hence λ is a constant function on M .

Now we give the components of $L \in \mathcal{L}_K(V)$ in the decomposition.

Theorem 5.4. *Let $L \in \mathcal{L}_K(V)$ and $K(L)$ be the Ricci tensor of L . Further let $\lambda = \text{Tr } K(L)$ and L_1, L_W, L_2 be the components of L in the decomposition of Theorem 5.1. Then*

$$(5.5) \quad L_1' = -\lambda(I \cdot I)/n(n+1),$$

$$(5.6) \quad L_2' = -2(K \cdot I)/(n+2) + 2\lambda(I \cdot I)/n(n+2),$$

$$(5.7) \quad L_W' = L' + 2(K \cdot I)/(n+2) - \lambda(I \cdot I)/(n+1)(n+2).$$

(Here $K = K(L)$ and $I = \text{identity endomorphism of } V^+.$)

Proof. Let R_1, R_2, R_W denote respectively the right-hand sides of (5.5), (5.6) and (5.7). We show that $R_1 \in \mathcal{L}_K^1(W)$, $R_W \in \mathcal{L}_K^W(V)$ and $R_2 \in \mathcal{L}_K^2(V)$. Clearly $R_1 \in \mathcal{L}_K^1(V)$.

$$K(R_W) = K(L) + 2K(L_{K \cdot I})/(n+2) - \lambda K(L_{I \cdot I})/(n+1)(n+2)$$

which is seen to be zero using (4.4). Hence $R_W \in \mathcal{L}_K^W(V)$.

It remains to show $R_2 \in \mathcal{L}_K^2(V)$.

First we observe that $\text{Tr } K(R_2) = 0$ (use (4.5)) hence $R_2 \in \mathcal{L}_K^W(V) \oplus \mathcal{L}_K^2(V)$. It suffices to show that R_2 is perpendicular to $\mathcal{L}_K^W(V)$. That is, $\text{Tr}(R_2' R') = 0$ for all $R \in \mathcal{L}_K^W(V)$.

We can choose an orthonormal basis $(e_1, \dots, e_n)$ of V^+ such that $K e_i = \lambda_i e_i$, $i = 1, 2, \dots, n$ (since $K \in s(V^+)$). Then $(K \cdot I)(e_i \cdot e_j) = (\lambda_i + \lambda_j)e_i \cdot e_j$, $i, j = 1, 2, \dots, n$, and $(I \cdot I)(e_i \cdot e_j) = 2e_i \cdot e_j$ and hence

$$(5.8) \quad R_2(e_i \cdot e_j) = (-2/(n+2))\{(\lambda_i + \lambda_j) - 2\lambda/n\}e_i \cdot e_j \quad \forall i, j.$$

For $R \in \mathcal{L}_K^W(V)$,

$$(5.9) \quad \sum_s R_{sjsl} = 0 \quad \text{for all } l, j$$

(since $K(R) = 0$). In particular

$$(5.10) \quad \sum_{s,j} R_{sjsj} = 0.$$

Then $(R_2, R) = \text{Tr}(R_2' R') = \sum_{i,j,s,t} (R_2')_{ijst} R'_{stij}$ ($1 < j, s < t$). Since $R'_{ijst} = 0$ unless $(i, j) = (s, t)$ we have

$$(5.11) \quad \text{Tr}(R_2' R') = \sum_{i,j} \{(\lambda_i + \lambda_j) - 2\lambda/n\} R'_{ijij} \quad (i < j).$$

Now using (5.9), (5.10) and skew-symmetry in i and j , we can show that each

term in (5.11) is separately zero, and hence $R_2 \in \mathfrak{L}_K^2(V)$. Since the decomposition is unique we have $R_1 = L'_1$, $R_2 = L'_2$, $R_W = L'_W$.

Remark. In the Kaehler case L'_W is given by

$$L'_W = L' + 2(K \cdot I)/(n+2) - (\text{Tr } K)(I \cdot I)/(n+1)(n+2).$$

Then if we compute the components $(L'_W)_{\alpha\beta\gamma\delta}$ w.r.t. an orthogonal coordinate (e_α) of V^+ we see that this is the formal tensor introduced by Bochner (which he denotes by K) up to sign (cf. [7, p. 161]). We call any $L_W \in \mathfrak{L}_K^W(V)$ a *generalized Bochner-Weyl tensor* and $\mathfrak{L}_K^W(V)$ is called the *Weyl subspace* of $\mathfrak{L}_K(V)$.

Now we can restate our Theorems 5.1 and 5.4 in terms of $s(V^+)$.

Theorem 5.12. Let $(V, J, \langle \cdot, \cdot \rangle)$ be a real $2n$ -dimensional hermitian vector space. Let $\mathfrak{L}_K(V) = \mathfrak{L}_K^1(V) \oplus \mathfrak{L}_K^W(V) \oplus \mathfrak{L}_K^2(V)$ be the decomposition of Theorem 5.1. Then

- (a) $\mathfrak{L}_K^1(V) \oplus \mathfrak{L}_K^2(V) = \{L \in \mathfrak{L}_K(V) \mid L' = A \cdot I, A \in s(V^+)\}$.
- (b) The Ricci map $K: \mathfrak{L}_K(V) \rightarrow s(V^+)$ is onto, $\ker K = \mathfrak{L}_K^W(V)$ and K is bijective from $\mathfrak{L}_K^1(V) \oplus \mathfrak{L}_K^2(V)$ onto $s(V^+)$. Furthermore, $K(\mathfrak{L}_K^1(V)) = \{A \in s(V^+) \mid A = \lambda I, \lambda \in \mathbb{R}\}$ and $K(\mathfrak{L}_K^2(V)) = \{A \in s(V^+) \mid \text{Tr } A = 0\}$.

Proof of (a). Let $L = L_{A \cdot I} \in \mathfrak{L}_K(V)$ with $A \in s(V^+)$. Then $L' = A \cdot I$. Let $L = L_1 + L_2 + L_W$. To show $L_W = 0$ but finding $K = K(L')$ and $\lambda = \text{Tr } K$ from $L' = A \cdot I$, and substituting for K and λ in L'_W we see easily L'_W is zero. Conversely, if $L = L_1 + L_2$ then $L'_W = 0$ and hence $L' = A \cdot I$ with $A = \lambda I/(n+1)(n+2) - 2K/(n+2) \in s(V^+)$.

Proof of (b). Let $B \in s(V^+)$. Write B uniquely as $B = B_0 + \mu I$ with $B_0 \in s(V^+)$ and $\text{Tr } B_0 = 0$. Then, from (4.4), $K(L_{(-2B_0 \cdot I)/(n+2)}) = B_0$. Also we have $K(L_{-\zeta I \cdot I}) = \zeta(n+1)I$ for any real number ζ ; in particular for $\zeta = -\mu/(n+1)$ we have $K(L_{-\mu/(n+1)I \cdot I}) = \mu I$.

Set $A = -((2B_0 \cdot I)/(n+2) + (\mu I \cdot I)/(n+1))$. Then $K(L_A) = B_0 + \mu I = B$. Hence K is onto and the other statements are all obvious.

Corollary 5.13. The real dimension of $\mathfrak{L}_K^2(V) = n^2 - 1$ where $n = \text{complex dimension of } V$.

This is immediate from the bijectivity of $K: \mathfrak{L}_K^1(V) \oplus \mathfrak{L}_K^2(V) \rightarrow s(V^+)$.

Corollary 5.14. Let V be a $2n$ -dimensional real vector space with a complex structure J and an hermitian inner product $\langle \cdot, \cdot \rangle$. Then the real dimension of $\mathfrak{L}_K^W(V) = n^2(n-1)(n+3)/4$ where $n = \dim \mathbb{C}V$.

Proof. Since $\dim_{\mathbb{R}} s(V^+ \cdot V^+) = (n(n+1)/2)^2$, $\dim_{\mathbb{R}} \mathcal{Q}_K(V) = n^2(n+1)^2/4$. Hence by Corollary 5.13, $\dim_{\mathbb{R}} \mathcal{Q}_K^W(V) = n^2(n+1)^2/4 - n^2 = n^2(n-1)(n+3)/4$. Note that the Weyl subspace $\mathcal{Q}_K^W(V)$ is of real dimension 5 when V is of complex dimension 2. In general to give some characterisation of the Weyl subspace $\mathcal{Q}_K^W(V)$ seems quite difficult.

REFERENCES

1. R. C. Gunning and H. Rossi, *Analytic functions of several complex variables*, Prentice-Hall, Englewood Cliffs, N. J., 1965. MR 31 #4927.
2. S. Kobayashi and K. Nomizu, *Foundations of differential geometry*. Vol. I, Interscience, New York, 1963. MR 27 #2945.
3. ———, *Foundations of differential geometry*. Vol. II, Interscience Tracts in Pure and Appl. Math., no. 15, Interscience, New York, 1969. MR 38 #6501.
4. A. Lascoux and M. Berger, *Variétés Kählériennes compactes*, Lecture Notes in Math., vol. 154, Springer-Verlag, Berlin and New York, 1970. MR 43 #3979.
5. K. Nomizu, *On the decomposition of generalized curvature tensor fields* (to appear).
6. I. M. Singer and J. A. Thorpe, *The curvature of 4-dimensional Einstein spaces*, Global Analysis (papers in Honor of K. Kodaira) (edited by D. C. Spencer and S. Iyanaga), Univ. of Tokyo Press, Tokyo, 1969, pp. 355–365. MR 41 #959.
7. K. Yano and S. Bochner, *Curvature and Betti numbers*, Ann. of Maths. Studies, no. 32, Princeton Univ. Press, Princeton, N. J., 1953. MR 15, 989.

DEPARTMENT OF MATHEMATICS, MADURAI UNIVERSITY, MADURAI-21, INDIA.