GENERALIZED SEMIGROUPS OF QUOTIENTS

BY

C. V. HINKLE, JR.

ABSTRACT. For S a semigroup with 0 and M_S a right S -set, certain classes of sub S -sets called right quotient filters are defined. A study of these right quotient filters is made and examples are given including the classes of intersection large and dense sub S -sets respectively. The general semigroup of right quotients Q corresponding to a right quotient filter on a semigroup S is developed and basic properties of this semigroup are noted. A nonzero regular semigroup S is called primitive dependent if each nonzero right ideal of S contains a 0-minimal right ideal of S . The theory developed in the paper enables us to characterize all primitive dependent semigroups having singular congruence the identity in terms of sub-direct products of column monomial matrix semigroups over groups.

1. Introduction. The classical ring of quotients of a ring R and its generalizations have been topics of major interest in ring theory over the past twenty years with much of the basic ground work laid by Asano [1], Johnson [9], and Utumi [15]. Each type of right quotient ring over R is developed from a given type of filter of right ideals in the ring R and the same general technique is applied in each case.

In this paper generalizations of these concepts are developed for semigroups and S -sets and an attempt is made to unify the theory as much as possible by considering certain classes of sub S -sets called right quotient filters. §3 is devoted to a study of these right quotient filters. Several examples which are analogues of concepts in ring theory are given including those of intersection large, dense, and strongly dense. We also consider a certain S -set extension (semigroup extension) of a semigroup S , called a $\mathcal{P}$ -quotient set (semigroup) where $\mathcal{P}$ is a right quotient filter on S , and we show how $\mathcal{P}$ induces a right quotient filter on such an extension. In certain cases the induced filter is of the same type as that of the original filter.

In §4 a generalization of the singular congruence and the torsion relation introduced by Feller and Gantos [6] is given. This generalization, which we call the $\mathcal{P}$ -torsion congruence on an S -set M , is defined in terms of an arbitrary right quo-

Received by the editors October 5, 1972.

AMS (MOS) subject classifications (1970). Primary 20M10.

Key words and phrases. Semigroups of quotients, singular congruences, Rees matrix semigroups, primitive dependent semigroups.

Copyright © 1973, American Mathematical Society

tient filter $\mathcal{P}$ on S , and we are particularly interested in the case where M is $\mathcal{P}$ -torsion free, i.e., where the $\mathcal{P}$ -torsion congruence on M is the identity.

§5 is devoted to the study of the general semigroup of right quotients corresponding to a right quotient filter $\mathcal{P}$ on an S -set. We begin by giving the basic construction of this semigroup. The main theorems of this section point out the fact that whenever S is $\mathcal{P}$ -torsion free, then $Q_{\mathcal{P}}(S)$ is a maximal $\mathcal{P}$ -torsion free $\mathcal{P}$ -quotient semigroup over S which contains an isomorphic copy over S of every such semigroup extension of S . Furthermore, whenever $\mathcal{P}$ is a special filter on S we find that the corresponding quotient semigroup of Q with respect to the induced filter on Q is itself.

As an application of this theory we note that whenever S is $\mathcal{P}_I$ -torsion free, then $Q_I(S)$ is the injective hull of S and is furthermore self-injective. We also restate the well-known theorems concerning the classical semigroup of quotients in terms of the theory developed herein.

In §6, we consider semigroups which are 0-direct unions of semigroups. If a right quotient filter is given on each of the component semigroups of S it is shown that a corresponding right quotient filter is induced on S in a natural way. Here we see that the semigroup of quotients of S is isomorphic to the direct product of the semigroups of quotients on the individual component semigroups. Furthermore, it is shown that every $\mathcal{P}$ -torsion free $\mathcal{P}$ -quotient semigroup over S is isomorphic to some subdirect product of $\mathcal{P}$ -torsion free $\mathcal{P}$ -quotient semigroups over the individual components.

A nonzero regular semigroup S is called *primitive dependent* if each nonzero idempotent of S lies above a primitive idempotent of S under the natural partial ordering on the idempotents of S . The main purpose of §7 is to characterize all primitive dependent semigroups for which the singular congruence is the identity. To this end we first consider completely 0-simple semigroups. A characterization of the singular congruence on a completely 0-simple semigroup expressed as a regular Rees matrix semigroup is given and it is noted that the semigroup of quotients of such a semigroup is isomorphic to a semigroup of column monomial matrices over a group with 0. The completely 0-simple semigroups for which the singular congruence is the identity are then characterized as certain subsemigroups, called $\mathcal{H}$ -semigroups, of these semigroups of column monomial matrices. The section is concluded by characterizing the primitive dependent semigroups for which the singular congruence is the identity as certain subdirect products of column monomial matrix semigroups over groups with 0.

Much of the basic notation and definitions used throughout the paper are given in §2. We assume that the reader is familiar with the basic terminology and results on algebraic semigroups employed herein, as presented in Clifford and Preston [4] and [5].

2. Preliminaries. Throughout this paper each semigroup will contain a zero (0) unless otherwise specified. Let S be a semigroup. A (centered right) S -set M_S is a set M , with an associative scalar operation on M by elements of S , which contains an element (necessarily unique) θ such that $\theta = \theta s = m0$ for all $m \in M$ and for all $s \in S$. The symbol θ will be called the zero of M . Since the distinction between the zero of M and the zero of S is clear from the context, we shall denote both by the same symbol 0. Note that if R is a right ideal of S then R becomes an S -set R_S under ordinary multiplication. Also, if S is contained in a semigroup T and if M_T is a T -set then M becomes an S -set M_S by restricting the scalar multiplication to the elements of S . A sub S -set N_S of an S -set M_S is a subset N of M such that $NS \subseteq N$. If $m, n \in M_S$ and if $E \subseteq S$ we shall say that mE is *pointwise equal* to nE when $ms = ns$ for each $s \in E$. This will be denoted as $mE = nE$.

Let M_S and N_S be S -sets. A function $f: M_S \rightarrow N_S$ is an S -homomorphism if, for each $m \in M$ and $s \in S$, $f(ms) = f(m)s$. The collection of all such S -homomorphisms will be denoted by $\text{Hom}_S(M, N)$. If there exists $f \in \text{Hom}_S(M, N)$ such f is 1-1 and onto then we say M_S is S -isomorphic to N_S and write $M_S \approx_S N_S$. If M_S and N_S are S -sets each containing A_S as a sub S -set and if there exists an S -isomorphism $\phi: M_S \rightarrow N_S$ such that the restriction $\phi|A$ of ϕ to A is the identity map on A then we say that " ϕ is an S -isomorphism over A_S " or " M_S and N_S are S -isomorphic over A_S ". Corresponding terminology will be used for semigroups and semigroup isomorphisms. All homomorphisms between semigroups will be considered as semigroup homomorphisms unless S -homomorphism is clearly indicated. If the semigroups S and T are isomorphic as semigroups we shall write $S \approx T$.

If f is an S -homomorphism the domain of f will be denoted by D_f and the range of f by R_f . The zero map from M_S will be denoted by 0 and the identity map on M by 1_M . If $f: M_S \rightarrow N_S$ and if $A_S \subseteq N_S$ then $f^{-1}(A) = \{m \in M; f(m) \in A\}$. The S -homomorphism f is called *0-restricted* if $f^{-1}(0) = 0$.

An S -congruence τ on M_S is an equivalence relation on M such that $(ms, ns) \in \tau$ whenever $(m, n) \in \tau$ for all $s \in S$. The S -congruence τ is said to be *0-restricted* if $(a, 0) \in \tau$ implies $a = 0$. If N_S is a sub S -set of M_S and if ρ is an S -congruence on M_S then $\rho|N = \rho \cap (N \times N)$ is an S -congruence on N_S .

If S has an identity 1 the S -set M_S is said to be *unital* when $m1 = m$ for each $m \in M$. For each semigroup S we shall define S^1 by $S^1 = S \cup \{1\}$ where 1 is a symbol not in S and where multiplication on S is extended to S^1 by defining $1x = x1 = x$ for each $x \in S^1$. With the operation so defined, S^1 is a semigroup. Note that this definition for S^1 differs from the standard one. However, with the definition given here each S -set M_S becomes a unital S^1 -set by defining $m1 = m$ for each $m \in M$.

The following definitions and theorems are taken from the paper by Berthiaume [3]. A sub S -set N_S of M_S is said to be *large (essential)* in M_S if for each $f \in \text{Hom}(M_S, K_S)$ such that $f|N$ is 1-1 then f is 1-1. In this case M_S is called an *essential extension* of N_S .

Lemma 2.1. N_S is large in M_S iff for every S -congruence ρ on M_S such that $\rho \neq \iota_M$ we have $\rho|N \neq \iota_N$.

An S -set M_S is *injective* if for each $A_S \subseteq B_S$ and for each $f \in \text{Hom}_S(A, M)$ there exists $f' \in \text{Hom}_S(B, M)$ such that $f'|A = f$. If $M_S \subseteq N_S$ and if N_S is injective then N_S is called an *injective extension* of M_S . The following theorem due to Berthiaume guarantees the existence of a minimal injective extension which is unique up to S -isomorphism.

Theorem 2.2. The S -set M_S is a maximal essential extension of N_S iff M_S is a minimal injective extension of N_S . Every S -set N_S has such an extension which is unique up to S -isomorphism over N_S .

The minimal injective extension of N_S given in the above theorem is called the *injective hull* of N_S . Note that M_S is the injective hull of N_S iff N_S is essential in M_S and M_S is injective.

A semigroup S will be called *self-injective* if S_S is injective.

3. Right quotient filters. The general theory developed throughout this paper is largely dependent on a certain type of filter of sub S -sets which will be called a *right quotient filter*. Several examples of these are given. These examples will later provide applications to the general theory developed herein.

The following definitions are generalizations of corresponding concepts in ring theory. Let $N_S \subseteq M_S$ where M_S is an S -set. Then N_S is said to be *intersection large* in M_S if for each $0 \neq m \in M$ there exists $s \in S^1$ such that $0 \neq ms \in N$. Note that N_S is intersection large in M_S iff the intersection of N with any non-zero sub S -set of M_S is always nonzero. A second definition taken from Utumi's paper [15] states that N_S is *dense* in M_S iff for each $0 \neq m, n \in M$ there exists $s \in S^1$ such that $ms \neq 0$ and $ms \in N$. We easily see that N_S is intersection large in M_S whenever N_S is dense in M_S . Less generally, N_S will be called *strongly dense* in M_S if for each $m_1, m_2, n \in M$ where $m_1 \neq m_2$ there exists $s \in S^1$ such that $m_1s \neq m_2s$ and $ns \in N$. This latter definition was suggested to the author by J. K. Luedeman for S -sets in general. In case $M = S$ and S has an identity this is equivalent to the definition of "dense" as given by F. R. McMorris in [10]. The fourth definition is a generalization of the vital ideal discussed by McMorris in [11]. Let C be any subsemigroup (possibly without zero) of S . Then $N_S \subseteq M_S$ will be called *C-vital* if for each $0 \neq m \in M$ there exists $c \in C^1$ such that

$0 \neq mc \in N$. Clearly, each C -vital sub S -set is intersection large in M_S and when $C = S$ these two definitions coincide.

The following lemma shows that each of the four concepts given above has the transitive property. In each case the proof follows from the corresponding definition and will be omitted.

Lemma 3.1. *Let " $\leq$ " denote any one of the terms "intersection large", "dense", "strongly dense", or "C-vital". Then $X_S \leq Y_S \leq Z_S$ iff $X_S \leq Z_S$ where $X_S \subseteq Y_S \subseteq Z_S$.*

Thus " $\leq$ " is a partial order relation on the class of S -sets for each definition given above. In the special case where " $\leq$ " represents "C-vital", the following theorem shows that " $\leq$ " is preserved under inverse images of S -homomorphisms.

Lemma 3.2. *Let M_S and N_S be S -sets and let $\phi \in \text{Hom}_S(M, N)$. If A_S is C -vital in N_S then $\phi^{-1}(A)$ is C -vital in M_S .*

Proof. Let $0 \neq m \in M$. If $\phi(m) = 0$ then $m1 = m \in \phi^{-1}(A)$. So, suppose $\phi(m) \neq 0$. Then there exists $c \in C^1$ such that $0 \neq \phi(m)c = \phi(mc) \in A$. Hence $0 \neq mc \in \phi^{-1}(A)$ and the result follows.

Note that if N_S is C -vital in M_S then $m^{-1}N = \{s \in S: ms \in N\}$ is C -vital in S_S for all $m \in M$. In order to show this, define $\phi_m: S \rightarrow M$ by $\phi_m(s) = ms$. Then $\phi_m \in \text{Hom}_S(S, M)$ and $\phi_m^{-1}(N) = m^{-1}N$ is C -vital in S_S by the lemma.

If we restrict ourselves to the class of sub S -sets of M_S a similar result is valid for the remaining definitions given.

Lemma 3.3. *Let " $\leq$ " denote any of the terms "intersection large", "dense", "strongly dense", or "C-vital". Let $A_S \leq M_S$ and $B_S \leq M_S$ and let $\phi \in \text{Hom}_S(A, M)$. Then $\phi^{-1}(B) \leq M_S$.*

Proof. By Lemma 3.2 we need only consider "dense" and "strongly dense". Let " $\leq$ " denote "dense" and let $0 \neq m, n \in M$. Then there exists $s_1 \in S^1$ such that $0 \neq ms_1$ and $ns_1 \in A$ and there exists $s_2 \in S^1$ such that $0 \neq ms_1s_2$ and $f(ns_1)s_2 \in B$. Therefore, $0 \neq m(s_1s_2)$ and $n(s_1s_2) \in f^{-1}(B)$ and it follows that $f^{-1}(B) \leq M_S$ in this case. The proof is similar when " $\leq$ " denotes "strongly dense".

Let $\mathcal{P}_I = \mathcal{P}_I(M_S)$, $\mathcal{P}_D = \mathcal{P}_D(M_S)$, $\mathcal{P}_{SD} = \mathcal{P}_{SD}(M_S)$, and $\mathcal{P}_{CV} = \mathcal{P}_{CV}(M_S)$ denote the classes of intersection large, dense, strongly dense, and C -vital sub S -sets of M_S respectively. In addition, $\mathcal{P}_B = \mathcal{P}_B(M_S)$ will denote the singleton class $\{M_S\}$. In case M is the semigroup S , the subscript " S " will generally be omitted in the notation above.

In general, a nonempty class of sub S -sets of M_S is called a *right quotient filter* (R.Q.F.) of M_S if

- (1) $A_S \in \mathcal{P}$ and $A_S \subseteq B_S \subseteq M_S$ imply $B_S \in \mathcal{P}$, and
- (2) $A_S, B_S \in \mathcal{P}$ and $f \in \text{Hom}_S(A, M)$ imply $f^{-1}(B) \in \mathcal{P}$.

It is easily seen from Lemma 3.1 and Lemma 3.3 that each of the classes $\mathcal{P}_I, \mathcal{P}_D, \mathcal{P}_{SD}, \mathcal{P}_{CV}$, and $\mathcal{P}_B$ is a right quotient filter on M_S . These specific filters will be called *standard filters*.

Lemma 3.4. *A right quotient filter $\mathcal{P}$ of M_S is closed under finite intersections.*

Proof. Let $A_S, B_S \in \mathcal{P}$ and let $\iota: A \rightarrow M$ be the identity map. Then $A \cap B = \iota^{-1}(B) \in \mathcal{P}$ by property (2) above.

Let $\mathcal{P}$ be a right quotient filter on M_S . A nonempty subclass $\mathcal{P}'$ of $\mathcal{P}$ is said to be a *base* for $\mathcal{P}$ if for each $A \in \mathcal{P}$ there exists $B \in \mathcal{P}'$ such that $B \subseteq A$. Note that a given class of sub S -sets of M_S can be a base for at most one right quotient filter on M_S . An arbitrary class $\mathcal{P}'$ of sub S -sets will be called a *right quotient filter base* if it is a base for some right quotient filter $\mathcal{P}$ on M_S . In this case it is easily seen that $\mathcal{P}'$ is a base for the right quotient filter $\mathcal{P} = \{A_S: A' \subseteq A \subseteq M \text{ for some } A' \in \mathcal{P}'\}$. This fact yields the following proposition.

Proposition 3.5. *A nonempty class $\mathcal{P}'$ of sub S -sets of M_S is a right quotient filter base if and only if for each $A', B' \in \mathcal{P}'$ and $f \in \text{Hom}_S(A', M)$ there exists C' in $\mathcal{P}'$ with $C' \subseteq f^{-1}(B')$.*

Note that if $\mathcal{P}$ is a right quotient filter on M_S and if $N_S \in \mathcal{P}$, $\mathcal{P}|N = \{N \cap A: A \in \mathcal{P}\}$ is a right quotient filter for N_S and is also a base for $\mathcal{P}$. In fact, if $\mathcal{P}$ denotes one of the standard filters on M_S then $\mathcal{P}|N$ is the corresponding standard filter on N_S .

Let S be a semigroup and let $\mathcal{P}$ be a right quotient filter on S_S . An S -set M_S containing S_S as a sub S -set is called a $\mathcal{P}$ -*quotient set* over S if $m^{-1}S = \{s \in S: ms \in S\} \in \mathcal{P}$ for each $m \in M$. In case M is a semigroup containing S as a sub-semigroup, M is called a $\mathcal{P}$ -*quotient semigroup* over S . Clearly, S is always a $\mathcal{P}$ -quotient semigroup over itself. Examples of other $\mathcal{P}$ -quotient sets are given in the easily proved proposition which follows.

Proposition 3.6. *If $\mathcal{P} = \mathcal{P}(M_S)$ denotes one of the standard filters $\mathcal{P}_I, \mathcal{P}_{CV}, \mathcal{P}_D$, or $\mathcal{P}_{SD}$ on M_S and if $S_S \in \mathcal{P}(M_S)$ then M_S is a $\mathcal{P}$ -quotient set over S .*

In the remainder of this section we shall give sufficient conditions for extending a right quotient filter $\mathcal{P}$ on S to a right quotient filter on any $\mathcal{P}$ -quotient semigroup over S . This is given below in the special cases where $\mathcal{P} = \mathcal{P}_I, \mathcal{P}_D$, or $\mathcal{P}_{SD}$.

Theorem 3.7. *Let $S \subseteq T$ where T is a semigroup.*

- (a) *If $S_S \in \mathcal{P}_D(T_S)$ then $\mathcal{P}' = \{ET^1: E \in \mathcal{P}_I(S)\}$ is a base for $\mathcal{P}_I(T)$.*
- (b) *If $S_S \in \mathcal{P}_D(T_S)$ then $\mathcal{P}' = \{ET^1: E \in \mathcal{P}_D(S)\}$ is a base for $\mathcal{P}_D(T)$.*
- (c) *If $S_S \in \mathcal{P}_{SD}(T_S)$ then $\mathcal{P}' = \{ET^1: E \in \mathcal{P}_{SD}(S)\}$ is a base for $\mathcal{P}_{SD}(T)$.*

Proof. We shall give the proof of (a). The same general techniques are applied to prove (b) and (c). Let $E \in \mathcal{P}_I(S)$ and let $0 \neq t \in T$. Since $E_S \in \mathcal{P}_I(T_S)$ by Lemma 3.1, there exists $s \in S^1 \subseteq T^1$ such that $0 \neq ts \in E \subseteq ET^1$. Thus $ET^1 \in \mathcal{P}_I(T)$ and it follows that $\mathcal{P}' \subseteq \mathcal{P}_I(T)$. Now let $E_T \in \mathcal{P}_I(T)$ and let $E' = E \cap S$. We claim that $E' \in \mathcal{P}_I(S)$. Let $0 \neq s \in S$. Since $E \in \mathcal{P}_I(T)$, there exists $t \in T^1$ such that $0 \neq st \in E$. The claim is evident if $t = 1$. Otherwise, since $S_S \in \mathcal{P}_D(T_S)$, there exists $s' \in S^1$ such that $0 \neq sts'$ and $ts' \in S$. Hence we have $0 \neq s(ts') \in E \cap S = E'$ where $(ts') \in S^1$. Thus, $E' \in \mathcal{P}_I(S)$ and it follows that $E'T^1 = (E \cap S)T^1 \subseteq ET^1 \subseteq E$. Therefore, $\mathcal{P}'$ is a base for $\mathcal{P}_I(T)$.

We can generalize Theorem 3.7 as follows.

Theorem 3.8. *Let $S \subseteq T$ where S and T are semigroups. Let $\mathcal{P}'(T_S)$ be a base for a right quotient filter $\mathcal{P}(T_S)$ on T_S . Then $\mathcal{P}'(T) = \{ET^1: E \in \mathcal{P}'(T_S)\}$ is a base for a right quotient filter $\mathcal{P}(T)$ on T_T . Furthermore, $\mathcal{P}(T)$ is independent of the base $\mathcal{P}'(T_S)$ taken for $\mathcal{P}(T_S)$.*

Proof. Let $E_1 T^1$ and $E_2 T^1 \in \mathcal{P}'(T)$ where E_1 and $E_2 \in \mathcal{P}'(T_S)$ and let $f \in \text{Hom}_T(E_1 T^1, T)$. Let $f' = f|_{E_1}$. Then $f' \in \text{Hom}_S(E_1, T)$ and since $\mathcal{P}'(T_S)$ is a base for $\mathcal{P}(T_S)$, there exists $E_3 \in \mathcal{P}'(T_S)$ such that $E_3 \subseteq f'^{-1}(E_2)$. It follows that $E_3 T^1 \subseteq f'^{-1}(E_2) T^1 \subseteq f^{-1}(E_2) T^1 \subseteq f^{-1}(E_2 T^1)$. Hence, by Proposition 3.5, $\mathcal{P}'(T)$ is a filter base on T_T . Now, let $\mathcal{P}'_1(T_S)$ and $\mathcal{P}'_2(T_S)$ be any two bases for $\mathcal{P}(T_S)$ and let $\mathcal{P}'_i(T) = \{ET^1: E \in \mathcal{P}'_i(T_S)\}$, $i = 1, 2$. Let $\mathcal{P}_1(T)$ and $\mathcal{P}_2(T)$ be the right quotient filters on T_T generated by $\mathcal{P}'_1(T)$ and $\mathcal{P}'_2(T)$ respectively. If $F \in \mathcal{P}_1(T)$ then there exists $E \in \mathcal{P}'_1(T_S)$ such that $ET^1 \subseteq F$. Since $\mathcal{P}'_2(T_S)$ is also a base for $\mathcal{P}(T_S)$, there exists $E' \in \mathcal{P}'_2(T_S)$ such that $E' \subseteq E$. Hence $E'T^1 \in \mathcal{P}_2(T)$ and $E'T^1 \subseteq ET^1 \subseteq F$. Therefore, $F \in \mathcal{P}_2(T)$ and it follows that $\mathcal{P}_1(T) \subseteq \mathcal{P}_2(T)$. A symmetrical argument gives the reverse inclusion.

A right quotient filter $\mathcal{P}$ on S_S is said to be *special* if it has the following property:

"If F is a right ideal of S , $E \in \mathcal{P}$, and $x^{-1}F = \{s: xs \in F\} \in \mathcal{P}$ for all $x \in E$, then $F \in \mathcal{P}$."

Lemma 3.9. *Let $\mathcal{P}$ be a right quotient filter on S_S . Then $\mathcal{P}$ is special iff $\mathcal{P}$ has the following property:*

"For each $E \in \mathcal{P}$ and for each $x \in E$, let E_x denote a member of $\mathcal{P}$. Then $E' = \bigcup_{x \in E} xE_x \in \mathcal{P}$."

Proof. ($\Rightarrow$) If $x \in E$ then $x^{-1}E' = \{s: xs \in E'\}$ which contains E_x and $E_x \in \mathcal{P}$. Hence $x^{-1}E' \in \mathcal{P}$ for each $x \in E$ and it follows that $E' \in \mathcal{P}$.

($\Leftarrow$) Let $E' = \bigcup_{x \in E} x(x^{-1}F)$. Then $E' \in \mathcal{P}$ by the hypothesis. However, $E' \subseteq F$. Therefore, $F \in \mathcal{P}$ and we have that $\mathcal{P}$ is special.

Before giving some examples of special quotient filters, we shall digress briefly and consider a semigroup S containing right cancellable elements. Let C be a subsemigroup of the right cancellable elements of S . The semigroup S has the *right common multiple property* (R.C.M.P.) with respect to C if for each $s \in S$ and for each $c \in C$, $cS \cap sC \neq \emptyset$. Let $\mathcal{P}_C = \{R: R \text{ is a right ideal of } S \text{ and } R \cap C \neq \emptyset\}$.

Proposition 3.10. $\mathcal{P}_C$ is a right quotient filter on S_S iff S has the R.C.M.P. with respect to C . In this case, $\mathcal{P}_C = \mathcal{P}_{CV}(S)$.

Proof. ($\Rightarrow$) Let $c \in C$ and $s \in S$. Define $\phi_s: S \rightarrow S$ by $\phi_s(t) = st$ for each $t \in S$. Then $\phi_s \in \text{Hom}_S(S, S)$. Since $c^2 \in cS \cap C$, we have $cS \in \mathcal{P}_C$. Hence by the definition of an R.Q.F. it follows that $\phi_s^{-1}(cS) = \{t \in S: st \in cS\} \in \mathcal{P}_C$. Let $c' \in \phi_s^{-1}(cS) \cap C$. Then $sc' \in cS$ and we have $sC \cap cS \neq \emptyset$. Therefore, S has the R.C.M.P. with respect to C .

($\Leftarrow$) Now, suppose S has the R.C.M.P. with respect to C . We claim that $\mathcal{P}_C = \mathcal{P}_{CV}(S)$. Let $R \in \mathcal{P}_C$ and let $0 \neq s \in S$. Then there exists $n \in R \cap C$ and there exists $c' \in C$ and $s' \in S$ such that $sc' = ns'$. Since $c' \in C$, $sc' \neq 0$ and it follows that $sC^1 \cap R \neq \emptyset$. Therefore, $R \in \mathcal{P}_{CV}(S)$. Now, let $R \in \mathcal{P}_{CV}(S)$ and let $c \in C$. Since $R \in \mathcal{P}_{CV}(S)$, there exists $c' \in C^1$ such that $cc' \in R$. However, $cc' \in C$ and we have $R \cap C \neq \emptyset$. Therefore, $R \in \mathcal{P}_C$ and the theorem follows.

Several examples of special right quotient filters can now be given as shown in the proposition below.

Proposition 3.11. Let S be a semigroup with zero.

(a) If S contains right cancellable elements and if S has the R.C.M.P. with respect to a subsemigroup C of the right cancellable elements of S then $\mathcal{P}_{CV}(S)$ is a special R.Q.F.

(b) $\mathcal{P}_B(S)$ is special iff $S^2 = S$.

(c) If $\{s \in S \mid sS = 0\} = \{0\}$ then $\mathcal{P}_D(S)$ is special.

(d) If $\{s \in S \mid sS = 0\} \neq \{0\}$ then $\mathcal{P}_D(S) = \mathcal{P}_B(S)$.

(e) If S is right reductive (i.e. $sS = tS$ implies $s = t$ for $s, t \in S$) then $\mathcal{P}_{SD}(S)$ is special.

(f) If S is not right reductive then $\mathcal{P}_{SD}(S) = \mathcal{P}_B(S)$.

Proof. (a) Let $E \in \mathcal{P}_{CV}(S)$ and let $E_x \in \mathcal{P}_{CV}(S)$ for each $x \in E$. Since $E \in \mathcal{P}_{CV}$ and since $\mathcal{P}_{CV} = \mathcal{P}_C$ by Proposition 3.10, there exists $c \in C \cap E$ and since

$E_c \in \mathcal{P}_C$, there exists $c' \in C \cap E_c$. Thus we have $cc' \in cE_c \subseteq E' = \bigcup_{x \in E} xE_x$ and it follows that $E' \in \mathcal{P}_C = \mathcal{P}_{C_V}$. Therefore, $\mathcal{P}_{C_V}$ is special by Lemma 3.9.

(b) Since $\mathcal{P}_B(S) = \{S\}$, $\mathcal{P}_B$ is special iff $S = \bigcup_{s \in S} sS = S^2$.

(c) Let $D \in \mathcal{P}_D(S)$ and for each $x \in D$ let $D_x \in \mathcal{P}_D(S)$. By Lemma 3.9 and Lemma 3.1, it will be sufficient to show that $D' = \bigcup_{x \in D} xD_x \in \mathcal{P}_D(D_S)$. Let $0 \neq m, n \in D$. By the hypothesis there exists $s \in S$ such that $ms \neq 0$. Since $D_n \in \mathcal{P}_D(S)$, there exists $s' \in S^1$ such that $mss' \neq 0$ and $ss' \in D_n$. Thus, $m(ss') \neq 0$ and $n(ss') \in nD_n \subseteq D'$. Therefore, $D' \in \mathcal{P}_D(D_S)$.

(d) Let $0 \neq m \in S$ such that $mS = 0$. Then for $D \in \mathcal{P}_D(S)$ and $s \in S$ there exists $s' \in S^1$ such that $0 \neq ms'$ and $ss' \in D$. However, since $mS = 0$ we must have $s' = 1$ and it follows that $D = S$. Therefore, $\mathcal{P}_D(S) = \{S\} = \mathcal{P}_B(S)$ in this case.

(e) Let $D \in \mathcal{P}_{SD}(S)$ and for each $x \in D$ let $D_x \in \mathcal{P}_{SD}(S)$. As in (c), it is sufficient to show that $D' = \bigcup_{x \in D} xD_x \in \mathcal{P}_{SD}(D_S)$. Let $m_1 \neq m_2, n \in D$. Since S is right reductive there exists $s \in S$ such that $m_1s \neq m_2s$. Hence since $D_n \in \mathcal{P}_{SD}(S)$, there exists $s' \in S^1$ such that $m_1ss' \neq m_2ss'$ and $ss' \in D_n$. Therefore, $m_1(ss') \neq m_2(ss')$ and $n(ss') \in nD_n \subseteq D'$ and we have $D' \in \mathcal{P}_{SD}(D_S)$.

(f) Let $m_1 \neq m_2 \in S$ such that $m_1S = m_2S$ and let $D \in \mathcal{P}_{SD}(S)$. Then for $s \in S$, there exists $s' \in S^1$ such that $m_1s' \neq m_2s'$ and $ss' \in D$. However, s' must be 1. Therefore $D = S$ and we have $\mathcal{P}_{SD}(S) = \{S\} = \mathcal{P}_B(S)$.

We will now return to the general problem of extending a right quotient filter $\mathcal{P}$ on S to a right quotient filter on any $\mathcal{P}$ -quotient semigroup over S .

Theorem 3.12. *Let $\mathcal{P}' = \mathcal{P}'(S)$ be a base for a special right quotient filter $\mathcal{P} = \mathcal{P}(S)$ on S_S . If M_S is a $\mathcal{P}$ -quotient set over S then $\mathcal{P}'$ is a base for a right quotient filter $\mathcal{P}(M_S)$ which is independent of the base taken for $\mathcal{P}(S)$.*

Proof. Let $A, B \in \mathcal{P}'$ and let $f \in \text{Hom}_S(A, M)$. By Proposition 3.5, it is sufficient to show that $f^{-1}(B)$ contains an element of $\mathcal{P}'$. For each $a \in A$, let $E_a = [f(a)]^{-1}S$. Then since M_S is a $\mathcal{P}$ -quotient set over S , $E_a \in \mathcal{P}$. Note that for each element $s \in E_a$, $f(as) = f(a)s \in S$. Let $E' = \bigcup_{a \in A} aE_a$. Then $E' \in \mathcal{P}$ since $\mathcal{P}$ is special. If $f' = f|_{E'}$ then $f' \in \text{Hom}_S(E', S)$. Hence $f'^{-1}(B) \in \mathcal{P}(S)$ and it follows that there exists $C \in \mathcal{P}'$ such that $C \subseteq f'^{-1}(B) \subseteq f^{-1}(B)$. An argument similar to that of Theorem 3.8 shows that $\mathcal{P}(M_S)$ is independent of the base $\mathcal{P}'$ taken for $\mathcal{P}(S)$.

Theorem 3.13. *Let $\mathcal{P}' = \mathcal{P}'(S)$ be a base for a special right quotient filter $\mathcal{P}(S)$ on S_S . Let T be a $\mathcal{P}(S)$ -quotient semigroup over S . Then $\mathcal{P}'(T) = \{ET^1: E \in \mathcal{P}'\}$ is a base for a special right quotient filter $\mathcal{P}(T)$ on T_T which is independent of the base $\mathcal{P}'$ taken for $\mathcal{P}(S)$.*

Proof. By Theorem 3.12 and Theorem 3.8, $\mathcal{P}'(T)$ is a base for a right quo-

right filter $\mathcal{P}(T)$ on T_T which is independent of the base $\mathcal{P}'$ taken for $\mathcal{P}(S)$. Hence we need only show that $\mathcal{P}(T)$ is special. Without loss of generality we may assume that $\mathcal{P}' = \mathcal{P}(S)$. Let $D \in \mathcal{P}(T)$ and let $D_x \in \mathcal{P}(T)$ for each $x \in D$. There exists $E \in \mathcal{P}(S)$ such that $E \subseteq ET^1 \subseteq D$ and, for each $e \in E$, there exists $E_e \in \mathcal{P}(S)$ such that $E_e \subseteq E_e T^1 \subseteq D_e$. Let $E' = \bigcup_{e \in E} eE_e$. Then $E' \in \mathcal{P}(S)$ and we have $E' T^1 = (\bigcup_{e \in E} eE_e) T^1 \subseteq (\bigcup_{x \in D} xD_x) T^1 \subseteq \bigcup_{x \in D} xD_x$. Therefore, $\bigcup_{x \in D} xD_x \in \mathcal{P}(T)$ and the theorem follows.

The class $\mathcal{P}(T)$ will be called the special right quotient filter induced by $\mathcal{P}(S)$ on T_T .

The next lemma gives a general analogue of the transitivity property given by Lemma 3.1.

Lemma 3.1.4. *Let $\mathcal{P}(S)$ be a special right quotient filter on S . Let T be a $\mathcal{P}(S)$ -quotient semigroup over S and let $\mathcal{P}(T)$ be the special right quotient filter on T_T induced by $\mathcal{P}(S)$. Let U_T be a $\mathcal{P}(T)$ -quotient T -set over T . Then U_S is a $\mathcal{P}(S)$ -quotient S -set over S .*

Proof. Let $u \in U$. We need to show that $(u^{-1}S)_S \in \mathcal{P}(S)$. Since U_T is a $\mathcal{P}(T)$ -quotient T -set over T , $(u^{-1}T)_T \in \mathcal{P}(T)$ and, since $\mathcal{P}(T)$ is induced by $\mathcal{P}(S)$, there exists $E \in \mathcal{P}(S)$ such that $E \subseteq ET^1 \subseteq (u^{-1}T)_T$. For every $s \in E$, let $E_s = [(us)^{-1}S]_S$. Then $E_s \in \mathcal{P}(S)$ for all $s \in E$, since $us \in T$ and T is a $\mathcal{P}(S)$ -quotient semigroup over S . Let $E' = \bigcup_{s \in E} sE_s$. Then $E' \in \mathcal{P}(S)$ and $E' \subseteq (u^{-1}S)_S$. Therefore, $(u^{-1}S)_S \in \mathcal{P}(S)$.

4. The torsion congruence induced by a right quotient filter. A generalization of the singular congruence and the torsion relation defined by Feller and Gantos [6] will now be given. We begin by defining the $\mathcal{P}$ -torsion relation on M_S and noting that it is an S -congruence when $\mathcal{P}$ is an R.Q.F. Several facts concerning the $\mathcal{P}$ -torsion congruence when $\mathcal{P}$ is one of the standard filters are then given.

In general, let $\mathcal{P}$ be any class of right ideals of S and let M_S be a right S -set. Let $\psi_{\mathcal{P}}(M_S) = \{(m_1, m_2) \in M \times M : m_1 P = m_2 P \text{ for some } P \in \mathcal{P}\}$. Then $\psi_{\mathcal{P}}(M_S)$ is called the $\mathcal{P}$ -torsion relation (or $\mathcal{P}$ -torsion congruence if $\psi_{\mathcal{P}}(M_S)$ is also an S -congruence). When $\mathcal{P}$ is an R.Q.F. on S_S this relation is indeed an S -congruence as shown in the following lemma which is easily proved using the properties of a right quotient filter.

Lemma 4.1. *Let $\mathcal{P}'$ be any base for a right quotient filter $\mathcal{P}$ on S_S . Then*

- (a) $\psi_{\mathcal{P}'}(M_S) = \psi_{\mathcal{P}}(M_S)$;
- (b) $\psi_{\mathcal{P}}(M_S)$ is an S -congruence on M_S ; and
- (c) if $M = S$ then $\psi_{\mathcal{P}}(S_S)$ is a 2-sided congruence on S .

If $\mathcal{P} = \mathcal{P}_I, \mathcal{P}_D, \mathcal{P}_{SD}, \mathcal{P}_{CV}$, or $\mathcal{P}_B$ then the corresponding $\mathcal{P}$ -torsion congruence will be denoted by $\psi_I = \psi_I(M_S)$, $\psi_D = \psi_D(M_S)$, $\psi_{SD} = \psi_{SD}(M_S)$, $\psi_{CV} = \psi_{CV}(M_S)$, or $\psi_B = \psi_B(M_S)$ respectively. In case $M = S$ the subscript S will be omitted. The congruence $\psi_I(M_S)$ is generally known as the *singular congruence* on M_S .

The following lemma shows that the order relation " $\subseteq$ " is preserved in passing from right quotient filters to $\mathcal{P}$ -torsion congruences.

Lemma 4.2. *Let $\mathcal{P}_1$ and $\mathcal{P}_2$ be any two right quotient filters on S_S . If $\mathcal{P}_1 \subseteq \mathcal{P}_2$ then $\psi_{\mathcal{P}_1} \subseteq \psi_{\mathcal{P}_2}$. In particular, we have $\psi_B \subseteq \psi_{SD} \subseteq \psi_D \subseteq \psi_I$ and $\psi_{CV} \subseteq \psi_I$.*

Let S be a semigroup with right cancellable elements and let C be a sub-semigroup of S consisting entirely of right cancellable elements of S such that S has the right common multiple property with respect to C . In Proposition 3.10, it was shown that $\mathcal{P}_{CV} = \{R_S \subseteq S_S : R \cap C \neq \emptyset\} = \mathcal{P}_C$. The next lemma shows that $\psi_{CV}(M_S)$ is the same as the torsion relation defined by Feller and Gantos in [6].

Lemma 4.3. *Let C be as above and let S have the R.C.M.P. with respect to C . Then $\psi_{CV}(M_S) = \{(m_1, m_2) : m_1 c = m_2 c \text{ for some } c \in C\}$.*

Proof. Let $(m_1, m_2) \in \psi_{CV}(M_S)$. Then there exists $E \in \mathcal{P}_{CV}$ such that $m_1 E = m_2 E$. Since $E \cap C \neq \emptyset$, it follows that (m_1, m_2) is a member of the right side. Now, let (m_1, m_2) be an element of the right side. Then there exists $c \in C$ such that $m_1 c = m_2 c$. Let $E = cS$. Then since $c^2 \in cS \cap C$ we have $cS \in \mathcal{P}_{CV}$. Clearly, $m_1 E = m_2 E$ and we have $(m_1, m_2) \in \psi_{CV}(M_S)$.

The following corollary is evident from the lemma.

Corollary 4.4. *Under the hypotheses of Lemma 4.3 $\psi_{CV}(S) = \iota$.*

It is clear from the definitions that $\mathcal{P}_D(M_S)$ is always contained in $\mathcal{P}_I(M_S)$. The next lemma gives a sufficient condition for equality of these two filters.

Lemma 4.5. *If $\psi_I(M_S)$ is 0-restricted then $\mathcal{P}_I(M_S) = \mathcal{P}_D(M_S)$.*

Proof. We need only show that $\mathcal{P}_I(M_S) \subseteq \mathcal{P}_D(M_S)$. Let $N_S \in \mathcal{P}_I(M_S)$ and let $0 \neq m, n \in M$. In a similar manner to the note following Lemma 3.2 we have $n^{-1}N = \{s \in S : ns \in N\} \in \mathcal{P}_I(S)$. If $m(n^{-1}N) = 0$ then $(m, 0) \in \psi_I(M_S)$. However, $m \neq 0$ and $\psi_I(M_S)$ is 0-restricted. Therefore, $m(n^{-1}N) \neq 0$ and it follows that there exists $s \in S^1$ such that $ms \neq 0$ and $ns \in N$. Hence, $N_S \in \mathcal{P}_D(M_S)$.

This lemma and the lemma which follows show that "dense" and "intersection large" are the same concepts on a regular semigroup.

Lemma 4.6. *If $S \subseteq T$ such that $S \in \mathcal{P}_I(T_S)$ and T is a regular semigroup, then $\psi_I(T_S)$ is 0-restricted.*

Proof. Let $E \in \mathcal{P}_I(S)$ and let $0 \neq t \in T$. Let t' be an inverse of t in T . Since

$S \in \mathcal{P}_I(T_S)$ we have $E \in \mathcal{P}_I(T_S)$. Thus there exists $s \in S^1$ such that $0 \neq t's \in E$. Hence $0 \neq t's = t't's \in E$ and it follows that $0 \neq t(t's) \in tE$. Therefore, $\psi_I(T_S)$ must be 0-restricted since E is an arbitrary element of $\mathcal{P}_I(T_S)$ and contains an element x such that $tx \neq 0$ for each $0 \neq t \in T$.

In general, $\mathcal{P}_I(S)$ is not a special right quotient filter. However, the next proposition gives necessary and sufficient conditions for this to be true.

Proposition 4.7. *The following are equivalent:*

- (a) $\mathcal{P}_I(S)$ is special.
- (b) $\psi_I(S)$ is 0-restricted.
- (c) $\mathcal{P}_I(S) = \mathcal{P}_D(S)$ and $\{s \in S: sS = 0\} = 0$.

Proof. (a) $\Rightarrow$ (b). Let $A = \{s \in S: (s, 0) \in \psi_I(S)\}$. We need to show that $A = 0$. By Zorn's lemma there exists a maximal sub S -set A'_S of S such that $A \cap A' = 0$. It easily follows that $B = A \cup A' \in \mathcal{P}_I(S)$. For each $b \in B$, let

$$D_b = \begin{cases} b^{-1}\{0\} & \text{if } b \in A, \\ B & \text{if } b \in A'. \end{cases}$$

Note that $b^{-1}\{0\} \in \mathcal{P}_I(S)$ for each $b \in A$. Hence $D = \bigcup_{b \in B} bD_b \in \mathcal{P}_I(S)$ since $\mathcal{P}_I(S)$ is special. However, we can write

$$D = \left[\bigcup_{b \in A} b(b^{-1}\{0\}) \right] \cup \left[\bigcup_{b \in A'} bB \right] = \bigcup_{b \in A'} bB \subseteq A'.$$

Thus it follows that $A' \in \mathcal{P}_I(S)$ and since $A \cap A' = 0$ we must have $A = 0$.

(b) $\Rightarrow$ (c). This follows immediately from Lemma 4.5.

(c) $\Rightarrow$ (a). This is essentially a restatement of Proposition 3.11(c).

Now let $\mathcal{P}$ denote one of $\mathcal{P}_I$, $\mathcal{P}_D$, or $\mathcal{P}_{SD}$. If S and T are semigroups such that $S \subseteq T$ then T has a $\mathcal{P}(S)$ -torsion congruence as well as a $\mathcal{P}(T)$ -torsion congruence. These are the same under the conditions given below.

Theorem 4.8. *Let S and T be semigroups such that $S \in \mathcal{P}_D(T_S)$. Then $\psi_D(T_S) = \psi_D(T)$ and $\psi_I(T_S) = \psi_I(T)$. Also, if $S \in \mathcal{P}_{SD}(T_S)$ then $\psi_{SD}(T_S) = \psi_{SD}(T)$.*

Proof. Let $\mathcal{P}$ denote any of $\mathcal{P}_D$, $\mathcal{P}_{SD}$, or $\mathcal{P}_I$ and let ψ denote the corresponding torsion congruence. If $(t_1, t_2) \in \psi(T)$ then there exists $E \in \mathcal{P}(T)$ such that $t_1E \doteq t_2E$. By Theorem 3.7 ((a), (b), or (c) depending on the choice of $\mathcal{P}$), there exists $E' \in \mathcal{P}(S)$ such that $E' \subseteq E'T^1 \subseteq E$. Thus $t_1E' \doteq t_2E'$ and we have $(t_1, t_2) \in \psi_{\mathcal{P}}(T_S)$. On the other hand if $(t_1, t_2) \in \psi_{\mathcal{P}}(T_S)$, there exists $E \in \mathcal{P}(S)$ such that $t_1E \doteq t_2E$. Hence $t_1(ET^1) \doteq t_2(ET^1)$ and $ET^1 \in \mathcal{P}(T)$ by the same

theorem mentioned above. Therefore, $(t_1, t_2) \in \psi(T)$ and the result follows.

The theorem can also be stated for any special filter and for any $\mathcal{P}$ -quotient semigroup. The proof is similar to that of the previous theorem.

Theorem 4.9. *If $\mathcal{P} = \mathcal{P}(S)$ is a special right quotient filter on S and T is a $\mathcal{P}$ -quotient semigroup over S then $\psi_{\mathcal{P}(S)}(T_S) = \psi_{\mathcal{P}(T)}(T)$ where $\mathcal{P}(T)$ is the right quotient filter induced on T by $\mathcal{P}$.*

Let $\mathcal{P}$ be an R.Q.F. on S . The S -set M_S will be called $\mathcal{P}$ -torsion free if $\psi_{\mathcal{P}}(M_S) = \iota$. We have already seen that if S has a subsemigroup C of right cancellable elements and has the R.C.M.P. with respect to C then S is $\mathcal{P}_{CV}(S)$ -torsion free. We observe below that S is $\mathcal{P}_{SD}$ -torsion free for a large class of semigroups which includes each semigroup with identity.

Lemma 4.10. *For all semigroups S , $\psi_{SD}(S) = \psi_B(S)$. Hence $\psi_{SD}(S)$ is the identity congruence iff S is right reductive.*

Proof. Clearly, $\psi_B(S) = \iota$ iff S is right reductive. Suppose $(s, t) \in \psi_{SD}(S)$ such that $s \neq t$. Then there exists $D \in \mathcal{P}_{SD}(S)$ such that $sD = tD$. If $s' \in S$ such that $ss' \neq ts'$ then there exists $t' \in S^1$ such that $ss't' \neq ts't'$ and $s't' \in D$, which is a contradiction. Thus $sS = tS$ and it follows that $\psi_B(S) = \psi_{SD}(S)$.

McMorris showed in [12] that $\mathcal{P}_I(S) = \mathcal{P}_{SD}(S)$ when $\psi_I(S) = \iota$. A similar proof gives this result for all S -sets.

Lemma 4.11. *If $\psi_I(M_S) = \iota$ then $\mathcal{P}_{SD}(M_S) = \mathcal{P}_D(M_S) = \mathcal{P}_I(M_S)$.*

We also have the following two lemmas involving $\mathcal{P}$ -quotient sets.

Lemma 4.12. *Let M_S be a $\mathcal{P}$ -torsion free S -set such that $S_S \subseteq M_S$ where $\mathcal{P}$ denotes one of the filters $\mathcal{P}_I, \mathcal{P}_D$, or $\mathcal{P}_{SD}$. Then M_S is a $\mathcal{P}$ -quotient set over S iff $S \in \mathcal{P}(M_S)$.*

Proof. If $S \in \mathcal{P}(M_S)$ then M_S is a $\mathcal{P}$ -quotient set over S by Proposition 3.6. Conversely, suppose that M_S is a $\mathcal{P}$ -quotient set over S . First, consider the case where $\mathcal{P} = \mathcal{P}_D$ and let $0 \neq m, n \in M$. Then $n^{-1}S \in \mathcal{P}(S)$ and $m(n^{-1}S) \neq 0$ since $\psi_D(M_S) = \iota$. Thus there exists $s \in S$ such that $0 \neq ms$ and $ns \in S$ and it follows that $S \in \mathcal{P}_D(M_S)$. Next, suppose that $\mathcal{P} = \mathcal{P}_I$. Since $\psi_I(M_S) = \iota$, we have $\mathcal{P}_I(M_S) = \mathcal{P}_D(M_S)$ by Lemma 4.11 and it also follows that $\psi_I(S) = \iota$. Thus $\mathcal{P}_I(S) = \mathcal{P}_D(S)$ by Lemma 4.11 again and the result follows from the previous case. The proof for the case $\mathcal{P} = \mathcal{P}_{SD}$ is similar to that of the first case and will be omitted.

Lemma 4.13. *Let $\mathcal{P}$ be an arbitrary right quotient filter on S such that S is $\mathcal{P}$ -torsion free and let M_S be any $\mathcal{P}$ -quotient set over S_S . Then M_S is $\mathcal{P}$ -torsion free iff S_S is large in M_S .*

Proof. ($\Rightarrow$) Let $\Phi \in \text{Hom}_S(M, K)$ such that $\Phi|_S$ is 1-1 where K_S is an arbitrary S -set. Let $m_1, m_2 \in M$ such that $\Phi(m_1) = \Phi(m_2)$. Let $D = m_1^{-1}S \cap m_2^{-1}S \in \mathcal{P}$. Since $\Phi(m_1s) = \Phi(m_2s)$ for all $s \in D$ and since $\Phi|_S$ is 1-1, we have $m_1D = m_2D$. Therefore, $m_1 = m_2$ since M_S is $\mathcal{P}$ -torsion free and it follows that S_S is large in M_S .

($\Leftarrow$) Since $\psi_{\mathcal{P}}(S) = \psi_{\mathcal{P}}(M_S)|_S = \iota$, it follows from Lemma 1.1 that $\psi_{\mathcal{P}}(M_S) = \iota$.

5. Right quotient semigroups. Let S be a semigroup, M_S be a right S -set and $\mathcal{P}$ be a right quotient filter on M_S . Let $F = F_{\mathcal{P}}(M_S) = \bigcup_{D \in \mathcal{P}} \text{Hom}_S(D, M)$ and define multiplication on F by $fg = b$ where $b: D_g \cap g^{-1}(D_f) \rightarrow M$ is given by $b(x) = f(g(x))$. Then under this multiplication F is a semigroup, called the *semigroup of partial S -homomorphisms of M_S with respect to $\mathcal{P}$* .

Lemma 5.1. Define a binary relation $\omega = \omega_{\mathcal{P}}(M_S)$ on the semigroup F by $(f, g) \in \omega$ iff there exists $D \in \mathcal{P}$ such that $f|_D = g|_D$. Then ω is a 2-sided congruence on F .

The semigroup $Q_{\mathcal{P}} = Q_{\mathcal{P}}(M_S) = F/\omega$ will be called the *semigroup of right quotients of M_S with respect to $\mathcal{P}$* . The elements of $Q_{\mathcal{P}}(M_S)$ will be denoted by $\bar{f}$ where $f \in F$. When $\mathcal{P} = \mathcal{P}_I, \mathcal{P}_D, \mathcal{P}_{SD}, \mathcal{P}_{CV}$, or $\mathcal{P}_B$ then $Q_{\mathcal{P}}$ will be denoted by Q_I, Q_D, Q_{SD}, Q_{CV} , or Q_B respectively. As before, the subscript S in $Q_{\mathcal{P}}(M_S)$ will be omitted when $M = S$.

Note that $Q_B(M_S) = \text{Hom}_S(M, M)$. More generally, the following lemma shows that if the R.Q.F. $\mathcal{P}$ on M_S has a least element D_S under set inclusion then there is a natural isomorphism between $Q_{\mathcal{P}}(M_S)$ and $\text{Hom}_S(D, D)$.

Lemma 5.2. Let $\mathcal{P}$ be a right quotient filter on M_S with a least element D_S . Then $Q_{\mathcal{P}}(M_S) \approx \text{Hom}_S(D, D)$ under the map $f \rightarrow \bar{f}$ where $f \in \text{Hom}_S(D, D)$ and multiplication on $\text{Hom}_S(D, D)$ is composition.

Proof. The isomorphism will clearly follow once it is shown that $\text{Hom}_S(D, M) = \text{Hom}_S(D, D)$. Let $f \in \text{Hom}_S(D, M)$. Then $f^{-1}(D) \in \mathcal{P}$ and since D is the least element of $\mathcal{P}$ we have $D \subseteq f^{-1}(D)$. Thus, $D = f^{-1}(D)$ and it follows that $f(D) \subseteq D$. Therefore, $f \in \text{Hom}_S(D, D)$. Since the reverse inclusion is obvious, we have equality.

In this paper we are mainly concerned with the case where M is the semigroup S . As noted below we see that there is a natural representation of S in $Q_{\mathcal{P}}(S)$.

Let $\mathcal{P}$ be a right quotient filter on S_S . For each $s \in S$, define $\phi_s: S \rightarrow S$ by $\phi_s(t) = st$. Then $\phi_s \in \text{Hom}_S(S, S)$. It is easily seen that the mapping $\phi: S \rightarrow Q = Q_{\mathcal{P}}(S)$ by $\phi(s) = \bar{\phi}_s$ is a representation of S in $Q_{\mathcal{P}}(S)$. The image of S under ϕ will be denoted by $\bar{S}$. Since $\bar{S}$ is a subsemigroup of Q , Q may be regarded as a

centered right $\bar{S}$ -set $Q_{\bar{S}}$ in a natural way. Also, Q becomes a centered right S -set Q_S by defining $\bar{f}s = \bar{f}\phi_s$ for each $\bar{f} \in Q$ and for each $s \in S$. It is easy to show that the following lemmas involving the representation of S in $Q_{\mathcal{P}}(S)$ are valid.

Lemma 5.3. $\psi_{\mathcal{P}}(S) = \phi^{-1} \circ \phi$.

Lemma 5.4. For each $f \in F_{\mathcal{P}}(S)$ and for each $s \in D_f$, $f\phi_s = \phi_{f(s)}$.

When $\psi_{\mathcal{P}}(S) = \iota$, we shall assume that S is embedded in $Q = Q_{\mathcal{P}}(S)$ under the identification $s \leftrightarrow \bar{\phi}_s$. From Lemma 5.4 we see that $\bar{f}s = f(s)$ for each $\bar{f} \in Q_{\mathcal{P}}(S)$ and for each $s \in D_f$ under the identification described above. Furthermore, since $D_f \subseteq (\bar{f})^{-1}S$ and $D_f \in \mathcal{P}$, $Q_{\mathcal{P}}(S)$ is also a $\mathcal{P}$ -quotient semigroup over S . In addition, the next lemma shows that Q_S is $\mathcal{P}(S)$ -torsion free.

Lemma 5.5. Let $\mathcal{P} = \mathcal{P}(S)$ be a right quotient filter on S and let $Q = Q_{\mathcal{P}}(S)$. If S is $\mathcal{P}$ -torsion free then Q_S is $\mathcal{P}$ -torsion free.

Proof. Let $(\bar{f}_1, \bar{f}_2) \in \psi_{\mathcal{P}}(Q_S)$. Then there exists $E \in \mathcal{P}$ such that $\bar{f}_1 E = \bar{f}_2 E$. Let $E' = E \cap D_{f_1} \cap D_{f_2} \in \mathcal{P}$. Then for each $s \in E'$, we have $f_1(s) = \bar{f}_1 s = \bar{f}_2 s = f_2(s)$ and it follows that $\bar{f}_1 = \bar{f}_2$.

The following corollary is immediate from Lemma 4.13 and the remarks preceding the above theorem.

Corollary 5.6. Let $\mathcal{P}$ be a right quotient filter on S and let $Q = Q_{\mathcal{P}}(S)$. If S is $\mathcal{P}$ -torsion free then S_S is large in Q_S .

The next theorem shows among other things that each $\mathcal{P}$ -torsion free $\mathcal{P}$ -quotient set (semigroup) can be embedded in $Q_{\mathcal{P}}(S)$ as an S -set (semigroup) such that the elements of S are fixed under the identification $s \leftrightarrow \bar{\phi}_s$.

Theorem 5.7. Let M_S be a $\mathcal{P}$ -quotient set (semigroup) over S . Then there exists an S -homomorphism (homomorphism) $\Phi: M_S \rightarrow Q_{\mathcal{P}}(S) = Q_S$ such that

- (a) $\Phi(s) = \bar{\phi}_s$ for each $s \in S$ and
- (b) $\Phi^{-1} \circ \Phi = \psi_{\mathcal{P}}(M_S)$.

Proof. Let $m \in M$. Since M_S is a $\mathcal{P}$ -quotient set over S , $m^{-1}S \in \mathcal{P}$. Define $\theta_m: m^{-1}S \rightarrow S$ by $\theta_m(s) = ms$. Then $\bar{\theta}_m \in Q = Q_{\mathcal{P}}(S)$. Let $\Phi: M \rightarrow Q$ by $\Phi(m) = \bar{\theta}_m$. For each $s \in S$, we have $D_{\theta_s} = s^{-1}S = S$. Thus $\Phi(s) = \bar{\theta}_s = \bar{\phi}_s$ and (a) follows. Furthermore, $\Phi(m_1) = \Phi(m_2)$ iff $\bar{\theta}_{m_1} = \bar{\theta}_{m_2}$ which is true iff there exists $E \in \mathcal{P}$ such that $m_1 s = \theta_{m_1}(s) = \theta_{m_2}(s) = m_2 s$ for all $s \in E$. Hence it follows that $\Phi^{-1} \circ \Phi = \psi_{\mathcal{P}}(M_S)$. Now let $m \in M$ and $s \in S$. Then $\Phi(ms) = \bar{\theta}_{ms}$ and $\Phi(m)s = \bar{\theta}_m s = \bar{\theta}_m \bar{\phi}_s = \bar{\theta}_{m\phi_s}$. Let $t \in D_{\theta_{ms}} \cap D_{\theta_{m\phi_s}} \in \mathcal{P}$. Then $\theta_{ms}(t) = (ms)t = m(st) = \theta_{m\phi_s}(st) = \theta_{m\phi_s}(t)$. Thus it follows that Φ is an S -homomorphism. Now suppose M is a $\mathcal{P}$ -quotient semigroup over S . Let $m_1, m_2 \in M$. In order to show that Φ is a

semigroup homomorphism it is sufficient to show that $\bar{\theta}_{m_1 m_2} = \bar{\theta}_{m_1} \bar{\theta}_{m_2}$. Let $t \in D_{\theta_{m_1 m_2}} \cap D_{\theta_{m_1} \theta_{m_2}} \in \mathcal{P}$. Then $\theta_{m_1 m_2}(t) = (m_1 m_2)t = m_1(m_2 t) = \theta_{m_1} \theta_{m_2}(t)$. Therefore, $\bar{\theta}_{m_1 m_2} = \bar{\theta}_{m_1} \bar{\theta}_{m_2}$.

As the following corollary shows, $Q_{\mathcal{P}}(S)$ is the maximal $\mathcal{P}$ -torsion free $\mathcal{P}$ -quotient semigroup over S (up to semigroup isomorphism over S) when S is $\mathcal{P}$ -torsion free.

Corollary 5.8. *If S is $\mathcal{P}$ -torsion free then $Q = Q_{\mathcal{P}}(S)$ is a $\mathcal{P}$ -torsion free $\mathcal{P}$ -quotient semigroup over S which contains an S -isomorphic (isomorphic) copy over S of every $\mathcal{P}$ -torsion free $\mathcal{P}$ -quotient set (semigroup) over S .*

Proof. By Lemma 5.5 and the remarks preceding it, Q is a $\mathcal{P}$ -torsion free $\mathcal{P}$ -quotient semigroup over S . Since $\psi_{\mathcal{P}}(S) = \iota$, the rest follows from Theorem 5.7.

Proposition 5.9. *If T is a $\mathcal{P}$ -quotient semigroup over S and Φ is a semigroup endomorphism on T such that $\Phi|_S = \iota_S$ then $\Phi \subseteq \psi_{\mathcal{P}}(T_S)$.*

Proof. Suppose $\Phi(t_1) = t_2$. Let $D = t_1^{-1}S \cap t_2^{-1}S \in \mathcal{P}$. Then for each $s \in D$, we have $t_1 s = \Phi(t_1 s) = \Phi(t_1) \Phi(s) = t_2 s$. Thus, it follows that $(t_1, t_2) \in \psi_{\mathcal{P}}(T_S)$.

Corollary 5.10. *If T is a $\mathcal{P}$ -torsion free $\mathcal{P}$ -quotient semigroup over S then the only semigroup endomorphism on T fixing S is the identity.*

Let $\mathcal{P} = \mathcal{P}(S)$ be a special right quotient filter on S such that S is $\mathcal{P}$ -torsion free. Let $\mathcal{P}(Q)$ be the special right quotient filter on $Q = Q_{\mathcal{P}}(S)$ regarded as a Q -set as given by Theorem 3.13. By Lemma 5.5, Q_S is $\mathcal{P}(S)$ -torsion free and it follows from Theorem 4.9 that Q_Q is also $\mathcal{P}(Q)$ -torsion free. Thus there exists a natural embedding of Q in $Q' = Q_{\mathcal{P}(Q)}(Q)$ and under this embedding Q'_Q is a $\mathcal{P}(Q)$ -quotient semigroup over Q . The next theorem shows that this embedding is actually onto Q' .

Theorem 5.11. *Let $\mathcal{P} = \mathcal{P}(S)$ be a special right quotient filter on S such that S is $\mathcal{P}$ -torsion free and let $Q = Q_{\mathcal{P}}(S)$. Then $Q' = Q_{\mathcal{P}(Q)}(Q) = Q$ with the identification given above where $\mathcal{P}(Q)$ is the special right quotient filter induced on Q_Q by $\mathcal{P}(S)$.*

Proof. By Lemma 5.5 with S replaced by Q , we have Q'_Q is $\mathcal{P}(Q)$ -torsion free. We claim that Q'_S is $\mathcal{P}(S)$ -torsion free. Let $(t_1, t_2) \in \psi_{\mathcal{P}(S)}(Q'_S)$. Then there exists $E \in \mathcal{P}(S)$ such that $t_1 E = t_2 E$. Hence $t_1(EQ) = t_2(EQ)$ and $EQ \in \mathcal{P}(Q)$. Thus $(t_1, t_2) \in \psi_{\mathcal{P}(Q)}(Q'_Q) = \iota$ and it follows that Q'_S is $\mathcal{P}(S)$ -torsion free. Furthermore, by Lemma 3.14, Q'_S is a $\mathcal{P}(S)$ -quotient semigroup over S . So, by Corollary 5.8 there exists a semigroup isomorphism Φ of Q' into Q which fixes S . Let $\Phi' = \Phi|_Q$. Then by Corollary 5.10, $\Phi' = \iota_Q$. Therefore, since $\Phi: Q' \rightarrow Q$ is an isomorphism which fixes Q it follows that $Q' = Q$.

From Proposition 3.11 we see that $\mathcal{P}_D(S)$ is special whenever S is $\mathcal{P}_D$ -torsion free. Also, from Lemma 4.10 and Proposition 3.11, $\mathcal{P}_{SD}(S)$ is special whenever S is $\mathcal{P}_{SD}$ -torsion free. Hence the corollary given below for these particular cases follows.

Corollary 5.12. *If S is $\mathcal{P}_D$ -torsion free then $Q_D(Q_D(S)) = Q_D(S)$ and if S is $\mathcal{P}_{SD}$ -torsion free then $Q_{SD}(Q_{SD}(S)) = Q_{SD}(S)$.*

In [8], the author noted that $Q_I(S)$ is the injective hull of S_S and is self-injective whenever S is $\mathcal{P}_I$ -torsion free. These results can also be obtained as applications of the preceding theory.

Theorem 5.13. *If S is $\mathcal{P}_I$ -torsion free then $Q = Q_I(S)$ is the injective hull of S_S .*

Proof. Since $\psi_I(S) = \iota$, we see from Corollary 5.6 that S_S is large in Q_S . Let M_S be the injective hull of S_S . Since S_S is large in M_S , $S_S \in \mathcal{P}_I(M_S)$ as shown by Feller and Gantos in [6] and we see that M_S is a $\mathcal{P}_I$ -quotient set over S . Also by Lemma 4.13, M_S is $\mathcal{P}_I$ -torsion free. Thus, from Corollary 5.8 we may assume that M_S is chosen such that $S_S \subseteq M_S \subseteq Q_S$. However, since S_S is large in Q_S and since M_S is a maximal essential extension of S_S we must have $M_S = Q_S$ and the result follows.

Theorem 5.14. *If S is $\mathcal{P}_I$ -torsion free then $Q = Q_I(S)$ is self-injective.*

Proof. By Lemma 5.5 we have $\psi_I(Q_S) = \iota$. Since $\mathcal{P}_I(S) = \mathcal{P}_D(S)$ by Lemma 4.5, $S \in \mathcal{P}_D(Q_S)$ by Lemma 4.12. Hence by Theorem 4.8, $\psi_I(Q) = \iota$. So, again by Lemma 4.5, $\mathcal{P}_I(Q) = \mathcal{P}_D(Q)$. Therefore, since $\mathcal{P}_D(S) = \mathcal{P}_I(S)$ and $\mathcal{P}_D(Q) = \mathcal{P}_I(Q)$ it follows from Corollary 5.12 that $Q_I(Q_I(S)) = Q_I(S)$. Thus Q_Q is injective by Theorem 5.13 where S is replaced by Q in the theorem.

Let C be a subsemigroup of the 2-sided cancellative elements of S . A semigroup Q with identity containing S as a subsemigroup is called a *classical semigroup of right quotients of S with respect to C* if (a) each element of C has a 2-sided inverse in Q , and (b) $Q = \{ab^{-1} : a \in S, b \in C\}$.

The following theorem is a generalization of a theorem in ring theory proved by Asano in [1]. A classical proof of this theorem was also given by Smith in [14].

Theorem 5.15. *S has a classical semigroup of right quotients with respect to C iff S has the R.C.M.P. with respect to C .*

Proof. ($\Rightarrow$) Let $S \subseteq Q$ where Q is a classical semigroup of right quotients over S with respect to C . Let $n \in C$ and $s \in S$. There exists $s_1 \in S$ and $n_1 \in C$ such that $n^{-1}s = s_1n_1^{-1}$. Hence $sn_1 = ns_1$ and it follows that $sC \cap nS \neq \emptyset$.

($\Leftarrow$) If S has the R.C.M.P. with respect to C then $\mathcal{P}_{CV}$ as defined in §3

is a special right quotient filter on S by Proposition 3.11. Also by Corollary 4.4, S_S is $\mathcal{P}_{CV}$ -torsion free. Let $Q = Q_{CV}(S)$. As before we can identify S with the semigroup $\Phi(S) = \bar{S}$ in Q . Hence without loss of generality $S \subseteq Q$ under the identification $s \leftrightarrow \bar{\phi}_s$. For each $n \in C$, $\phi_n: S \rightarrow nS$ is 1-1 since n is left cancellative. Hence $\phi_n^{-1}: nS \rightarrow S$ exists and $\phi_n \phi_n^{-1} = \bar{\phi}_n^{-1} \bar{\phi}_n = \bar{\tau}_S$. Thus each element of C has a 2-sided inverse in Q . Now let $\bar{f} \in Q$ and let $n \in D_f \cap C$. Then $\bar{f}n = f(n)$. Therefore $\bar{f} = f(n)n^{-1} \in SC^{-1}$ and the theorem follows.

A semigroup S' (without 0) is said to be *left reversible* if $s_1 S' \cap s_2 S' \neq \emptyset$ for every $s_1, s_2 \in S'$. The following well-known theorem due to Ore (see Clifford and Preston [4]) now follows as a corollary of Theorem 5.15.

Corollary 5.16. *A cancellative semigroup S' (without 0) is embeddable in a group of right quotients of S' iff S' is left reversible.*

Proof. Let $S = S'^0$ and let $C = S'$. If S' is left reversible then S has the R.C.M.P. with respect to $C = S'$. Hence S has a semigroup of right quotients Q with respect to S' . The properties of Q immediately yield the fact that Q is a group with 0 adjoined. Let $A' = Q \setminus \{0\}$. Then Q' is a group of right quotients of S' . The proof of the converse is easy and will be omitted.

Lemma 5.17. *Let $S \subseteq T \subseteq U$ where T and U are classical semigroups of quotients over S with respect to C . Then $T = U$.*

Theorem 5.18. *If S has a classical semigroup of quotients T with respect to C then T is unique up to isomorphism.*

Proof. Let T be any classical semigroup of quotients of S with respect to C and let $Q = Q_{CV}(S)$ be the one obtained in Theorem 5.15. If $t \in T$ then $t = sn^{-1}$ for some $s \in S$ and some $n \in C$. Thus $n \in t^{-1}S$ and it readily follows that $t^{-1}S \in \mathcal{P}_{CV}(S)$. Since each element of C has an inverse in T , it is immediate from Lemma 4.3 that T_S is $\mathcal{P}_{CV}$ -torsion free. Thus T is $\mathcal{P}_{CV}$ -torsion free and a $\mathcal{P}_{CV}$ -quotient semigroup over S . Hence by Corollary 5.8, T can be embedded in Q such that $S \subseteq T \subseteq Q$ and the result follows from Lemma 5.17.

6. Right quotient semigroups of 0-direct unions of semigroups. A right ideal R of a semigroup S is called a *null right ideal* if $RS = 0$. Throughout this section we shall let $\{S_\alpha\}_{\alpha \in \Omega}$ be a collection of 0-disjoint semigroups each having no nonzero null right ideals. The semigroup S will denote the 0-direct union of $\{S_\alpha\}_{\alpha \in \Omega}$ (i.e., S is the union of the S_α 's and $S_\alpha S_\beta = 0$ for $\alpha \neq \beta$). Note that a natural embedding of S in $\Pi_{\alpha \in \Omega} S_\alpha$ is given by $\Lambda: s \rightarrow \hat{s}$ where

$$\hat{s}(\alpha) = \begin{cases} s & \text{if } s \in S_\alpha, \\ 0 & \text{otherwise.} \end{cases}$$

When convenient, we shall identify S with its image in $\prod_{\alpha \in \Omega} S_\alpha$.

For each $\alpha \in \Omega$, let $\mathcal{P}_\alpha$ be a right quotient filter on S_α . The next theorem shows that the collection $\{\mathcal{P}_\alpha\}_{\alpha \in \Omega}$ induces a right quotient filter $\mathcal{P}$ on S .

Theorem 6.1. *For each $\alpha \in \Omega$, let $\mathcal{P}'_\alpha$ be a base for the right quotient filter $\mathcal{P}_\alpha$ on S_α . Let $\mathcal{P}' = \{A_S \subseteq S : A \cap S_\alpha \in \mathcal{P}'_\alpha \text{ for each } \alpha \in \Omega\}$. Then $\mathcal{P}'$ is a base for a right quotient filter $\mathcal{P}$ on S which is independent of the bases $\mathcal{P}'_\alpha$ taken for $\mathcal{P}_\alpha$, $\alpha \in \Omega$. Furthermore, if $\mathcal{P}'_\alpha = \mathcal{P}_\alpha$ for each $\alpha \in \Omega$ then $\mathcal{P}'$ is the right quotient filter $\mathcal{P}$.*

Proof. Let $A, B \in \mathcal{P}'$. Let $A_\alpha = A \cap S_\alpha$ and let $B_\alpha = B \cap S_\alpha$ for each $\alpha \in \Omega$. Let $f \in \text{Hom}_S(A, S)$ and let $f_\alpha = f|_{A_\alpha}$. We claim that $f_\alpha \in \text{Hom}_{S_\alpha}(A_\alpha, S_\alpha)$. It is sufficient to show that $f_\alpha(A_\alpha) \subseteq S_\alpha$. Suppose $f_\alpha(s) = t \in S_\beta$ where $\beta \neq \alpha$. Then $f_\alpha(s)S = tS = 0$. Since S has no nonzero null right ideals it follows that $t = 0$. Since $A_\alpha, B_\alpha \in \mathcal{P}'_\alpha$ and since $\mathcal{P}'_\alpha$ is a base for $\mathcal{P}$ on S_α , there exists $C_\alpha \in \mathcal{P}'_\alpha$ such that $C_\alpha \subseteq f_\alpha^{-1}(B_\alpha)$. Let $C = \bigcup_\alpha C_\alpha$. Then $C \in \mathcal{P}'$ and clearly $C \subseteq f^{-1}(B)$. Therefore, $\mathcal{P}'$ is a base for a right quotient filter $\mathcal{P}$ on S . Now suppose that $\mathcal{P}''_\alpha$ is also a base for $\mathcal{P}_\alpha$, $\alpha \in \Omega$ and let $\mathcal{P}'' = \{A_S \subseteq S : A \cap S_\alpha \in \mathcal{P}''_\alpha \text{ for each } \alpha \in \Omega\}$. We claim that each element A of $\mathcal{P}'$ contains an element of $\mathcal{P}''$. If $A_\alpha = A \cap S_\alpha$ then $A_\alpha \in \mathcal{P}'_\alpha \subseteq \mathcal{P}_\alpha$ and since $\mathcal{P}''_\alpha$ is also a base for $\mathcal{P}_\alpha$ there exists $A'_\alpha \in \mathcal{P}''_\alpha$ such that $A'_\alpha \subseteq A_\alpha$. Let $A' = \bigcup_{\alpha \in \Omega} A'_\alpha$. Then $A' \in \mathcal{P}''$ and we have $A' \subseteq A$. A symmetrical argument shows that each element of $\mathcal{P}''$ contains an element of $\mathcal{P}'$. Thus it follows that $\mathcal{P}'$ and $\mathcal{P}''$ are both bases for the same R.Q.F. $\mathcal{P}$ on S . Finally, suppose that each $\mathcal{P}_\alpha$ is an R.Q.F. on S_α and let $A, B \in \mathcal{P}'$ and $f \in \text{Hom}_S(A, S)$. Then

$$f_\alpha = f|_{A \cap S_\alpha} \in \text{Hom}_{S_\alpha}(A \cap S_\alpha, S_\alpha)$$

and

$$f^{-1}(B) = f^{-1} \left[\bigcup_{\alpha \in \Omega} (B \cap S_\alpha) \right] = \bigcup_{\alpha \in \Omega} [f^{-1}(B \cap S_\alpha)] = \bigcup_{\alpha \in \Omega} [f_\alpha^{-1}(B \cap S_\alpha)] \in \mathcal{P}'.$$

Therefore, in this case $\mathcal{P}'$ is an R.Q.F. on S and the theorem follows.

The filter $\mathcal{P}$ on S given by the above theorem will be called the *right quotient filter induced on S by $\{\mathcal{P}_\alpha\}_{\alpha \in \Omega}$* . For the remainder of this section, $\mathcal{P}$ will always denote this induced filter, unless a specific statement to the contrary is made. If each $\mathcal{P}_\alpha$ is a standard filter on S_α of a given type, the next result shows that $\mathcal{P}$ is the standard filter on S of that type.

Lemma 6.2. *If $\mathcal{P}_\alpha$ is $\mathcal{P}_D(S_\alpha)$, $\mathcal{P}_I(S_\alpha)$, or $\mathcal{P}_B(S_\alpha)$ respectively for each $\alpha \in \Omega$ then $\mathcal{P}$ is $\mathcal{P}_D(S)$, $\mathcal{P}_I(S)$, or $\mathcal{P}_B(S)$ in the same order.*

Proof. Let $A \in \mathcal{P}_D(S)$ and let $A_\alpha = A \cap S_\alpha$ for each $\alpha \in \Omega$. Let $0 \neq m, n \in S_\alpha \subseteq S$. Then there exists $s \in S^1$ such that $ms \neq 0$ and $ns \in A$. However, $m \in S_\alpha$ and $ms \neq 0$. Hence $s \in S_\alpha^1$ and it follows that $ms \neq 0$ and $ns \in A \cap S_\alpha = A_\alpha$. So, we have $A_\alpha \in \mathcal{P}_D(S_\alpha)$ for each α and it follows that $A \in \mathcal{P}$. Now suppose that $A \in \mathcal{P}$. Then $A_\alpha = A \cap S_\alpha \in \mathcal{P}_\alpha = \mathcal{P}_D(S_\alpha)$ for each $\alpha \in \Omega$. Let $0 \neq m, n \in S$, say $m \in S_\alpha$ and $n \in S_\beta$. If $\alpha = \beta$ then there exists $s \in S_\alpha^1$ such that $ms \neq 0$ and $ns \in A_\alpha \subseteq A$. Suppose $\alpha \neq \beta$. Since S has no nonzero null right ideals there exists $s \in S$ such that $ms \neq 0$. Thus $s \in S_\alpha$ and we have $ns = 0 \in A$. Hence in either case there exists $s \in S^1$ such that $ms \neq 0$ and $ns \in A$. Therefore $A \in \mathcal{P}_D(S)$ and it follows that $\mathcal{P} = \mathcal{P}_D(S)$. The proof for the remaining cases follows in a similar way from the corresponding definitions and will be omitted.

Returning now to right quotient filters in general, let us see how the $\mathcal{P}$ -torsion congruence on S is determined entirely from the individual $\mathcal{P}_\alpha$ -torsion congruences on the S_α 's.

Proposition 6.3. $\psi_{\mathcal{P}}(S) = \bigcup_{\alpha} \psi_{\mathcal{P}_\alpha}(S_\alpha) \cup \{(a, b) : (a, 0) \in \psi_{\mathcal{P}_\alpha}(S_\alpha) \text{ and } (0, b) \in \psi_{\mathcal{P}_\beta}(S_\beta) \text{ for some } \alpha, \beta \in \Omega\}$.

Proof. Let $(a, b) \in \psi_{\mathcal{P}}(S)$. Then there exists $E \in \mathcal{P}$ such that $as = bs$ for all $s \in E$. If a and b are both in the same S_α then $as = bs$ for all $s \in E \cap S_\alpha \in \mathcal{P}_\alpha$. Hence $(a, b) \in \psi_{\mathcal{P}_\alpha}(S_\alpha)$. Suppose that $a \in S_\alpha$ and $b \in S_\beta$ where $\alpha \neq \beta$. Then $as = bs = 0$ for all $s \in E \cap S_\alpha$ and $bs = as = 0$ for all $s \in E \cap S_\beta$. Therefore, in this case $(a, 0) \in \psi_{\mathcal{P}_\alpha}(S_\alpha)$ and $(0, b) \in \psi_{\mathcal{P}_\beta}(S_\beta)$. Now, if $(a, b) \in \psi_{\mathcal{P}_\alpha}(S_\alpha)$ for some $\alpha \in \Omega$ then there exists $A_\alpha \in \mathcal{P}_\alpha$ such that $aA_\alpha = bA_\alpha$. Let $A = A_\alpha \cup (\bigcup_{\beta \neq \alpha} S_\beta)$. Then $A \in \mathcal{P}$ and $aA = bA$. Thus we have $(a, b) \in \psi_{\mathcal{P}}(S)$ and it follows that $\psi_{\mathcal{P}_\alpha}(S_\alpha) \subseteq \psi_{\mathcal{P}}(S)$. The remaining inclusion is immediate from this and the transitivity of $\psi_{\mathcal{P}}(S)$.

Corollary 6.4. S is $\mathcal{P}$ -torsion free iff each S_α is $\mathcal{P}_\alpha$ -torsion free.

The next theorem shows that the semigroup of right quotients of S with respect to $\mathcal{P}$ is determined by the semigroups $\{Q_{\mathcal{P}_\alpha}(S_\alpha)\}_{\alpha \in \Omega}$.

Theorem 6.5. *There exists an isomorphism Φ from $\prod_{\alpha \in \Omega} Q_{\mathcal{P}_\alpha}(S_\alpha)$ onto $Q_{\mathcal{P}}(S)$, under which S is invariant when $\psi_{\mathcal{P}}(S) = \iota$.*

Proof. Let $(\bar{f}_\alpha)_{\alpha \in \Omega} \in \prod_{\alpha \in \Omega} Q_{\mathcal{P}_\alpha}(S_\alpha)$ which we abbreviate as $(\bar{f}_\alpha)$. Let $D_f = \bigcup_{\alpha} D_{f_\alpha} \in \mathcal{P}$ and define $f: D_f \rightarrow S$ by $f(x) = f_\alpha(x)$ where $x \in D_{f_\alpha}$. Then $f \in \text{Hom}_S(D_f, S)$. Define $\Phi: \prod_{\alpha \in \Omega} Q_{\mathcal{P}_\alpha}(S_\alpha) \rightarrow Q_{\mathcal{P}}(S)$ by $\Phi((\bar{f}_\alpha)) = \bar{f}$. It is easily seen that Φ is the desired isomorphism.

The invariance of S under Φ when $\psi_{\mathcal{P}}(S) = \iota$ is shown in the diagram below for an arbitrary element s of S .

$$\begin{array}{ccccccc}
 s & \longleftrightarrow & \bar{\phi}_s & \longleftrightarrow & \hat{\phi}_s & \xrightarrow{\Phi} & \bar{\phi}_s \longleftrightarrow s \\
 \text{in} & & \text{in} & & \text{in} & & \text{in} \\
 S_\alpha \subseteq S & & Q_{\mathcal{P}_\alpha}(S_\alpha) & & \prod_{\alpha \in \Omega} Q_{\mathcal{P}_\alpha}(S_\alpha) & & Q_{\mathcal{P}}(S) \subseteq S
 \end{array}$$

For each $\alpha \in \Omega$, let T_α be a semigroup such that $S_\alpha \subseteq T_\alpha$. A semigroup T containing S as a subsemigroup is called a *subdirect product* of $\{T_\alpha\}_{\alpha \in \Omega}$ over S if there exists an isomorphism θ from T into $\prod_{\alpha \in \Omega} T_\alpha$ such that

(a) $\theta|_S = \iota_S$ (under the identification given at the beginning of this section), and

(b) $\pi_\alpha \theta(T) = T_\alpha$ for each $\alpha \in \Omega$, where π_α is the projection mapping of $\prod_{\alpha \in \Omega} T_\alpha$ onto T_α .

We will now show that subdirect products over S of $\mathcal{P}_\alpha$ -torsion free $\mathcal{P}_\alpha$ -quotient semigroups over S_α , $\alpha \in \Omega$, determine up to isomorphism all $\mathcal{P}$ -torsion free $\mathcal{P}$ -quotient semigroups over S . The following lemma which can easily be shown will be useful in the proof. It is valid for any semigroup S with zero and any right quotient filter $\mathcal{P}$ on S .

Lemma 6.6. *If T is a $\mathcal{P}$ -quotient semigroup over S and if θ is an isomorphism from T onto a semigroup T' containing S such that $\theta|_S = \iota_S$ then T' is a $\mathcal{P}$ -quotient semigroup over S . Furthermore, T' is $\mathcal{P}$ -torsion free iff T is $\mathcal{P}$ -torsion free.*

Theorem 6.7. *Let S be $\mathcal{P}$ -torsion free and let T be a semigroup containing S . Then T is a $\mathcal{P}$ -torsion free $\mathcal{P}$ -quotient semigroup over S iff there exist semigroups T_α , $\alpha \in \Omega$, such that each T_α is a $\mathcal{P}_\alpha$ -torsion free $\mathcal{P}_\alpha$ -quotient semigroup over S_α and T is a subdirect product over S of $\{T_\alpha\}_{\alpha \in \Omega}$.*

Proof. ($\Rightarrow$) By Corollary 5.8 there exists a semigroup T' isomorphic to T such that $S \subseteq T' \subseteq Q_{\mathcal{P}}(S)$ and S is invariant under the given isomorphism. Let $T'' = \Phi^{-1}(T')$ where Φ is given in Theorem 6.5. Hence we have $S \subseteq T'' \subseteq \prod_{\alpha \in \Omega} Q_{\mathcal{P}_\alpha}(S_\alpha)$. For each $\alpha \in \Omega$, let $T_\alpha = \pi_\alpha(T'')$ where π_α is the projection of $\prod_{\alpha \in \Omega} Q_{\mathcal{P}_\alpha}(S_\alpha)$ onto $Q_{\mathcal{P}_\alpha}(S_\alpha)$. Thus we have $S_\alpha \subseteq T_\alpha \subseteq Q_{\mathcal{P}_\alpha}(S_\alpha)$ for each $\alpha \in \Omega$. Since $Q_{\mathcal{P}_\alpha}(S_\alpha)$ is a $\mathcal{P}_\alpha$ -torsion free $\mathcal{P}_\alpha$ -quotient semigroup over S_α , it clearly follows that each T_α is also a $\mathcal{P}_\alpha$ -torsion free $\mathcal{P}_\alpha$ -quotient semigroup over S_α . Furthermore, it is straightforward to see that T is a subdirect product over S of the $\{T_\alpha\}_{\alpha \in \Omega}$.

($\Leftarrow$) By Lemma 6.6, we may assume that $S \subseteq T \subseteq \prod_{\alpha \in \Omega} T_\alpha$ such that $\pi_\alpha(T) = T_\alpha$.

Recall that in this context we are regarding S as a subset of $\prod_{\alpha \in \Omega} S_\alpha \subseteq$

$\prod_{\alpha \in \Omega} T_\alpha$ under the identification $s \leftrightarrow \hat{s}$. Let $t \in T$ be arbitrary. Since T_α is a $\mathcal{P}_\alpha$ -quotient semigroup over S_α for each α , it follows that $t(\alpha)^{-1}S_\alpha \in \mathcal{P}_\alpha$. Let $E = \bigcup_{\alpha \in \Omega} t(\alpha)^{-1}S_\alpha$. Then $E \in \mathcal{P}$ and for each $s \in E \cap S_\alpha$, $\alpha \in \Omega$, we have

$$ts(\beta) = \begin{cases} 0 & \text{if } \beta \neq \alpha, \\ t(\alpha)s \in S_\alpha & \text{if } \beta = \alpha. \end{cases}$$

Hence it follows that $s \in t^{-1}S$. Thus $E \subseteq t^{-1}S$ and $t^{-1}S \in \mathcal{P}$ and we see that T is a $\mathcal{P}$ -quotient semigroup over S . In order to show that T is $\mathcal{P}$ -torsion free, suppose that $t, t' \in T$ and $E \in \mathcal{P}$ such that $tE \neq t'E$. We claim that $t(\alpha) = t'(\alpha)$ for each $\alpha \in \Omega$. Let $E_\alpha = E \cap S_\alpha \in \mathcal{P}_\alpha$. Then for each $s \in E_\alpha$, we have $t(\alpha)s = t(\alpha)\hat{s}(\alpha) = (t\hat{s})(\alpha) = (t'\hat{s})(\alpha) = t'(\alpha)\hat{s}(\alpha) = t'(\alpha)s$. Hence $(t(\alpha), t'(\alpha)) \in \psi_{\mathcal{P}_\alpha}(S_\alpha) = \iota$ and the claim follows. Therefore, $t = t'$ and it follows that T is $\mathcal{P}$ -torsion free.

7. Primitive dependent semigroups. Let S be a semigroup and let $E = E(S)$ be the set of idempotents in S . A partial ordering $\leq$, called the *natural partial ordering on E* , is defined by $e \leq f$ iff $e = fe = ef$. An element $0 \neq e \in E$ is said to be primitive if $0 \leq f \leq e$, $f \in E$, implies that $f = 0$ or $f = e$. The set of primitive idempotents of S will be denoted by $E' = E'(S)$. A regular semigroup S is said to be *primitive regular* if $E' = E \setminus \{0\}$. An important subclass of the class of primitive regular semigroups is the collection of all completely 0-simple semigroups. These semigroups have been characterized as regular Rees matrix semigroups over groups with 0 as shown in Chapter III of Clifford and Preston [4].

A nonzero regular semigroup S will be called *primitive dependent* if for each $0 \neq e \in E = E(S)$ there exists $e' \in E' = E'(S)$ such that $e' \leq e$. This class of semigroups contains the primitive regular semigroups, all finite regular semigroups, and, as will be shown, all completely semisimple semigroups with principal series.

In this section we shall develop several theorems involving the singular congruence on a completely 0-simple semigroup. In addition, a characterization of the completely 0-simple $\mathcal{P}_I$ -torsion free semigroups will be given. These facts along with the general theorems given in §6 will lead to a complete characterization of all primitive dependent $\mathcal{P}_I$ -torsion free semigroups as regular subdirect products of column monomial matrix semigroups over groups.

We begin with a few definitions and lemmas which can be stated for S -sets in general. A nonzero sub S -set N_S of an S -set M_S is said to be 0-minimal if N_S contains no proper nonzero sub S -sets. Such S -sets are characterized as follows.

Lemma 7.1. *Let $0 \neq N_S \subseteq M_S$. Then N_S is 0-minimal iff for each $0 \neq x, y \in N$, there exists $s \in S^1$ such that $y = xs$ (i.e., iff $N = xS^1$ for each $0 \neq x \in N$).*

Proof. ($\Rightarrow$) Suppose N is 0-minimal. Let $0 \neq x \in N$. Then $0 \subsetneq xS^1 \subseteq N$. Hence $xS^1 = N$ since N is 0-minimal.

($\Leftarrow$) Let $0 \subsetneq N' \subseteq N$ and let $n \in N$ and $0 \neq n' \in N'$. Then there exists $s \in S^1$ such that $n = n's \in N'$. Thus $N' = N$ and the result follows.

The *right socle* $\Sigma = \Sigma_r(M)$ of an S -set M is the union of $\{0\}$ and all 0-minimal sub S -sets of M_S . The *left socle* $\Sigma_l(M)$ is defined dually for left S -sets.

Lemma 7.2. $\Sigma_r(M) \subseteq N$ for every $N \in \mathcal{P}_l(M_S)$.

Proof. Let $0 \neq m \in \Sigma_r(M)$. Then $mS^1 \cap N \neq 0$ and since mS^1 is 0-minimal it follows that $m \in mS^1 \subseteq N$. Therefore, $\Sigma_r(M) \subseteq N$.

Lemma 7.3. $M = \Sigma_r(M)$ iff $\mathcal{P}_l(M_S) = \{M\}$.

Proof. ($\Rightarrow$) This is immediate from Lemma 7.2.

($\Leftarrow$) Let $0 \neq m \in M$. We claim that mS^1 is 0-minimal. Let $0 \neq ms_1, ms_2 \in mS^1$. As in the proof of Proposition 4.7, there exists a sub S -set N of M such that $ms_1S^1 \cap N = 0$ and $ms_1S \cup N \in \mathcal{P}_l(M_S)$. Hence, since $\mathcal{P}_l(M_S) = \{M\}$, we have $ms_1S^1 \cup N = M$. If $m \in N$ then $ms_1 \in ms_1S^1 \cap N = \{0\}$ which is a contradiction. Thus, $m \in ms_1S^1$ and it follows that $ms_2 \in ms_1S^1$. Therefore, mS^1 is 0-minimal by Lemma 7.1 and $m \in mS^1 \subseteq \Sigma_r(M)$.

Note that if $\Sigma = \Sigma_r(M) \in \mathcal{P}_l(M_S)$ then, by Lemma 5.2, $Q_l(M_S) \approx \text{Hom}_S(\Sigma, \Sigma)$ under the map $\Phi(f) = \bar{f}$ where $f \in \text{Hom}_S(\Sigma, \Sigma)$. Hereafter, we shall regard this isomorphism as an identity.

The theorem which follows determines all regular semigroups S for which $\Sigma_r(S) = S$. Parts (a), (b), (c), and (d) of the theorem are taken from Theorem 6.39 of Clifford and Preston [5]. The equivalence of (c) and (e) is given by Lemma 7.3.

Theorem 7.4. *The following are equivalent where $S = S^0$ is a semigroup.*

- (a) S is a 0-direct union of completely 0-simple semigroups.
- (b) S is a union of 0-minimal right ideals of the form eS where $e^2 = e \in S$.
- (c) S is regular and $\Sigma_r(S) = S$.
- (d) S is primitive regular.
- (e) S is regular and $\mathcal{P}_l(S) = \{S\}$.

From part (e) of the above theorem we note that the terms "right reductive" and " $\mathcal{P}_l$ -torsion free" are equivalent on a primitive regular semigroup. The next theorem, which describes the socle of any regular semigroup, is essentially a restatement of Theorem 7.59 in Clifford and Preston [5].

Theorem 7.5. *Let S be a regular semigroup with 0. Then $\Sigma_r(S) = \Sigma_l(S)$ and the socle $\Sigma (= \Sigma_r(S) = \Sigma_l(S))$, if nonzero, is the largest primitive regular ideal of S and it contains all the primitive idempotents of S .*

Note that if $E' = E'(S) \neq \square$ then $\Sigma(S) = SE'S = E'S = SE'$ for any regular semigroup S .

The preceding two theorems indicate that completely 0-simple semigroups are some of the basic "building blocks" for all regular semigroups with nonzero socle. Furthermore, the well-known Rees theorem given on p. 94 of [4] describes each completely 0-simple semigroup as a regular Rees matrix semigroup $\mathfrak{M}^0(G, I, J; P)$ over a group with 0. For this reason we begin our study of the singular congruence and the essential semigroup of quotients on a regular semigroup by considering first the case where S is a regular Rees matrix semigroup.

Let $S = \mathfrak{M}^0(G, I, J; P)$ be the regular Rees matrix semigroup over the group G with zero, having indices I and J and $J \times I$ sandwich matrix P . The nonzero elements of S will be written in the form $A = (a)_{ij}$ where $a \in G$ is the single nonzero entry in the $I \times J$ matrix A which appears in the i th row and j th column. The element in the (j, i) position of P will be denoted by p_{ji} . Row j_1 of P is said to be *left proportional* to row j_2 of P if there exists $c \in G$ such that $p_{j_1 i} = cp_{j_2 i}$ for all $i \in I$. Dually, column i_1 is *right proportional* to column i_2 if there exists $c \in G$ such that $p_{j i_1} = p_{j i_2}c$ for all $j \in J$. The next theorem determines the singular congruence on S .

Theorem 7.6. For $S = \mathfrak{M}^0(G, I, J; P)$, $\psi_I(S) = \{[(a)_{i_1 j_1}, (b)_{i_2 j_2}]: \text{row } j_1 \text{ is left proportional to row } j_2, \text{ say } p_{j_1 i} = cp_{j_2 i} \text{ for each } i \in I, \text{ and } a, b \in G \text{ such that } a^{-1}b = c\} \cup \{(0, 0)\}$.

Proof. Since S is regular, $\psi_I(S)$ is 0-restricted by Lemma 4.6. Let $[(a)_{i_1 j_1}, (b)_{i_2 j_2}] \in \psi_I(S)$. Since $\mathcal{P}_I(S) = \{S\}$, we have $(a)_{i_1 j_1}(x)_{ij} = (b)_{i_2 j_2}(x)_{ij}$ for each $(x)_{ij} \in S$. Hence $(ap_{j_1 i}x)_{i_1 j} = (bp_{j_2 i}x)_{i_2 j}$ for all $(x)_{ij} \in S$. Thus, it follows that $i_1 = i_2$ and $ap_{j_1 i} = bp_{j_2 i}$ for each $i \in I$. Therefore, $p_{j_1 i} = a^{-1}bp_{j_2 i} = cp_{j_2 i}$ for each $i \in I$, where $c = a^{-1}b$, and we see that $[(a)_{i_1 j_1}, (b)_{i_1 j_2}]$ is a member of the right side. The opposite inclusion is immediate from retracing the above steps.

Corollary 7.7. If P consists entirely of 0's and 1's then $\psi_I(S) = \{[(x)_{i_1 j_1}, (x)_{i_2 j_2}]: \text{rows } j_1 \text{ and } j_2 \text{ of } P \text{ are equal}\} \cup \{(0, 0)\}$.

Corollary 7.8. $\psi_I(S) = \iota$ iff no two rows of P are left proportional.

Corollary 7.9. If S is a primitive inverse semigroup then $\psi_I(S) = \iota$.

Proof. By Theorem 7.4, S is a 0-direct union of $\{S_\alpha\}_{\alpha \in \Omega}$ where each S_α is completely 0-simple. Since $S_\alpha \subseteq S$, it follows that S_α is a Brandt semigroup for each $\alpha \in \Omega$. Hence by Theorem 3.9 of Clifford and Preston [4], S_α is isomorphic to a regular Rees matrix semigroup $\mathfrak{M}^0(G_\alpha, I_\alpha, I_\alpha; \Delta_\alpha)$ over a group G_α with 0 and with the $I_\alpha \times I_\alpha$ identity matrix Δ_α as sandwich matrix. Thus, from Corollary 7.8 we see that $\psi_I(S_\alpha) = \iota$ for each $\alpha \in \Omega$. Therefore, the result follows by Lemma 6.2 and Corollary 6.4.

Gordon L. Bailes [2] has defined a *right inverse semigroup* as a regular semigroup S in which each element has a unique left unit (i.e., $|xV(x)| = 1$ where $V(x)$ is the set of inverses of x in S). A left inverse semigroup is defined dually. Corollary 4 of [2] shows that S is an inverse semigroup iff S is both a left and a right inverse semigroup.

Theorem 7.10. *Let S be a completely 0-simple semigroup. Then the following are equivalent.*

(a) S is right inverse.

(b) S is isomorphic to a regular Rees matrix semigroup $\mathfrak{M}^0(G, I, J; P)$ where P consists entirely of 0's and 1's and each column contains exactly one nonzero entry.

(c) $\psi_I(S) = \iota$ and S is orthodox.

Proof. (a) $\Leftrightarrow$ (b) is essentially a restatement of Theorem 44 in [2] and (b) $\Leftrightarrow$ (c) is immediate from Corollary 7.8 and Theorem 6 of [7].

Corollary 7.11. *A primitive regular semigroup is a right inverse semigroup iff $\psi_I(S) = \iota$ and S is orthodox.*

Proof. By Theorem 7.4, S is a 0-direct union of completely 0-simple semigroups $\{S_\alpha\}_{\alpha \in \Omega}$. Clearly, by the definition S is a right inverse semigroup iff each S_α is a right inverse semigroup. By the theorem this is true iff each S_α is orthodox and $\psi_I(S_\alpha) = \iota$. Finally, by Corollary 6.4, this follows iff S is orthodox and $\psi_I(S) = \iota$.

Corollary 7.12. *An orthodox primitive regular semigroup is inverse iff $\psi_I(S) = \iota$ and ${}_I\psi(S) = \iota$, where ${}_I\psi(S)$ is the left-right dual of $\psi_I(S)$.*

Let $S = \mathfrak{M}^0(G, I, J; P)$ be a regular Rees matrix semigroup and let $\mathcal{C}^0(G, I)$ denote the semigroup of all $I \times I$ column monomial matrices over G^0 . If $A \in \mathcal{C}^0(G, I)$ then $A(i, j)$ will denote the element in the (i, j) position of A . Multiplication in S will be indicated by juxtaposition whereas ordinary multiplication between matrices as defined on p. 87 of Clifford and Preston [4] will be denoted by " $\circ$ ". Hence, if $A, B \in S$ then $AB = A \circ P \circ B$. Since $S = \Sigma(S)$ in this case, $\mathcal{P}_I(S) = \{S\}$ and we have $\mathcal{Q}_I(S) = \text{Hom}_S(S, S)$ which is precisely the semigroup $\Lambda(S)$ of all left translations of S . This has been characterized by Petrich [13] as a wreath product over a group. Another characterization of $\Lambda(S)$ is also given in Proposition 4.5.6 of [13] which is restated below without proof.

Theorem 7.13. *Let $S = \mathfrak{M}^0(G, I, J; P)$ be a regular Rees matrix semigroup. Then there exists an isomorphism $\Phi: \text{Hom}_S(S, S) \xrightarrow{\text{onto}} \mathcal{C}^0(G, I)$, say $\Phi(f) = C_f$, such that $f(X) = C_f \circ X$ for each $f \in \text{Hom}_S(S, S)$ and for each $X \in S$.*

Let ϕ denote the representation of S in $Q_I(S) = \text{Hom}_S(S, S)$ given in §5. Then $\Phi\phi$ is a representation of S in $\mathcal{C}^0(G, I)$. The image of this representation is given by the following proposition.

Proposition 7.14. *Let $\Phi\phi$ be the representation of S in $\mathcal{C}^0(G, I)$ given above. For each $0 \neq X = (x)_{ij} \in S$ let C_X denote the element of $\mathcal{C}^0(G, I)$ given by*

$$C_X = (i', k) = \begin{cases} 0 & \text{if } i \neq i', k \in I, \\ xp_{jk} & \text{if } i = i', k \in I. \end{cases}$$

Then $\Phi\phi(X) = C_X$.

Proof. Let $f \in Q_I(S)$ such that $\Phi(f) = C_X$. We claim that $f = \phi_X$. Let $Y = (y)_{i'j'}$ be an arbitrary element of S . Then

$$\begin{aligned} f(Y) &= C_f \circ Y = C_X \circ Y = \begin{pmatrix} 0 & & 0 \\ & xp_{ji'} & \\ 0 & & 0 \end{pmatrix} \circ \begin{pmatrix} 0 & 0 \\ 0 \cdots 0 & y0 \cdots 0 \\ 0 & 0 \end{pmatrix} \\ &= (xp_{ji'}y)_{ij'} = XY = \phi_X(Y). \end{aligned}$$

Therefore, $f = \phi_X$ and the proposition follows.

Note that if $S = \mathfrak{M}^0(G, I, I; \Delta)$ where Δ is the $I \times I$ identity matrix then $\Phi\phi(X) = X$ for each $X \in S$, i.e., the representation of S is actually the identity map.

The above proposition leads us to the following definition.

Let G be a group, I and J be index sets, and let P be a regular $J \times I$ matrix over G^0 such that no two rows of P are left proportional. Let $\mathcal{H}^0 = \mathcal{H}^0(G, I, J; P)$ denote the subset of all elements "A" in $\mathcal{C}^0(G, I)$ such that for some $i \in I, j \in J$, and $g \in G^0$,

$$A(i', k) = \begin{cases} 0 & \text{if } i' \neq i, k \in I, \\ gp_{jk} & \text{if } i' = i, k \in I. \end{cases}$$

If S is the semigroup $\mathfrak{M}^0(G, I, J; P)$ then S is $\mathcal{P}_I$ -torsion free since no two rows of P are left proportional. Hence $\mathcal{H}^0$ is the isomorphic image of S under the map $\Phi\phi$ into the semigroup $\mathcal{C}^0(G, I)$. Thus since $\phi(S)$ is a left ideal of $Q_I(S) = \text{Hom}_S(S, S)$, we see that $\mathcal{H}^0$ is a left ideal of $\mathcal{C}^0(G, I)$. Furthermore, a trace of the isomorphisms involved shows that $\mathcal{C}^0(G, I)$ is isomorphic over $\mathcal{H}^0$ to $Q_I(\mathcal{H}^0)$. The semigroup $\mathcal{H}^0$ will be called an $\mathcal{H}$ -semigroup. The next theorem follows from the discussion above and §5.

Theorem 7.15. *Let $\mathcal{H}^0 = \mathcal{H}^0(G, I, J; P)$ be an arbitrary $\mathcal{H}$ -semigroup. Then*

- (a) A semigroup S is completely 0-simple right reductive iff S is isomorphic to some $\mathcal{H}$ -semigroup.
- (b) $\mathcal{H}^0$ is a left ideal of $\mathcal{C}^0(G, I)$.
- (c) $\mathcal{C}^0(G, I)$ is isomorphic over $\mathcal{H}^0$ to $Q_I(\mathcal{H}^0)$.
- (d) $\mathcal{C}^0(G, I)$ is the injective hull of $\mathcal{H}^0$.
- (e) $\mathcal{C}^0(G, I)$ is self-injective.

These results can also be applied to primitive regular semigroups as stated below. The proofs are immediate from the general theory developed in §6 since each primitive regular semigroup is a 0-direct union of completely 0-simple semigroups.

Theorem 7.16. *Let S be a primitive regular semigroup. Then*

- (a) S is primitive regular right reductive iff S is isomorphic to a 0-direct union of $\mathcal{H}$ -semigroups.
- (b) $Q_I(S) = \text{Hom}_S(S, S)$ and is isomorphic to a direct product of column monomial matrix semigroups over groups.

We shall now consider primitive dependent semigroups. Certainly any primitive regular semigroup is primitive dependent, as well as any finite regular semigroup. A more general class of primitive dependent semigroups is given in the theorem which follows. The reader is referred to §§2.6 and 6.6 of Clifford and Preston [4] and [5] for the definitions of the terms used.

Theorem 7.17. *Let $S = S^0$ be a completely semisimple semigroup with a principal series. Then S is a primitive dependent semigroup.*

Proof. Let $S = S_1 \supset S_2 \supset \dots \supset S_n = \{0\}$ be a principal series for S . We shall proceed by induction on n . If $n = 2$, then $S = S_1$ is completely 0-simple and the result follows. Suppose the theorem is true for all completely semisimple semigroups having a principal series of length k where $2 \leq k < n$, $n > 2$, and let S have a principal series of length n as above. Then S_2 has a principal series of length $n - 1$. Let $0 \neq e = e^2 \in S = S_1$. If $e \in S_2$ then by the induction hypothesis there exists an idempotent $f' \in S_2$ such that $f' \leq e$ and such that f' is primitive in S_2 . Since S_2 is an ideal of S , it follows that f' is also primitive in S_1 . Hence the result is true in this case. Now suppose $e \in S \setminus S_2$. If e is primitive in S , we are through. So, suppose e is not primitive in S . Then there exists $0 \neq f \in E(S)$ such that $0 \neq f < e$. If $f \in S \setminus S_2$ then $0 \neq f < e$ in $S/S_2 = S_1/S_2$ which is completely 0-simple. But this is impossible. Hence $f \in S_2$ and by the induction hypothesis there exists f' primitive in S_2 (hence in S) such that $f' \leq f$. Therefore $f' < e$ and the theorem is proved.

Let R be a subset of a semigroup S . The left annihilator of R in S is ${}_R A =$

$\{x \in S: xR = 0\}$. The right annihilator of R is defined dually and is denoted by A_R . Several equivalences to the definition of primitive dependent can now be stated.

Theorem 7.18. *Let S be a regular semigroup and let Σ be the socle of S . Then the following are equivalent.*

- (a) $\Sigma A = 0$.
- (b) $\Sigma \in \mathcal{P}_I(S_\Sigma)$.
- (c) $\Sigma \in \mathcal{P}_I(S)$.
- (d) Every nonzero right ideal of S contains a 0-minimal right ideal of S .
- (e) S is primitive dependent.

Proof. (a) $\Rightarrow$ (b). Let $0 \neq s \in S$. Since $\Sigma A = 0$, there exists $x \in \Sigma$ such that $sx \neq 0$ and since Σ is a 2-sided ideal of S , we have that $0 \neq sx \in \Sigma$. Thus it follows that $\Sigma \in \mathcal{P}_I(S_\Sigma)$.

(b) $\Rightarrow$ (c). This is immediate from Theorem 3.7.

(c) $\Rightarrow$ (d). Let R be a nonzero right ideal of S . Since $\Sigma \in \mathcal{P}_I(S)$, there exists $0 \neq x \in R \cap \Sigma$, and since $x \in \Sigma$, xS^1 is 0-minimal. Furthermore, $xS^1 \subseteq R$.

(d) $\Rightarrow$ (e). Let $0 \neq e \in E$. By the hypothesis eS contains a 0-minimal right ideal R of S . Since S is regular, R is of the form $R = fS$ for some $0 \neq f \in E$, and since $f \in \Sigma$, we have that $f \in E'$. Furthermore, $ef = f$. Let $f' = fe$. Then $f' \in E$ since $f'^2 = (fe)(fe) = f(ef)e = ffe = fe = f'$, and $f'f = fef = ff = f \neq 0$. Hence $f' \neq 0$. Furthermore, $f' = fe \in fS \subseteq \Sigma$. Thus it follows that $f' \in E'$. Finally, $f'e = fee = fe = f' = fe = efe = ef'$. Therefore, $0 \neq f' \leq e$ and $f' \in E'$ and it follows that S is primitive dependent.

(d) $\Rightarrow$ (a). Let $0 \neq x \in S$ and let x' be an inverse of x . Then there exists $f \in E'$ such that $f \leq x'x$. Hence $x'xf = f \in E' \subseteq \Sigma$. Therefore, $0 \neq xf \in x\Sigma$ and it follows that $\Sigma A = 0$.

Note that the left-right duals of (a), (b), (c) and (d) respectively are equivalent to (e) also.

The essential semigroup of quotients of a primitive dependent semigroup is given by the following lemma.

Lemma 7.19. *Let S be primitive dependent and let Σ be its socle. Then $Q_I(S)$ is semigroup isomorphic to $\text{Hom}_\Sigma(\Sigma, \Sigma) = Q_I(\Sigma)$.*

Proof. From a note following Lemma 7.3 and from the above theorem we see that $Q_I(S) \approx \text{Hom}_S(\Sigma, \Sigma)$. Hence it is sufficient to show that every Σ -homomorphism on Σ is an S -homomorphism. Let $f \in \text{Hom}_\Sigma(\Sigma, \Sigma)$, $t \in \Sigma$, and $s \in S$. Then for t' , an inverse of t in Σ , we have $f(ts) = f(tt'ts) = f(t)t'ts = f(tt't)s = f(t)s$ and the result follows.

Another characterization of the singular congruence can be given for a primitive dependent semigroup.

Lemma 7.20. *If S is primitive dependent then $\psi_I(S) = \{(x, y): xe = ye \text{ for all } e \in E'\}$.*

Proof. By Lemma 7.2 and Theorem 7.18, $\{\Sigma\}$ is a base for $\mathcal{P}_I(S)$. Thus by Lemma 4.1, $\psi_I(S) = \{(x, y): xs = ys \text{ for all } s \in \Sigma\}$. Since $E' \subseteq \Sigma$, it follows that $\psi_I(S)$ is contained in the right side above. Now, let (x, y) be in the right side and let $s \in \Sigma$. Then there exists $0 \neq e = e^2 \in \Sigma$ such that $s = es$ and, since $e \in \Sigma$, $e \in E'$. Therefore $xs = xes = yes = ys$ and $(x, y) \in \psi_I(S)$.

Lemma 7.21. *Let $S \subseteq T$ be regular semigroups such that S is a left (right, 2-sided) ideal of T . Then $\Sigma(S) \subseteq \Sigma(T)$.*

Proof. Let $e \in E'(S)$. We claim that $e \in E'(T)$. Suppose there exists $f^2 = f \in T$ such that $f \leq e$. Then $f = fe = ef \in S$ in any case above. Hence, since e is primitive in S , it follows that $f = 0$ or $f = e$. Therefore, $e \in E'(T)$ and we see that $\Sigma(S) = E'(S)S \subseteq E'(T)T = \Sigma(T)$ by the note following Theorem 7.5.

The next theorem shows that certain regular $\mathcal{P}_I$ -quotient semigroups over a primitive dependent semigroup are also primitive dependent.

Theorem 7.22. *Let S be a primitive dependent semigroup and let T be any regular $\mathcal{P}_I(S)$ -quotient semigroup over S such that S is a left (right, 2-sided) ideal of T and $\psi_I(T_S)$ is 0-restricted. Then T is primitive dependent with socle $\Sigma(T) = \Sigma(S)T^1$.*

Proof. By Lemma 7.21, $\Sigma(S) \subseteq \Sigma(T)$. Hence $\Sigma(S)T^1 \subseteq \Sigma(T)$. We claim that $S \in \mathcal{P}_I(T_S)$. Let $0 \neq t \in T$. Then since T is a $\mathcal{P}_I$ -quotient semigroup over S , $t^{-1}S \in \mathcal{P}_I(S)$. Furthermore, $t(t^{-1}S) \neq 0$ since $\psi_I(T_S)$ is 0-restricted and the claim follows. Thus, since $\Sigma(S) \in \mathcal{P}_I(S)$, we see from Lemma 4.5 and Theorem 3.7(a) that $\Sigma(S)T^1 \in \mathcal{P}_I(T)$. So, $\Sigma(T) \subseteq \Sigma(S)T^1$ by Lemma 7.2 and equality follows. Furthermore, since $\Sigma(T) = \Sigma(S)T^1 \in \mathcal{P}_I(T)$ we see from Theorem 7.18 that T is primitive dependent.

Corollary 7.23. *Let S be a primitive dependent $\mathcal{P}_I$ -torsion free semigroup. Then $Q_I(S)$ is primitive dependent.*

Proof. Since $Q_I(\Sigma(S)) \approx Q_I(S)$ by Lemma 7.19, and $\Sigma(S)$ is primitive dependent, we may assume without loss of generality that $S = \Sigma(S)$. Hence S is embedded in $Q = Q_I(S)$ and Q is a $\mathcal{P}_I$ -torsion free $\mathcal{P}_I$ -quotient semigroup over S . Furthermore, since $Q = \text{Hom}_S(S, S)$ we have that $fs = f(s)$ for each $f \in Q$, $s \in S$. Therefore, S is a left ideal of Q . Furthermore, by Theorem 4.2 of [8], Q is regular. Hence the result follows from Theorem 7.22.

Our final theorem gives a complete characterization of all $\mathcal{P}_I$ -torsion free primitive dependent semigroups.

Theorem 7.24. *Let Ω be an index set and for each $\alpha \in \Omega$, let $\mathcal{H}_\alpha^0 = \mathcal{H}^0(G_\alpha, I_\alpha, J_\alpha; P_\alpha)$ be an $\mathcal{H}$ -semigroup. Let S be the 0-direct union of $\{\mathcal{H}_\alpha^0\}_{\alpha \in \Omega}$. Let T_α be any regular semigroup extension of $\mathcal{H}_\alpha^0$ in $\mathcal{C}_\alpha = \mathcal{C}_\alpha^0(G_\alpha, I_\alpha)$, $\alpha \in \Omega$, and let T be any regular subdirect product of the $\{T_\alpha\}_{\alpha \in \Omega}$ over S . Then T is a primitive dependent $\mathcal{P}_I$ -torsion free semigroup. Conversely, every primitive dependent $\mathcal{P}_I$ -torsion free semigroup is isomorphic to a semigroup obtained in this manner.*

Proof. Since $\mathcal{C}_\alpha \approx \mathcal{Q}_I(\mathcal{H}_\alpha^0)$ over $\mathcal{H}_\alpha^0$ for each $\alpha \in \Omega$, we have that $\mathcal{C}_\alpha$ is a $\mathcal{P}_I(\mathcal{H}_\alpha^0)$ -quotient semigroup over $\mathcal{H}_\alpha^0$ which is $\mathcal{P}_I(\mathcal{H}_\alpha^0)$ -torsion free. Thus T_α is a $\mathcal{P}_I(\mathcal{H}_\alpha^0)$ -torsion free $\mathcal{P}_I(\mathcal{H}_\alpha^0)$ -quotient semigroup for each $\alpha \in \Omega$. Hence by Theorem 6.7, T is a $\mathcal{P}_I(S)$ -torsion free $\mathcal{P}_I(S)$ -quotient semigroup over S and by Lemma 4.12 we see that $S \in \mathcal{P}_I(T_S)$. Thus, T is also $\mathcal{P}_I(T)$ -torsion free by Lemma 4.5 and Theorem 4.8. Since $\mathcal{H}_\alpha^0$ is a left ideal of $\mathcal{C}_\alpha$ for each $\alpha \in \Omega$, we have $\mathcal{H}_\alpha^0$ is a left ideal of T_α . Let $t \in T$ and $s \in \mathcal{H}_\alpha^0 \subseteq S$. Then

$$ts(\beta) = \begin{cases} 0 & \text{if } \beta \neq \alpha, \\ t(\alpha)s \in \mathcal{H}_\alpha^0 & \text{if } \beta = \alpha. \end{cases}$$

Hence $t\hat{s} = t(\alpha)s$ and it follows that $S \approx \hat{S}$ is a left ideal of T . Therefore, the result follows by Theorem 7.22.

Conversely, let T be a $\mathcal{P}_I$ -torsion free primitive dependent semigroup and let $S = \Sigma(T)$. Since $S \in \mathcal{P}_I(T)$ by Theorem 7.18(c), S is $\mathcal{P}_I(S)$ -torsion free. Hence by Theorem 7.16 we may assume that S is a 0-direct union of $\mathcal{H}$ -semigroups $\mathcal{H}_\alpha^0 = \mathcal{H}_\alpha^0(G_\alpha, I_\alpha, J_\alpha; P_\alpha)$, $\alpha \in \Omega$. By Theorem 7.18(b) and Proposition 3.6, T is a $\mathcal{P}_I(S)$ -quotient semigroup over S which is $\mathcal{P}_I(S)$ -torsion free. Thus, by Theorem 6.7, there exist semigroups T_α , $\alpha \in \Omega$, such that each T_α is a $\mathcal{P}_I(\mathcal{H}_\alpha^0)$ -torsion free $\mathcal{P}_I(\mathcal{H}_\alpha^0)$ -quotient semigroup over $\mathcal{H}_\alpha^0$ and T is a subdirect product of the $\{T_\alpha\}_{\alpha \in \Omega}$ over S . Since $\mathcal{C}_\alpha = \mathcal{C}_\alpha^0(G_\alpha, I_\alpha)$ is isomorphic over $\mathcal{H}_\alpha^0$ to $\mathcal{Q}_I(\mathcal{H}_\alpha^0)$, we may choose the T_α such that $\mathcal{H}_\alpha^0 \subseteq T_\alpha \subseteq \mathcal{C}_\alpha^0(G_\alpha, I_\alpha)$ by Corollary 5.8. Furthermore, each T_α is regular since it is the homomorphic image under the projection map of the regular semigroup T . Therefore, the theorem follows.

This paper constitutes part of the author's doctoral dissertation written under the direction of Professor D. R. LaTorre. The author wishes to thank Professor LaTorre and Professor John K. Luedeman for their valuable advice and encouragement.

Research for this work was supported by a National Science Foundation Traineeship and by a Faculty Fellowship from the Virginia Military Institute.

REFERENCES

1. K. Asano, *Über die Quotientenbildung von Schieftringen*, J. Math. Soc. Japan 1 (1949), 73–78. MR 11, 154.

2. Gordon L. Bailes, *Right inverse semigroups*, Thesis, Clemson University, Clemson, S.C., 1972.
3. P. Berthiaume, *The injective envelope of S -sets*, *Canad. Math. Bull.* **10** (1967), 261–273. MR 35 #4321.
4. A. H. Clifford and G. B. Preston, *The algebraic theory of semigroups*, 2nd ed., Vol. I, *Math. Surveys*, no. 17, Amer. Math. Soc., Providence, R.I., 1961. MR 24 #A2627.
5. ———, *The algebraic theory of semigroups*. Vol. II, *Math. Surveys*, no. 17, Amer. Math. Soc., Providence, R.I., 1967. MR 36 #1558.
6. E. H. Feller and R. L. Gantos, *Indecomposable and injective S -systems with zero*, *Math. Nachr.* **41** (1969), 37–48. MR 40 #7374
7. T. E. Hall, *On regular semigroups whose idempotents form a subsemigroup*, *Bull. Austral. Math. Soc.* **1** (1969), 195–208. MR 40 #2772.
8. C. V. Hinkle, Jr., *The extended centralizer of an S -set* (submitted).
9. R. E. Johnson, *The extended centralizer of a ring over a module*, *Proc. Amer. Math. Soc.* **2** (1951), 891–895. MR 13, 618.
10. F. R. McMorris, *On quotient semigroups* (submitted).
11. ———, *Vital injective S -systems*, *Math. Nachr.* **47** (1970), 121–125. MR 44 #342.
12. ———, *The singular congruence and the maximal quotient semigroup*, *Canad. Math. Bull.* (to appear).
13. Mario Petrich, *Topics in semigroups*, Pennsylvania State University, 1967.
14. David A. Smith, *On semigroups, semirings, and rings of quotients*, *J. Sci. Hiroshima Univ. Ser. A-I Math.* **30** (1966), 123–130. MR 34 #7692.
15. Yuzo Utumi, *On quotient rings*, *Osaka Math. J.* **8** (1956), 1–18. MR 18, 7.

DEPARTMENT OF MATHEMATICS, CLEMSON UNIVERSITY, CLEMSON, SOUTH CAROLINA 29631

DEPARTMENT OF MATHEMATICS, VIRGINIA MILITARY INSTITUTE, LEXINGTON, VIRGINIA 24450 (Current address)