ON THE ISOTROPIC GROUP OF A HOMOGENEOUS POLYNOMIAL

BY

SIU-MING HO⁽¹⁾

ABSTRACT. Let G be the linear group leaving a homogeneous polynomial of degree k fixed. The author shows that either the polynomial is a polynomial in fewer than the assigned number of variables or that the $(k-1)$ st prolongation of G is 0. The author also shows that this result is optimal.

Let $P^k[X_1, \dots, X_n]$ be the space of homogeneous polynomials in $X_1, \dots, X_n$ of degree k . Note that the linear group $GL(n)$ acts on $P^k[X_1, \dots, X_n]$. If $A = (a_i^j) \in GL(n)$ and $f(X_1, \dots, X_n) \in P^k[X_1, \dots, X_n]$, we denote the action of A on f by Af where

$$(1) \quad Af(X_1, \dots, X_n) = f\left(\sum_{j=1}^n a_1^j X_j, \dots, \sum_{j=1}^n a_n^j X_j\right).$$

The Lie algebra $\mathfrak{gl}(n)$ of $GL(n)$ is the set of $n \times n$ matrices which also acts on $P^k[X_1, \dots, X_n]$. If $a = (a_i^j) \in \mathfrak{gl}(n)$, $f \in P^k[X_1, \dots, X_n]$ then

$$(2) \quad a \cdot f(X_1, \dots, X_n) = \sum_{i,j} \frac{\partial f}{\partial X_i} a_i^j X_j.$$

This is obtained by writing $A = \exp(ta)$ and differentiating (1) with respect to t .

Let G be the Lie subgroup of $GL(n)$ which leaves $f \in P^k[X_1, \dots, X_n]$ invariant, and $\mathfrak{g}$ the Lie algebra of G . By formula (2) we have the following.

Lemma. The Lie algebra $\mathfrak{g}$ of G consists of those matrices (a_i^j) for which $\sum_{i,j} \frac{\partial f}{\partial X_i} a_i^j X_j = 0$.

For each element $a = (a_i^j) \in \mathfrak{gl}(n)$ we can think of $(A^1, \dots, A^n)$ as an n -tuple of homogeneous polynomials of first order, where $A^i = \sum_{j=1}^n a_i^j X_j$. From now on we identify $\mathfrak{gl}(n)$ with the space of n -tuple of homogeneous polynomials of first order. So we can also think of the subspace $\mathfrak{g}$ of $\mathfrak{gl}(n)$ as a subspace of the above space.

Received by the editors November 3, 1972.

AMS (MOS) subject classifications (1970). Primary 53C10.

Key words and phrases. G -structure, prolongation.

⁽¹⁾ This research was partially supported by a grant from NSF, P22927X1.

For any Lie subalgebra g (see [3]) of $gl(n)$ we define the *first prolongation* $g^{(1)}$ of g as the set of all n -tuples $(A^1(X), \dots, A^n(X))$, where A^i is a homogeneous polynomial of degree two in $X_1, \dots, X_n$ and all the first derivatives, $i = 1, \dots, n$, $(\partial A^1/\partial X_i, \dots, \partial A^n/\partial X_i)$ lie in g . Similarly we define the k th prolongation $g^{(k)}$ of g as the set of all n -tuples $(A^1(X), \dots, A^n(X))$ where $A^i(X)$ is a homogeneous polynomial of degree $k+1$ in $X_1, \dots, X_n$, and all the k th derivatives $(\partial^k A^1/\partial X^\alpha, \dots, \partial^k A^n/\partial X^\alpha)$, $|\alpha| = k$, lie in g .

Definition. The algebra g is of finite type if $g^{(k)} = 0$ for some k (and, therefore for all large k) G is said to be of type k if $g^{(k-1)} \neq 0$ and $g^{(k)} = 0$.

Definition. $f(X)$ is a nonsingular homogeneous polynomial if f cannot be written as a polynomial in fewer than n variables.

We have the following result:

Theorem. If $f \in P^k[X_1, \dots, X_n]$ and nonsingular then $g^{(k-1)} = 0$.

Proof. If $(A^1(X), \dots, A^n(X)) \in G^{(k-1)}$, by definition

$$(\partial^{k-1} A^1/\partial X^\alpha, \dots, \partial^{k-1} A^n/\partial X^\alpha), \quad |\alpha| = k-1, \in g.$$

then

$$\sum_{i=1}^n \frac{\partial f}{\partial X_i} \frac{\partial^{k-1}}{\partial X^\alpha} A^i = 0 \quad \text{for all } |\alpha| = k-1.$$

By repeated application of Euler's identity for homogeneous functions we can rewrite this as

$$\sum_{i=1}^n \frac{\partial^{k-1}}{\partial X^\alpha} \left(\frac{\partial f}{\partial X_i} \right) A^i = 0 \quad \text{for all } |\alpha| = k-1.$$

However $\partial^{k-1}/\partial X^\alpha (\partial f/\partial X_i)$ are constants, so choosing any X_0 such that the $A^i(X_0)$ are not all zero, and setting $b_i = A^i(X_0)$ we get

$$\sum \frac{\partial^{k-1}}{\partial X^\alpha} \left(\frac{\partial f}{\partial X_i} \right) b_i = 0 \quad \text{for all } |\alpha| = k-1.$$

This implies $\sum b_i \partial f/\partial X_i = 0$, in other words f can be written as a polynomial in $n-1$ variables which contradicts the hypotheses. Q.E.D.

In what follows we construct some examples which show that the above theorem is the best possible result we can get. Let

$$f(X_1, X_2, X_3, X_4, X_5) = \begin{vmatrix} X_3 & X_4 & X_5 \\ X_1 & X_2 & 0 \\ 0 & X_1 & X_2 \end{vmatrix} = X_3X_2^2 - X_4X_1X_2 + X_5X_1^2.$$

It is easily seen that f cannot be written as a polynomial fewer than 5 variables.

Let $(A^1, A^2, A^3, A^4, A^5)$ be a 5-tuple of homogeneous polynomials of second degree where $A^1 = 0$, $A^2 = 0$, $A^3 = X_1^2$, $A^4 = 2X_1X_2$, $A^5 = X_2^2$. Then

$$\sum_{i=1}^n \frac{\partial f}{\partial X_i} \frac{\partial A^i}{\partial X_j} = 0 \quad \text{for all } j = 1, 2, 3, 4, 5.$$

This implies $(A^1, A^2, A^3, A^4, A^5) \in g^{(1)}$, so $g^{(1)}$ is nontrivial.

Let $f \in P^{k-2}[X_1, X_2, \dots, X_k]$ be defined as follows:

$$\begin{aligned} A^1 &= 0, & A^4 &= X_1^{k-4}X_2, \\ A^2 &= 0, & A^i &= X_1^{k-i}X_2^{i-3}, \\ A^3 &= X_1^{k-3}, & A^k &= X_2^{k-3}; \end{aligned}$$

$$f(X_1, X_2, \dots, X_k) = \begin{vmatrix} \frac{\partial^{k-4}A^3}{\partial X_1^{k-4}} & \frac{\partial^{k-4}A^4}{\partial X_1^{k-4}} & \dots & \frac{\partial^{k-4}A^k}{\partial X_1^{k-4}} \\ \frac{\partial^{k-4}A^3}{\partial X_1^{k-5}\partial X_2} & \frac{\partial^{k-4}A^4}{\partial X_1^{k-5}\partial X_2} & \dots & \frac{\partial^{k-4}A^k}{\partial X_1^{k-5}\partial X_2} \\ \frac{\partial^{k-4}A^3}{\partial X_2^{k-4}} & \frac{\partial^{k-4}A^4}{\partial X_2^{k-4}} & \dots & \frac{\partial^{k-4}A^k}{\partial X_2^{k-4}} \end{vmatrix}$$

It is easily seen that f is nonsingular and also

$$\sum_{i=1}^n \frac{\partial f}{\partial X_i} \frac{\partial^{k-4}A^i}{\partial X^\alpha} = 0 \quad \text{for all } |\alpha| = k-4$$

so we have g^{k-4} of f is nontrivial.

Note that the first example of $g^{(1)} \neq 0$ occurs in dimension 5. Using a result of Gordan-Noether (see [1]) one can show that, for dimensions ≤ 4 , $g^{(1)} = 0$ and, for dimension 5, $g^{(2)} = 0$.

The result above has interesting consequences for the problem of when a partial differential operator can be transformed to a constant coefficient operator see [2].

BIBLIOGRAPHY

1. P. Gordan and M. Noether, *Ueber die algebraischen Formen deren Hesse'sche Determinante identisch verschwindet*, Math. Ann. 10 (1876), 547–568.
2. V. Guillemin and I. M. Singer, *Differential equations and G-structures*, U. S.-Japan Seminar in Differential Geometry (Kyoto, 1965), Nippon Hyoronsha, Tokyo, 1966, pp. 34–36. MR 35 #4840.
3. I. M. Singer and S. Sternberg, *On the infinite groups of Lie and Cartan. I, The transitive groups*, J. Analyse Math. 15 (1965), 1–114. MR 36 #911.

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MASSACHUSETTS 02139