λ CONNECTED PLANE CONTINUA

BY

CHARLES L. HAGOPIAN⁽¹⁾

ABSTRACT. A continuum M is said to be λ connected if any two distinct points of M can be joined by a hereditarily decomposable continuum in M . Recently this generalization of arcwise connectivity has been related to fixed point problems in the plane. In particular, it is known that every λ connected nonseparating plane continuum has the fixed point property. The importance of arcwise connectivity is, to a considerable extent, due to the fact that it is a continuous invariant. To show that λ connectivity has a similar feature is the primary purpose of this paper. Here it is proved that if M is a λ connected continuum and f is a continuous function of M into the plane, then $f(M)$ is λ connected. It is also proved that every semiaposyndetic plane continuum is λ connected.

Introduction. A nondegenerate metric space that is both compact and connected is called a *continuum*. It is known that every plane continuum that has a hereditarily decomposable boundary and does not separate the plane has the fixed point property [1]. Recently the author [4] proved that every arcwise connected nonseparating plane continuum has a hereditarily decomposable boundary. Hence all arcwise connected nonseparating plane continua have the fixed point property. In [7] it is pointed out that the author's theorem remains true if the word "arcwise" is replaced by " λ ". In fact, in [7] it is proved that a plane continuum that does not have infinitely many complementary domains is λ connected if and only if its boundary does not contain an indecomposable continuum.

This paper is primarily concerned with the following questions:

- (1) What other theorems about arcwise connected continua also hold for λ connected continua?
- (2) Are there general properties, other than arcwise connectivity for plane continua, that imply λ connectivity?

Received by the editors September 13, 1971 and, in revised form, September 6, 1973.
AMS (MOS) subject classifications (1970). Primary 54C05, 54F25, 54F60, 57A05;
Secondary 54F15.

Key words and phrases. Arcwise connectivity, λ connected continua, aposyndesis, semiaposyndesis, planar continuous images of hereditarily decomposable continua.

(¹) The author gratefully acknowledges conversations about these results with D. P. Bellamy, F. B. Fuller, E. E. Grace, F. B. Jones, and J. S. Rosasco.

Copyright © 1974, American Mathematical Society

Here, in response to the first question, we prove that every planar continuous image of a λ connected continuum is λ connected.

To deal with the second question, we must consider general properties that do not imply arcwise connectivity. Examples have been given [13, Example 4], [10, Example 6], and [3, Example 1], which indicate that, for plane continua, arcwise connectivity is not a consequence of the aposyndetic property defined by F. Burton Jones. Jones' property can be generalized as follows.

Definition. A continuum M is said to be *semiaposyndetic* if for each pair of distinct points x and y in M there exists a subcontinuum F of M such that the sets $M - F$ and the interior of F relative to M each contain a point of $\{x, y\}$.

Although not all semiaposyndetic plane continua are arcwise connected, these properties are related. If X is a semiaposyndetic plane continuum and for any positive real number ϵ there are at most a finite number of complementary domains of X of diameter greater than ϵ , then X is arcwise connected [6]. For another arc theorem involving semiaposyndesis see [5]. In the last section of this paper it is proved that all semiaposyndetic plane continua are λ connected.

Throughout this paper S^2 is a 2-sphere. The closure and the boundary of a given set Z are denoted by $\text{Cl } Z$ and $\text{Bd } Z$ respectively. The union of the elements of Z is denoted by $\text{St } Z$.

Definitions. Let X be a continuum in S^2 . A continuum L in X is said to be a *link* in X if L is either the boundary of a complementary domain of X or the limit of a convergent sequence of complementary domains of X . A continuum T in X is said to be a *2-link* in X if T is the union of two (not necessarily distinct) links in X . An indecomposable subcontinuum I of X is said to be *terminal* in X if there exists a composant C of I such that each subcontinuum of X that meets both C and $X - I$ contains I .

Preliminary results.

Theorem 1. *If X is a continuum in S^2 and I is an indecomposable subcontinuum of X that is contained in the union of finitely many links in X , then every subcontinuum of X that contains a nonempty open subset of I contains I .*

Proof. Assume I is contained in the union of α (α is a natural number) links in X . Suppose Z is a collection of $\alpha + 1$ disjoint circular regions in S^2 such that each element of Z intersects I . There exist points c and d belonging to distinct elements of Z such that $\{c, d\}$ is contained in a complementary domain of X . The theorem now follows directly from [4, Theorem 1 (proof)].

Theorem 2. *Suppose M is a hereditarily decomposable continuum and f is a continuous function of M into S^2 . Then no indecomposable subcontinuum of $f(M)$ is terminal in $f(M)$.*

Proof. Assume there exists an indecomposable subcontinuum I of $f(M)$ that is terminal in $f(M)$. There exists a composant C of I such that each subcontinuum of $f(M)$ that meets both C and $f(M) - I$ contains I . Note that since $f(M)$ is decomposable, I is a proper subcontinuum of $f(M)$. Let p be a point of $f^{-1}(C)$ and let Z be the p -component of $f^{-1}(I)$.

For each positive integer n , let G_n be an open set in M such that (1) $Z \subset G_n$, (2) $\text{Bd } G_n \cap f^{-1}(I) = \emptyset$, (3) $M - G_n$ is not the empty set, and (4) the distance from each point of G_n to Z is less than n^{-1} . For each n , let Y_n be the p -component of $\text{Cl } G_n$. Note that, for each n , the continuum Y_n meets $\text{Bd } G_n$ [12, Theorem 50, p. 18] and consequently $f(Y_n)$ is a continuum in $f(M)$ that meets both C and $f(M) - I$. Hence for each n , the continuum $f(Y_n)$ contains I . It follows from the continuity of f that $f(Z)$ is I . Since Z is hereditarily decomposable, this is a contradiction. Hence no indecomposable subcontinuum of $f(M)$ is terminal in $f(M)$.

Theorem 3. *If X is a continuum in S^2 and no finite collection of links in X contains an indecomposable continuum in its union, then X is λ connected.*

Proof. Let p_1 and q_1 be distinct points of X . We shall construct a hereditarily decomposable continuum Q in X that contains p_1 and q_1 .

Let A be an arc in S^2 from p_1 to q_1 ordered by $<$.

Notation. For points x and y of A ($x \leq y$) let $[x, y]$, $[x, y)$, $(x, y]$, and (x, y) denote the sets $\{z \in A \mid x \leq z \leq y\}$, $\{z \in A \mid x \leq z < y\}$, $\{z \in A \mid x < z \leq y\}$, and $\{z \in A \mid x < z < y\}$ respectively.

Let $\{G_n\}$ be the countable collection consisting of all complementary domains of X . If no element of $\{G_n\}$ intersects A , then A is in X and the construction is trivial. Assume this is not the case. Suppose without loss of generality that $G_1 \cap A \neq \emptyset$.

1. The construction of a continuum Z_1 in $(X \cup A) - G_1$ from p_1 to q_1 . Let H_0 be a link in X whose intersection with A is p_1 (if no such link exists let H_0 be $\emptyset$). Let L_0 be a link in X that meets A only at q_1 (if no such link exists let L_0 be $\emptyset$). The links H_0 and L_0 will be referred to as selected 2-links.

With H_0 , L_0 , the arc $[p_1, q_1]$, and the complementary domain G_1 as sets of reference, we now construct either a continuum K_1 or a set T_1 as follows.

If $H_0 \cap L_0 \neq \emptyset$, define $K_1 = H_0 \cup L_0$.

Assume that H_0 and L_0 are disjoint sets. Let x_1 and y_1 be the first and last points respectively of $[p_1, q_1] \cap \text{Bd } G_1$ (with respect to the order of A). Let W_1 be the collection consisting of all links in X that meet $[p_1, x_1] \cup [y_1, q_1]$. Let V_1 be the collection of all 2-links L in X such that L is the union of two (not necessarily distinct) elements of W_1 .

If L is an element of V_1 that is the union of elements H and K (not necessarily distinct) of W_1 , then we call any point set $\{x, y\}$ such that x belongs to $H \cap ([p_1, x_1] \cup [y_1, q_1])$ and y belongs to $K \cap ([p_1, x_1] \cup [y_1, q_1])$ a *basic set* in L .

Suppose no element of V_1 meets both $[p_1, x_1] \cup H_0$ and $[y_1, q_1] \cup L_0$ and no element of V_1 meets both $[p_1, x_1] \cup H_0$ and $(y_1, q_1] \cup L_0$. Define

$$K_1 = [p_1, x_1] \cup \text{Bd } G_1 \cup [y_1, q_1].$$

In the last part of this proof, $\text{Bd } G_1$ will be referred to as a selected 2-link.

Assume that V_1 does not have this property.

Suppose there exists an element E of V_1 in $X - \{p_1, q_1\}$ that meets both H_0 and L_0 . Define $K_1 = H_0 \cup E \cup L_0$. Later E will be referred to as a selected 2-link.

Assume no element of V_1 misses $\{p_1, q_1\}$ and meets both H_0 and L_0 .

Suppose there exists an element of V_1 that meets both H_0 and $[y_1, q_1]$. Let d be the least upper bound (with respect to $<$) of the set of points $[y_1, q_1]$ that can be joined to H_0 by an element of V_1 . Since the limit of a convergent sequence of links is a link, there is an element of V_1 that joins d to H_0 [12, Theorem 59, p. 24]. If possible, select an element H of V_1 joining d to p_1 and define $K_1 = H \cup [d, q_1]$. If no such element exists, select an element E of V_1 missing p_1 that joins d to H_0 and define $K_1 = H_0 \cup E \cup [d, q_1]$.

Assume that no element of V_1 meets both H_0 and $[y_1, q_1]$.

Suppose an element of V_1 meets L_0 and $[p_1, x_1]$. Let a be the first point of $[p_1, x_1]$ that can be joined to L_0 by an element of V_1 . If possible, select an element L of V_1 that joins a to q_1 and define $K_1 = [p_1, a] \cup L$. If no such element exists, select an element E of V_1 missing q_1 that joins a to L_0 and define $K_1 = [p_1, a] \cup E \cup L_0$.

Assume that no element of V_1 meets both L_0 and $[p_1, x_1]$. Let a_1 be the first point of $[p_1, x_1]$ that can be joined to $[y_1, q_1]$ by an element of V_1 . There exist a continuum L_1 and a point q_2 of $[y_1, q_1]$ such that (1) L_1 is a selected element of V_1 that joins q_2 to a_1 , and (2) q_2 is the first point of $[y_1, q_1]$ that can be joined to a_1 by an element of V_1 .

Let d_1 be the last point of $[y_1, q_1]$ that can be joined to $[p_1, x_1]$ by an element of V_1 . There exist a continuum H_1 and a point p_2 such that (1) H_1 is a selected element of V_1 that joins p_2 to d_1 , and (2) p_2 is the last point of $[p_1, x_1]$ that can be so joined to d_1 .

Suppose an element E of V_1 contains $\{a_1, d_1\}$. Define $K_1 = [p_1, a_1] \cup E \cup [d_1, q_1]$.

Assume $\{a_1, d_1\}$ is not contained in an element of V_1 .

Suppose there exists an element E of V_1 that does not contain p_1 and meets

both H_0 and H_1 . Define $K_1 = H_0 \cup E \cup H_1 \cup [d_1, q_1]$.

Assume that each element of V_1 that meets both H_0 and H_1 contains p_1 .

Suppose an element E of V_1 in $X - \{q_1\}$ meets L_0 and L_1 . Define $K_1 = [p_1, a_1] \cup L_1 \cup E \cup L_0$.

Assume that no element of V_1 in $X - \{q_1\}$ meets both L_0 and L_1 .

Suppose an element of V_1 meets $[p_1, a_1]$ and H_1 . Let a be the first point of $[p_1, a_1]$ that can be joined to H_1 by an element of V_1 . Select an element E of V_1 that joins a to H_1 . Define $K_1 = [p_1, a] \cup E \cup H_1 \cup [d_1, q_1]$.

Assume that no element of V_1 meets both $[p_1, a_1]$ and H_1 .

Suppose an element of V_1 meets L_1 and $[d_1, q_1]$. Let d be the last point of $[d_1, q_1]$ that can be joined to L_1 by an element of V_1 . Select an element E of V_1 that joins d to L_1 . Define $K_1 = [p_1, a_1] \cup L_1 \cup E \cup [d, q_1]$.

Assume that no element of V_1 meets both L_1 and $[d_1, q_1]$.

Suppose an element of V_1 meets H_0 and $[p_2, x_1]$. Let a be the first point of $[p_2, x_1]$ that can be joined to H_0 by an element of V_1 . If possible, select an element H of V_1 that joins a to p_1 and define $K_1 = H \cup [p_2, a] \cup H_1 \cup [d_1, q_1]$. If no such element exists, select an element E of V_1 missing p_1 that joins a to H_0 and define $K_1 = H_0 \cup E \cup [p_2, a] \cup H_1 \cup [d_1, q_1]$.

Assume that no element of V_1 meets both H_0 and $[p_2, x_1]$.

Suppose an element of V_1 meets $[y_1, q_2]$ and L_0 . Let d be the last point of $[y_1, q_2]$ that can be joined to L_0 by an element of V_1 . If possible, select an element L of V_1 that joins d to q_1 and define $K_1 = [p_1, a_1] \cup L_1 \cup [d, q_2] \cup L$. If no such element exists, select an element E of V_1 missing q_1 that joins d to L_0 and define $K_1 = [p_1, a_1] \cup L_1 \cup [d, q_2] \cup E \cup L_0$.

Assume that no element of V_1 meets both $[y_1, q_2]$ and L_0 .

Suppose an element of V_1 meets $[p_1, a_1]$ and $[p_2, x_1]$. Let a be the first point of $[p_1, a_1]$ that can be joined to $[p_2, x_1]$ by an element of V_1 . There exist a continuum E and a point b such that (1) E is a selected element of V_1 that joins b to a , and (2) b is the first point of $[p_2, x_1]$ that can be so joined to a . Define $K_1 = [p_1, a] \cup E \cup [p_2, b] \cup H_1 \cup [d_1, q_1]$.

Assume no element of V_1 meets both $[p_1, a_1]$ and $[p_2, x_1]$.

Suppose an element of V_1 meets $[y_1, q_2]$ and $[d_1, q_1]$. Let d be the last point of $[d_1, q_1]$ that can be joined to $[y_1, q_2]$ by an element of V_1 . There exist a continuum E and a point c such that (1) E is a selected element of V_1 that joins c to d , and (2) c is the last point of $[y_1, q_2]$ that can be so joined to d . Define $K_1 = [p_1, a_1] \cup L_1 \cup [c, q_2] \cup E \cup [d, q_1]$.

Assume no element of V_1 meets both $[y_1, q_2]$ and $[d_1, q_1]$.

If an element E of V_1 can be selected that meets L_1 and H_1 , define $K_1 = [p_1, a_1] \cup L_1 \cup E \cup H_1 \cup [d_1, q_1]$.

Assume no element of V_1 meets both L_1 and H_1 .

Suppose an element of V_1 meets L_1 and $[p_2, x_1]$. Let a be the first point of $[p_2, x_1]$ that can be joined to L_1 by an element of V_1 . Let E be a selected element of V_1 that joins a to L_1 . Define $K_1 = [p_1, a_1] \cup L_1 \cup E \cup [p_2, a] \cup H_1 \cup [d_1, q_1]$.

Assume no element of V_1 meets both L_1 and $[p_2, x_1]$.

Suppose an element of V_1 meets $[y_1, q_2]$ and H_1 . Let d be the last point of $[y_1, q_2]$ that can be joined to H_1 by an element of V_1 . Let E be a selected element of V_1 that joins d to H_1 . Define $K_1 = [p_1, a_1] \cup L_1 \cup [d, q_2] \cup E \cup H_1 \cup [d_1, q_1]$.

Assume no element of V_1 meets both $[y_1, q_2]$ and H_1 . Define $T_1 = [p_1, a_1] \cup L_1 \cup H_1 \cup [d_1, q_1]$.

If at this stage of the construction K_1 is defined, then let $Z_1 = K_1$. If K_1 is not defined, then repeat the process, using the continua H_1 , L_1 , the arc $[p_2, q_2]$ and the complementary domain G_1 as reference sets to define K_2 or T_2 . Note that no 2-link in X meets two distinct elements of $\{[p_1, a_1], (p_2, q_2), [d_1, q_1]\}$. Hence the resulting set, K_2 or T_2 , meets T_1 only at the points p_2 and q_2 . If K_2 is defined, let $Z_1 = K_2 \cup T_1$. If K_2 is not defined, repeat the process using H_2 , L_2 , $[p_3, q_3]$, and G_1 as reference sets.

If after this process is repeated n (finitely many) times K_n is defined, then let $Z_1 = K_n \cup \bigcup_{i=1}^{n-1} T_i$. If, for each positive integer n , the set T_n is defined, let $Z_1 = \text{Cl} \bigcup_{i=1}^{\infty} T_i$. Note that Z_1 is a continuum in $S^2 - G_1$ that contains $\{p_1, q_1\}$.

2. The properties of Z_1 . Let M_1 be the collection consisting of all selected 2-links that appear in the definition of Z_1 . The continua that appear in the definition of Z_1 have the *linear separating property*; that is, for any two distinct continua L and E in $S^2 - \{p_1\}$ appearing in the definition of Z_1 (each of L and E is either a selected 2-link or an arc in $A \cap Z_1$ that is maximal with respect to not having an interior point in common with $\text{Cl}(\text{St } M_1)$), either L separates p_1 from $E - L$ or E separates p_1 from $L - E$ in Z_1 but not both. Note that only one continuum appearing in the definition of Z_1 contains p_1 .

Note that Z_1 has the *linear linking condition* with respect to M_1 ; that is, no 2-link in X meets more than one component of $Z_1 - \text{Cl}(\text{St } M_1)$ and if J is a component of $Z_1 - \text{Cl}(\text{St } M_1)$ and r is an endpoint of J , there is at most one point v that is an endpoint of a component of $Z_1 - (J \cup \text{Cl}(\text{St } M_1))$ that can be joined to r by a 2-link in X .

3. The second phase of the construction of Q . If no element of $\{G_n\}$ intersects Z_1 , define $Q = Z_1$.

Assume that some element of $\{G_n\}$ meets Z_1 . Define $Q_1 = Z_1$. Suppose without loss of generality that $G_2 \cap Q_1 \neq \emptyset$. Since no 2-link in X meets more than

one component of $Q_1 - \text{Cl}(\text{St } M_1)$, the domain G_2 intersects only one component (p_1^2, q_1^2) of $Q_1 - (\text{Cl}(\text{St } M_1) \cup \{p_1, q_1\})$. Using the construction methods of §1, define a continuum Z_2 such that $Q_2 = (Q_1 - (p_1^2, q_1^2)) \cup Z_2$ is a continuum in $S^2 - (G_1 \cup G_2)$ that contains $\{p_1, q_1\}$. Here we start with two selected 2-links in Z_1 , the arc $[p_1^2, q_1^2]$, and the domain G_2 as sets of reference for the construction of Z_2 . Note that each basic set in any 2-link that is selected in the construction of Z_2 is by definition in $[p_1^2, q_1^2] - G_2$.

Assume without loss of generality that the continua used to define Q_2 (i.e., all continua except $[p_1^2, q_1^2]$ appearing in the definitions of Q_1 and Z_2) have the linear separating property. Note that it may be necessary to delete an element of M_1 in the definition of Q_2 to get this property.

Let M_2 be the collection of selected 2-links used to define Q_2 . The continuum Q_2 has the linear linking condition with respect to M_2 .

4. Continuing until no complementary domain meets the constructed set. If no element of $\{G_n\}$ meets Q_2 , define $Q = Q_2$. If this is not the case, we continue the process.

Assume the continuum Q_n is defined and M_n is the collection of all selected 2-links in Q_n . If Q_n is contained in X , define $Q = Q_n$. Assume Q_n intersects some element of $\{G_n\}$. There exists an integer i greater than n such that $Q_n \cap G_i \neq \emptyset$ and $Q_n \cap \bigcup_{j=1}^{i-1} G_j = \emptyset$. The domain G_i intersects only one component (p_1^{n+1}, q_1^{n+1}) of $Q_n - (\text{Cl}(\text{St } M_n) \cup \{p_1, q_1\})$. Using the construction methods of §1, define Z_{n+1} such that $Q_{n+1} = (Q_n - (p_1^{n+1}, q_1^{n+1})) \cup Z_{n+1}$ is a continuum in $S^2 - \bigcup_{j=1}^i G_j$ containing $\{p_1, q_1\}$. We can assume without loss of generality that the continua used to define Q_{n+1} have the linear separating property. The continuum Q_{n+1} has the linear linking condition with respect to M_{n+1} the collection of all selected 2-links used to define Q_{n+1} .

Suppose that for each positive integer n , a continuum Q_n is defined. Define Q to be the limit of $\{Q_n\}$.

Since in all cases Q is a continuum in X that contains $\{p_1, q_1\}$, the construction is complete. Now we must show that Q is hereditarily decomposable.

5. Derived subcontinua of Q . Suppose there exist an arc $[r, v]$ in A and links F_1, F_2 , and F_3 contained in $X - [r, v]$ such that for each $i = 1, 2$, and 3 , F_i is the limit of a convergent sequence $\{E_n^i\}$ of links in $X - \bigcup_{j=1}^3 F_j$ with the following properties:

- (1) For each n , the link E_n^i is contained in a selected 2-link L and a point v_n^i of a basic set in L is in $[r, v] \cap E_n^i$.
- (2) The sequence $\{v_n^i\}$ converges to v .

We now show that one of F_1, F_2 , and F_3 is contained in the union of the other two.

Assume that for $j = 1, 2$, and 3 , there exists a point w_j of $F_j - \bigcup_{i \neq j} F_i$. There exist a point s of (r, v) , a positive integer m , an element E of $\{E_n^3\}$, and circular regions U_1, U_2 , and U_3 whose closures are disjoint containing w_1, w_2 , and w_3 respectively such that (1) $\text{Cl}(U_1 \cup U_2 \cup U_3)$ is in $S^2 - [r, v]$, (2) E meets both $[r, s]$ and U_3 and does not intersect $\text{Cl}(U_1 \cup U_2)$, and (3) for each n greater than m ,

$$\begin{aligned}(F_1 \cup E_n^1) \cap \text{Cl}(U_2 \cup U_3) &= (F_2 \cup E_n^2) \cap \text{Cl}(U_1 \cup U_3) \\ &= (F_3 \cup E_n^3) \cap \text{Cl}(U_1 \cup U_2) = \emptyset.\end{aligned}$$

Let X_1 be a continuum in $S^2 - E$ that is the union of an arc in (s, v) , an element of $\{E_n^1\}$ in $X - \text{Cl}(U_2 \cup U_3)$ that meets both (s, v) and U_1 , and an element of $\{E_n^2\}$ in $X - \text{Cl}(U_1 \cup U_3)$ that meets both (s, v) and U_2 .

There exists a continuum X_2 in $S^2 - (E \cup X_1)$ that is the union of an arc in (s, v) , an element of $\{E_n^1\}$ in $X - \text{Cl}(U_2 \cup U_3)$ that meets both (s, v) and U_1 , and an element of $\{E_n^2\}$ in $X - \text{Cl}(U_1 \cup U_3)$ that meets both (s, v) and U_2 .

There exists a point t of (s, v) such that no element of $\{E_n^3\}$ meets both (t, v) and $X_1 \cup X_2$. Let T be the continuum $F_1 \cup F_2 \cup X_1 \cup X_2 \cup \text{Cl}(U_1 \cup U_2)$. There is an element H of $\{E_n^3\}$ in $S^2 - T$ that meets both (t, v) and U_3 . The disjoint continua $F_1 \cup F_2, X_1$, and X_2 can be closely approximated by arcs in such a way that the existence of the continuum $E \cup H \cup \text{Cl } U_3$ contradicts [12, Theorem 116, p. 247]. Hence one of F_1, F_2 , and F_3 is contained in the union of the other two.

It follows that the collection of all links having the properties that F_1, F_2 , and F_3 have with respect to $[r, v]$ when partially ordered by inclusion has at most two maximal elements. Hence the union of this collection is a 2-link F in X . Since r precedes v with respect to the order of A , we shall refer to F as a continuum in Q that is *derived from the left at v* . Similarly continua that are derived from the right at a point of A are 2-links in X . A continuum that is derived from the left or from the right at a point z of A will be referred to as a *derived continuum* in Q with *base point z* .

6. Definition of the collection C . If E is an element of M_i , for some i , that is not deleted at a later stage of the construction to get the linear separating property or if E is an arc in $A \cap Q$ that is maximal with respect to not having an interior point in common with $\bigcup \text{Cl}(\text{St } M_i)$, then E is said to be a continuum used to define Q . Let B denote the collection of all continua used to define Q . Let D denote the collection consisting of all derived continua in Q .

Define C to be the collection consisting of all continua Y that satisfy one of the following conditions:

(1) Y is the union of an element E of B with all elements of D that have their base points in E .

(2) There is an element F of D that does not have a base point in an element of B . The union of F with all elements of D that meet F is Y .

Note that $Q - \text{St } C$ is a totally disconnected set in A .

For each continuum E belonging to B , there are at most two elements of D that have base points in E . Furthermore, for each element F of D at most one element of B contains a base point of F . Since the continua at all stages of the construction have the linear linking condition, each element of D meets at most one other element of D . It follows that for each element Y of C , there exist points r and t (not necessarily distinct) of A and a collection N , consisting of finitely many links in X and possibly one arc in $A \cap Q$ that is maximal with respect to not having an interior point in $\bigcup \text{Cl}(\text{St } M_i)$, such that each element of N meets $\{r, t\}$ and the union of N is Y . The set $\{r, t\}$ will be referred to as the *core set* of Y .

7. An order relation in C in terms of separating. Define the binary relation $\ll$ in C as follows. For distinct elements Y and Z of C , $Y \ll Z$ if Y contains p_1 or Y separates p_1 from the set $Z - Y$ in Q . Since the continua at all stages of the construction have the linear linking condition and the sets used to define these continua have the linear separating property, $\ll$ is an order relation in C (i.e., $\ll$ is transitive and satisfies the law of trichotomy). When checking the properties of $\ll$, it is helpful to note that for any two elements Y and Z of C , if $Y \ll Z$, then the p_1 -component of $Q - Z$ meets $Y - Z$.

Note that if V , Y , and Z are elements of C such that $V \ll Y$ and $Y \ll Z$, then V and Z are mutually exclusive.

8. All continua in Q are decomposable. Suppose there exists an indecomposable continuum I in Q . Since I is not contained in the union of finitely many links and arcs in X and $Q - \text{St } C$ is totally disconnected, there exist mutually exclusive elements V , Y , and Z of C such that $V \ll Y$, $Y \ll Z$, and each component of I meets both V and Z . Note that since I is not separated by any of its proper subcontinua, Y cannot be an arc in A .

Suppose that the core set of Y consists of two distinct points r and t . The continuum Y separates V from Z in Q . Hence there exist six disjoint continua $\{P_i\}_{i=1}^6$ in $I - \{r, t\}$, each intersecting V , Z , and a link in X that meets $\{r, t\}$. Let R and U be disjoint circular regions in $S^2 - (V \cup Z \cup \bigcup_{i=1}^6 P_i)$ that are centered on r and t respectively. For each integer i ($1 \leq i \leq 6$), let e_i be a point of P_i that can be joined to either r or t by a link in X and let R_i be a circular region centered on e_i in $S^2 - (V \cup Z \cup \bigcup_{j \neq i} P_j)$. For each i , since e_i

belongs to a link in X that meets $\{r, t\}$, there exists an arc J_i in a complementary domain of X that meets both R_i and a component of $R \cup U$.

It follows that some three of $\{J_i\}_{i=1}^6$ meet one component of $R \cup U$. Assume without loss of generality that J_1 , J_2 , and J_3 each meet R . The continua V , Z , P_1 , P_2 , and P_3 can be closely approximated by arcs in such a way that the existence of J_1 , J_2 , and J_3 contradicts [12, Theorem 116, p. 247].

Using an argument similar to the preceding, we can show that assuming the core set of Y consists of one point also involves a contradiction. Hence Q is hereditarily decomposable.

Continuity.

Theorem 4. *Suppose M is a hereditarily decomposable continuum, f is a continuous function of M into the plane, and R is the union of finitely many links in $f(M)$. Then R does not contain an indecomposable continuum.*

Proof. Assume there exists an indecomposable continuum I in R . According to Theorem 1, every subcontinuum of $f(M)$ that contains a nonempty open subset of I contains I . It follows that I is terminal in $f(M)$ [8, Theorem 2]. Since M is hereditarily decomposable, this contradicts Theorem 2. Hence R does not contain an indecomposable continuum.

Theorem 5. *If M is a λ connected continuum and f is a continuous function of M into the plane, then $f(M)$ is λ connected.*

Proof. Suppose p and q are distinct points of $f(M)$. Since M is λ connected, there exists a hereditarily decomposable continuum H in M that meets both $f^{-1}(p)$ and $f^{-1}(q)$. According to Theorems 3 and 4, $f(H)$ is λ connected. Hence there exists a hereditarily decomposable continuum in $f(H)$ that joins p and q . It follows that $f(M)$ is λ connected.

Comment. Theorem 5 cannot be generalized to include all continuous images in Euclidean 3-space. There are continuous images of the topologist's sine curve in E^3 that are not λ connected.

Aposyndesis.

Theorem 6. *If X is a semiaposyndetic plane continuum, then X is λ connected.*

Proof. Assume there exists an indecomposable continuum I that is contained in the union of finitely many links in X . Let x and y be distinct points of I . According to Theorem 1, every subcontinuum of X that contains a point of $\{x, y\}$ in its interior relative to X contains I . This contradicts the assumption that X is semiaposyndetic. It follows from Theorem 3 that X is λ connected.

Corollary. *Every aposyndetic plane continuum is λ connected.*

Example. An aposyndetic continuum in Euclidean 3-space need not be λ connected. To see this let M be the product of the pseudo arc [11] with itself. M is aposyndetic [9, Theorem 7] and not λ connected. In fact, M does not contain a hereditarily decomposable continuum. Since the pseudo arc is chainable, M is embeddable in Euclidean 3-space [2].

BIBLIOGRAPHY

1. H. Bell, *On fixed point properties of plane continua*, Trans. Amer. Math. Soc. 128 (1967), 539–548. MR 35 #4888.
2. R. Bennett, *Embedding products of chainable continua*, Proc. Amer. Math. Soc. 16 (1965), 1026–1027. MR 31 #6216.
3. C. L. Hagopian, *Concerning arcwise connectedness and the existence of simple closed curves in plane continua*, Trans. Amer. Math. Soc. 147 (1970), 389–402. MR 40 #8030.
4. ———, *A fixed point theorem for plane continua*, Bull. Amer. Math. Soc. 77 (1971), 351–354. MR 42 #8469.
5. ———, *Arcwise connectedness of semi-aposyndetic plane continua*, Trans. Amer. Math. Soc. 158 (1971), 161–165. MR 44 #2205.
6. ———, *Arcwise connectivity of semi-aposyndetic plane continua*, Pacific J. Math. 37 (1971), 683–686. MR 46 #6322.
7. ———, *Another fixed point theorem for plane continua*, Proc. Amer. Math. Soc. 31 (1972), 627–628. MR 44 #3309.
8. ———, *Planar images of decomposable continua*, Pacific J. Math. 42 (1972), 329–331.
9. F. B. Jones, *Concerning non-aposyndetic continua*, Amer. J. Math. 70 (1948), 403–413. MR 9, 606.
10. ———, *Concerning aposyndetic and non-aposyndetic continua*, Bull. Amer. Math. Soc. 58 (1952), 137–151. MR 14, 71.
11. E. E. Moise, *An indecomposable plane continuum which is homeomorphic to each of its nondegenerate subcontinua*, Trans. Amer. Math. Soc. 63 (1948), 581–594. MR 10, 56.
12. R. L. Moore, *Foundations of point set theory*, rev. ed., Amer. Math. Soc. Colloq. Publ., vol. 13, Amer. Math. Soc., Providence, R. I., 1962. MR 27 #709.
13. G. T. Whyburn, *Semi-locally connected sets*, Amer. J. Math. 61 (1939), 733–749. MR 1, 31.

DEPARTMENT OF MATHEMATICS, CALIFORNIA STATE UNIVERSITY, SACRAMENTO,
CALIFORNIA 95819 (Current address)

DEPARTMENT OF MATHEMATICS, ARIZONA STATE UNIVERSITY, TEMPE, ARIZONA
85281