CONE BUNDLES

BY

CLINT MCCRORY⁽¹⁾

ABSTRACT. A theory of normal bundles for locally knotted codimension two embeddings of PL manifolds is developed. The classifying space for this theory is Cappell and Shaneson's space BRN_2 .

Cone bundles are a generalization of blockbundles [9], allowing local knotting of the base in the total space. They are a type of "mockbundle" [8], [2], closely related to the theory of stratified polyhedra [12], and designed to provide a simple foundation for Cappell and Shaneson's theory of singularities of PL embeddings [3], [4]. A similar definition has been given by Matumoto and Matsumoto [6].

§1 contains the basic definitions. A classifying space for cone bundles is constructed in §2. §3 contains a proof that the total space of a cone bundle over a manifold is a manifold. Finally, cone bundles are related to the topology of stratified polyhedra in §4.

The geometric idea for cone bundles comes from my paper [7] on cone complexes. I thank Sylvain Cappell for encouraging me to develop this idea.

I will work in the category of polyhedra and piecewise linear maps [11]. In particular, all manifolds and homeomorphisms will be piecewise linear.

1. Thickenings. Let M be a compact n -manifold. A codimension q *thickening* of M is a compact $(n + q)$ -manifold W containing M as a subpolyhedron, such that W collapses to M . Furthermore, $\partial W \cap M = \partial M$, and there is a collar of ∂W in W which restricts to a collar of ∂M in M . (This is called a "very proper" thickening in [4].)

The thickenings V and W of M are *equivalent* if there is a homeomorphism between V and W which is the identity on M .

If $q > 2$, a codimension q thickening W of M is an "abstract regular neighborhood" of M [9, p. 14], since M is locally flat in W . But if $q = 2$, M can be locally knotted in W . M can be locally knotted in a codimension one thickening if and only if the PL Schoenflies conjecture is false.

Received by the editors November 20, 1975.

AMS (MOS) subject classifications (1970). Primary 57C40, 57C50; Secondary 57C45.

Key words and phrases. Mockbundle, blockbundle, locally knotted embedding, thickening, cone complex, piecewise linear stratification.

⁽¹⁾ Supported in part by NSF Grant GP-43128.

© American Mathematical Society 1977

Let K be a (PL) cell complex. A q -cone bundle ξ/K consists of a polyhedron $E(\xi)$ containing $|K|$ such that

- (i) For each p -cell $\sigma_i \in K$ there is a $(p+q)$ -ball $\beta_i \subset E(\xi)$, containing σ_i , such that (β_i, σ_i) is homeomorphic with the cone on a sphere pair. (N.B. this sphere pair may be nonlocally flat or knotted.) β_i is called the *block* over σ_i .
- (ii) $E(\xi)$ is the union of the blocks β_i .
- (iii) The interiors of the blocks are disjoint.
- (iv) $\beta_i \cap \beta_j$ is the union of the blocks over the cells contained in $\sigma_i \cap \sigma_j$.

By the Zeeman unknotting theorem, a q -cone bundle is a blockbundle [9] if $q > 2$, i.e., (β_i, σ_i) is homeomorphic with a standard sphere pair for each $\sigma_i \in K$. The following results show that cone bundles bear essentially the same relation to thickenings that blockbundles bear to abstract regular neighborhoods.

LEMMA 1. *Let N be a manifold containing the manifold M as a subpolyhedron. Suppose that $\partial N \cap M = \partial M$ and $(\partial N, \partial M)$ is collared in (N, M) . Then there is a cone bundle ξ/K with $|K| = M$ such that $E(\xi)$ is a regular neighborhood of M in N .*

PROOF. ξ is constructed in the same manner as a normal blockbundle for a locally flat submanifold, using dual cells. (The usual construction must be slightly modified near the boundary; cf. [7, p. 284].)

LEMMA 2. *If ξ/K is a cone bundle, and $|K|$ is a compact manifold, then $E(\xi)$ is a thickening of K .*

PROOF. It is easy to see that $E(\xi)$ collapses to $|K|$, since $E(\xi)$ can be triangulated as a stellar neighborhood of $|K|$, by induction on the dimension of ξ (cf. [7, p. 274]). The fact that $E(\xi)$ is a *manifold* will be proved in §3.

LEMMA 3. *If W is a thickening of the compact manifold M , there is a cone bundle ξ/K , $|K| = M$, such that the thickening $E(\xi)$ of M is equivalent to W .*

PROOF. This follows from Lemma 1 and the uniqueness of regular neighborhoods [11, p. 33].

REMARKS. (1) If ξ/K and η/L are cone bundles with $|K| = |L| = M$, and the thickenings $E(\xi)$ and $E(\eta)$ of M are equivalent, one might expect (by analogy with blockbundles) that ξ and η have isomorphic "subdivisions". However, this is not true—one has to introduce the weaker relation of *concordance* (§2) in order to get a bijection between classes of bundles and classes of thickenings.

(2) In Matumoto and Matumoto's definition of " RN_2 -bundles" [6], condition (i) in the definition of a 2-cone bundle is weakened to the condition that (β_i, σ_i) is an arbitrary ball pair. Lemma 2 is not true for their bundles, since the total space need not collapse to the base space.

(3) A 2-cone bundle ξ/K has a canonical "Noguchi characteristic class" $n \in H^2(K; \gamma)$ (twisted coefficients), where γ is the Fox-Milnor cobordism group (cf. [4]). n is represented by the cocycle which assigns to each 2-cell $\sigma \in K$, the cobordism class of the knot $(\partial\beta, \partial\sigma)$, where β is the block over σ . Thus n is the primary obstruction to making ξ a blockbundle. (The analogous higher obstructions are not defined *a priori* since $\partial\sigma_i$ is not necessarily locally flat in $\partial\beta_i$ if $\dim \sigma_i > 2$.)

2. A classifying space. The following definitions come from [9].

If ξ/K is a cone bundle and L is a subcomplex of K , the *restriction* $\xi|L$ is defined by putting $\beta_i(\xi|L) = \beta_i(\xi)$ for each $\sigma_i \in L$.

The cone bundles $\xi_0, \xi_1/K$ are *isomorphic* if there is a homeomorphism $h: E(\xi_0) \rightarrow E(\xi_1)$ such that h is the identity on $|K|$ and $h(\beta_i(\xi_0)) = \beta_i(\xi_1)$ for each $\sigma_i \in K$.

The cone bundles $\xi_0, \xi_1/K$ are *concordant* if there is a cone bundle $\eta/(K \times I)$ such that $\eta|(K \times \{i\})$ is isomorphic with ξ_i , $i = 0, 1$. Here $I = [0, 1]$ and $K \times I$ is the usual product complex. (Two blockbundles are concordant if and only if they are isomorphic [9, p. 6]. This is not true for 2-cone bundles.)

We will construct a classifying space for concordance classes of 2-cone bundles analogous to the classifying space $B\widetilde{PL}_q$ for q -blockbundles. (The same construction also works for 1-cone bundles.)

Let $\mathcal{K}(K)$ be the set of concordance classes of 2-cone bundles over K . $\mathcal{K}$ is a contravariant functor from the category with objects PL cell complexes and morphisms generated by isomorphisms and inclusions of subcomplexes, to the category of (based) sets. (The base point of $\mathcal{K}(K)$ is the class of the trivial bundle over K .)

THEOREM 1. $\mathcal{K}$ has a unique extension to the category of CW complexes and homotopy classes of maps.

PROOF. This is a corollary of the "mockbundle" recipe for homotopy functors [2, I]. It is clear that cone bundles can be glued (axiom G [2, p. 15]), so we only have to verify the extension axiom (E, [2, p. 15]). That is, if $e: K_0 \rightarrow K$ is an elementary expansion, and ξ_0/K_0 is a cone bundle, we must construct a cone bundle ξ/K such that $\xi|K_0 = \xi_0$. We follow [2, p. 21]. $K = K_0 \cup \{\sigma, \tau\}$, where σ is a principal cell of K and τ is a free face of σ . Let J be the subcomplex of $\partial\sigma$ consisting of all the faces of σ except τ . Now $|J|$ is a ball, and $E(\xi_0|J)$ is a thickening of $|J|$ by Lemma 2, so $E(\xi_0|J)$ is a ball. Let (B, C) denote the ball pair $(E(\xi_0|J), |J|)$. Identifying (σ, C, τ) with $(C \times I, C \times \{0\}, (\partial C \times I) \cup (C \times \{1\}))$, we define the extension ξ/K as follows. The block of ξ over σ is the cone on the boundary of the ball $B \times I$, and the block over τ is the cone on the boundary of the ball

$$\text{cl}[\partial(B \times I) \setminus (B \cup (B^* \times I))],$$

where $B^* = \text{cl}[\partial B \setminus \bigcup \beta_i(\xi_0)]$, union over all i such that $\sigma_i \subset \partial C$.

THEOREM 2. $\mathcal{H}$ is a representable functor.

PROOF. Let G be the (based) Δ -set whose k -simplexes are 2-cone bundles over Δ^k (the standard k -simplex) which are embedded blockwise in $\Delta^k \times R^\infty$, and let γ be the canonical cone bundle on G (cf. [8, p. 131] and [2, p. 37]). G is a Kan Δ -set by the extension axiom (see the proof of Theorem 1) and general position. It follows that if $\mathcal{G}$ is the realization of G , pulling back the class of γ induces a bijection

$$\mathcal{H}(X) \cong [X, \mathcal{G}]$$

for all CW complexes X , where $[,]$ denotes homotopy classes of maps (cf. [10, §6] and [9, §2]).

Theorems 1 and 2 are also true for Matumoto and Matsumoto's RN_2 -bundles, by the same proofs. As they have pointed out to me, any RN_2 -bundle is concordant to a cone bundle by the Alexander trick, so the corresponding homotopy functors are the same.

The thickenings W_0 and W_1 of the n -manifold M are *concordant* if there is a thickening Q of $M \times I$ such that W_i is a regular neighborhood of $M \times \{i\}$ rel $\partial M \times \{i\}$ in ∂Q , $i = 0, 1$ (cf. [4]).

THEOREM 3. If M is a compact n -manifold, $\xi \mapsto E(\xi)$ induces a bijection between $\mathcal{H}(M)$ and concordance classes of codimension 2 thickenings of M .

PROOF. Every thickening of M is in fact *equivalent* to $E(\xi)$ for some ξ over M , by Lemma 3. On the other hand, given ξ_0 and ξ_1 , and a concordance Q between $E(\xi_0)$ and $E(\xi_1)$, a concordance between ξ_0 and ξ_1 can be constructed as a regular neighborhood of $M \times I$ in Q , by the relative version of Lemma 1.

It follows that the classifying space $\mathcal{G}$ is (canonically homotopy equivalent with) Cappell and Shaneson's classifying space BRN_2 [3], [4]. In the same way, "oriented" 2-cone bundles are classified by $BSRN_2$, and 2-cone bundles which are blockbundles on the $(k - 1)$ -skeleton are classified by $BRN_{2,k}$.

REMARKS. (1) The Noguchi obstruction n can be viewed as a natural transformation from $\mathcal{H}(\cdot)$ to $H^2(\cdot; \gamma)$, since $n(\xi)$ depends only on the concordance class of ξ . Furthermore, $n(\xi) = 0$ if and only if ξ is concordant to a cone bundle which is a blockbundle on the 2-skeleton (cf. [4, §3]).

(2) In [4], Cappell and Shaneson completely determine the homotopy type of $BSRN_2$. An interesting problem is to give a geometric description of the resulting H -space structure on $BSRN_2$.

3. Collared complexes. A *collared complex* $\mathcal{C}$ on a polyhedron $X = |\mathcal{C}|$ is a locally finite covering of X by compact subpolyhedra, together with a subpolyhedron $\delta\alpha$ of each element α of $\mathcal{C}$ such that

- (i) for each $\alpha \in \mathcal{C}$, $\delta\alpha$ is a union of elements of $\mathcal{C}$,
- (ii) if α and β are distinct elements of $\mathcal{C}$, $\alpha^\circ \cap \beta^\circ$ is empty, where $\alpha^\circ = \alpha \setminus \delta\alpha$,
- (iii) $\delta\alpha$ is collared in α for each $\alpha \in \mathcal{C}$. ((i) and (ii) imply that $\alpha \cap \beta$ is a union of elements of $\mathcal{C}$.)

Collared complexes are Akin's "general complexes" [1]. Examples of collared complexes are cell complexes, manifold complexes [5], and cone complexes [7].

The usefulness of collared complexes comes from the following proposition, derived from the proof of a lemma of Cohen and Sullivan [5, p. 142]. (See also [2, p. 21] and [7, p. 278].)

If $\mathcal{C}$ is a collared complex and $\alpha \in \mathcal{C}$, let $L(\alpha)$ be the geometric realization of the nerve of the finite partially ordered set $\{\beta \in \mathcal{C}, \alpha < \beta\}$, where $\alpha < \beta$ means $\alpha \subset \delta\beta$.

PROPOSITION 1. *If $\mathcal{C}$ is a collared complex on X , $\alpha \in \mathcal{C}$, and $x \in \alpha^\circ$, then*

$$lk(x; X, \alpha) \cong (L(\alpha) * lk(x; \alpha), lk(x; \alpha)),$$

where lk denotes the link, and $*$ denotes the join.

PROOF. Use induction on the "depth" of α , i.e. the length of a maximal chain $\alpha < \alpha_1 < \dots < \alpha_n$ in $\mathcal{C}$.

With this proposition, we can prove Lemma 2, by induction on the dimension of the base. Let ξ/K be a cone bundle, with $|K|$ a manifold (with boundary). If $\sigma_i \in K$, and β_i is the block of ξ over σ_i , let $\delta\beta_i = E(\xi|_{\partial\sigma_i})$. By induction hypothesis, $\delta\beta_i$ is a codimension 0 submanifold of $\partial\beta_i$, so $\delta\beta_i$ is collared in β_i . Thus the set of blocks of ξ forms a collared complex $\mathcal{C}$ on $E(\xi)$. The map $\beta_i \mapsto \sigma_i$ is an incidence preserving bijection between $\mathcal{C}$ and K . Therefore $L(\beta_i) = L(\sigma_i)$ for all $\sigma_i \in K$. Thus the proposition implies $E(\xi)$ is a manifold (with boundary), since each block β_i is a manifold and $|K|$ is a manifold.

4. Geometry of codimension 2 thickenings. Let W be a codimension 2 thickening of the compact n -manifold M . If $x \in M$, the *intrinsic dimension* $d(x; W, M)$ is the smallest integer k such that x is in the k -skeleton of every (PL) cell complex on W which has M as a subcomplex. The k th *intrinsic stratum* S_k of M in W is

$$\{x \in M \setminus \partial M, d(x; W, M) = k\} \cup \{x \in \partial M, d(x; \partial W, \partial M) = k - 1\}.$$

(Cf. [12, p. 13]. Recall that $(\partial W, \partial M)$ is collared in (W, M) .) S_k is a k -dimensional submanifold of M , and $\text{cl}(S_k) = \bigcup_{j \leq k} S_j$. S_n is the set of

locally flat points of M in W , and S_k can be thought of as the points at which the “degree of local knottedness” of M in W is $n - k$. Let $\mathfrak{S} = \{S_k\}$ denote this *intrinsic stratification* of M in W .

By Lemma 3, we can assume that $W = E(\xi)$ for some 2-cone bundle ξ/K , $|K| = M$. Now for each block β of ξ , choose a cellular subdivision of the “rim” $\beta^* = \text{cl}(\partial\beta \setminus \delta\beta)$. These cells, together with the blocks themselves, form a cell complex $\mathcal{C}$ on W . Choose a cone structure for each cell of $\mathcal{C}$ so that σ_i is a subcone of β_i for each cell $\sigma_i \in K$. (N.B. K is not a subcomplex of $\mathcal{C}$.) Then the *dual* cone complex $\mathcal{C}^*$ on W [7] will have the complex K^* on M as a subcomplex. (Note that the cones of $\mathcal{C}^*$ are cells, but if $\alpha \in \mathcal{C}^*$ and $\alpha \cap \partial W \neq \emptyset$, the apex of α lies in ∂W .) It follows that $\text{cl}(S_k)$ is a subcomplex of K^* for all k , i.e. *the cells of K are transverse to the intrinsic stratification of M in W* . (See [7, p. 287] for a discussion of transversality to a stratification.)

If K' is a subdivision of K , the cone bundle ξ'/K' is a *subdivision* of ξ/K if for each $\sigma_i \in K$, $\beta_i(\xi) = \bigcup \beta_j(\xi')$, where the union is taken over all blocks $\beta_j(\xi')$ over cells $\tau_j \in K'$ such that $\tau_j \subset \sigma_i$.

It is not hard to see that a subdivision ξ' of ξ will exist over the subdivision K' of K if and only if $(K')^*$ extends to a cell complex on $W = E(\xi)$ (for some cone structuring of K'). This is equivalent to the condition that K' be transverse to the intrinsic stratification $\mathfrak{S}$. Therefore, ξ can be restricted to precisely those subpolyhedra of M which are transverse to $\mathfrak{S}$.

Thus the fact that concordance classes of cone bundles can be “pulled back” is a consequence of the geometric fact that any subpolyhedron X of the manifold M can be moved transverse to $\mathfrak{S}$. (In fact, Stone’s transversality theorem [12] can be easily proved from the mockbundle viewpoint—cf. [7, p. 287].)

The following result is important in [4].

PROPOSITION 2. *Let W be a codimension 2 thickening of M , and let N be a locally flat codimension q submanifold of M , with $\partial M \cap N = \partial N$. Suppose that N is transverse to the intrinsic stratification $\mathfrak{S}$ of M in W . Then there is a cone bundle ξ over M with $E(\xi) = W$, and a normal blockbundle ν of N in M such that $E(\nu)$ is transverse to $\mathfrak{S}$, and $E(\xi|E(\nu))$ is a codimension q thickening of $E(\xi|N)$ equivalent to $E(q^*\nu)$, where $q: E(\xi|N) \rightarrow N$ is a homotopy inverse of the inclusion.*

PROOF. N is transverse to $\mathfrak{S}$ implies there is a cone bundle η/L , $|L| = M$, with $E(\eta) = W$ and N a subcomplex of L . Let K be the canonical “full” subdivision of L constructed in [7, p. 276], and let ξ be a subdivision of η over K . (It is easy to construct ξ explicitly.) Then the union of the cells in K which meet N is a regular neighborhood of N , and so this neighborhood equals $E(\nu)$ for some blockbundle ν over N . $E(\nu)$ is transverse to $\mathfrak{S}$ since it is a

subcomplex of K . $E(\xi|E(\nu))$ is a manifold by Lemma 2, and it collapses to $E(\xi|N)$ since $E(\nu)$ collapses to N . Thus $E(\xi|E(\nu))$ is a thickening of $E(\xi|N)$. $E(\xi|N)$ is locally flat in $E(\xi|E(\nu))$ by Proposition 1, since the given collared complexes on $E(\xi|E(\nu))$ and $E(\nu)$ are abstractly isomorphic. Thus $E(\xi|E(\nu)) \supset E(\xi|N)$ is equivalent to $E(q^*\nu) \supset E(\xi|N)$ by the uniqueness of regular neighborhoods.

REFERENCES

1. E. Akin, *Transverse cellular mappings of polyhedra*, Trans. Amer. Math. Soc. **169** (1972), 401–438. MR **48** #5088.
2. S. Buoncristiano, C. P. Rourke and B. J. Sanderson, *A geometric approach to homology theory*, London Math. Soc. Lecture Notes Series, no. 18, Cambridge Univ. Press, London, 1976.
3. S. E. Cappell and J. L. Shaneson, *Nonlocally flat embeddings, smoothings, and group actions*, Bull. Amer. Math. Soc. **79** (1973), 577–582. MR **47** #9635.
4. ———, *Piecewise linear embeddings and their singularities*, Ann. of Math. **103** (1976), 163–228.
5. M. M. Cohen and D. Sullivan, *On the regular neighborhood of a two-sided submanifold*, Topology **9** (1970), 141–147. MR **41** #9262.
6. T. Matumoto and Y. Matsumoto, *The unstable difference between homology cobordism and piecewise linear bundles* (to appear).
7. C. McCrory, *Cone complexes and PL transversality*, Trans. Amer. Math. Soc. **207** (1975), 269–291.
8. C. P. Rourke, *Block structures in geometric and algebraic topology*, Actes Congres Internat. Math. (Nice, 1970), Vol. 2, Gauthier-Villars, Paris, 1971, pp. 127–132.
9. C. P. Rourke and B. J. Sanderson, *Block bundles*. I, Ann. of Math. (2) **87** (1968), 1–28. MR **37** #2234a.
10. ———, Δ -sets. I: *Homotopy theory*, Quart. J. Math. Oxford Ser. (2) **22** (1971), 321–338. MR **45** #9327.
11. ———, *Introduction to piecewise-linear topology*, Ergebnisse Math., Band 69, Springer-Verlag, Berlin and New York, 1972. MR **50** #3236.
12. D. Stone, *Stratified polyhedra*, Lecture Notes in Math., vol. 252, Springer-Verlag, Berlin and New York, 1972. MR **51** #9074.

DEPARTMENT OF MATHEMATICS, BROWN UNIVERSITY, PROVIDENCE, RHODE ISLAND 02912