THE COHOMOLOGY OF THE SYMMETRIC GROUPS

BY

BENJAMIN MICHAEL MANN

ABSTRACT. Let $\mathfrak{S}_n$ be the symmetric group on n letters and SG the limit of the sets of degree $+1$ homotopy equivalences of the $n-1$ sphere. Let p be an odd prime. The main results of this paper are the calculations of $H^*(\mathfrak{S}_n, \mathbb{Z}/p)$ and $H^*(SG, \mathbb{Z}/p)$ as algebras, determination of the action of the Steenrod algebra, $\mathcal{Q}(p)$, on $H^*(\mathfrak{S}_n, \mathbb{Z}/p)$ and $H^*(SG, \mathbb{Z}/p)$ and integral analysis of $H^*(\mathfrak{S}_n, \mathbb{Z}, p)$ and $H^*(SG, \mathbb{Z}, p)$.

0. Introduction. Let K and L be discrete groups with L abelian. The groups $H^n(K, L)$ have been of interest for years. [12] and [11] first considered these cohomology groups algebraically and their relation with topological problems. The algebraic groups $H^n(K, L)$ are isomorphic to $H^n(B_K, L)$ where B_K is the topological classifying space for the group K .

Suppose K is $\mathfrak{S}_n$, the symmetric group on n letters. Then $H^*(\mathfrak{S}_n, L)$ is especially important. In the 1950's, work on cohomology operations, [29] and [30], showed the necessity for knowledge of $H^*(\mathfrak{S}_p, \mathbb{Z}/p)$. The construction of the mod p Steenrod operations depends on properties of $\mathfrak{S}_p$. Furthermore the Adem relations were derived using the structure of $H^*(\mathfrak{S}_p, \mathbb{Z}/p)$.

If L is a ring then $H^*(K, L)$ is a graded ring. The homology of symmetric products, [9], [17], [20], [21], and [28], computed the groups $H^i(\mathfrak{S}_n, \mathbb{Z}/p)$ as $\mathbb{Z}/p$ vector spaces. The graded ring structure, which was not analyzed, becomes important in later problems.

There is an interesting link that ties $\mathfrak{S}_n$ to SG . Recall $Q(S^0) = \text{dir lim } \Omega^n S^n$ is the space of "infinite loops of S^∞ " and $SG = \text{dir lim } SG_n$ where SG_n is the space of degree $+1$ homotopy equivalences of S^{n-1} . SG is homotopy equivalent to the $+1$ component of $Q(S^0)$.

THEOREM. (1) *There is a canonical map $\omega: B_{\mathfrak{S}_\infty} = \text{dir lim } B_{\mathfrak{S}_n} \rightarrow Q(S^0)_0$ inducing integral and mod p homology isomorphisms.*

(2) *The inclusions $\mathfrak{S}_n \times \mathfrak{S}_m \rightarrow \mathfrak{S}_{n+m}$ give $H_*(\mathfrak{S}_\infty)$ the structure of an algebra. ω_* is an algebra isomorphism and a Hopf algebra isomorphism mod p where $H_*(Q(S^0)_0)$ is an algebra under the loop sum product.*

Received by the editors October 21, 1975 and, in revised form, March 29, 1977.

AMS (MOS) subject classifications (1970). Primary 18H10; Secondary 55F40.

Key words and phrases. Cohomology of groups, classifying spaces, Steenrod algebra.

The above theorem is contained in the work of many people including [10], [16], [22], [24], [25].

Thus B_{S_∞} properly interpreted is a model for SG .

In all that follows let p be an *odd* prime. We will write $H^*(K)$ for $H^*(K, \mathbb{Z}/p)$. $H^*(K, \mathbb{Z}, p)$ is, by definition, [5], the p -primary component of $H^*(K, \mathbb{Z})$. In [4] the algebra structure of $H^*(S_{p^2})$ is computed but the arguments do not generalize to S_{p^i} ; $i \geq 3$. The main results of this paper are the calculations of $H^*(S_n)$ and $H^*(SG)$ as algebras, determination of the action of the Steenrod algebra, $\mathcal{Q}(p)$, on $H^*(S_n)$ and $H^*(SG)$ and integral analysis of $H^*(S_n, \mathbb{Z}, p)$ and $H^*(SG, \mathbb{Z}, p)$.

This paper is essentially my Stanford University Ph. D. thesis written under the direction of R. James Milgram, whom I would like to thank for his advice and encouragement. I would also like to thank the referee for his numerous helpful comments including shorter proofs for two of the propositions in §II. In addition after submission of this paper I learned that Benjamin Cooper [35] and Hùynh Mùì [36] have also studied $H^*(S_{p^i})$.

I. Statement of results. It is well known that a p -Sylow subgroup K_p of a finite group K contains all the p -primary homology information; more precisely, $H^*(K)$ and $H^*(K, \mathbb{Z}, p)$ are isomorphic to subrings of $H^*(K_p)$ and $H^*(K_p, \mathbb{Z}, p)$ respectively, which are invariant under the action of certain automorphisms. It is also well known, [6], that a p -Sylow subgroup of S_{p^i} is isomorphic to $wr^i \mathbb{Z}/p$, the i -fold wreath product of $\mathbb{Z}/p$. In the next section we examine a specific embedding of $wr^i \mathbb{Z}/p$ in S_{p^i} and show the existence of an $H^*()$ detecting family consisting of subgroups of the form $\times^m \mathbb{Z}/p$. In fact we have the following subgroups and natural inclusions: $k_{j,i}: T_{j,i} \rightarrow S_{p^i}$ for $1 \leq j \leq i$ and the map $k_i^* = \prod_{j=1}^i k_{j,i}^*: H^*(S_{p^i}) \rightarrow \prod_{j=1}^i H^*(T_{j,i})$, where $T_{j,i} = \times^{p^{i-j}} (\times^j \mathbb{Z}/p)$.

The first theorems compute the images of $k_{j,i}^*$'s and the map k_i^* . We show that k_i^* detects a set of multiplicative generators for $H^*(S_{p^i})$ whose relations are trivial to compute. Hence the map k_i^* determines $H^*(S_{p^i})$. Later for simplicity we will want to identify $u \in H^*(S_{p^i})$ with its natural image $k_{j,i}^*(u) \in H^*(T_{j,i})$ but we must wait until Theorems A–D have been stated to avoid possible confusion.

Recall $H^*(\times^k \mathbb{Z}/p) = E(e_1, \dots, e_k) \otimes P(b_1, \dots, b_k)$ with $\text{degree}(e_m) = 1$, $\text{degree}(b_m) = 2$ for all m . Furthermore $\beta_p(e_m) = b_m$, where β_p is the Bockstein operator associated with the exact coefficient sequence $0 \rightarrow \mathbb{Z}/p \rightarrow \mathbb{Z}/p^2 \rightarrow \mathbb{Z}/p \rightarrow 0$.

Consider the following classes in $H^*(\times^i \mathbb{Z}/p)$: (a matrix cohomology class will always mean the cohomology class given by the formal determinant of that matrix)

$$L_i = \begin{vmatrix} b_1^{p^{i-1}} & \cdots & b_i^{p^{i-1}} \\ \vdots & \vdots & \vdots \\ b_1^{p^r} & \cdots & b_i^{p^r} \\ \vdots & \vdots & \vdots \\ b_1 & \cdots & b_i \end{vmatrix} \quad \begin{array}{l} \text{i.e. the } k, j \text{ entry of } L_i \\ \text{is } b_j^{p^k} \text{ (} 0 \leq r \leq i-1 \text{).} \end{array}$$

$$Q_{j,i} = \frac{\begin{vmatrix} b_1^{p^j} & \cdots & b_i^{p^j} \\ \vdots & \vdots & \vdots \\ \widehat{b_1^{p^j}} & \cdots & \widehat{b_i^{p^j}} \\ \vdots & \vdots & \vdots \\ b_1 & \cdots & b_i \end{vmatrix}}{L_i} \quad \begin{array}{l} \text{i.e. the } b^{p^j} \text{ row of the} \\ \text{numerator is omitted} \\ \text{(} 1 \leq j \leq i-1 \text{).} \end{array}$$

$$\underline{L}_i = \begin{vmatrix} b_1^{p^{i-1}} & \cdots & b_i^{p^{i-1}} \\ \vdots & \vdots & \vdots \\ b_1^p & \cdots & b_i^p \\ e_1 & \cdots & e_i \end{vmatrix} \quad \begin{array}{l} \text{i.e. } \underline{L}_i \text{ is the } L_i \text{ determinant} \\ \text{with the } b_1 \cdots b_i \text{ row replaced} \\ \text{by the row } e_1 \cdots e_i. \end{array}$$

$$M_{j,i} = \begin{vmatrix} b_1^{p^{i-1}} & \cdots & b_i^{p^{i-1}} \\ \vdots & \vdots & \vdots \\ \widehat{b_1^{p^j}} & \cdots & \widehat{b_i^{p^j}} \\ \vdots & \vdots & \vdots \\ b_1 & \cdots & b_i \\ e_1 & \cdots & e_i \end{vmatrix} \quad \begin{array}{l} \text{i.e. the } b^{p^j} \text{ row is} \\ \text{omitted (} 1 \leq j \leq i-1 \text{).} \end{array}$$

Note. (i) If $i = 1$ then $L_1 = b_1$ and $\underline{L}_1 = e_1$ are the only two classes defined.

(ii) [19] proved $Q_{j,i}$ is integral, not merely rational, mod p . See appendix for proof.

$\mathcal{S}_{p^i}$ can be thought of as the permutations of the point set $\Pi^i Z/p$. Let $k_{i,i}$: $T_{i,i} = \times^i Z/p \rightarrow \{\text{permutations of } \Pi^i Z/p\}$ be defined by: $k_{i,i}(a_1, \dots, a_i)$ sends $(b_1, \dots, b_i)$ to $(a_1 + b_1, \dots, a_i + b_i)$ where Z/p is written additively. Then $k_{i,i}$ is seen to be equivalent to the adjoint representation (2.5) and

includes $T_{i,i}$ in $\mathcal{S}_{p^i}$. The normalizer N of $k_{i,i}(T_{i,i})$ in $\mathcal{S}_{p^i}$ maps onto $GL(i, Z/p)$ (2.10) and induces an action on $H^*(T_{i,i})$ as follows. If $\cup_x$ in $GL(i, Z/p)$ represents the coset $xT_{i,i}$ in N then the homomorphism $\text{ad}_x: H^*(T_{i,i}) \rightarrow H^*(T_{i,i})$ operates as follows: $\text{ad}_x(e_m) = \cup_x e_m$, $\text{ad}_x(b_m) = \cup_x b_m$ where e_m , b_m are treated as the vectors $(0, \dots, e, \dots, 0)$ and $(0, \dots, b, \dots, 0)$ in $H^*(T_{i,i})$ with nonzero entries in the m th place. Hence ad_x operates on the above determinant classes via the determinant function; that is, $\text{ad}_x(L_i) = \det(\cup_x L_i)$. By 2.13 image $k_{i,i}^*$ is contained in $H^*(T_{i,i})^{GL(i, Z/p)}$.

Let $\mathcal{W}_i$ be the algebra $E(L_1 L_1^{p-2}) \otimes P(L_1^{p-1})$. For i greater than 1 let $\mathcal{W}_i$ be the subalgebra of $H^*(T_{i,i})$ generated by: $1, L_i^{p-1}, Q_{j,i}, L_i L_i^{p-2}, M_{j,i} L_i^{p-2}, M_{j,i} L_i L_i^{p-3}, M_{j,i} M_{h,i} L_i^{p-3}$ with $1 \leq j, h \leq i-1$ and $j < h$. $\mathcal{W}_i$ is contained in $H^*(T_{i,i})^{GL(i, Z/p)}$ (2.12). Then $\mathcal{W}_i$ contains the polynomial algebra $P(L_i^{p-1}, Q_{1,i}, Q_{2,i}, \dots, Q_{i-1,i})$ and all other generators of $\mathcal{W}_i$ are exterior. However the algebra they generate is not an exterior subalgebra as there are zero products. The multiplication of these exterior products is determined by the relations:

$$(1) \underline{L}_i^2 = M_{j,i}^2 = 0, 1 \leq j \leq i-1,$$

$$(2) \underline{L}_i M_{1,i} M_{2,i} \cdots M_{i-1,i} \neq 0.$$

For example $(M_{2,i} M_{3,i} L_i^{p-3})(M_{2,i} M_{5,i} L_i^{p-3}) = 0$.

THEOREM A. image $k_{i,i}^* \cong \mathcal{W}_i$.

EXAMPLES. (i) If $i = 1$ then $0 \rightarrow H^*(\mathcal{S}_p) \rightarrow k_{1,1}^* H^*(Z/p)$ where $H^*(Z/p) \cong E(L_1) \otimes P(L_1)$ and $H^*(\mathcal{S}_p) \cong E(L_1 L_1^{p-2}) \otimes P(L_1^{p-1})$.

(ii) If $i = 2$ the results of [4] are obtained.

(iii) Let $p = 3, i = 3$ then $k_{3,3}^*: H^*(\mathcal{S}_{27}) \rightarrow H^*(Z/3 \times Z/3 \times Z/3)$ and image $k_{3,3}^*$ is generated by:

(1) polynomial generators

$$L_3^2 = \begin{vmatrix} b_1^9 & b_2^9 & b_3^9 \\ b_1^3 & b_2^3 & b_3^3 \\ b_1 & b_2 & b_3 \end{vmatrix}^2, \quad Q_{1,3} = \frac{\begin{vmatrix} b_1^{27} & b_2^{27} & b_3^{27} \\ b_1^9 & b_2^9 & b_3^9 \\ b_1 & b_2 & b_3 \end{vmatrix}}{L_3},$$

$$Q_{2,3} = \frac{\begin{vmatrix} b_1^{27} & b_2^{27} & b_3^{27} \\ b_1^3 & b_2^3 & b_3^3 \\ b_1 & b_2 & b_3 \end{vmatrix}}{L_3}.$$

(2) exterior generators

$$M_{1,3}M_{2,3} = \begin{vmatrix} b_1^9 & b_2^9 & b_3^9 \\ b_1 & b_2 & b_3 \\ e_1 & e_2 & e_3 \end{vmatrix} \begin{vmatrix} b_1^3 & b_2^3 & b_3^3 \\ b_1 & b_2 & b_3 \\ e_1 & e_2 & e_3 \end{vmatrix},$$

$$M_{1,3}\underline{L}_3, M_{1,3}L_3, M_{2,3}\underline{L}_3, M_{2,3}L_3, \underline{L}_3L_3.$$

(3) the relations that any product of exterior generators is zero except

$$(a) (M_{1,3}M_{2,3})(\underline{L}_3L_3) = - (M_{1,3}\underline{L}_3)(M_{2,3}L_3) = (M_{2,3}\underline{L}_3)(M_{1,3}L_3),$$

$$(b) (M_{1,3}L_3)(\underline{L}_3L_3) = (M_{1,3}\underline{L}_3)L_3^2,$$

$$(c) (M_{2,3}L_3)(\underline{L}_3L_3) = (M_{2,3}\underline{L}_3)L_3^2,$$

$$(d) (M_{1,3}L_3)(M_{2,3}L_3) = (M_{1,3}M_{2,3})L_3^2.$$

The proof of Theorem A depends, in part, on [17] and a counting argument. As noted above the classes in image $k_{i,i}^*$ are $GL(i, Z/p)$ invariant. A calculation and [8] show $P(b_1, \dots, b_i)^{GL(i, Z/p)}$ is isomorphic to the polynomial subalgebra of image $k_{i,i}^*$. For $i = 2$, [4] shows

$$(E(e_1, e_2) \otimes P(b_1, b_2))^{GL(2, Z/p)} \cong H^*(Z/p \times Z/p)^{GL(2, Z/p)} \cong \text{image } k_{2,2}^*.$$

If $p \geq 5$, $i \geq 3$ then $(E(e_1, \dots, e_i) \otimes P(b_1, \dots, b_i))^{GL(i, Z/p)}$ properly contains image $k_{i,i}^*$; for example, $M_{1,i}M_{2,i}\underline{L}_iL_i^{p-4}$ is not in image $k_{i,i}^*$. For $p = 3$, $i \geq 3$, it is unknown if image $k_{i,i}^*$ equals the ring of invariants.

Consider the inclusion $\times_{m=1}^p (\mathfrak{S}_{p^{i-1}})_m \rightarrow \mathfrak{S}_{p^i}$ where $(\mathfrak{S}_{p^{i-1}})_m$ permutes the p^{i-1} letters $((m-1)p^{i-1} + 1, \dots, mp^{i-1})$. Then let $k_{i-1,i}: T_{i-1,i} \rightarrow \mathfrak{S}_{p^i}$ be the composition $I_{i-1}(\times_{m=1}^p (k_{i-1,i-1})_m)$. More generally let $k_{j,i}: T_{j,i} \rightarrow \mathfrak{S}_{p^i}$ be the composition $I_j(\times_{m=1}^{p^{i-j}} (k_{j,i})_m)$ where I_j is the inclusion $\times_{m=1}^{p^{i-j}} (\mathfrak{S}_{p^j})_m \rightarrow \mathfrak{S}_{p^i}$ given by letting $(\mathfrak{S}_{p^j})_m$ permute the m th block of p^j letters.

Let $1 \leq j \leq i$, then $\mathfrak{S}_{p^{i-j}}$ operates on $T_{j,i}$ and on the algebra $\bigotimes_{m=1}^{p^{i-j}} (\mathcal{U}_j)_m$ contained in $H^*(T_{j,i}) \cong \bigotimes_{m=1}^{p^{i-j}} (H^*(\times^j Z/p))_m$ by permuting the p^{i-j} copies of $\times^j Z/p$.

THEOREM B. For $1 \leq j \leq i$ image $k_{j,i}^*$ is isomorphic to the algebra of $\mathfrak{S}_{p^{i-j}}$ invariant classes of $\bigotimes_{m=1}^{p^{i-j}} (\mathcal{U}_j)_m$.

Notation. Let $u_m \in (\mathcal{U}_j)_m$ then $\mathfrak{S}\langle u_1, u_2, \dots, u_{p^{i-j}} \rangle$ is the $\mathfrak{S}_{p^{i-j}}$ invariant class generated by $u_1 u_2 \dots u_{p^{i-j}}$ (u_m is allowed to be $1 \in H^0(\times^j Z/p)$). If u_1 is odd dimensional then $\mathfrak{S}\langle u_1, u_1, \dots, u_{p^{i-j}} \rangle = 0$.

EXAMPLES. (i) image $k_{1,1}^*$ is generated by:

$$\begin{aligned} A_{k,i} &= \sum_{m=1}^{p^{i-1}} (\underline{L}_1 L_1^{(p-2)+k(p-1)})_m \\ &= \mathfrak{S}\langle (\underline{L}_1 L_1^{(p-2)+k(p-1)}), 1, \dots, 1 \rangle, \text{ for } 0 \leq k \leq p^{i-1} - 1, \end{aligned}$$

and

$$B_{k,i} = \sum (L_1^{p-1})_{m_1} (L_1^{p-1})_{m_2} \dots (L_1^{p-1})_{m_k},$$

where $1 \leq k \leq p^{i-1}$ and the sum runs over all sequences $1 \leq m_1 < m_2$

$\langle \dots \langle m_k \leq p^{i-1}$. Thus $B_{k,i} = \mathfrak{S} \langle L_1^{p-1}, L_1^{p-1}, \dots, L_1^{p-1}, 1, \dots, 1 \rangle$ where L_1^{p-1} appears k times.

(ii) Let $p = 3$, then $k_{2,3}^*$: $H^*(\mathfrak{S}_{27}) \rightarrow H^*(T_{2,3})$ and image $k_{2,3}^*$ is generated by:

$$\begin{array}{lll} \mathfrak{S} \langle \text{ext}, 1, 1 \rangle & \mathfrak{S} \langle \text{poly}, 1, 1 \rangle & \mathfrak{S} \langle \text{ext}, \text{poly}, 1 \rangle \\ \mathfrak{S} \langle M_{1,2} \underline{L}_2, M_{1,2} \underline{L}_2, M_{1,2} \underline{L}_2 \rangle & & \mathfrak{S} \langle \text{ext}, \text{poly}, \text{poly} \rangle \\ \mathfrak{S} \langle \text{poly}, \text{poly}, 1 \rangle & \mathfrak{S} \langle \text{poly}, \text{poly}, \text{poly} \rangle & \mathfrak{S} \langle \text{ext}, \text{ext}, \text{poly} \rangle \end{array}$$

where

(a) ext runs through $M_{1,2} \underline{L}_2$, $M_{1,2} L_2$, and $\underline{L}_2 L_2$.

(b) poly runs through L_2^2 and $Q_{1,2}$.

(c) As $M_{1,2} L_2$ and $\underline{L}_2 L_2$ are odd dimensional neither can appear twice in any $\mathfrak{S} \langle -, -, - \rangle$. For example $\mathfrak{S} \langle \underline{L}_2 L_2, \underline{L}_2 L_2, 1 \rangle = 0$. Note that $\mathfrak{S} \langle M_{1,2} \underline{L}_2, 1, 1 \rangle$ has height 3 while $\mathfrak{S} \langle M_{1,2} L_2, 1, 1 \rangle$ is exterior.

(iii) In image $k_{2,i}^*$ the classes

$$\mathfrak{S} \langle M_{1,2} \underline{L}_2 L_2^{p-3}, 1, \dots, 1 \rangle$$

and

$$\mathfrak{S} \langle (M_{1,2} \underline{L}_2 L_2^{p-3})_1, \dots, (M_{1,2} \underline{L}_2 L_2^{p-3})_p, 1, \dots, 1 \rangle$$

have height p while $\mathfrak{S} \langle M_{1,2} \underline{L}_2 L_2^{p-3}, \dots, M_{1,2} \underline{L}_2 L_2^{p-3} \rangle$ is exterior. This pattern generalizes to image $k_{j,i}^*$, $3 \leq j \leq i-1$, in the obvious way.

Note. Example (iii) shows how all even dimension exterior generators in $\mathfrak{W}_j$ build classes in $H^*(T_{j,i})$ which are the images under $k_{j,i}^*$ of classes $u \in H^*(\mathfrak{S}_{p^i})$ where each u generates a truncated polynomial algebra of height p in $H^*(\mathfrak{S}_n)$. These are the truncated polynomial algebras described in [22].

Let $u \in H^*(\mathfrak{S}_{p^i})$ then $k_i^*(u) = (k_{1,i}^*(u), \dots, k_{i,i}^*(u))$ and the algebra structure restricted to these detecting groups is compatible with component-wise projection. Clearly to calculate $H^*(\mathfrak{S}_{p^i})$ we must know when a class $u \in H^*(\mathfrak{S}_{p^i})$ has nontrivial image under more than one $k_{i,j}^*$.

DEFINITION. $u \in H^*(\mathfrak{S}_{p^i})$ is a multiple image class if and only if $k_{j,i}^*(u) \neq 0$ for at least two different values of j .

Given $u_1, u_2 \in H^*(\mathfrak{S}_{p^i})$ with u_1 detected only by $k_{j_1,i}^*$ and u_2 detected only by $k_{j_2,i}^*$ with $j_1 \neq j_2$ then $u_1 + u_2$ is a multiple image class. However this type of multiple image class is decomposable as a sum of classes and thus is a "trivial" multiple image class. The next three definitions and following theorem give all "nontrivial"; i.e., indecomposable, multiple image classes.

DEFINITION. $\mathfrak{N}_i$ is the subalgebra contained in $\mathfrak{W}_i$ generated by $1, M_{g,i} M_{h,i} L_i^{p-3}, Q_{h,i}, 1 \leq g, h \leq i-1, g < h$.

DEFINITION. Given $x_{m,j} \in \mathfrak{N}_j$ we define $x_{m,j-1} \in \mathfrak{W}_{j-1}$ as follows:

(a) If $x_{m,j} = 1$ then $x_{m,j-1} = 1$.

(b) If $x_{mj} = Q_{hj}$ then $x_{m,j-1} = Q_{h-1,j-1}$, for $2 \leq j \leq i$ and $1 \leq h \leq j-1$ with the convention $Q_{0,j-1} = L_{j-1}^{p-1}$.

(c) If $x_{mj} = M_{gj}M_{hj}L_j^{p-3}$ then $x_{m,j-1} = -M_{g-1,j-1}M_{h-1,j-1}L_{j-1}^{p-3}$, for $3 \leq j \leq i$, $0 < g, h < j$ and $g < h$ with the convention $M_{0,j-1} = L_{j-1}$.

(d) If $x_{mj} = x'_{mj}x''_{mj}$ then $x_{m-1,j-1} = x'_{m-1,j-1}x''_{m-1,j-1}$.

Note. (a) through (d) define a unique class $x_{m,j-1}$ for every $x_{mj} \in \mathfrak{N}_j$.

DEFINITION. $u \in H^*(\mathfrak{S}_n)$ is sum indecomposable if and only if $u = u_1 + u_2$ for $u_1, u_2 \in H^*(\mathfrak{S}_n)$ implies u_1 or u_2 is zero.

THEOREM C. Suppose $u \in H^*(\mathfrak{S}_{p^i})$ is both sum indecomposable and a multiple image class. Further suppose j is the largest integer such that $k_{j,i}^*(u) \neq 0$. Then

$$k_{j,i}^*(u) = \mathfrak{S} \langle x_{1,j}, \dots, x_{p^{i-j},j} \rangle$$

with $x_{m,j} \in \mathfrak{N}_j$ for $1 \leq m \leq p^{1-j}$, and

$$k_{j-1,i}^*(u) = \mathfrak{S} \langle x_{1,j-1}, \dots, x_{1,j-1}, \dots, x_{p^{i-j,j-1}}, \dots, x_{p^{i-j,j-1}} \rangle$$

where each $x_{m,j-1}$ is as defined above and appears p times in $k_{j-1,i}^*(u)$. If $j-1 \geq 2$ and each $x_{m,j-1} \in \mathfrak{N}_{j-1}$ (not just $\mathfrak{N}_{j-1}$) then $k_{j-2,i}^*(u) \neq 0$ and may be obtained from $k_{j-1,i}^*(u)$ precisely as $k_{j-1,i}^*(u)$ was obtained from $k_{j,i}^*(u)$. In fact this iteration continues r times until either $j-r=2$ or $x_{m,j-r} \notin \mathfrak{N}_{j-r}$ when $k_{j-(r+1),i}^*(u) = 0$ for all $t > 0$. Thus u has $r+1$ nontrivial images in the detecting groups: $k_{j-s,i}^*$ for $0 \leq s \leq r$.

EXAMPLE. For $H^*(\mathfrak{S}_{27}, \mathbb{Z}/3)$ the only sum-indecomposable multiple image classes of k_0^* occurring as generators in the examples after Theorems A and B are:

$$\begin{aligned} & (B_9, (Q_{1,2})_1(Q_{1,2})_2(Q_{1,2})_3, Q_{2,3}), \\ & (0, (L_2^2)_1(L_2^2)_2(L_2^2)_3, Q_{1,3}), \\ & (0, (M_{1,2}L_2)_1(M_{1,2}L_2)_2(M_{1,2}L_2)_3, -M_{1,3}M_{2,3}), \\ & (B_3, \mathfrak{S} \langle Q_{1,2}, 1, 1 \rangle, 0), \\ & (B_6, \mathfrak{S} \langle Q_{1,2}, Q_{1,2}, 1 \rangle, 0). \end{aligned}$$

Consider $u_1 u_2$ in $H^*(\mathfrak{S}_{p^3})$ where $k_3^*(u_1) = (\mathfrak{S} \langle L_1^{p-1}, 1, \dots, 1 \rangle, 0, 0)$ and $k_3^*(u_2) = (0, \mathfrak{S} \langle L_2^{p-1}, 1, \dots, 1 \rangle, 0)$. Then $k_3^*(u_1 u_2) = 0$ but in fact $u_1 u_2 \neq 0$ in $H^*(\mathfrak{S}_{p^3})$ and $u_1 u_2$ is detected by subgroups of the form $T_1 \times T_2 \times \dots \times T_p$ where $T_n = T_{1,2}$ or $T_{2,2}$ and both $T_{1,2}$ and $T_{2,2}$ must occur at least once. These detecting groups are included in $\mathfrak{S}_{p^3}$ through $\times^p(\mathfrak{S}_{p^2})$. More generally a nonsymmetric detecting group, $\times_{n=1}^p (\times_{m=1}^t (T_{r_m, s_m}))_n$ of $\mathfrak{S}_{p^i}$ is a product of detecting groups of $\mathfrak{S}_{p^{i-1}}$ included in $\mathfrak{S}_{p^i}$ through $\times^p(\mathfrak{S}_{p^{i-1}})$ where $T_{r_1, s_1} \neq T_{r_2, s_2}$ for some r_1, r_2, s_1 and s_2 . These nonsymmetric detecting groups detect all classes $u \in H^*(\mathfrak{S}_{p^i})$ not detected by the map k_i^* as stated in Theorem D. First we need

DEFINITION. Let $u \in H^*(\mathcal{S}_{p^i})$ and $n < p^i$. Then we have the natural inclusion $I_{p^i,n}: \mathcal{S}_n \hookrightarrow \mathcal{S}_{p^i}$. We say u restricts nonzero to $\mathcal{S}_n$ if and only if $I_{p^i,n}^*(u) \neq 0$. For notational convenience we write u for both the class in $H^*(\mathcal{S}_{p^i})$ and the restriction in $H^*(\mathcal{S}_n)$.

THEOREM D. (1) The classes in $H^*(\mathcal{S}_{p^i})$ not detected by k_i^* are products of classes that are detected by k_i^* .

(2) Let $u_m \in H^*(\mathcal{S}_{p^i})$. Suppose $k_i^*(u_m) \neq 0$, $\prod_{m=1}^r k_i^*(u_m) = 0$ and let n_m be the smallest power of p such that u_m restricts nonzero to $H^*(\mathcal{S}_{n_m})$. Then $\prod_{m=1}^r u_m \neq 0$ in $H^*(\mathcal{S}_{p^i})$ unless:

(a) $u_{m_1} = u_{m_2}$ is an odd dimensional exterior class in $H^*(\mathcal{S}_{n_{m_1}})$, for some $1 \leq m_1 < m_2 \leq r$.

(b) $u_{m_1} = u_{m_2} = \cdots = u_{m_p}$ is an even dimensional exterior class in $H^*(\mathcal{S}_{n_{m_1}})$ for some $1 \leq m_1 < m_2 < \cdots < m_p \leq r$ or

(c) $\mathcal{S}_{n_1} \times \cdots \times \mathcal{S}_{n_r}$ is not contained in $\mathcal{S}_{p^i}$.

Note. The classes u_{m_1} appearing in condition (b) are the generators for the truncated polynomial algebras described in example (iii) after Theorem B.

Thus every $u \in H^*(\mathcal{S}_{p^i})$ is expressible as a sum of monomials $\sum a(u_1, \dots, u_r) u_1 \otimes \cdots \otimes u_r$ where $a(u_1, \dots, u_r) \in \mathbb{Z}/p$, $u_i \in H^*(\mathcal{S}_{p^i})$ with $k_i(u_i) \neq 0$ for all i .

DEFINITION. $u \in H^*(\mathcal{S}_{p^i})$ is proper if and only if $u = \sum a(u_1, \dots, u_r) u_1 \otimes \cdots \otimes u_r$ with $k_i^*(u_1 \otimes \cdots \otimes u_r) \neq 0$ for each monomial in the sum.

Thus Theorems A through D compute $H^*(\mathcal{S}_{p^i})$ and from this point on we will identify elements of $H^*(\mathcal{S}_{p^i})$ with their image under k_i^* . That is $L_i^{p-1} Q_{j,i} \in H^*(\mathcal{S}_{p^i})$ is the unique proper class $u \in H^*(\mathcal{S}_{p^i})$ such that $k_i^*(u) = (0, \dots, 0, L_i^{p-1} Q_{j,i})$. Care must be taken with multiple image classes under this identification. Notice, by Theorem C, that $Q_{1,i} \in H^*(\mathcal{S}_{p^i})$ is the unique proper class $u \in H^*(\mathcal{S}_{p^i})$ such that $k_i^*(u) = (0, \dots, 0, \langle L_{i-1}^{p-1}, \dots, L_{i-1}^{p-1} \rangle, Q_{1,i})$.

Since

$$\mathcal{P}^j(b^{p^k}) = \begin{cases} b^{p^k} & \text{if } j = 0, \\ b^{p^{k+1}} & \text{if } j = p^k, \\ 0 & \text{otherwise,} \end{cases}$$

it is easy to determine the action of the Steenrod algebra $\mathcal{Q}(p)$ on $H^*(\mathcal{S}_{p^i})$. Consider $M_{1,3}L_3$ in $H^{47}(\mathcal{S}_{27}, \mathbb{Z}/3)$. Then

$$\mathcal{P}^1 \left[\begin{vmatrix} b_1^9 & b_2^9 & b_3^9 \\ b_1 & b_2 & b_3 \\ e_1 & e_2 & e_3 \end{vmatrix} \middle| \begin{vmatrix} b_1^9 & b_2^9 & b_3^9 \\ b_1^3 & b_2^3 & b_3^3 \\ b_1 & b_2 & b_3 \end{vmatrix} \right] = \begin{vmatrix} b_1^9 & b_2^9 & b_3^9 \\ b_1^3 & b_2^3 & b_3^3 \\ e_1 & e_2 & e_3 \end{vmatrix} \middle| \begin{vmatrix} b_1^9 & b_2^9 & b_3^9 \\ b_1^3 & b_2^3 & b_3^3 \\ b_1 & b_2 & b_3 \end{vmatrix} = \underline{L}_3 L_3.$$

This computation involved use of the Cartan formula; however, all terms except the first are zero. The next theorem describes the $\mathcal{Q}(p)$ action on $\mathcal{W}_i$. Note the polynomial subalgebra of $\mathcal{W}_i$ is closed under the $\mathcal{Q}(p)$ action while a class in the ideal generated by the exterior generators of $\mathcal{W}_i$ may be "bocksteined" into the polynomial algebra; e.g., $\beta \mathcal{P}^1(M_{1,i}L_i^{p-2}) = L_1^{p-1}$ for $i > 1$. Using the Cartan formula and the following theorem it is trivial to compute the $\mathcal{Q}(p)$ action on all the detecting groups.

THEOREM E. *The following relations and the Cartan formula describe the $\mathcal{Q}(p)$ action on $\mathcal{W}_i$.*

- (1) $\mathcal{P}^{p^{h-1}}(M_{j,i}M_{h,i}L_i^{p-3}) = M_{j,i}M_{h-1,i}L_i^{p-3}$, $j > h$ and $M_{0,i} = \underline{L}_i$,
- (2) $\mathcal{P}^{p^{j-1}}(M_{j,i}M_{h,i}L_i^{p-3}) = M_{j-1,i}M_{h,i}L_i^{p-3}$, $j > h$ and $M_{0,i} = \underline{L}_i$,
- (3) $\beta(\underline{L}_i) = L_i$,
- (4) $\mathcal{P}^{p^{h-1}}(Q_{h,i}) = Q_{h-1,i}$, with $Q_{0,i} = L_i^{p-1}$,
- (5) $\mathcal{P}^{p^{i-1}}(L_i^{p-1}) = -Q_{i-1,i}L_i^{p-1}$ for $i > 1$ while $\mathcal{P}^j(L_1^{p-1}) = (p^{-1})L_1^{(p-1)(j+1)}$ for $j \leq p-1$.
- (6) $\mathcal{P}^{p^{i-1}}(M_{i-1,i}\underline{L}_iL_i^{p-3}) = (p-2)(M_{i-1,i}\underline{L}_iL_i^{p-3})(Q_{i-1,i})$,
 $\mathcal{P}^{p^{i-1}}(M_{i-1,i}L_i^{p-2}) = (p-2)(M_{i-1,i}L_i^{p-2})(Q_{i-1,i})$.

The following diagram is conceptually helpful.

$$\begin{array}{ccccccc}
 & & & & & & \underline{LL}^{p-2} \xrightarrow{\beta} L^{p-1} \\
 & & & & & & \uparrow \mathcal{P}^1 \quad \uparrow \mathcal{P}^1 \\
 & & & & & & M_1 \underline{LL}^{p-3} \xrightarrow{-\beta} M_1 L^{p-2} \quad Q_1 \\
 & & & & & & \uparrow \mathcal{P}^p \quad \uparrow \mathcal{P}^p \\
 & & & & & & M_2 M_1 L^{p-3} \xrightarrow{\mathcal{P}^1} M_2 \underline{LL}^{p-3} \xrightarrow{-\beta} M_2 L^{p-2} \quad Q_2 \\
 & & & & & & \uparrow \mathcal{P}^{p^2} \quad \uparrow \mathcal{P}^{p^2} \quad \uparrow \mathcal{P}^{p^2} \quad \uparrow \mathcal{P}^{p^2} \\
 & & & & & & \vdots \quad \vdots \quad \vdots \quad \vdots \\
 & & & & & & M_{i-3} M_{i-4} L^{p-3} \xrightarrow{\mathcal{P}^{p^{i-5}}} \vdots \quad \vdots \\
 & & & & & & \uparrow \mathcal{P}^{p^{i-3}} \quad \uparrow \mathcal{P}^{p^{i-3}} \quad \uparrow \mathcal{P}^{p^{i-3}} \quad \uparrow \mathcal{P}^{p^{i-3}} \\
 & & & & & & M_{i-2} M_{i-3} L^{p-3} \xrightarrow{\mathcal{P}^{p^{i-4}}} M_{i-2} M_{i-4} L^{p-3} \xrightarrow{\mathcal{P}^{p^{i-5}}} \dots \xrightarrow{\mathcal{P}^1} M_{i-2} \underline{LL}^{p-3} \xrightarrow{-\beta} M_{i-2} L^{p-2} \quad Q_{i-2} \\
 & & & & & & \uparrow \mathcal{P}^{p^{i-2}} \quad \uparrow \mathcal{P}^{p^{i-2}} \quad \uparrow \mathcal{P}^{p^{i-2}} \quad \uparrow \mathcal{P}^{p^{i-2}} \\
 M_{i-1} M_{i-2} L^{p-3} \xrightarrow{\mathcal{P}^{p^{i-3}}} M_{i-1} M_{i-3} L^{p-3} \xrightarrow{\mathcal{P}^{p^{i-4}}} M_{i-1} M_{i-4} L^{p-3} \xrightarrow{\mathcal{P}^{p^{i-5}}} \dots \xrightarrow{\mathcal{P}^1} M_{i-1} \underline{LL}^{p-3} \xrightarrow{-\beta} M_{i-1} L^{p-2} \quad Q_{i-1}
 \end{array}$$

THE ACTION OF $\mathcal{Q}(p)$ ON THE GENERATORS OF $\mathcal{W}_i$

EXAMPLES. (i) Consider $A = (0, \mathcal{S} \langle M_{1,2}\underline{L}_2, M_{1,2}\underline{L}_2, M_{1,2}\underline{L}_2 \rangle, -M_{2,3}M_{1,3})$ in $H^{30}(\mathbb{S}_{27}, \mathbb{Z}/3)$. Then

$$\mathcal{P}^1 \beta(A) = (0, -\mathcal{S} \langle \underline{L}_2 L_2, M_{1,2}\underline{L}_2, M_{1,2}\underline{L}_2 \rangle, 0)$$

while

$$\beta \mathcal{P}^1(A) = (0, 0, M_{2,3}L_3).$$

(ii)

$$\begin{aligned}\mathcal{P}^{p^{i-2}}\mathcal{P}^{p^{i-1}}(M_{i-1,i}L_i^{p-2}) &= \mathcal{P}^{p^{i-2}}((p-2)(M_{i-1,i}L_i^{p-2})Q_{i-1,i}) \\ &= (p-2)[(M_{i-2,i}L_i^{p-2})Q_{i-1,i} + (M_{i-1,i}L_i^{p-2})Q_{i-2,i}].\end{aligned}$$

Let n be an arbitrary integer. Then n may be written uniquely as follows: $n = \sum_{j=0}^i a_j p^j$ with $0 \leq a_j \leq p-1$, $a_i \neq 0$. A p -Sylow subgroup K_p of S_n is isomorphic to

$$K_p = \times^{a_i} \left(\text{wr } Z/p \right) \times \times^{a_{i-1}} \left(\text{wr } Z/p \right) \times \cdots \times \times^{a_1} (Z/p).$$

To compute $H^*(\mathcal{S}_n)$ consider the following diagram of inclusions

$$\begin{array}{ccccc}\bar{K}_p = \times^{a_i} (\mathcal{S}_{p^i}) \times \cdots \times \times^{a_1} (\mathcal{S}_p) & \xrightarrow{J_n} & \mathcal{S}_n & \xrightarrow{I_{p^{i+1},n}} & \mathcal{S}_{p^{i+1}} \\ & & \uparrow & & \uparrow \\ & & K_p & \xrightarrow{i+1} & \text{wr } Z/p\end{array}$$

THEOREM F. (1) $I_{p^{i+1},n}^*$ is surjective.

(2) J_n^* is injective.

(3) $v \in \text{Image } J_n^*$ if and only if there exists a $u \in H^*(\mathcal{S}_{p^{i+1}})$ such that

$$\begin{aligned}(I_{p^{i+1},n} \circ J_n)^*(u) &= v \\ &= \sum \mathcal{S}\langle u_{i,1}, \dots, u_{i,a_i} \rangle \otimes \cdots \otimes \mathcal{S}\langle u_{1,1}, \dots, u_{1,a_1} \rangle \in H^*(\bar{K}_p)\end{aligned}$$

with $u_{t,r} \in H^*(\mathcal{S}_{p^t})$ for each r .

IMPORTANT EXAMPLE. Let $n = 2p^i$. We have

$$\begin{array}{ccccc}\mathcal{S}_{p^i} \times \mathcal{S}_{p^i} & \nearrow & \mathcal{S}_{2p^i} & \xrightarrow{I_{p^{i+1},2p^i}} & \mathcal{S}_{p^{i+1}} \\ & & \uparrow & & \uparrow \\ & & \text{wr } Z/p \times \text{wr } Z/p & \xrightarrow{i+1} & \text{wr } Z/p\end{array}$$

Recall the definition of $A_{k,i}$ and $B_{k,i}$ (see example (i) after Theorem B). Then $I_{p^{i+1},2p^i}^*(A_{k,i+1}) = A_{k,i} \otimes 1 + 1 \otimes A_{k,i} = \mathcal{S}\langle A_{k,i}, 1 \rangle$ for $1 \leq k \leq p^i$, while for

$p^i < k \leq p^{i+1}$, $I_{p^{i+1}, 2p^i}^*(A_{k,i+1}) = A'_{k,i} \otimes 1 + 1 \otimes A'_{k,i}$ where $A'_{k,i}$ is expressible in terms of $A_{r,i}$ and $B_{r,i}$ for $r < p^i$.

$$I_{p^{i+1}, 2p^i}^*(B_{k,i+1}) = \sum_{n+m=k} B_{n,i} \otimes B_{m,i} = \sum_{n=0}^{p^i} \mathbb{S} \langle B_{n,i}, B_{2p^i-n,i} \rangle,$$

where $0 \leq n, m \leq p^i$, $0 \leq k \leq 2p^i$, and $B_{0,i} = 1$. Similar restrictions occur on the other detecting groups. Thus the natural inclusions $\mathbb{S}_n \rightarrow \mathbb{S}_{n+1} \rightarrow \cdots \rightarrow \text{dir lim } \mathbb{S}_n$ are easily analyzed. Clearly

$$\mathbb{S}_{p^i} \rightarrow \mathbb{S}_{p^{i+1}} \rightarrow \cdots \rightarrow \text{dir lim } \mathbb{S}_{p^i}$$

is a cofinal direct limit and we have $H^*(\text{dir lim } \mathbb{S}_n) \cong H^*(\text{dir lim } \mathbb{S}_{p^i}) \cong \text{inv lim } H^*(\mathbb{S}_{p^i})$. Notice Theorem F implies $\text{inv lim } H^t(\mathbb{S}_{p^i})$ is attained for each t at a finite stage.

Recall the theorem stated in the introduction that ties $\text{dir lim } B_{\mathbb{S}_n}$ to $Q(S^0) = \text{dir lim } \Omega^n S^n$. Furthermore, if G_n is the set of homotopy equivalences of S^{n-1} then $G = \text{dir lim } G_n$ is homotopy equivalent to the union of the $+1$ and -1 components of $Q(S^0)$. Thus $\text{dir lim } B_{\mathbb{S}_n}$ properly interpreted is a model for G and we have:

$$\text{inv lim } H^*(\mathbb{S}_{p^i}) \cong H^*(Q(S^0)_0) \cong H^*(SG)$$

as algebras. Thus $H^*(SG)$ can be identified with "infinite symmetric sums" in the $\mathcal{W}_i$ algebras with the proper identifications; i.e., $\mathbb{S} \langle Q_{j,i}, 1, \dots \rangle \leftrightarrow \mathbb{S} \langle Q_{j-1,i-1}, \dots, Q_{j-1,i-1}, 1, \dots \rangle$. The $\mathcal{Q}(p)$ action on $H^*(SG)$ restricts to that on $B_{\mathbb{S}_{p^i}}$ for each i and there is a unique action which has this property. Theorem E describes the restriction of this action. Recall, [22] and [24], $H^*(\text{dir lim } \mathbb{S}_{p^i})$ is a Hopf algebra isomorphic to $H^*(Q(S^0)_0)$ with the coalgebra product on $H^*(\text{dir lim } \mathbb{S}_{p^i})$ induced by the inclusions $\mathbb{S}_{p^i} \times \mathbb{S}_{p^i} \rightarrow \mathbb{S}_{2p^i}$. Thus Theorem F gives the loop sum coalgebra map on $H^*(Q(S^0)_0)$.

As $Q(S^0)_0$ is an H -space it is possible to obtain integral information about $H^*(SG, Z, p)$ on $H^*(\mathbb{S}_{p^i}, Z, p)$ (see [14]). [2] gives a Hopf algebra Bockstein spectral sequence with

$$E_1 \cong H^*(\text{dir lim } \mathbb{S}_{p^i}, Z/p),$$

$$E_\infty \cong H^*(\text{dir lim } \mathbb{S}_{p^i}, Z, p)/\text{Torsion}.$$

Let $x, y \in \mathcal{W}_j$ and let

$$L_{n,j}(x: y_{n+1}, \dots, y_m, 1, \dots) = \mathbb{S} \langle xL_j^{p-1}, \dots, xL_j^{p-1}, y_{n+1}, \dots, y_m, 1, \dots \rangle$$

and

$$\begin{aligned} \underline{L}_{n,j}(x: y_{n+1}, \dots, y_m, 1, \dots) \\ = \mathbb{S} \langle x\underline{L}_j L_j^{p-2}, x\underline{L}_j^{p-1}, \dots, x\underline{L}_j^{p-1}, y_{n+1}, \dots, y_m, 1, \dots \rangle \end{aligned}$$

where $y_r \neq xL_j^{p-1}$ or $x\underline{L}_j L_j^{p-2}$. Note a class in $H^*(\text{dir lim } \mathbb{S}_{p^i})$ may have

more than one representation as $L_{n,j}(\dots)$ or $\underline{L}_{n,j}(\dots)$; for example,

$$\mathfrak{S}\langle xL_j^{p-1}, xL_j^{p-1}, yL_j^{p-1}, 1, \dots \rangle = L_{2,j}(x: y, 1, \dots) = L_{1,j}(y: x, 1, \dots).$$

THEOREM G. Let $k_{j,\infty}^* = \text{dir lim}_i k_{j,i}^*$ and let $u \in H^*(\text{dir lim } \mathfrak{S}_{p^i})$ be a proper class. Then there exists a smallest positive integer j such that $k_{j,\infty}^*(u) \neq 0$. Then $k_{j,\infty}^*(u) = \mathfrak{S}\langle x_1, \dots, x_m, 1, \dots \rangle$ and

(1) If some x_n contains an odd number of $M_{g,j}$ factors or if $k_{j,\infty}^*(u) = \underline{L}_{n,j}(\dots)$ or $L_{n,j}(\dots)$ for n not divisible by p then u is in the image or domain of β_p .

(2) Let $r \geq 2$. If $d_{r-1}(v) = u$ in E_{r-1} of the Bockstein spectral sequence and $k_{j,\infty}^*(u) = \mathfrak{S}\langle x_1, \dots, x_m, 1, \dots \rangle$ with no x_n containing an odd number of $M_{g,h}$ terms or the factor $\underline{L}_j$ then there exist v' and u' such that $d_r(v') = u'$ where $k_{j,\infty}^*(u') = \mathfrak{S}\langle x_1, \dots, x_1, \dots, x_m, \dots, x_m, 1, \dots \rangle + \Sigma u''$. Each x_h appears p times in $\mathfrak{S}\langle x_1, \dots, x_1, \dots, x_m, \dots, x_m, 1, \dots \rangle$ and each $u'' = \mathfrak{S}\langle x_1, \dots, x_t, 1, \dots \rangle$ with $t < pm$.

COROLLARY 1. Let $r \geq 2$ then

$$d_r(\underline{L}_{p^{r-1},j}(x: 1, \dots)) = L_{p^{r-1},j}(x: 1, \dots)$$

where x satisfies the same conditions as the x_n 's in (2) of Theorem G.

Let R_i be the inclusion $\mathfrak{S}_{p^i} \rightarrow \text{dir lim } \mathfrak{S}_{p^i}$ then R_i^* gives the Bockstein structure of $H^*(\mathfrak{S}_{p^i}, Z, p)$.

COROLLARY 2. $Q_{j,i} \in H^*(\mathfrak{S}_{p^i}, Z, p)$ has order p^{j+1} .

EXAMPLES. (i) $L_{p^r,j}(M_{1,j}\underline{L}_jL_j^{p-3}; 1, \dots)$ is a class of order p in $H^*(SG, Z, p)$, while $L_{p^r,j}(M_{1,j}M_{2,j}L_j^{p-3}; 1, \dots)$ is a class of order p^{r+1} .

(ii) $(B_6, \mathfrak{S}\langle Q_{1,2}, Q_{1,2}, 1 \rangle, 0) \in H^{24}(\mathfrak{S}_{27}, Z, 3)$ has order 9.

Finally the results of this paper have an application to cobordism theory. Although [3], [13] and [18] completely compute the PL and TOP cobordism ring at the prime 2, the odd case still has unanswered questions, notably the odd torsion in Ω^{PL} . Using results of [3], [15], [26], [27], [32], [34], [37], [38], [39] and this paper one may calculate the E^2 term of the Adams spectral sequence converging to $\Omega^{\text{PL}} \otimes Z_{(p)}$. Current joint work with H. Ligaard, J. P. May and R. J. Milgram computes this E^2 term and gives infinite families of nontrivial differentials of all orders in the spectral sequence.

II. The embedding and the detecting family.

2.1. DEFINITION. Let K be a finite group and L a subgroup of $\mathfrak{S}_n$ then $K \text{ wr } L$ is defined to be the group whose elements are

$$\{(f, g): f \text{ is a mapping of } (1, 2, \dots, n) \text{ into } K, g \in L\}$$

and whose multiplication is given by $(f, g)(f', g') = (ff'_g, gg')$, where $f_g(g(i)) = f(i)$ and $ff'(i) = f(i)f'(i)$.

2.2. DEFINITION. Let X be a space and $\{A_i\}$ a collection of subspaces of X . $\{A_i\}$ is a $\mathbb{Z}/p$ cohomology detecting family for X if the inclusion map $H^*(X) \rightarrow \prod H^*(A_i)$ is an injection.

2.3. LEMMA. Let K_p be a p -Sylow subgroup of K , then the transfer $t(K, K_p): H^*(K_p) \rightarrow H^*(K)$ is an epimorphism and the inclusion $i(K_p, K): H^*(K) \rightarrow H^*(K_p)$ is a monomorphism whose image consists of stable elements of $H^*(K_p)$. Furthermore we have the direct sum decomposition $H^*(K_p) \cong \text{Im } i(K_p, K) \oplus \text{Ker } t(K, K_p)$.

PROOF. See [5, Chapter XII, p. 257] for the definition of stable and p. 259 for a proof of the lemma.

Recalling that a p -Sylow subgroup of $\mathfrak{S}_{p^i}$ is isomorphic to $\text{wr}^i \mathbb{Z}/p$, [6] gives

2.4. COROLLARY. If $\{A_j\}$ is a $\mathbb{Z}/p$ detecting family for $\text{wr}^i \mathbb{Z}/p$ then it is one for $\mathfrak{S}_{p^i}$ also.

2.5. DEFINITION. Let G be a finite group of order n . Then the adjoint representation $A: G \rightarrow \mathfrak{S}_n$ is defined as follows: Let $A(g)$ be the permutation $\{g_i \mapsto gg_i\}$ where $\mathfrak{S}_n$ is thought of as the permutations on the n elements of G .

The adjoint representation is obviously a monomorphism and includes G in $\mathfrak{S}_n$. Let $G = \times^i \mathbb{Z}/p$, then the adjoint representation of $\times^i \mathbb{Z}/p$ in $\mathfrak{S}_{p^i}$ is clearly equivalent to the map $k_{i,i}: \times^i \mathbb{Z}/p \rightarrow \mathfrak{S}_{p^i}$ defined in §I. (The two maps differ by at most a reordering of the elements of $\times^i \mathbb{Z}/p$; that is, an inner automorphism of $\mathfrak{S}_{p^i}$.)

Again considering $\mathfrak{S}_{p^i}$ as the permutations on the set $\prod^i \mathbb{Z}/p$ the map $I_{i-1}: \times_{m=1}^p (\mathfrak{S}_{p^{i-1}})_m \rightarrow \mathfrak{S}_{p^i}$ defined in the introduction is realized by letting $(\mathfrak{S}_{p^{i-1}})_m$ permute the set $\prod^{i-1} \mathbb{Z}/p \times \{m\}$ contained in $\prod^i \mathbb{Z}/p$.

Note that under the specific embeddings $k_{i,i}$ and I_{i-1} the subgroup $\times^{i-1} \mathbb{Z}/p \times \{0\} \rightarrow \times^i \mathbb{Z}/p \xrightarrow{k_{i,i}} \mathfrak{S}_{p^i}$ is contained in the subgroup $\times_{m=1}^p (\mathfrak{S}_{p^{i-1}})_m \xrightarrow{I_{i-1}} \mathfrak{S}_{p^i}$. Any p -Sylow subgroup of $\times_{m=1}^p (\mathfrak{S}_{p^{i-1}})_m$ that contains $\times^{i-1} \mathbb{Z}/p \times \{0\}$ is isomorphic to $\times_{m=1}^p (\text{wr}^{i-1} \mathbb{Z}/p)_m$. Then $\times_{m=1}^p (\text{wr}^{i-1} \mathbb{Z}/p)_m$ and $\times^i \mathbb{Z}/p$ generate a p -Sylow subgroup of $\mathfrak{S}_{p^i}$ which must be isomorphic to $\text{wr}^i \mathbb{Z}/p$. Thus we have the following commutative diagram with the above mentioned inclusions:

$$\begin{array}{ccccc}
 & & \times_{m=1}^p (\mathfrak{S}_{p^{i-1}})_m & \xrightarrow{I_{i-1}} & \mathfrak{S}_{p^i} \\
 & \nearrow \tilde{k}_{i-1,i} & \uparrow & & \nwarrow k_{i,i} \\
 \times_{m=1}^p \left(\times^{i-1} \mathbb{Z}/p \right)_m & \longrightarrow & \times_{m=1}^p \left(\text{wr}^{i-1} \mathbb{Z}/p \right)_m & \longrightarrow & \text{wr}^i \mathbb{Z}/p \longleftarrow \times^i \mathbb{Z}/p
 \end{array}$$

where $\tilde{k}_{i-1,i} = \times_{m=1}^p (k_{i-1,i-1})_m$. The specific form of $k_{i,i}$ and $\tilde{k}_{i-1,i}$ guarantees $\times_{m=1}^p (\times^{i-1} Z/p)_m$ factors through $\times_{m=1}^p (\text{wr}^{i-1} Z/p)_m$.

More generally if $I_{m_1, \dots, m_n}: \mathbb{S}_{p^{m_1}} \times \dots \times \mathbb{S}_{p^{m_n}} \rightarrow \mathbb{S}_{p^i}$ is defined by letting $\mathbb{S}_{p^{m_r}}$ permute the p^{m_r} letters $(p^{m_1} + \dots + p^{m_{r-1}} + 1, \dots, p^{m_1} + \dots + p^{m_r})$ then the map $I_{m_1, \dots, m_n} \circ (\prod_{r=1}^n k_{m_r, m_r})$ includes $\prod_{r=1}^n (\times^{m_r} Z/p)$ in $\mathbb{S}_{p^i}$.

If $m_1 = m_2 = \dots = m_{p^{i-j}} = j$ then $\prod_{r=1}^{p^{i-j}} (\times^j Z/p) \rightarrow \mathbb{S}_{p^i}$ has the form

$$k_{j,i} = I_{j, \dots, j} \circ \prod_{r=1}^{p^{i-j}} (k_{j,j})_r: \times^{p^{i-j}} \left(\times^j Z/p \right) \rightarrow \mathbb{S}_{p^i}.$$

2.6. DEFINITION. Let $T_{j,i} = \times^{p^{i-j}} (\times^j Z/p)$. Let $k_{j,i}: T_{j,i} \rightarrow \mathbb{S}_{p^i}$ be the above inclusion. Then $T_{j,i}$ is called a totally symmetric detecting group.

Notice $T_{j,i}$ and $k_{j,i}$ are defined for $1 \leq j \leq i$. The following lemmas are established in the proofs of Theorems A through D:

2.7. LEMMA. The set $\{I_{m_1, \dots, m_n} \circ (\prod_{r=1}^n k_{m_r, m_r}): \prod_{r=1}^n \times^{m_r} (Z/p) \rightarrow \mathbb{S}_{p^i}\}$ forms a Z/p detecting family for $\mathbb{S}_{p^i}$.

2.8. LEMMA. The totally symmetric detecting groups $T_{j,i}$, $1 \leq j \leq i$, detect a set of multiplicative generators for $H^*(\mathbb{S}_{p^i})$. (This is the first part of Theorem D.)

2.9. LEMMA. In Z/p cohomology, $\text{Ker } k_{i,i}^* \cap \text{Ker } I_{i-1}^* = 0$.

These lemmas may be proved directly using [27], induction on i , and 3.1.

We now examine the normalizers of the detecting subgroups in $\mathbb{S}_{p^i}$. Consider $k_{i,i}: T_{i,i} \rightarrow \mathbb{S}_{p^i}$. Let $a_r \in \mathbb{S}_{p^i}$ generate $k_{i,i}(0 \times 0 \times \dots \times (Z/p)_r \times \dots \times 0)$ and let N_i be the normalizer of $k_{i,i}(T_{i,i})$ in $\mathbb{S}_{p^i}$. Define a homomorphism $\psi: N_i \rightarrow \text{GL}(i, Z/p)$ as follows: If $x \in N_i$ then $xa_r x^{-1} = a_1^{s_{1,r}} a_2^{s_{2,r}} \dots a_i^{s_{i,r}}$. Then let $\psi(x)$ be the matrix whose (m, n) th entry is $s_{m,n}$. Clearly $\psi(x)$ is nonsingular.

2.10. PROPOSITION. The sequence $1 \rightarrow k_{i,i}(T_{i,i}) \rightarrow N_i \xrightarrow{\psi} \text{GL}(i, Z/p) \rightarrow 1$ is exact.

PROOF. Preceding $k_{i,i}$ by any automorphism $\varphi: T_{i,i} \rightarrow T_{i,i}$ is just a reordering of the underlying set of $T_{i,i}$. This reordering, considered as an element of $\mathbb{S}_{p^i}$, conjugates $k_{i,i}$ to $k_{i,i} \circ \varphi$. This implies ψ is onto. The remainder of the proposition follows trivially.

For $x \in \mathbb{S}_{p^i}$ the homomorphism $\text{ad}_x: H^*(T_{i,i}) \rightarrow H^*(xT_{i,i}x^{-1})$ is induced by the inner automorphism $y \rightarrow xyx^{-1}$. Let $E = \sum_{m=1}^i a_m e_m$ and $B = \sum_{m=1}^i a'_m b_m$ in $H^*(T_{i,i})$ then it follows directly from the definition of ψ that

2.11. PROPOSITION. For $x \in N_i$, $\text{ad}_x(E) = \psi(x)E$ and $\text{ad}_x(B) = \psi(x)B$.

Since ad_x is a ring homomorphism 2.11 determines ad_x on all of $H^*(T_{i,i})$.

Since the p th power homomorphism, $a \mapsto a^p$, is the identity on Z/p we have $P(x_1^p, \dots, x_i^p) = (P(x_1, \dots, x_i))^p$ for all polynomials P . This fact and direct computation yield

2.12. PROPOSITION. ad_x operates on the classes $L_i, Q_{j,i}, M_{j,i}, \underline{L}_i$ via multiplication by the determinant function.

2.13. COROLLARY. The algebra $\mathcal{W}_i$ is contained in $H^*(T_{i,i})^{\text{GL}(i, Z/p)}$.

2.14. LEMMA. If G is a finite group, K a subgroup, and $N_{K,G}$ the normalizer of K in G then the image of $H^*(G)$ in $H^*(K)$ is contained in $H^*(K)^{N_{K,G}}$.

PROOF. Any inner automorphism of G induces the identity on $H^*(G)$. Hence we have the following commutative diagram:

$$\begin{array}{ccc} H^*(G) & \xrightarrow{\text{id}} & H^*(G) \\ i(K, G) \downarrow & & \downarrow i(xKx^{-1}, G) \\ H^*(K) & \xrightarrow{\text{ad}_x} & H^*(xKx^{-1}) \end{array}$$

Allowing x to run through $N_{K,G}$ gives the lemma.

2.15. COROLLARY. Let $u \in H^*(\mathfrak{S}_{p'})$ then $k_{i,i}^*(u) \in H^*(T_{i,i})^{\text{GL}(i, Z/p)}$.

PROOF. Immediate from 2.10 and 2.14.

Let $N_{j,i}$ be the normalizer of $k_{j,i}: T_{j,i} \rightarrow \mathfrak{S}_{p'}$ in $\mathfrak{S}_{p'}$.

2.16. PROPOSITION. The sequence

$$1 \rightarrow \times_{p^{i-j}} N_j \rightarrow N_{j,i} \xrightleftharpoons[\psi]{\varphi} \mathfrak{S}_{p^{i-j}} \rightarrow 1$$

is exact.

PROOF. Both $N_{j,i}$ and $\times_{p^{i-j}} N_j$ act on $T_{j,i}$ via conjugation. But $x \in N_{j,i}$ permutes the p^{i-j} orbits of $\times_{p^{i-j}} N_j$. This gives a homomorphism $\varphi: N_{j,i} \rightarrow \mathfrak{S}_{p^{i-j}}$ which is clearly onto and has an obvious section ψ . Notice $\psi(\varphi(x)^{-1}) \cdot x \in \times_{p^{i-j}} N_j$ as $\psi(\varphi(x)^{-1}) \in N_{j,i}$ and $\psi(\varphi(x)^{-1}) \cdot x \in \times_{p^{i-j}} \mathfrak{S}_{p'}$. The proposition follows.

Let $N_{m_1, \dots, m_n}$ be the normalizer of $I_{m_1, \dots, m_n}(\prod_{r=1}^n (k_{m_r, m_r}))$: $\prod_{r=1}^n (\times^{m_r} Z/p) \rightarrow \mathfrak{S}_{p'}$ in $\mathfrak{S}_{p'}$ and let $\mathfrak{S}_{(m_1, \dots, m_n)}$ be the subgroup of $\mathfrak{S}_n$ generated by the transpositions (a, c) where $m_a = m_c$. Minor modification of 2.16 yields the following three propositions.

2.17. PROPOSITION. The sequence $1 \rightarrow \times_{r=1}^n N_{m_r} \rightarrow N_{m_1, \dots, m_n} \xrightleftharpoons{\varphi} \mathfrak{S}_{(m_1, \dots, m_n)} \rightarrow 1$ is exact.

2.18. PROPOSITION. Let $\bar{N}_j$ be the normalizer of $I_j: \times_{p^{i-j}} \mathfrak{S}_{p'} \rightarrow \mathfrak{S}_{p'}$ in $\mathfrak{S}_{p'}$. Then the sequence $1 \rightarrow \times_{p^{i-j}} \mathfrak{S}_{p'} \rightarrow \bar{N}_j \xrightleftharpoons{\varphi} \mathfrak{S}_{p^{i-j}} \rightarrow 1$ is exact.

2.19. PROPOSITION. Let $\bar{N}_{m_1, \dots, m_n}$ be the normalizer of $I_{m_1, \dots, m_n}: \times_{r=1}^n \mathcal{S}_{m_r} \rightarrow \mathcal{S}_{p^i}$ in $\mathcal{S}_{p^i}$. Then the sequence $1 \rightarrow \times_{r=1}^n \mathcal{S}_{m_r} \rightarrow \bar{N}_{m_1, \dots, m_n} \rightrightarrows \mathcal{S}_{(m_1, \dots, m_n)} \rightarrow 1$ is exact.

2.20. LEMMA. If G is a finite group and K a subgroup then $i(K, G)^*t(G, K) = \sum_{x \in G/K} t_x i_x \text{ad}_x$ where $\text{ad}_x: H^*(K) \rightarrow H^*(xKx^{-1})$ is the homomorphism induced by $y \mapsto xyx^{-1}$ for $y \in K$, i_x is the inclusion map $H^*(xKx^{-1}) \rightarrow H^*(xKx^{-1} \cap K)$ and t_x is the transfer $H^*(xKx^{-1} \cap K) \rightarrow H^*(K)$.

PROOF. [5, XII. 9.1, p. 257].

2.21. PROPOSITION. If K is a proper subgroup of $\times^m \mathbb{Z}/p$ then the transfer $t: H^*(K) \rightarrow H^*(\times^m \mathbb{Z}/p)$ is zero.

PROOF. [4, I.2.1].

III. Some properties of $\mathcal{Q}(p)$ and the proof of Theorem E. In this section we state facts about the Steenrod algebra needed to prove Theorems A through D and give a proof of Theorem E.

First recall the construction of the Steenrod p th powers ([31] gives the complete treatment and we quote it frequently in what follows). Let X be a finite regular cell complex then we have the following spaces and maps:

$$X^p \xrightarrow{j} W_{\mathbb{Z}/p} \times_{\mathbb{Z}/p} X^p \xleftarrow{1 \times \Delta} W_{\mathbb{Z}/p} \times_{\mathbb{Z}/p} X = B_{\mathbb{Z}/p} \times X$$

where j is the inclusion and Δ is the diagonal map. Given any $u \in H^*(X)$ there exists a unique natural class $\mathcal{P}(u)$ in $H^*(W_{\mathbb{Z}/p} \times_{\mathbb{Z}/p} X^p)$ such that:

$$(1) j^*(\mathcal{P}(u)) = u \otimes \dots \otimes u = u^{\otimes p}.$$

(2) $(1 \times \Delta)^*(\mathcal{P}(u))$ in $H^*(B_{\mathbb{Z}/p} \times X)$ can be expanded by the Künneth theorem. $(1 \times \Delta)^*(\mathcal{P}(u)) = \sum w_k \otimes D_k(u)$ where w_k generates $H^k(\mathbb{Z}/p)$ and $D_k: H^q(X) \rightarrow H^{pq-k}(X)$ are homomorphisms which define the elements of $\mathcal{Q}(p)$.

$$(3) \beta D_{2k}(u) = D_{2k-1}(u), \beta D_{2k-1}(u) = 0 \text{ and } \beta D_0(u) = 0.$$

3.1. THEOREM [31]. If $z \in H^*(W_{\mathbb{Z}/p} \times_{\mathbb{Z}/p} X^p)$, then z is of the form $z = tz_1 + z_2 \cdot \mathcal{P}(z_3)$ with $z_1 \in H^*(X^p)$, $z_2 \in H^*(B_{\mathbb{Z}/p})$ and $z_3 \in H^*(X)$, where t is the transfer. Furthermore the sequence

$$H^*(X^p) \xrightarrow{t} H^*(W_{\mathbb{Z}/p} \times_{\mathbb{Z}/p} X^p) \xrightarrow{(1 \times \Delta)^*} H^*(B_{\mathbb{Z}/p} \times X)$$

is exact.

PROOF. [31, VII. 4.1, p. 104 and VIII. 3.6, p. 126].

3.2. DEFINITION [31]. Let $u \in H^q(X)$ then

$$\mathcal{P}^j(u) = a_{j,q} D_{(q-2j)(p-1)}(u),$$

$$\beta \mathcal{P}^j(u) = a_{j,q} D_{(q-2j)(p-1)-1}(u),$$

where $a_{j,q}$ is a nonzero constant in Z/p dependent on j and q . If $k \neq (q-2j)(p-1)$ or $(q-2j)(p-1)-1$ for some j then $D_k(u) = 0$.

3.3. PROPOSITION. If q is even, say $q = 2n$, then $a_{j,2n} = (-1)^{j+n}$.

PROOF. Follows directly from [31, VII. 6.1 and VII. 6.3] (note correction of the formula in VII. 6.1 on the first page of the appendix to [31]).

The following is well known:

3.4. LEMMA. I. Let p be a prime and $a = \sum_{i=0}^m a_i p^i$, $c = \sum_{i=0}^m c_i p^i$ ($0 \leq a_i, c_i \leq p-1$). Then

$$\binom{c}{a} \equiv \prod_i \binom{c_i}{a_i} \pmod{p}.$$

II. $\mathcal{P}^j(e) = 0$ for all $j > 0$.

III. $\mathcal{P}^j(b^k) = \binom{k}{j} b^{k+(p-1)j}$.

IV. (Cartan formula) $\mathcal{P}^j(uv) = \sum_{m+n=j} \mathcal{P}^m(u) \mathcal{P}^n(v)$.

V.

$$\mathcal{P}^j(b^{p^m}) = \binom{p^m}{j} b^{p^m+(p-1)j} = \begin{cases} b^{p^m} & \text{if } j = 0, \\ b^{p^{m+1}} & \text{if } j = p^m, \\ 0 & \text{otherwise.} \end{cases}$$

PROOF. [31, see I.2.6, V. 1, VII. 2.2 and VI. 2.3].

The proof of Theorem E follows from direct calculation and Lemma 3.4. Note: To prove relation (4) of Theorem E, just expand $\mathcal{P}^{p^k-1}(Q_{k,i} L_i^{p-1})$.

IV. Symmetric products and image $k_{i,i}^*$. In this chapter we summarize results of [17] which give $H^*(\mathfrak{S}_n)$ as Z/p vector spaces and give an upper bound on the size of image $k_{i,i}^*$.

Recall the monomial $\mathcal{P}^I = \beta^{\epsilon_k} \mathcal{P}^{s_k} \cdots \beta^{\epsilon_1} \mathcal{P}^{s_1} \in \mathcal{Q}(p)$ is called admissible if $s_i \geq ps_{i-1} + \epsilon_{i-1}$ for each $i \geq 1$, and the excess of $\mathcal{P}^I = 2s_k + \epsilon_k - \sum_{j=1}^{k-1} (2s_j(p-1) + \epsilon_j)$. The excess of any admissible monomial is nonnegative. Let $\mathcal{Q}(p)_n$ be the subvector space of $\mathcal{Q}(p)$ spanned by those monomials of excess $< n$.

Let $\text{SP}^k(S^{2n})$ be the k symmetric product of S^{2n} (see [17] for the definition and properties of the symmetric products of a space).

4.1. THEOREM [17]. (1) $H_*(\text{SP}^k(S^{2n})) = \sum_{m=1}^k H_*(\text{SP}^m(S^{2n}), \text{SP}^{m-1}(S^{2n}))$.

(2) $\mathcal{R}(S^{2n}, Z/p) = \sum_{m=1}^\infty H_*(\text{SP}^m(S^{2n}), \text{SP}^{m-1}(S^{2n}))$ is isomorphic to $H_*(K(Z, 2n))$.

There is a bigrading of $\mathcal{R}(S^{2n}, Z/p)$ given by

$$\mathcal{R}_{i,m}(S^{2n}, Z/p) = H_i(\text{SP}^m(S^{2n}), \text{SP}^{m-1}(S^{2n})).$$

(3) For $\mathcal{R}(S^{2n}, Z/p)$ the generators q_I in homology are in 1-1 correspondence with admissible monomials $\mathcal{P}^I = \beta^{\epsilon_k} \mathcal{P}^{s_k} \cdots \beta^{\epsilon_1} \mathcal{P}^{s_1}$ in $\mathcal{Q}(p)_{2n}$ and the bidegree

of this generator is $(|\mathcal{P}^I| + 2n, p^i)$. Moreover $\langle q_I, \mathcal{P}^I(i) \rangle = 1$ under the isomorphism in (2).

PROOF. [17].

REMARKS. (1) is due to N. E. Steenrod. [8] and [21] also studied (1) and (2).

The next theorem follows from the fact that the singular locus of $(S^{2n})^{p^i}$ under $\mathfrak{S}_{p^i}$ has dimension $2n(p^i - 1)$.

4.2. THEOREM [17]. For $k < 2n - 1$, $H^k(\mathfrak{S}_{p^i}) \cong H_{2n(p^i)-k}(\mathrm{SP}^{p^i}(S^{2n}))$.

Since $H_j(\mathrm{SP}^{p^i}(S^{2n})) \cong H_j(\mathrm{SP}^{p^j}(S^{2n}), \mathrm{SP}^{p^{i-1}}(S^{2n}))$ for $j > 2n(p^i - 1) + 1$ we may identify $H^k(\mathfrak{S}_{p^i})$ with elements in $\mathcal{R}(S^{2n}, \mathbb{Z}/p)$ of bidegree $(2n(p^i) - k, p^i)$. Thus for $k < 2n - 1$ classes in $H^k(\mathfrak{S}_{p^i})$ correspond to classes Σa ; with each $a \in \mathcal{R}(S^{2n}, \mathbb{Z}/p)$ having bidegree $(-, p^i)$. This gives $H^k(\mathfrak{S}_{p^i})$ as $\mathbb{Z}/p$ vector spaces. Recall there are two types of classes in $\mathcal{R}(S^{2n}, \mathbb{Z}/p)$ having bidegree $(-, p^i)$:

(1) a corresponds to $\mathcal{P}^I$ of bidegree $(|\mathcal{P}^I| + 2n, p^i)$,

(2) $a = \prod b_k$ where b_k has bidegree $(-, p^j)$, for some $j < i$ and occurs in $H_*(\mathrm{SP}^{p^j}(S^{2n}), \mathrm{SP}^{p^{j-1}}(S^{2n}))$.

On the other hand the multiplication map $M: \mathrm{SP}^{p^{i-1}}(S^{2n}) \times \cdots \times \mathrm{SP}^{p^{i-1}}(S^{2n}) \rightarrow \mathrm{SP}^{p^i}(S^{2n})$ and 4.2 give a map $m: \bigotimes^p H^*(\mathfrak{S}_{p^{i-1}}) \rightarrow H^*(\mathfrak{S}_{p^i})$.

4.3. LEMMA [21]. m is the transfer map induced by the inclusion

$$I_{i-1}: \bigotimes^p \mathfrak{S}_{p^{i-1}} \rightarrow \mathfrak{S}_{p^i}.$$

PROOF. [21].

4.4. LEMMA. Let $u \in H^*(\mathfrak{S}_{p^i})$ correspond to $a \in \mathcal{R}(S^{2n}, \mathbb{Z}/p)$. If a is of type 2 then $k_{i,i}^*(u) = 0$.

PROOF. Suppose a is of type 2 then a is in the image of M_* . By 4.3, u is in the image of the transfer $t: H^*(\bigotimes^p \mathfrak{S}_{p^{i-1}}) \rightarrow H^*(\mathfrak{S}_{p^i})$. But 3.1 implies $k_{i,i}^* t = 0$. Hence $k_{i,i}^*(u) = 0$.

Let $\mathcal{R}'_{2n(p^i)-k, p^i}(S^{2n}, \mathbb{Z}/p)$ be the subspace of $\mathcal{R}_{2n(p^i)-k, p^i}(S^{2n}, \mathbb{Z}/p)$ spanned by elements of type 1. Then 4.4 yields:

4.5. THEOREM [17]. As $\mathbb{Z}/p$ vector spaces

$$\dim((\text{image } k_{i,i}^*)_k) \leq \dim(\mathcal{R}'_{2n(p^i)-k, p^i}(S^{2n}, \mathbb{Z}/p)).$$

V. The proof of Theorem A. We now proceed with the proof of Theorem A.

5.1. LEMMA. $\mathcal{W}_i$ is contained in image $k_{i,i}^*$.

PROOF. By induction on i . The lemma is classically true for $i = 1$ and [4] proves the lemma for $i = 2$. Assume $\mathcal{W}_{i-1}$ is contained in image $k_{i-1, i-1}^*$. The next four lemmas establish 5.1.

5.2. LEMMA. *There exists $u \in H^*(\mathfrak{S}_{p^i})$ such that*

$$k_{i-1,i}^*(u) = (M_{i-2,i-1}M_{i-3,i-1}L_{i-1}^{p-3})^{\otimes p} \in H^*(T_{i-1,i}).$$

PROOF. Recall the following commutative diagram containing the construction of the Steenrod powers on $\mathfrak{S}_{p^{i-1}}$:

$$\begin{array}{ccccc}
 & & B_{\mathfrak{S}_{p^i}} & & \\
 & & \uparrow & & \\
 & & B_{N_p \times (\mathfrak{S}_{p^{i-1}}, \mathfrak{S}_{p^i})} & & \\
 & & \uparrow & & \\
 B_p \times_{\mathfrak{S}_{p^{i-1}}} & \xrightarrow{j} & W_{Z/p} \times_{Z/p} B_p \times_{\mathfrak{S}_{p^{i-1}}} & = & B_{\mathfrak{S}_{p^{i-1}} \text{ wr } Z/p} \\
 \uparrow & & \uparrow & & \uparrow \\
 & & 1 \times \Delta & & \\
 & & W_{Z/p} \times_{Z/p} B_{\mathfrak{S}_{p^{i-1}}} & = & B_{Z/p} \times B_{\mathfrak{S}_{p^{i-1}}} \\
 & & \uparrow & & \\
 & & B_{T_{i,i}} & &
 \end{array}$$

Of course the composition $B_{T_{i-1,i}} \rightarrow B_{\mathfrak{S}_{p^i}}$ is $Bk_{i-1,i}$ and the composition $B_{T_{i,i}} \rightarrow B_{\mathfrak{S}_{p^i}}$ is $Bk_{i,i}$.

Let $u' \in H^*(\mathfrak{S}_{p^{i-1}})$ be such that $k_{i-1,i-1}^*(u') = M_{i-2,i-1}M_{i-3,i-1}L_{i-1}^{p-3}$ then $\mathfrak{P}(u') = u'' \in H^*(\mathfrak{S}_{p^{i-1}} \text{ wr } Z/p)$. Let $A = \mathfrak{S}_{p^{i-1}} \text{ wr } Z/p$. Then 2.20 gives

$$i(A, \mathfrak{S}_{p^i})^* t(\mathfrak{S}_{p^i}, A) = \sum_{x \in \mathfrak{S}_{p^i}/A} t_x i_x \text{ad}_x$$

and we have the following commutative diagram:

$$\begin{array}{ccccccc}
 H^*(A) & \xrightarrow{\text{ad}_x} & H^*(xAx^{-1}) & \xrightarrow{i_x} & H^*(xAx^{-1} \cap A) & \xrightarrow{t_x} & H^*(A) \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \Sigma H^*(T') & \xrightarrow{\Sigma \text{ad}_x} & \Sigma H^*(xT'x^{-1}) & \xrightarrow{\Sigma i_x} & \Sigma H^*(xT'x^{-1} \cap T_{i-1,i}) & \xrightarrow{\Sigma t_x} & H^*(T_{i-1,i})
 \end{array}$$

where T' runs through all inclusions $\times^m Z/p$ in A . (The last square commutes by 2.21 and [31, V. 7.2], as $xT_{i-1,i}x^{-1} \subset A$ implies $x \in A$.)

Thus 2.16, 2.18 and 2.21 show

$$k_{i-1,i}^* t(A, \mathfrak{S}_{p^i})(u'') = \sum_x (M_{i-2,i-1}M_{i-3,i-1}L_{i-1}^{p-3})^{\otimes p}$$

where the sum runs over a coset representation $\bar{N}_{i-1} = N_{\times^p \mathfrak{S}_{p^{i-1}}, \mathfrak{S}_{p^i}} \bmod A$. As

A contains a p -Sylow subgroup of $\mathfrak{S}_{p^i}$, $[\bar{N}_{i-1}; A] = c \not\equiv 0 \pmod{p}$. Let $u = t(A, \mathfrak{S}_{p^i})(c^{-1}u'')$; then $k_{i-1,i}^*(u) = (M_{i-2,i-1}M_{i-3,i-1}L_{i-1}^{p-3})^{\otimes p}$.

5.3. LEMMA. *There exists $u \in H^*(\mathfrak{S}_{p^i})$ such that*

$$k_{i-1,i}^*(u) = (Q_{i-2,i-1})^{\otimes p} \in H^*(T_{i-1,i}).$$

PROOF. Identical to that of 5.2.

5.4. LEMMA. *There exists $u \in H^*(\mathfrak{S}_{p^i})$ such that $k_{i,i}^*(u) = M_{i-1,i}M_{i-2,i}L_i^{p-3}$.*

PROOF. Let $u' \in H^*(\mathfrak{S}_{p^{i-1}})$ be such that $k_{i-1,i-1}^*(u') = M_{i-2,i-1}M_{i-3,i-1}L_{i-1}^{p-3}$ and $u \in H^*(\mathfrak{S}_{p^i})$ be such that $k_{i-1,i}^*(u) = (M_{i-2,i-1}M_{i-3,i-1}L_{i-1}^{p-3})^{\otimes p} \in H^*(T_{i-1,i})$. Recall 3.1 implies image $k_{i,i}^*$ is contained in the $H^*(Z/p)$ module generated by $\text{image}(1 \times \Delta)^*\mathcal{P}$. A simple dimension check shows that the only classes in $H^*(\mathfrak{S}_{p^{i-1}} \text{ wr } Z/p)$ that could project to $k_{i-1,i}^*(u)$ are $\mathcal{P}(u')$ and $b_1^x + \mathcal{P}(u')$, where $x = \frac{1}{2} \text{dimension}(u)$. By 2.15, $k_{i,i}^*(u)$ is $\text{GL}(i, Z/p)$ invariant. As $(u')^p = 0$ in $H^*(\mathfrak{S}_{p^{i-1}})$ the class $b_1^x + \mathcal{P}(u')$ is not $\text{GL}(i, Z/p)$ invariant (there cannot be a pure b_r^x term in $(1 \times \Delta)^*(\mathcal{P}(u'))$ for $r \geq 1$). Hence $k_{i,i}^*(u) = (1 \times \Delta)^*(\mathcal{P}(u'))$. It is easy to see that $\text{dimension}(u) = 2(p^{i-1} - p^{i-2} - p^{i-3}) = 2n$. Thus

$$\begin{aligned} k_{i,i}^*(u) &= (1 \times \Delta)^*(\mathcal{P}(u')) = \sum_k w_k \otimes D_k(M_{i-2,i-1}M_{i-3,i-1}L_{i-1}^{p-3}) \\ &= (-1)^n \left[\sum_j w_{(2n-2j)(p-1)} \otimes (-1)^j \mathcal{P}^j(M_{i-2,i-1}M_{i-3,i-1}L_{i-1}^{p-3}) \right. \\ &\quad \left. + \sum_j w_{(2n-2j)(p-1)-1} \otimes (-1)^j \beta \mathcal{P}^j(M_{i-2,i-1}M_{i-3,i-1}L_{i-1}^{p-3}) \right]. \end{aligned}$$

Consider $M_{i-1,i}M_{i-2,i}L_i^{p-3}$. Expanding along the e_1, b_1 columns we have

$$\begin{aligned} M_{i-1,i}M_{i-2,i}L_i^{p-3} &= \sum_{\substack{A \\ B \\ C_k}} (-1)^{\varphi} b_1^r (ABC_1 \cdots C_{p-3}) \\ &\quad + \sum_{\substack{D \\ E \\ C_k}} (-1)^{\varphi} e_1 b_1^s (DEC_1 \cdots C_{p-3}) \end{aligned}$$

where A runs over all $i-1 \times i-1$ minors of $M_{i-1,i}$ eliminating the $b_1^{p''}$ ($0 \leq u \leq i-2$) row and column, B runs over all $i-1 \times i-1$ minors of $M_{i-2,i}$ eliminating the $b_1^{p''}$ ($0 \leq v \leq i-3$, or $v = i-1$) row and column, C_k ($k = 1, \dots, p-3$) is any $i-1 \times i-1$ minor of L_i eliminating the $b_1^{p''k}$ ($0 \leq z_k \leq i-1$) row and column, r satisfies the relation $\dim(M_{i-1,i}M_{i-2,i}L_i^{p-3}) = 2r + \dim(A) + \dim(B) + \sum_{k=1}^{p-3} \dim(C_k)$, and $\varphi \equiv u + v + \sum_{k=1}^{p-3} z_k \pmod{2}$ if $v \neq i-1$, and $\equiv (i-u) + \sum_{k=1}^{p-3} z_k \pmod{2}$ if

$v = i - 1$. D and E are $i - 1 \times i - 1$ minors of $M_{i-1,i}$ and $M_{i-2,i}$ respectively with exactly one minor eliminating the e_1 row and column, the other eliminating a $b_1^{p'}$ row and column.

If C_k is the minor eliminating the $b_1^{p^k}$ row and column then $C_k = \mathcal{P}^{m_{z_k}}(L_{i-1})$ where $m_{z_k} = p^{z_k} + p^{z_k+1} + \dots + p^{i-2}$ ($= 0$ if $z_k = i - 1$).

Case 1. Suppose $v = i - 1$. Then the minor of $M_{i-2,i}$ eliminating the $b_1^{p^v}$ row and column is $M_{i-2,i-1}$. If A is an $i - 1 \times i - 1$ minor of $M_{i-1,i}$ eliminating the $b_1^{p^u}$ row and column and $AM_{i-2,i-1} \neq 0$ then $u \neq i - 2$. Thus $A = \mathcal{P}^{j_1}(M_{i-3,i-1})$ where $j_1 = p^u + p^{u+1} + \dots + p^{i-4}$ (if $u = i - 3$ then $j_1 = 0$). Thus if $v = i - 1$ we have

$$ABC_1 \cdots C_{p-3} = (-1)^{\mathcal{P}^0(M_{i-2,i-1})} \mathcal{P}^{j_1}(M_{i-3,i-1}) \mathcal{P}^{m_{z_1}}(L_{i-1}) \cdots \mathcal{P}^{m_{z_{p-3}}}(L_{i-1}).$$

Case 2. Suppose $0 \leq v \leq i - 3$. Then $A = \mathcal{P}^{j_1}(M_{i-2,i-1})$ where $j_1 = p^u + p^{u+1} + \dots + p^{i-3}$ unless $u = i - 2$ in which case $j_1 = 0$ and $B = \mathcal{P}^{j_2}(M_{i-3,i-1})$ where $j_2 = p^v + p^{v+1} + \dots + p^{i-4} + p^{i-2}$ unless $v = i - 3$ in which case $j_2 = p^{i-2}$. Then we have

$$ABC_1 \cdots C_{p-3} = \mathcal{P}^{j_1}(M_{i-2,i-1}) \mathcal{P}^{j_2}(M_{i-3,i-1}) \mathcal{P}^{m_{z_1}}(L_{i-1}) \cdots \mathcal{P}^{m_{z_{p-3}}}(L_{i-1}).$$

Note. In Case 1 we have terms involving $(-1)^{\mathcal{P}^0(M_{i-2,i-1})} \mathcal{P}^{j_1}(M_{i-3,i-1})$ and in Case 2 if $u = i - 2$ we have terms involving $\mathcal{P}^0(M_{i-2,i-1}) \mathcal{P}^{j_2}(M_{i-3,i-1})$ but it is clear that j_1 can never equal j_2 in these cases.

Thus if $ABC_1 \cdots C_{p-3} \neq 0$ we have written $ABC_1 \cdots C_{p-3}$ uniquely as $\mathcal{P}^{j_1}(M_{i-2,i-1}) \mathcal{P}^{j_2}(M_{i-3,i-1}) \mathcal{P}^{m_{z_1}}(L_{i-1}) \cdots \mathcal{P}^{m_{z_{p-3}}}(L_{i-1})$ for certain $j_1, j_2, m_{z_1}, \dots, m_{z_{p-3}}$. 3.4 clearly shows if

$$Y = \mathcal{P}^{s_1}(M_{i-2,i-1}) \mathcal{P}^{s_2}(M_{i-3,i-1}) \mathcal{P}^{s_{z_1}}(L_{i-1}) \cdots \mathcal{P}^{s_{z_{p-3}}}(L_{i-1}) \neq 0$$

then $Y = ABC_1 \cdots C_{p-3}$ for a suitable choice of $A, B, C_1, \dots, C_{p-3}$ and is thus analyzed in Case 1 or Case 2 above.

Let $j = j_1 + j_2 + \sum_{k=1}^{p-3} m_{z_k}$. For both $v = i - 3$ and $v < i - 3$ it is trivial to see that $\varphi = j \pmod{2}$. Hence the Cartan formula and the above facts yield the following decomposition of $M_{i-1,i} M_{i-2,i} L_i^{p-3}$ where the first sum runs over all integers j .

$$\begin{aligned} M_{i-1,i} M_{i-2,i} L_i^{p-3} &= \sum_j b_1^{(n-j)(p-1)} \otimes (-1)^j \mathcal{P}^j(M_{i-2,i-1} M_{i-3,i-1} L_{i-1}^{p-3}) \\ &\quad + \sum_{\substack{D \\ E \\ C_k}} (-1)^{\varphi} e_1 b_1^s \otimes DEC_1 \cdots C_{p-3}. \end{aligned}$$

Let $U = k_{i,i}^*(u) - (-1)^n (M_{i-1,i} M_{i-2,i} L_i^{p-3})$. U is clearly $\text{GL}(i, \mathbb{Z}/p)$ invariant. Any monomial term in U must contain the factor $e_1 e_j$ ($j \neq 1$) but as there is no monomial in U with an $e_2 e_3$ factor symmetry implies $U = 0$. As

$n = p^{i-1} - p^{i-2} - p^{i-3}$ we have

$$k_{i,i}^*(u) = -M_{i-1,i}M_{i-2,i}L_i^{p-3}.$$

This proves 5.4.

Note. By keeping careful track of D , E , and $\beta(\mathcal{P}^{j_1}(M_{i-2,i-1})\mathcal{P}^{j_2}(M_{i-3,i-1}))$ it is possible to see directly that

$$\sum_{\substack{D \\ E \\ D_k}} (-1)^{\varphi} e_1 b_1^s \otimes DEC_1 \cdots C_{p-3} \\ = - \sum_j e_1 b_1^{(n-j)(p-1)-1} \otimes (-1)^j \beta \mathcal{P}^j(MML^{p-3})$$

where $MML^{p-3} = M_{i-2,i-1}M_{i-3,i-1}L_{i-1}^{p-3}$.

5.5. LEMMA. *There exists $u \in H^*(\mathcal{S}_{p'})$ such that $k_{i,i}^*(u) = Q_{i-1,i}$.*

PROOF. The proof is similar to that of 5.4. We let $u' \in H^*(\mathcal{S}_{p'-1})$ be such that $k_{i-1,i-1}^*(u') = Q_{i-2,i-1}$ and $u \in H^*(\mathcal{S}_{p'})$ be such that $k_{i-1,i}^*(u) = (Q_{i-2,i-1})^{\otimes p} \in H^*(T_{i-1,i})$. Then $k_{i,i}^*(u)$ is the $\text{GL}(i, \mathbb{Z}/p)$ invariant class containing $(1 \times \Delta)^*(\mathcal{P}(u'))$. But [8] proved $Q_{i-1,i}$ is the only $\text{GL}(i, \mathbb{Z}/p)$ invariant polynomial in this dimension. Thus $k_{i,i}^*(u) = cQ_{i-1,i}$, where c is a constant. Note $(1 \times \Delta)^*(\mathcal{P}(u'))$ contains the term $w_0 \otimes D_0(Q_{i-2,i-1}) = (Q_{i-2,i-1})^p \neq 0$. Hence $c \neq 0$.

The naturality of the Steenrod algebra implies image $k_{i,i}^*$ contains $\mathcal{Q}(p)(M_{i-1,i}M_{i-2,i}L_i^{p-3}, Q_{i-1,i})$. By Theorem E any generator $\mathcal{W}_i$ is contained in $\mathcal{Q}(p)(M_{i-1,i}M_{i-2,i}L_i^{p-3}, Q_{i-1,i})$ (see the diagram after Theorem E). This completes the proof of Lemma 5.1.

By 4.5, to complete the proof of Theorem A it suffices to construct a 1-1 correspondence between nonzero monomials in $\mathcal{W}_i$ and admissible monomials in $\mathcal{Q}(p)$.

5.6. LEMMA. $M_{i-1,i}M_{i-2,i} \cdots M_{1,i}L_i \neq 0$.

PROOF. The term $e_1 e_2 \cdots e_i (b_1^{p^{i-1}})^{i-1} (b_2^{p^{i-2}})^{i-1} \cdots (b_i)^{i-1}$ appears with coefficient 1 in the term-by-term expansion of $M_{i-1,i}M_{i-2,i} \cdots M_{1,i}L_i$.

The only admissible monomials of length 1 in $\mathcal{Q}(p)_{2n}$ are $\mathcal{P}^{n-j}(u_{2n})$ and $\beta \mathcal{P}^{n-j}(u_{2n})$ which correspond to $(L_1^{p-1})^j$ and $(L_1 L_1^{p-2})(L_1^{p-1})^{j-1}$ in $\mathcal{W}_i$. Thus we may assume, by induction, that an $i-1$ length admissible monomial in $\mathcal{Q}(p)_{2n}$ starting with $\mathcal{P}^{n-j}(u_{2n})$ corresponds to a j -fold product monomial in $\mathcal{W}_{i-1}$ ($j < n$). Let A be an admissible monomial in $\mathcal{Q}(p)_{2n}$.

Case 1. $e_1 = 0$; that is, $A = \beta^{e_1 \mathcal{P}^{s_1}} \cdots \beta^{e_2 \mathcal{P}^{s_2}} \mathcal{P}^{n-j}(u_{2n})$. The dimension of $\mathcal{P}^{n-j}(u_{2n})$ is $2p(n-j) + 2j$ and hence $s_2 = p(n-j) + k$, $0 \leq k < j$, if $A(u_{2n})$ is nonzero and admissible. Consider

$$A' = \beta^{e_1 \mathfrak{P}^{s_1}} \cdots \beta^{e_2 \mathfrak{P}^{s_2}} (\bar{u}_{2(p(n-j)+j)}) \quad \text{where } \bar{u}_{2(p(n-j)+j)} = \mathfrak{P}^{n-j}(u_{2n}).$$

A' is an admissible monomial of length $i-1$ and $s_2 = (p(n-j)+j) - (j-k)$. Thus A' corresponds to a $(j-k)$ -fold product monomial in $\mathcal{W}_{i-1}$, call it U_{j-k} . Identify A with $\bar{U}_{j-k}(Q_{i-1,i})^k$ in $\mathcal{W}_i$. $\bar{U}_{j-k}$ comes from U_{j-k} by changing the detecting index from $i-1$ to i ; i.e., $Q_{m,i-1} \rightarrow Q_{m,i}$.

Case 2. $e_1 = 1$; that is, $A = \beta^{e_1 \mathfrak{P}^{s_1}} \cdots \beta^{e_2 \mathfrak{P}^{s_2}} \beta^{\mathfrak{P}^{n-j}}(u_{2n})$. Then consider that part of A until a second Bockstein occurs.

$$A = \beta^{e_1 \mathfrak{P}^{s_1}} \cdots \beta^{\mathfrak{P}^{s_k} \mathfrak{P}^{s_{k-1}}} \cdots \beta^{p(p(n-j)+m_1)+m_2 \mathfrak{P}^{p(n-j)+m_1}} \beta^{\mathfrak{P}^{n-j}}(u_{2n})$$

with $m_1 \geq 1$.

Further suppose $k < i$. Then

$$s_k = p(p(p(\cdots(p(n-j)+m_1)+m_2)+\cdots+m_{k-2}))+m_{k-1}$$

and $\mathfrak{P}^{s_k} \cdots \beta^{\mathfrak{P}^{n-j}}(u_{2n})$ has dimension $2p^k(n-j) + 2p^{k-1}m_1 + 2p^{k-2}m_2 + \cdots + 2pm_{k-1} + 2(j-m_1-m_2-\cdots-m_{k-1}) + 1$. For A to be admissible and nonzero we must also have $j-m_1-m_2-\cdots-m_{k-1} \geq 0$ and $j-m_1-m_2-\cdots-m_{k-1}+1 \geq 0$. Then

$$A' = \beta^{e_1 \mathfrak{P}^{s_1}} \cdots \mathfrak{P}^{s_{k+1}} (\beta^{\mathfrak{P}^{s_k}} \cdots \beta^{\mathfrak{P}^{n-j}}(u_{2n})) = A'' (\beta^{\mathfrak{P}^{s_k}} \cdots \beta^{\mathfrak{P}^{n-j}}(u_{2n}))$$

and A'' corresponds to a $j-m_1-m_2-\cdots-m_k+1$ fold product monomial in $\mathcal{W}_{i-k}$, call it $U_{A''}$. Identify A with the monomial

$$\bar{U}_{A''} (M_{i-k,i} M_{i-1,i} L_i^{p-3}) (Q_{i-k,i})^{m_{k-1}-1} (Q_{i-k-1,i})^{m_{k-2}} \cdots (Q_{i-2,i})^{m_2} (Q_{i-1,i})^{m_1-1}$$

where $\bar{U}_{A''}$ comes from $U_{A''}$ by changing the detecting index from $i-k$ to i ; i.e., $Q_{m,i-k} \rightarrow Q_{m,i}$. If $k=i$ or no second Bockstein occurs assign to A the monomial

$$(M_{i-1,i} L_i^{p-3}) (L_i^{p-3})^{m_i} (Q_{1,i})^{m_{i-1}} \cdots (Q_{i-2,i})^{m_2} (Q_{i-1,i})^{m_1-1} \quad \text{or} \\ (M_{i-1,i} L_i^{p-2}) (L_i^{p-3})^{m_i} (Q_{1,i})^{m_{i-1}} \cdots (Q_{i-2,i})^{m_2} (Q_{i-1,i})^{m_1-1} \quad \text{respectively}$$

where $m_i = j-m_1-m_2-\cdots-m_{i-1}$.

Let $U_{A(u_{2n})}$ be the above constructed monomial in $\mathcal{W}_i$ corresponding to $A(u_{2n})$. It is routine to verify that for $U_{A''}$ in $\mathcal{W}_{i-k}$ and $\bar{U}_{A''}$ in $\mathcal{W}_i$ constructed above we have $\dim(U_{A''}) + 2j(p^i - p^{i-k}) = \dim(U_{A(u_{2n})})$. This fact and induction on i show that if $A(u_{2n})$ has dimension $2n(p^i) - k$ then $U_{A(u_{2n})}$ has dimension k . Lemma 5.6 shows $U_{A(u_{2n})} \neq 0$. Hence, by Theorem 4.5, $(\mathcal{W}_i)_k$ must fill out $(\text{image } k_{i,i}^*)_k$ for $k \ll n$. This finishes the proof of Theorem A.

VI. Proof of Theorems B, C, D, and F. Consider the following commutative diagram:

$$\begin{array}{ccccc}
& & & H^*(T_{i,i}) & \\
& & & \uparrow \text{incl}^* & \\
& & & H^*(\mathbb{S}_{p^{i-1}} \times Z/p) & \\
& & & \uparrow (1 \times \Delta)^* & \\
& & & H^*(\mathbb{S}_{p^{i-1}} \text{ wr } Z/p) & \\
& & \nearrow k_{i,i}^* & \nearrow \bar{k}_{i,i}^* & \nearrow h^* \\
H^*(\times^p \mathbb{S}_{p^{i-1}}) & \xrightarrow{t} & H^*(\mathbb{S}_{p^i}) & \xrightarrow{h^*} & H^*(\mathbb{S}_{p^{i-1}} \text{ wr } Z/p) \\
\downarrow t'' & & \downarrow t' & & \\
H^*(\mathbb{S}_{p^{i-1}} \text{ wr } Z/p) & \xrightarrow{t'} & H^*(\mathbb{S}_{p^i}) & \xrightarrow{h^*} & H^*(\mathbb{S}_{p^{i-1}} \text{ wr } Z/p)
\end{array}$$

where $h = i(\mathbb{S}_{p^{i-1}} \text{ wr } Z/p, \mathbb{S}_{p^i})$.

6.1. PROPOSITION. Let $u \in H^*(\mathbb{S}_{p^i})$. If $k_{i,i}^*(u) = 0$ then there exists $z \in H^*(\times^p \mathbb{S}_{p^{i-1}})$ such that $t(z) = u$.

PROOF. By 4.4 and Theorem A, $k_{i,i}^*(u) = 0$ implies $\bar{k}_{i,i}^*(u) = 0$. Hence $(1 \times \Delta)^* h^*(u) = 0$ and $h^*(u) \in \ker(1 \times \Delta)^*$. By 3.1 there exists $z \in H^*(\times^p (\mathbb{S}_{p^{i-1}}))$ such that $t''(z) = h^*(u)$. Then $t(z) = t' t''(u) = t'(h^*(u)) = [\mathbb{S}_{p^i} : \mathbb{S}_{p^{i-1}} \text{ wr } Z/p] u = u \pmod{p}$.

Let $u_{s,i-1} \in H^*(\mathbb{S}_{p^{i-1}})$, then, by induction, $u_{s,i-1}$ pulls back to a $\mathbb{S}_{p^{i-1}}$ detecting subgroup $\prod_{i=1}^q T_{s_i, s_i} \rightarrow \mathbb{S}_{p^{i-1}}$ (recall §II gives these subgroups and their inclusions into $\mathbb{S}_{p^{i-1}}$). Thus to complete the computation of $H^*(\mathbb{S}_{p^i})$ it suffices to compute the map $I_{i-1}^* t$. First consider the maps $\Phi_{m_1, \dots, m_n} = (I_{m_1, \dots, m_n} \circ (\prod_{r=1}^n (k_{m_r, m_r})))^* t_{m_1, \dots, m_n} : H^*(\times_{r=1}^n \mathbb{S}_{p^{m_r}}) \rightarrow H^*(\mathbb{S}_{p^i}) \rightarrow \bigotimes_{r=1}^n H^*(T_{m_r, m_r})$ for all $(m_1, \dots, m_r)$ such that $\sum_{r=1}^n p^{m_r} = p^i$, with $n \geq 2$ and $t_{m_1, \dots, m_n}$ the transfer $H^*(\times_{r=1}^n \mathbb{S}_{p^{m_r}}) \rightarrow H^*(\mathbb{S}_{p^i})$.

6.2. LEMMA. Let $u = u_{1, m_1} \otimes \dots \otimes u_{n, m_n} \in H^*(\times_{r=1}^n \mathbb{S}_{p^{m_r}})$ and $k_{m_r, m_r}^*(u_{r, m_r}) = v_r$. Then

$$\Phi_{m_1, \dots, m_n}(u) = \sum_{\sigma \in \mathcal{S}_{(m_1, \dots, m_n)}} v_{\sigma(1)} \otimes \dots \otimes v_{\sigma(n)}.$$

PROOF. As in the proof of 5.2, 2.16 through 2.21 and the following commutative diagram give the proposition:

$$\begin{array}{ccccccc}
H^*(A) & \xrightarrow{\text{ad}_x} & H^*(xAx^{-1}) & \xrightarrow{i_x} & H^*(xAx^{-1} \cap A) & \xrightarrow{t_x} & H^*(A) \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
\Sigma H^*(T') & \xrightarrow{\Sigma \text{ad}_x} & \Sigma H^*(xT'x^{-1}) & \xrightarrow{\Sigma i_x} & \Sigma H^*(xT'x^{-1} \cap T) & \xrightarrow{\Sigma t_x} & H^*(T)
\end{array}$$

where $A = \times_{r=1}^n (\mathbb{S}_{p^{m_r}})$, T' runs through all inclusions of $\times^m Z/p$ in A and $T = \times_{r=1}^n T_{m_r, m_r}$.

The only $\mathcal{S}_{(m_1, \dots, m_n)}$ invariant classes not in image $\Phi_{m_1, \dots, m_n}$ are classes $u' = u_{1, m_1} \otimes \dots \otimes u_{n, m_n}$ containing $(u_{r_0, m_{r_0}})^{\otimes p} \in \bigotimes^p H^*(T_{m_{r_0}, m_{r_0}})$ as a factor. Recall $u^{\otimes p} \xleftarrow{(1 \times \Delta)^*} \mathcal{P}(u) \xrightarrow{(1 \times \Delta)^*} (1 \times \Delta)^*(\mathcal{P}(u))$. Thus u' is in the

$$\text{image} \left(I_{m_1, \dots, m_n} \circ \left(\prod_{r=1}^n k_{m_r, m_r} \right) \right)^* : H^*(\mathfrak{S}_p) \rightarrow \bigotimes_{r=1}^n H^*(T_{m_r, m_r}).$$

Hence we have

6.3. LEMMA. $\text{Image}(I_{m_1, \dots, m_n} \circ (\prod_{r=1}^n k_{m_r, m_r}))^* \cong \mathfrak{S}_{(m_1, \dots, m_n)}$ invariant classes of $\bigotimes_{r=1}^n H^*(T_{m_r, m_r})$.

This proves Theorems B and D. A trivial modification of 6.2 and 6.3 proves Theorem F. As 3.1 shows the only multiple image classes are generated by the $\mathcal{P}(\cdot)$'s, Theorem C follows, up to constants. Using the notation of Theorem C if $x_{m,i-1} = M_{i-2,i-1} M_{i-3,i-1} L_{i-1}^{p-3}$ then 5.4 gives $x_{m,i} = -1(M_{i-1,i} M_{i-2,i} L_i^{p-3})$. If $x_{m,i-1} = Q_{i-2,i-1}$ then direct computation shows the constant c in 5.5 is 1 hence $x_{m,i} = Q_{i-1,i}$. It is easy to see that application of the Steenrod p th powers or direct computation yield that the constant is $+1$ for multiple image polynomial generators and -1 for even dimensional multiple image exterior generators.

VII. Proof of Theorem G.

PROOF OF (1). Let $k_{j,\infty}^*(u) = \mathfrak{S} \langle x_1, \dots, x_m, 1, \dots \rangle$. As j is the smallest integer such that $k_{j,\infty}^*(u) \neq 0$ it follows that at least one x_h contains a factor equal to L_j^{p-1} , $\underline{L}_j L_j^{p-2}$, $M_{g,j} \underline{L}_j L_j^{p-3}$, or $M_{g,j} L_j^{p-2}$. If $k_{j,\infty}^*(u)$ has at least one representative of the form $\underline{L}_{n,j}(\dots)$ with p not dividing n then $\beta_p(k_{j,\infty}^*(u)) = \sum n L_{n,j}(\dots) + B \neq 0$ (where B cannot contain terms in the first sum). Similarly if some $x_h = M_{g,j} \underline{L}_j L_j^{p-3} Y$ and no $x_{h'} = M_{g,j} L_j^{p-2} Y$ then $\beta_p(k_{j,\infty}^*(u)) \neq 0$. Suppose every time the term $M_{g,j} \underline{L}_j L_j^{p-3} Y$ appears the term $M_{g,j} L_j^{p-2} Y$ also appears; then if $k_{j,\infty}^*(u) \neq \underline{L}_{n,j}(\dots) Y$ must be a product of $Q_{h,j}$'s. It is then easy to construct a class u' such that $\beta_p(u') = u$ (just replace one $M_{g,j} L_j^{p-2} Y$ by $M_{g,j} \underline{L}_j L_j^{p-3} Y$). If $\beta_p(u) = 0$ and $M_{g,j} L_j^{p-2} Y$ appears a similar construction yields u' such that $\beta_p(u') = u$. The only possibility left is $\beta_p(u) = 0$, and $k_{j,\infty}^*(u) = L_{n,j}(\dots)$. Then $\beta_p(u') = u$ where $k_{j,\infty}^*(u') = \underline{L}_{n,j}$.

PROOF OF (2). We need the following

THEOREM [2]. Let $r \geq 2$. In homology with the loop sum multiplication if $d^{r-1}(a) = b$ then $d^r(a^p) = a^{p-1}b$.

PROOF. Theorem 5.4 of [2].

The homology and cohomology Bockstein spectral sequences are Hopf algebra duals and Theorem F gives the loop sum coalgebra map in cohomology. If a, b in $H_*(Q(S^0)_0)$ are dual to u, v respectively then Theorem F gives $\langle u', a^p \rangle = 1$. Now u' is not dual to a^p on the E_1 level; in fact $(u')^* = a^p + \sum a_i$. It is easy to see however that the a_i are all dual to classes u'' where $k_{j,\infty}^*(u'') = \mathfrak{S} \langle x_1, \dots, x_t, 1, \dots \rangle$ with $t < pm$.

Many times it is easy to see that the a_i classes do not live to E_r . Such is the case with Corollary 1 as induction on r and the fact that $\{L_{p^{m_j}}(x: 1, \dots)\}_{m=1}^{r-1}$ generate the subalgebra $\{L_{n_j}(x: 1, \dots)\}$ (where $n = 1, \dots, p^r - 1$) prove the corollary.

PROOF OF COROLLARY 2. The reduction homomorphism $j_r: H^*(, Z/p^r) \rightarrow E_r$ is onto and if $k_{i,i}^*(u) = Q_{j,i}$ then $k_{j,i}^*(u) = R_i^*(L_{p^r/j}(1: 1, \dots))$.

Appendix. We give a proof that the quotient determinants, $Q_{j,i} \in \mathbb{Q}_i$ are integral mod p . L_i has an explicit factorization first discovered by E. H. Moore in 1896

LEMMA [19]. $L_i = \prod_{(m_1, \dots, m_i)} (m_1 b_1 + \dots + m_i b_i)$ where $(m_1, \dots, m_i)$ runs over all elements of $T_{i,i}$ with first nonzero coefficient equal to one.

PROOF. (Compare with [8, p. 76].) L_i is invariant under the special linear group $SL(i, Z/p)$ which acts transitively on the nonzero elements of $T_{i,i}$. Since b_1 is a factor of L_i it follows that $\alpha(b_1) = m_1 b_1 + \dots + m_i b_i$ is a factor as well. Hence the product above divides L_i (the factors are all relatively prime). But both sides have the same degree, hence they differ only up a constant factor. But the diagonal term $b_1^{p^i-1} b_2^{p^i-2} \dots b_i$ occurs in both sides only once and each time with coefficient 1.

More generally b_1 is a factor of the numerator of $Q_{j,i}$ for every j , so L_i is also a factor of the numerator of $Q_{j,i}$ by the above argument. This gives:

LEMMA. $Q_{j,i}$ is a nontrivial polynomial invariant under $GL(i, Z/p)$.

REFERENCES

1. S. Araki and T. Kudo, *Topology of H_n -spaces and H -squaring operations*, Mem. Fac. Sci. Kyusyu Univ. Ser. A **10** (1956), 85–120.
2. W. Browder, *Homotopy commutative H -spaces*, Ann. of Math. (2) **75** (1962), 283–311. MR **27** #765.
3. G. Brumfiel, I. Madsen, and R. J. Milgram, *PL-characteristic classes and cobordism*, Ann. of Math. (2) **97** (1973), 82–159.
4. H. Cárdenas, *El algebra de cohomologia del grupo simétrico de grado p^2* , Bol. Soc. Mat. Mexicana **10** (1965), 1–30.
5. H. Cartan and S. Eilenberg, *Homological algebra*, Princeton Univ. Press, Princeton, N. J., 1956. MR **17**, 1040.
6. A. Cauchy, *Exercices d'analyse et de physique mathématique*. III, Paris, 1844.
7. L. E. Dickson, *The Madison Colloquium Lectures on Mathematics*, Amer. Math. Soc. Colloq. Publ., vol. 4, Amer. Math. Soc., Providence, R. I., 1913, pp. 33–40.
8. ———, *A fundamental system of invariants of the general modular linear group with a solution of the form problem*, Trans. Amer. Math. Soc. **12** (1911), 75–98.
9. A. Dold, *Homology of symmetric products and other functors of complexes*, Ann. of Math. (2) **68** (1958), 54–80. MR **20** #3537.

10. E. Dyer and R. Lashof, *Homology of iterated loop-spaces*, Amer. J. Math. **84** (1962), 35–88. MR **25** #4523.
11. S. Eilenberg and S. Mac Lane, *On the group $H(\pi, n)I$* , Ann. of Math. (2) **58** (1953), 55–106. MR **15**, 54.
12. H. Hopf, *Über die Bettischen Gruppen, die zu einer beliebigen Gruppe gehören*, Comment. Math. Helv. **17** (1945), 39–79. MR **6**, 279.
13. I. Madsen and R. J. Milgram, *The universal smooth surgery class*, Comment. Math. Helv. **50** (1975), 281–310. MR **52** #4285.
14. ———, *Higher torsion in SG and BSG*, Math. Z. **143** (1975), 55–80. MR **51** #11503.
15. J. P. May, *Homology operations on infinite loop spaces*, Algebraic topology (Proc. Sympos. Pure Math., Vol. 22, Univ. Wisconsin, Madison, Wis., 1970), Amer. Math. Soc., Providence, R. I., 1971, pp. 171–185. MR **47** #7740.
16. R. J. Milgram, *The mod 2 spherical characteristic classes*, Ann. of Math. (2) **92** (1970), 238–261. MR **41** #7705.
17. ———, *The homology of symmetric products*, Trans. Amer. Math. Soc. **138** (1969), 251–265. MR **39** #3483.
18. ———, *Surgery with coefficients*, Ann. of Math. (2) **100** (1974), 194–248. MR **50** #14801.
19. E. H. Moore, *A two-fold generalization of Fermat's theorem*, Bull. Amer. Math. Soc. **2** (1896), 189–199.
20. M. Morse, *The calculus of variations in the large*, Amer. Math. Soc. Colloq. Publ., vol. 18, Amer. Math. Soc., Providence, R. I., 1934.
21. M. Nakaoka, *Decomposition theorem for homology groups of symmetric groups*, Ann. of Math. (2) **71** (1960), 16–42. MR **22** #2989.
22. ———, *Homology of the infinite symmetric group*, Ann. of Math. (2) **73** (1961), 229–257. MR **24** #A1721.
23. G. Nishida, *Cohomology operations in iterated loop spaces*, Proc. of the Japan Acad. **44** (1968), 104–109. MR **39** #2156.
24. S. Priddy, *On $\Omega^\infty S^\infty$ and the infinite symmetric group*, Algebraic topology (Proc. Sympos. Pure Math., Vol. 22, Univ. Wisconsin, Madison, Wis., 1970), Amer. Math. Soc., Providence, R. I., 1971, pp. 217–220. MR **50** #11226.
25. D. Quillen, Unpublished.
26. ———, *On the cohomology and K-theory of the general linear groups over a finite field*, Ann. of Math. (2) **96** (1972), 552–586. MR **47** #3565.
27. ———, *The Adams conjecture*, Topology **10** (1971), 67–80.
28. P. A. Smith and M. Richardson, *Periodic transformations of complexes*, Ann. of Math. (2) **39** (1938), 611–633.
29. N. E. Steenrod, *Products of cocycles and extensions of mappings*, Ann. of Math. (2) **48** (1947), 290–320. MR **9**, 154.
30. ———, *Homology groups of symmetric groups and reduced power operations*, Proc. Nat. Acad. Sci. U.S.A. **39** (1953), 213–217. MR **14**, 1005.
31. N. E. Steenrod and D. B. A. Epstein, *Cohomology operations*, Ann. of Math. Studies, No. 50, Princeton Univ. Press, Princeton, N. J., 1962. MR **26** #3056.
32. J. Tornehave, *Developing the Quillen map*, Thesis, M. I. T., 1971.
33. A. Tsuchiya, *Characteristic classes for spherical fiber spaces*, Nagoya Math. J. **43** (1971), 1–39. MR **45** #7736.
34. ———, *Characteristic classes for PL micro bundles*, Nagoya Math. **43** (1971), 169–198. MR **47**, 2614.
35. B. Cooper, *Cohomology of symmetric groups*, Augsburg College, 1975 (mimeo).
36. H. Mui, *Modular invariant theory and cohomology algebras of symmetric groups*, J. Fac. Sci. Univ. Tokyo **22** (1975), 319–369.
37. F. Cohen, T. Lada and J. P. May, *The homology of iterated loop spaces*, Lecture Notes in Math., vol. 533, Springer-Verlag, Berlin and New York, 1976.

38. H. Ligaard, *On the Adams spectral sequence for $\Pi_*(MSTOP)$ and infinite loop maps from SF to $BO_{\otimes}$ at the prime 2*, Thesis, University of Chicago, 1977.

39. B. Mann and R. J. Milgram, *On the Chern classes of the regular representations of some finite groups*, Stanford Univ., 1975 (mimeo).

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, STANFORD, CALIFORNIA 94305

Current address: Department of Mathematics, Harvard University, Cambridge, Massachusetts 02138