ON A SUFFICIENT CONDITION FOR PROXIMITY

BY

KA-SING LAU¹

ABSTRACT. A closed subspace M in a Banach space X is called U -proximal if it satisfies: $(1 + \rho)S \cap (S + M) \subseteq S + \varepsilon(\rho)(S \cap M)$, for some positive valued function $\varepsilon(\rho)$, $\rho > 0$, and $\varepsilon(\rho) \rightarrow 0$ as $\rho \rightarrow 0$, where S is the closed unit ball of X . One of the important properties of this class of subspaces is that the metric projections are continuous. We show that many interesting subspaces are U -proximal, for example, the subspaces with the 2-ball property (semi M -ideals) and certain subspaces of compact operators in the spaces of bounded linear operators.

1. Introduction. We call a closed subspace M of a real Banach space X an M -ideal if the annihilator $M^\perp$ of M is an L -summand in X^* . This notion was formulated and studied by Alfsen and Effros [1]. It was proved that if M is an M -ideal, then M is a proximal subspace of X [1], [5]. In [4], Hennefeld showed that the space of compact operators on l^p (or c_0), $1 < p < \infty$, is an M -ideal in the space of bounded linear operators on l^p (or c_0 respectively). This theorem is also true for operators from l^p into l^q , $1 < p < q < \infty$ [11]. (It is well known that if $1 < q < p < \infty$, then every bounded linear operator from l^p into l^q is compact.) M -ideal theory provides a convenient tool to study the approximation of operators by the space of compact operators and has been investigated by many authors [3], [4], [5], [9], [11], [15], [16]. However, in some cases, the class of M -ideals appears to be too restricted; for example, the space of compact operators on l^1 is not an M -ideal in the space of bounded linear operators on l^1 [16]. It is our attempt to consider another sufficient condition for proximity which preserves certain important properties of M -ideals and also includes some other interesting classes of proximal subspaces.

Motivated by a lemma of Holmes in [5], we call a closed subspace M of a Banach space X U -proximal if there exists a positive function $\varepsilon(\rho)$, $\rho > 0$, with $\varepsilon(\rho) \rightarrow 0$ as $\rho \rightarrow 0$ and satisfies

$$(1 + \rho)S \cap (S + M) \subseteq S + \varepsilon(\rho)(S \cap M), \quad \rho > 0,$$

Received by the editors February 3, 1978 and, in revised form, July 24, 1978.

AMS (MOS) subject classifications (1970). Primary 41A65; Secondary 46B99.

Key words and phrases. Compact operators, measurable functions, M -ideals, proximity, uniformly convex.

¹Supported in part by the Faculty of Arts and Sciences research grant of the University of Pittsburgh.

where S denotes the closed unit ball of X . Examples of U -proximal subspaces are:

- (i) X is uniformly convex and M is a closed subspace of X ;
- (ii) $X = B(K)$, the space of bounded functions on a topological space K and $M = C(K)$, the space of bounded continuous functions on K ;
- (iii) M is a semi M -ideal or semi L -summand in a Banach space X ;
- (iv) $X = L^\infty(\Omega, L^1)$, the space of bounded Bochner measurable functions from a σ -finite measure space Ω into L^1 and M is the subspace of f in X such that $f(\Omega)$ is weakly precompact. In particular, if we let $L(E, F)$ ($K(E, F)$) denote the space of bounded linear operators (compact operators, respectively), then $K(L^1(\Omega), l^1)$ is proximal in $L(L^1(\Omega), l^1)$.

Our paper is divided into six sections. In §2, we define some basic terminologies and give several reformulations of the definition of U -proximity. The metric projection from X into a proximal subspace M is the map P which sends $x \in X$ to the set of best approximations from M to x . The study of the continuity of metric projections is an important component of the theory of best approximation. In §3, we show that if M is U -proximal, then the metric projection P is continuous (with respect to the Hausdorff metric in the range). We also give a condition for P to be Lipschitz continuous. In §§4–6, we show that the examples listed above are in fact U -proximal subspaces.

ACKNOWLEDGEMENT. The author is indebted to Professor T. A. Metzger for some suggestions in the first draft. He would also like to thank the referee for bringing his attention to several recent publications in this subject.

2. Definitions and preliminaries. Let X be a real Banach space, let $S_r(X)$ (or S_r) denote the closed ball of radius r centered at 0; $S_1(X) = S(X)$ ($S_1 = S$). If M is a closed subspace of X , then for $x \in X$, we denote the subspace generated by M and x by $\langle M, x \rangle$, and denote $S_r(\langle M, x \rangle)$ by $S_r[x]$.

Let $F(X)$ be a family of nonempty bounded closed subsets of X . For any $A, B \in F(X)$, we define

$$d_H(A, B) = \inf\{r: A \subseteq B + S_r \text{ and } B \subseteq A + S_r\}.$$

Then d_H is a metric on $F(X)$ and is called the *Hausdorff metric*.

For $x \in X$ and for any subset A in X , we define another distance function $d(x, A) = \inf\{\|x - z\|: z \in A\}$. A point y in a closed subspace M of X is called a *best approximation* from M to x if $\|x - y\| = d(x, M)$. M is called a *proximal subspace* of X if every $x \in X$ has a best approximation from M . Propositions 2.2 and 2.3 will give the motivation of a simple sufficient condition for M to be a proximal subspace.

LEMMA 2.1. Let g be a concave function defined on $[0, 1]$ with $g(0) > 0$ and $g(1) = 0$. Let h be a function defined on $[0, a]$, $a > 1$, with $h(x) = ag(x/a)$, $x \in [0, a]$. Then $h(1) > h(x) - g(x)$ for $0 < x < 1$.

PROOF. Note that the derivatives $h'(x)$, $g'(x)$ exist and decrease almost everywhere. Hence

$$h'(x) - g'(x) = g'\left(\frac{x}{a}\right) - g'(x) > 0 \quad \text{a.e.}$$

and

$$h(1) = h(1) - g(1) > h(x) - g(x), \quad 0 < x < 1. \quad \square$$

Let X be a two dimensional normed linear space. For $\rho > 0$, let L be a line which is tangent to S at w and cuts the sphere $\{z: \|z\| = 1 + \rho\}$ at x and y . Let $(x': y')$ be any open line segment which is parallel to L , with $\|x'\| = 1 + \rho$, $\|y'\| = 1$ and does not intersect S . Lemma 2.1 implies that $\|x' - y'\| < \max\{\|x - w\|, \|y - w\|\}$.

Let M be a proximal subspace in X , let $x \in X \setminus M$ and let $X_1 = \langle M, x \rangle$. Let $f \in X_1^*$ be such that $f^{-1}(0) = M$, $\|f\| = 1$ and define

$$\alpha(x, \rho) = d_H((1 + \rho)S[x] \cap f^{-1}(1), S[x] \cap f^{-1}(1)), \quad \rho > 0.$$

It follows from simple geometry that for each $x \in X \setminus M$,

$$|\alpha(x, \rho) - \alpha(x, \rho')| < \frac{2|1 + \rho + \rho'|}{\rho} |\rho - \rho'| \quad \text{for } \rho > 0.$$

Hence $\alpha(x, \cdot)$ is continuous on $\mathbb{R}^+$.

PROPOSITION 2.2. Let M be a proximal subspace of a Banach space X and let $\alpha(x, \rho)$ be defined as above. Then

$$\alpha(x, \rho) = \inf\{r > 0: (1 + \rho)S \cap (S[x] + M) \subseteq S + rS(M)\}.$$

PROOF. Without loss of generality, we may assume that $X = \langle M, x \rangle$. It is clear from the definition of $\alpha(x, \rho)$ that

$$\alpha(x, \rho) < \inf\{r > 0: (1 + \rho)S \cap (S + M) \subseteq S + rS(M)\}.$$

To prove the reverse inequality, let $0 < \varepsilon < 1$ and let $y \in (1 + \rho)S \cap (S + M)$. We will consider the case $y \notin M$ first. Assume $f(y) > 0$ (the case $f(y) < 0$ is similar), and let $y_1 = ty \in f^{-1}(1)$ for some $t > 0$. There exists $w_1 \in S \cap f^{-1}(1)$ such that

$$\|y_1 - w_1\| < d(y_1, S \cap f^{-1}(1)) + \varepsilon.$$

If $\|y_1\| < 1 + \rho$, we define $x_1 = y_1$, otherwise we define x_1 to be the point on the line segment of $(y_1 : w_1)$ with $\|x_1\| = 1 + \rho$. In either case, it can be shown that $\|x_1 - w_1\| < \alpha(x, \rho) + \varepsilon$. By the remark before the proposition,

we can choose a $z \in f^{-1}(f(y)) \cap \{w: \|w\| = 1\}$ such that $\|y - z\| < \alpha(x, \rho) + \varepsilon$. Hence

$$y = z + (y - z) \in S + (\alpha(x, \rho) + \varepsilon)S(M). \quad (1)$$

If $y \in M$, we can choose $y' \in (1 + \rho)S \setminus M$ with $\|y' - y\| < \varepsilon/2(1 + \rho)$ and $z' \in f^{-1}(f(y')) \cap S$ satisfies (1). Let

$$z'' = \|z' - (y' - y)\|^{-1}(z' - (y' - y)).$$

Then z'' is in M since

$$f(z' - (y' - y)) = f(z') - f(y') = 0.$$

That

$$\|y - z''\| \leq \varepsilon + (1 + \varepsilon)(\alpha(x, \rho) + \varepsilon) = \beta$$

implies that

$$y = z'' + (y - z'') \in S + \beta S(M).$$

Since this is also true for the $y \notin M$ as in (1) and since ε is arbitrary, we conclude that

$$\alpha(x, \rho) \geq \inf\{r > 0: (1 + \rho)S \cap (S + M) \subseteq S + rS(M)\}. \quad \square$$

The following proposition is the foundation of this paper; the proof is similar to [5, Lemma 2].

PROPOSITION 2.3. *Let M be a closed subspace of X and let $x \in X \setminus M$. Suppose there exists a function $\varepsilon: \mathbf{R}^+ \rightarrow \mathbf{R}^+$ (depending on x) such that $\varepsilon(\rho) \rightarrow 0$ as $\rho \rightarrow 0$ and*

$$(1 + \rho)S \cap \overline{(S[x] + M)} \subseteq \overline{S + \varepsilon(\rho)S(M)}, \quad \rho > 0.$$

Then x has a best approximation from M .

Furthermore, if $\|x\| \leq 1 + \rho_0$ and if $\inf\{\|x - z\|: z \in M\} < 1$. Then given any $r > \varepsilon(\rho_0)$, there exists a best approximation z_0 in M such that $\|z_0\| < r$.

PROOF. Without loss of generality, assume that $\|x\| = 1 + \rho_0$ and $\inf\{\|x - z\|: z \in M\} = 1$. We claim that $x \in \overline{S[x] + M}$. Indeed, a sequence $\{z_n\}$ in M can be chosen such that $\|x - z_n\| \rightarrow 1$. Let

$$x_n = \|x - z_n\|^{-1}(x - z_n) + z_n,$$

then the sequence $\{x_n\}$ is contained in $S[x] + M$ and $x_n \rightarrow x$.

Choose a sequence of positive numbers $\{\rho_n\}$ such that $\rho_n \rightarrow 0$ and $\sum_{n=0}^{\infty} \varepsilon(\rho_n) < r$ where $r > \varepsilon(\rho_0)$. By hypothesis there exists $z_1 \in \varepsilon(\rho_0)S(M)$ with $\|x - z_1\| \leq 1 + \rho_1$. Note that $x - z_1$ is also in $\overline{S[x] + M}$, the same argument yields a z_2 in $\varepsilon(\rho_1)S(M)$ with $\|x - z_1 - z_2\| \leq 1 + \rho_2$. Inductively, we can find a sequence $\{z_n\}$ such that $z_n \in \varepsilon(\rho_{n-1})S(M)$ and $\|x - \sum_{k=1}^n z_k\| \leq 1 + \rho_n$. Let $z_0 = \sum_{n=1}^{\infty} z_n$, then $\|z_0\| < \sum_{n=0}^{\infty} \varepsilon(\rho_n) < r$ and $\|x - z_0\| = 1$. This completes the proof. $\square$

PROPOSITION 2.4. *Let M be a closed subspace in X and let $x \in X \setminus M$. Then the following conditions are equivalent:*

(i) *there exists an $\varepsilon: \mathbf{R}^+ \rightarrow \mathbf{R}^+$ such that $\varepsilon(\rho) \rightarrow 0$ as $\rho \rightarrow 0$ and*

$$(1 + \rho)S \cap (\overline{S[x] + M}) \subseteq \overline{S + \varepsilon(\rho)S(M)}, \quad \rho > 0.$$

(ii) *there exists an $\varepsilon': \mathbf{R}^+ \rightarrow \mathbf{R}^+$ such that $\varepsilon'(\rho) \rightarrow 0$ as $\rho \rightarrow 0$ and*

$$(1 + \rho)S \cap (S[x] + M) \subseteq S + \varepsilon'(\rho)S(M), \quad \rho > 0.$$

(iii) *there exists an $\varepsilon: \mathbf{R}^+ \rightarrow \mathbf{R}^+$ such that $\tilde{\varepsilon}$ is increasing, continuous, $\tilde{\varepsilon}(\rho) \rightarrow 0$ as $\rho \rightarrow 0$ and*

$$(1 + \rho)S \cap (S[x] + M) \subseteq S + \tilde{\varepsilon}(\rho)S(M), \quad \rho > 0.$$

Moreover, if one of the three conditions holds, the function $\alpha(x, \rho)$ defined in Proposition 2.2 converges to zero as $\rho \rightarrow 0$.

PROOF. (i) $\Rightarrow$ (ii). Let $y \in (1 + \rho)S \cap (S[x] + M)$. By Proposition 2.3 if $r = 2\varepsilon(\rho)$, then there exists a $z \in M$ such that $\|y - z\| < 1$ and $\|z\| < 2\varepsilon(\rho)$. Let $\varepsilon'(\rho) = 2\varepsilon(\rho)$, then $y \in S + \varepsilon'(\rho)S(M)$ and (ii) follows. (ii) $\Rightarrow$ (i) is clear. To prove (ii) $\Rightarrow$ (iii), we can take $\tilde{\varepsilon}(\rho) = \alpha(x, \rho) + \rho$, then $\tilde{\varepsilon}$ is an increasing, continuous function. Proposition 2.2 implies that $\alpha(x, \rho) < \tilde{\varepsilon}(\rho)$. Hence $\tilde{\varepsilon}(\rho) \rightarrow 0$ as $\rho \rightarrow 0$. That (iii) $\Rightarrow$ (ii) is obvious. $\square$

DEFINITION 2.5. *Let M be a closed subspace of a Banach space X , we say that M is locally U -proximal if there exists a function $\varepsilon: (X \setminus M) \times \mathbf{R}^+ \rightarrow \mathbf{R}^+$ such that for each fixed x , $\varepsilon(x, \cdot)$ is continuous, increasing on ρ , $\varepsilon(x, \rho) \rightarrow 0$ as $\rho \rightarrow 0$ and*

$$(1 + \rho)S \cap (S[x] + M) \subseteq S + \varepsilon(x, \rho)S(M), \quad x \in M, \quad \rho > 0.$$

M is called U -proximal if the function ε can be chosen independent of x and

$$(1 + \rho)S \cap (S + M) \subseteq S + \varepsilon(\rho)S(M).$$

It follows from Proposition 2.3 that locally U -proximal and U -proximal subspaces are proximal.

3. Metric projections. Let M be a proximal subspace of X , for each $x \in X$, we define $P(x)$ to be the set of best approximations from M to x . We call the map P from X into $F(M)$, the family of bounded closed subsets in M , as the *metric projection* from X into M .

LEMMA 3.1. *Let M be a locally U -proximal subspace in X . For each $x \in X \setminus M$, let $r_x = d(x, M)$. Then*

$$M \cap (S_{(1+\rho)r_x} + x) \subseteq P(x) + r_x \varepsilon(x, \rho)S(M), \quad \rho > 0.$$

PROOF. Without loss of generality, we assume that $r_x = 1$. For $y \in M \cap (S_{(1+\rho)} + x)$, it follows that $\|x - y\| < 1 + \rho$. That

$$x - y = (x - z) + (z - y)$$

where z is the best approximation of x , implies that $x - y \in S + M$. Hence

$$(x - y) \in (1 + \rho)S \cap (S + M).$$

Since M is locally U -proximal, $x - y = w + v$ for some w in S and v in $\varepsilon(x, \rho)S(M)$. Note that $(x - w)$ is in M and, in fact, a best approximation to x (for $1 \leq \|x - (x - w)\| = \|w\| \leq 1$). Hence

$$y = (x - w) - v \in P(x) + \varepsilon(x, \rho)S(M). \quad \square$$

LEMMA 3.2. *Let M be a locally U -proximal subspace in X . Then for any $x, y \in X \setminus M$ with $\|x - y\| \leq \rho$,*

$$d_H(P(x), P(y)) \leq \max \left\{ r_x \varepsilon \left(x, \frac{2\rho}{r_x} \right), r_y \varepsilon \left(y, \frac{2\rho}{r_y} \right) \right\}.$$

PROOF. Since $r_y \leq r_x + \rho$ and

$$(S_y + y) \subseteq (S_{r_x+2\rho} + x),$$

it follows that

$$P(y) \subseteq M \cap (S_{r_x+2\rho} + x).$$

By Lemma 3.1, we have

$$P(y) \subseteq P(x) + r_x \varepsilon \left(x, \frac{2\rho}{r_x} \right) S(M).$$

A similar argument yields that

$$P(x) \subseteq P(y) + r_y \varepsilon \left(y, \frac{2\rho}{r_y} \right) S(M),$$

and the lemma follows easily from these inclusions and the definition of the Hausdorff metric on $F(M)$. $\square$

THEOREM 3.3. *Let M be a locally U -proximal subspace in X . Suppose for each $\rho > 0$, $\varepsilon(\cdot, \rho)$ is an upper semicontinuous function on $X \setminus M$. Then P is a continuous function from X into $F(M)$.*

PROOF. It is easy to show that P is continuous for $x \in M$. Let $x \in X \setminus M$, for any $\delta > 0$, there exists an η in $(0, 1)$ such that $\varepsilon(x, \eta) < \delta/2$. Since $\varepsilon(\cdot, \eta)$ is upper semicontinuous, there exists a ρ , $0 < \rho < (\eta \cdot r_x)/4$, such that for $\|x - y\| \leq \rho$, $\varepsilon(y, \eta) < \varepsilon(x, \eta) + \delta/2$. Hence for $\|x - y\| \leq \rho$, we have

$$\begin{aligned} d_H(P(x), P(y)) &\leq \max \left\{ r_x \varepsilon \left(x, \frac{2\rho}{r_x} \right), r_y \varepsilon \left(y, \frac{2\rho}{r_y} \right) \right\} \\ &\leq 2r_x \max \left\{ \varepsilon \left(x, \frac{2\rho}{r_x} \right), \varepsilon \left(y, \frac{4\rho}{r_x} \right) \right\} \\ &\leq 2r_x \delta. \end{aligned}$$

This shows that P is continuous at x . $\square$

THEOREM 3.4. *Let M be a U -proximal subspace of X . Then the metric projection $P: X \rightarrow F(M)$ is continuous.*

PROOF. In this case, we have $\epsilon(x, \rho) = \epsilon(y, \rho) = \epsilon(\rho)$ for any $x, y \in X \setminus M$. Theorem 3.3 implies that P is continuous. $\square$

COROLLARY 3.5. *Let M be a U -proximal subspace of X . Then the metric projection P admits a continuous selection $s: X \rightarrow M$ (i.e., $s(x) \in P(x)$).*

The following also follows easily from Lemma 3.2:

THEOREM 3.6. *Let M be a subspace of X . Suppose there exists a $k > 0$ such that*

$$(1 + \rho)S \cap (S + M) \subseteq S + k\rho S(M), \quad \rho > 0.$$

Then the metric projection $P: X \rightarrow F(M)$ is a Lipschitz continuous function with Lipschitz constant not greater than $2k$.

In the next three sections, we will see that many interesting examples will satisfy the above inclusion. We remark that the converse of the theorem is not true; for example, let M be a closed subspace of a Hilbert space X , then the metric projection $P: X \rightarrow M$ is a Lipschitz function but $\epsilon(\rho) > \sqrt{\rho^2 + 2\rho}$. We also remark that a U -proximal subspace may not have a uniformly continuous metric projection. Examples (certain closed subspaces in some uniformly convex spaces) can be found in [7], [14]. In general, a set valued function which satisfies the Lipschitz condition does not admit Lipschitz selection; it will be interesting to investigate this question for the metric projections in Theorem 3.6.

4. Some examples. In this section, we will give some simple examples of locally U -proximal and U -proximal subspaces.

PROPOSITION 4.1. *Let M be a finite dimensional subspace in a Banach space X . Then M is locally U -proximal.*

PROOF. Let $x \in X \setminus M$ and assume that $X = \langle M, x \rangle$. Let $\epsilon(x, \rho) = \alpha(x, \rho) + \rho$, then it follows from the compactness of the unit ball that $\epsilon(x, \rho) \rightarrow 0$ as $\rho \rightarrow 0$ and by Proposition 2.2,

$$(1 + \rho)S \cap (S + M) \subseteq S + \epsilon(x, \rho)S(M). \quad \square$$

A Banach space X is called *locally uniformly convex* if for each $x \in X$ with $\|x\| = 1$ and for any $\eta > 0$, there exists a $\delta > 0$ such that for any $y \in X$ with $\|y\| \leq 1$ and $\|x - y\| \geq \eta$, $\|x + y\| \leq 2(1 - \delta)$. X is called *uniformly convex* if the δ can be chosen independent of x .

PROPOSITION 4.2. *Let X be a locally uniformly convex space. Then every proximal subspace is locally U -proximal.*

PROOF. Let M be a proximal subspace of X and let $x \in X \setminus M$. Let $X_1 = \langle M, x \rangle$ and let $f \in X_1^*$ with $f^{-1}(0) = M$ and $\|f\| = 1$. If $y \in S$ with $f(y) = 1$ and $\alpha(x, \rho)$ is defined as in Proposition 2.2, then

$$\begin{aligned}\alpha(x, \rho) &= d_H((1 + \rho)S[x] \cap f^{-1}(1), \{y\}) \\ &< \text{diam}((1 + \rho)S[x] \cap f^{-1}(1)).\end{aligned}$$

By the locally uniformly convexity of the norm on X_1 , we can show that the last term approaches to zero as $\rho \rightarrow 0$. Let $\varepsilon(x, \rho) = \alpha(x, \rho) + \rho$; it follows that M is locally U -proximal. $\square$

PROPOSITION 4.3. *Let X be a uniformly convex space. Then every closed subspace is U -proximal.*

PROOF. Let $|L|$ denote the length of a line segment L and define

$$\varepsilon(\rho) = \sup\{|L|: L \text{ is in } (1 + \rho)S \setminus S\} + \rho.$$

It is clear that $\varepsilon(\rho) \rightarrow 0$ as $\rho \rightarrow 0$ and $\varepsilon(\rho) > \alpha(x, \rho) + \rho$ where $\alpha(x, \rho)$ is defined as in Proposition 2.2. Hence

$$(1 + \rho)S \cap (S + M) \subseteq S + \varepsilon(\rho)S(M)$$

and the proof is completed. $\square$

Let K be a topological space, let $B(K)$ be the space of bounded continuous functions on K with the supremum norm and let $C(K)$ be the subspace of bounded continuous functions in $B(K)$. It is well known that $C(K)$ is a proximal subspace in $B(K)$ [6].

PROPOSITION 4.4. *Let M be a closed subspace in $X = B(K)$ such that for each $h \in M$, $h \wedge \rho$, $h \vee (-\rho)$, $\rho > 0$, are also in M . Then M satisfies*

$$(1 + \rho)S \cap (S + M) \subseteq S + \rho S(M), \quad \rho > 0.$$

Thus M is U -proximal, and in particular, $C(K)$ is U -proximal.

PROOF. For any $f \in (1 + \rho)S \cap (S + M)$, we can write $f = g + h$ where $\|g\| \leq 1$ and $h \in M$. Let $h' = (h \wedge \rho) \vee (-\rho)$; then $h' \in M$ and $\|h'\| \leq \rho$. Let $g' = f - h'$, we need only show that $\|g'\| \leq 1$ and hence

$$f = g' + h' \in S + \rho S(M).$$

If $|h(x)| \leq \rho$, then $|g'(x)| = |g(x)| \leq 1$. If $h(x) > \rho$, then three cases arise:

(i) if $g(x) \geq 0$, then $|g'(x)| = f(x) - \rho \leq 1$;

(ii) if $g(x) < 0$ and $f(x) \geq 0$, then

$$|g'(x)| = |f(x) - \rho| \leq (1 + \rho) - \rho = 1;$$

(iii) if $g(x) < 0$ and $f(x) < 0$, then $|g(x)| = |f(x)| + h(x)$. This implies $|f(x)| < 1 - \rho$ and

$$|g'(x)| = |f(x) - \rho| \leq |f(x)| + \rho \leq 1.$$

If $h(x) < -\rho$, a similar proof shows that $|g'(x)| \leq 1$. Hence $\|g'\| \leq 1$ and the proof is completed. $\square$

5. Semi- M -ideals. A convex subset F in a convex set K is a *face* of K if $\lambda y + (1 - \lambda)z \in F, y, z \in K, 0 < \lambda < 1$ implies $y, z \in F$. Let X be a Banach space and let $x \in S$ with $\|x\| = 1$, we use $\text{face}(x)$ to denote the maximal proper face in S containing x .

Let J be a closed subspace of a Banach space X and let

$$J' = \left\{ x \in X: x = 0 \text{ or } \text{face}\left(\frac{x}{\|x\|}\right) \cap J = \emptyset \right\}.$$

J is called a *semi- L -summand* if each $x \in X$ has a unique decomposition as $x = y + z$ with $y \in J, z \in J'$ and $\|x\| = \|y\| + \|z\|$. J is called an *L -summand* if J' is a closed linear subspace of X . For detailed discussion of these and the semi- M -ideals, M -ideals, which we will define later, we refer the readers to [1], [12]. It follows directly from the definition that

$$S = \text{conv}((S \cap J) \cup (S \cap J')).$$

THEOREM 5.1. *Every semi- L -summand J in a Banach space X is U -proximal.*

PROOF. Let $x \in (1 + \rho)S \cap (S + J)$ and $\|x\| = 1 + \rho$; then $x = (1 + \rho)(y + z)$ with $y \in S \cap J$ and $z \in S \cap J'$. Note that $x \in S + J, d(x, J) \leq 1$. This implies $\|(1 + \rho)z\| \leq 1$. Also note that $(1 + \rho) - \rho/\|y\| > 0$ (for otherwise, $(1 + \rho)\|y\| < \rho$ will imply $(1 + \rho)\|y\| + (1 + \rho)\|z\| < 1 + \rho$). Let $y' = ((1 + \rho) - \rho/\|y\|)y$ and write $x = (y' + (1 + \rho)z) + \rho y/\|y\|$. Then $\|y' + (1 + \rho)z\| = 1$ and hence $x \in S + \rho(S \cap J)$. $\square$

Let M be a closed subspace of a Banach space X . M is called a *semi- M -ideal* (*M -ideal*) if $M^\perp = \{x^* \in S^*: x^*(x) = 0 \ \forall x \in M\}$ is a semi- L -summand (L -summand, respectively) in X^* . M is said to have the *n -ball property* if for any $\varepsilon > 0$ and for any n intersecting balls $S_i(a_i, r_i) = \{x: \|x - a_i\| \leq r_i\}$ with $S_i(a_i, r_i) \cap M \neq \emptyset$, then $\bigcap_{i=1}^n S_i(a_i, r_i + \varepsilon) \cap M \neq \emptyset$. It was proved that M is a semi- M -ideal (M -ideal) if and only if M has the 2-ball (n -ball, $n \geq 3$, respectively) property [1], [12]. In [5], Holmes showed that M -ideals are U -proximal. In the following, we will show that his theorem also holds for semi- M -ideals.

LEMMA 5.2. *Let X be a Banach space and let $e^* \in X^*$ with $\|e^*\| = 1$. Let $F = \{x \in S: e^*(x) = 1\}$. Suppose $S = \text{conv}(F \cup -F)$. Then $M = \{x: e^*(x) = 0\}$ satisfies*

$$(1 + \rho)S \cap (S + M) \subseteq S + \rho S(M).$$

PROOF. Let $x \in (1 + \rho)S \cap (S + M)$ with $\|x\| = 1 + \rho$. By assumption, there exist $a, b \in (1 + \rho)F$ such that $x \in [a : -b]$. Let $y = (\rho/(1 + \rho))(a - b)/2$. It is easy to show that $x = (x - y) + y \in S + \rho S(M)$. $\square$

LEMMA 5.3. *Let J be a one dimensional semi- L -summand in X . Let $e \in J$ with $\|e\| = 1$. Then e is an extreme point of S . Moreover, for any $x \in S$, $\|x\| = 1$, at least one of the line segments joining x with e and $-e$ is contained in the boundary of S .*

PROOF. Suppose $e = \frac{1}{2}(x + y)$, $x, y \in S$, $x, y \neq e$. By the definition of semi- L -summand, there exists $x_1, y_1 \in J'$, $0 < \alpha, \beta < 1$ such that

$$x = \alpha x_1 + (1 - \alpha)e, \quad y = \beta y_1 + (1 - \beta)e.$$

Hence $e = (\alpha x_1 + \beta y_1)/(\alpha + \beta)$. This contradicts that J is a semi- L -summand. The second part is clear by observing that the unit sphere of the subspace generated by e and x is a parallelogram. $\square$

THEOREM 5.4. *Let M be a semi- M -ideal in X . Then M satisfies*

$$(1 + \rho)S \cap \overline{(S + M)} \subseteq \overline{S + \rho S(M)}$$

and hence it is a U -proximal subspace in X .

PROOF. Let $x \in (1 + \rho)S \cap (S + M)$ with $\|x\| = 1 + \rho$. Note that $\langle M, x \rangle$ also has the 2-ball property. We may assume, without loss of generality, that $X = \langle M, x \rangle$. Then $J = M^\perp$ is a one dimensional semi- L -summand in X^* . Let $e^* \in J \cap S$. It follows from Lemma 5.3 and [13, p. 44, Theorem 4.7] that for any extreme point x^{**} in $S(X^{**})$, $x^{**}(e^*) = 1$. Let

$$F = \{x^{**} \in S(X^{**}) : x^{**}(e^*) = 1\};$$

$F \cup -F$ contains all extreme points of $S(X^{**})$. By the Krein-Milman theorem and the fact that F is w^* -compact,

$$\text{conv}(F \cup -F) = \overline{\text{conv}}^{w^*}(F \cup -F) = S(X^{**}).$$

Hence Lemma 5.2 applies and

$$\begin{aligned} x \in (S(X^{**}) + \rho(S(X^{**}) \cap M^{**})) \cap X &= \overline{S(X) + \rho(S(X) \cap M)}^{w^*} \cap S \\ &\subseteq \overline{S(X) + \rho(S(X) \cap M)}. \end{aligned}$$

It follows that $(1 + \rho)S \cap \overline{(S + M)} \subseteq \overline{S + \rho S(M)}$. Proposition 2.4 implies M is a U -proximal subspace with $\epsilon(\rho) = 2\rho$. $\square$

6. Approximation by compact operators. Let E, F be Banach spaces, we will use $L(E, F)$ ($K(E, F)$) to denote the space of bounded linear operators (compact operators) from E into F . In [4], [11], it is shown that if $E = l^p$, $F = l^q$, $1 < p, q < \infty$, then $K(E, F)$ is an M -ideal in $L(E, F)$. By Theorem 5.4, $K(E, F)$ is actually U -proximal in $L(E, F)$. For the case $E = F = l^1$,

Smith and Ward [16] and Mach and Ward [15] showed that $K(l^1, l^1)$ is not an M -ideal in $L(l^1, l^1)$; however, it is a proximal subspace. In the following, we will consider a more general setting and that $K(l^1, l^1)$ is a U -proximal subspace of $L(l^1, l^1)$ comes as a corollary.

LEMMA 6.1. *Let $(W, \mathfrak{B}, \sigma)$ be a positive measure space and suppose $f, g, h \in L^1(W)$ satisfy $f = g + h$, $\|f\| \leq 1 + \rho$, $\|g\| \leq 1$. Then there exist $g', h' \in L^1(W)$ such that $f = g' + h'$ with $\|g'\| \leq 1$, $\|h'\| \leq 8\rho$ and $|h'(x)| \leq |h(x)|$ for all $x \in W$.*

PROOF. We will assume that $1 < \|f\| \leq 1 + \rho$ (otherwise, we can take $g' = f$, $h' = 0$) and divide the measure space W into three parts:

$$D_1 = \{x: g(x) > 0, h(x) > 0\} \cup \{x: g(x) < 0, h(x) < 0\},$$

$$D_2 = \{x: g(x) > 0, h(x) < 0, f(x) < 0\} \\ \cup \{x: g(x) < 0, h(x) > 0, f(x) > 0\},$$

and

$$D_3 = \{x: g(x) > 0, h(x) < 0, f(x) > 0\} \\ \cup \{x: g(x) < 0, h(x) > 0, f(x) < 0\}.$$

Let f/D denote the restriction of f to D . It is clear that

$$\|f/D_1\| = \|g/D_1\| + \|h/D_1\|,$$

$$\|f/D_2\| = \|h/D_2\| - \|g/D_2\|,$$

and

$$\|f/D_3\| = \|g/D_3\| - \|h/D_3\|.$$

Three cases arise:

Case (i). If $\|h/D_1\| > \rho$, we let $h' = \rho \|h\chi_{D_1}\|^{-1} h\chi_{D_1}$ and let $g' = f - h'$. Then it is easy to check that $\|g'\| \leq 1$, $\|h'\| = \rho$ and $|h'(x)| \leq |h(x)|$ for all $x \in W$.

Case (ii). If $\|h/D_1\| < \rho$ and $\|h/D_2\| < 3\rho$, then we have

$$\|f/D_1\| \leq \rho + \|g/D_1\|, \|f/D_2\| < 3\rho$$

and

$$\|f/D_3\| = \|g/D_3\| - \|h/D_3\|.$$

Thus $1 < \|f\| \leq 1 + 4\rho - \|h/D_3\|$, which implies $\|h/D_3\| < 4\rho$ and hence $\|h\| < 8\rho$. For this case, we let $g' = g$, $h' = h$.

Case (iii). If $\|h/D_1\| < \rho$ and $\|h/D_2\| > 3\rho$, then $\|f/D_2\| > \rho$. For otherwise, $\|f/D_2\| < \rho$ implies that $\|g/D_2\| > 2\rho$. Thus $\|f/D_2\| < \rho < \|g/D_2\| - \rho$ and it follows that

$$\|f\| \leq (\|g/D_1\| + \rho) + (\|g/D_2\| - \rho) + \|g/D_3\| \leq 1,$$

which contradicts our assumption that $1 < \|f\|$ and proves the claim. We

define $h' = \rho \|f\chi_{D_2}\|^{-1} f\chi_{D_2}$ and let $g' = f - h'$. Then for $x \in D_2$, $|h'(x)| < |f(x)| < |h(x)|$ and hence $|h'(x)| < |h(x)|$ for all $x \in W$. It is clear that $\|g'\| = \|f\| - \rho < 1$. $\square$

Let (Ω, Σ, μ) be a σ -finite measure space and let Y be a Banach space. A function $f: \Omega \rightarrow Y$ is called a *Bochner measurable function* if there exists a sequence of functions $\{f_n\}$ of the form $\sum_{i=1}^{\infty} y_i \chi_{E_i}$, where $y_i \in Y$, $\{E_i\}$ is a measurable partition of Ω and $\{f_n\}$ converges to F uniformly except on a zero set. Let $L^\infty(\Omega, Y)$ denote the space of bounded Bochner measurable functions from Ω into Y with norm defined by the essential supremum norm. It is clear from the definition that the set of countable valued Bochner measurable functions is dense in $L^\infty(\Omega, Y)$.

Let (Ω, Σ, μ) be a σ -finite measure space and let $(W, \mathfrak{B}, \sigma)$ be a positive measure space. We use $L_{wc}^\infty(\Omega, L^1(W))$ to denote the subspace of $L^\infty(\Omega, L^1(W))$ consists of those functions f such that $f(\Omega \setminus N)$ is a weak precompact set for some zero set N in Ω .

LEMMA 6.2. *Let $H \in L_{wc}^\infty(\Omega, L^1(W))$ and let $H' \in L^\infty(\Omega, L^1(W))$ such that for almost all $\omega \in \Omega$, $|H'(\omega)| < |H(\omega)|$. Then $H' \in L_{wc}^\infty(\Omega, L^1(W))$.*

PROOF. It is well known that for any subset K in $L^1(W, \mathfrak{B}, \sigma)$, K is weakly precompact if and only if it is bounded and for any decreasing sequence $\{E_n\} \subseteq \mathfrak{B}$ such that $\cap E_n = \emptyset$, $\{\int_{E_n} f d\sigma\}$ converges uniformly to zero for all $f \in K$ [2, p. 292, p. 430]. This is also equivalent to $\{|f|: f \in K\}$ is weakly precompact. If $H \in L_{wc}^\infty(\Omega, L^1(W))$, then there exists a zero set N_1 such that $H(\Omega \setminus N_1)$ is weakly precompact. By hypothesis, $|H'(\omega)| < |H(\omega)|$ for $\omega \in \Omega \setminus N_2$ where N_2 is a zero set, it follows from the above remark that $H'(\Omega \setminus (N_1 \cup N_2))$ is also weakly precompact; therefore $H' \in L_{wc}^\infty(\Omega, L^1(W))$. $\square$

THEOREM 6.3. *$L_{wc}^\infty(\Omega, L^1(W))$ is a U -proximal subspace of $L^\infty(\Omega, L^1(W))$.*

PROOF. Let $X = L^\infty(\Omega, L^1(W))$ and let $M = L_{wc}^\infty(\Omega, L^1(W))$, in view of Proposition 2.4(i), it suffices to show that

$$(1 + \rho)S \cap (S_D + M_D) \subseteq S + 8\rho S(M)$$

where S_D and M_D denote the dense subsets of functions with countable values in S and M , respectively. Let $F \in (1 + \rho)S \cap (S_D + M_D)$; we can write $F = G + H$ with $\|G\| < 1$, $H \in M$ and

$$G(\omega)(y) = \sum_{n=1}^{\infty} g_n(y) \cdot \chi_{E_n}(\omega),$$

$$H(\omega)(y) = \sum_{n=1}^{\infty} h_n(y) \cdot \chi_{E_n}(\omega),$$

where $\omega \in \Omega$, $y \in W$ and $\{E_n\}$ is a measurable partition of E . By Lemma 5.1, for each n , there exist g'_n and h'_n such that $\|g'_n\| < 1$, $\|h'_n\| < 8\rho$, $|h'_n| < |h|$ and

$g_n + h_n = g'_n + h'_n$. Define G', H' for $\omega \in \Omega, y \in W$ by

$$G'(\omega)(y) = \sum_{n=1}^{\infty} g'_n(y) \cdot \chi_{E_n}(\omega),$$

$$H'(\omega)(y) = \sum_{n=1}^{\infty} h'_n(y) \cdot \chi_{E_n}(\omega).$$

It follows that $\|G'\| \leq 1$, $\|H'\| \leq 8\rho$ and $|H'(\omega)| \leq |H(\omega)|$, $\omega \in \Omega$. Hence $H' \in M$ (Lemma 6.2) and $F = G' + H' \in S + 8\rho S(M)$. $\square$

Let (Ω, Σ, μ) be a σ -finite measure space and let X be a Banach space with separable dual. For any operator $T: L^1(\Omega) \rightarrow X^*$, there exists a bounded Bochner measurable function $F: \Omega \rightarrow X^*$ such that

$$\langle T(f), x \rangle = \int_{\Omega} f(\omega) \langle F(\omega), x \rangle d\mu(\omega), \quad x \in X,$$

and $\|T\| = \|F\|_{\infty}$. Conversely, for any given bounded Bochner measurable function, we can define an integral operator T with F as kernel [2, p. 506]. We will identify $L(L^1(\Omega), X^*)$ and $L^{\infty}(\Omega, X^*)$. Moreover, it is known that T is weakly compact if and only if the corresponding F is in $L^{\infty}_{wc}(\Omega, X^*)$.

THEOREM 6.4. *Let (Ω, Σ, μ) be a σ -finite measure space. Then $K(L^1(\Omega), l^1)$ is a U -proximal subspace of $L(L^1(\Omega), l^1)$.*

PROOF. As is well known, weak sequential convergence and norm convergence are equivalent in l^1 . Hence the Eberlein-Smulian theorem [2, p. 430] implies that weak compactness and norm compactness in l^1 are identical. Upon identifying $K(L^1(\Omega), l^1)$ with $L^{\infty}_{wc}(\Omega, l^1)$, the result follows from Proposition 5.3. $\square$

COROLLARY 6.5. *$K(l^1, l^1)$ is a U -proximal subspace of $L(l^1, l^1)$.*

To conclude this section, we remark that little is known about the proximality of the subspace of compact operators on L^p , $1 < p < \infty$, $p \neq 2$ or $C(K)$. In [10] it was proved that if (i) $E = L^1(\mu)$ where μ is a σ -finite measure and F is uniformly convex or (ii) E^* is uniformly convex and $F = C(K)$ for some topological space X , then $K(E, F)$ is a proximal subspace in $L(E, F)$. For operators between l^p and l^q , $1 < p, q < \infty$, the only remaining unanswered case is $K(l^{\infty}, l^p)$, $1 < p < \infty$.

REFERENCES

1. E. Alfsen and E. Effros, *Structure in real Banach spaces*, Ann. of Math. (2) **96** (1972), 98–173.
2. N. Dunford and J. Schwartz, *Linear operators*. I, Interscience Publishers, New York, 1958.
3. H. Fakhoury, *Sur les M -idéaux dans certains espaces d'opérateurs et l'approximation par des opérateurs compacts* (to appear).
4. J. Hennefeld, *A decomposition of $B(X)^*$ and unique Hahn-Banach extensions*, Pacific J. Math. **46** (1973), 197–199.

5. R. Holmes, *M-ideals in approximation theory*, Approximation Theory. II, Academic Press, New York, 1976, pp. 391–396.
6. R. Holmes and B. Kripke, *Approximation of bounded functions by continuous functions*, Bull. Amer. Math. Soc. **71** (1965), 896–897.
7. ———, *Smoothness of approximation*, Michigan Math. J. **15** (1968), 225–248.
8. ———, *Best approximation by compact operators*, Indiana Univ. Math. J. **21** (1971), 255–263.
9. R. Holmes, B. Scranton and J. Ward, *Approximation from the space of compact operators and other M-ideals*, Duke Math. J. **42** (1975), 259–269.
10. K. Lau, *Approximation by continuous vector valued functions*, Studia Math. (to appear).
11. ———, *M-ideals and approximation by compact operators* (unpublished).
12. A. Lima, *Intersection of balls and subspaces in Banach spaces*, Trans. Amer. Math. Soc. **227** (1977), 1–62.
13. J. Lindenstrauss, *Extension of compact operators*, Mem. Amer. Math. Soc. No. 48, 1964.
14. ———, *On nonlinear projections in Banach spaces*, Michigan Math. J. **11** (1964), 263–287.
15. J. Mach and J. Ward, *Approximation by compact operators on certain Banach spaces*, J. Approximation Theory (to appear).
16. R. Smith and J. Ward, *M-ideal structure in Banach algebra*, J. Functional Analysis **27** (1978), 337–349.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PITTSBURGH, PITTSBURGH, PENNSYLVANIA 15260