THE DIRICHLET NORM AND THE NORM OF SZEGÖ TYPE¹

BY

SABUROU SAITOH

ABSTRACT. Let S be a smoothly bounded region in the complex plane. Let $g(z, t)$ denote the Green's function of S with pole at t . We show that

$$\iint_S |f'(z)|^2 dx dy < \frac{1}{2} \int_{\partial S} |f'(z)|^2 \left(\frac{\partial g(z, t)}{\partial n_z} \right)^{-1} |dz|$$

holds for any analytic function $f(z)$ on $S \cup \partial S$. This curious inequality is obtained as a special case of a much more general result.

1. Introduction and preliminary facts. Let S denote an arbitrary compact bordered Riemann surface with boundary contours $\{C_\nu\}_{\nu=1}^{2g+m}$ and of genus g . Let $\{C_\nu\}_{\nu=1}^{2g+m-1}$ be a canonical homology basis for S . Let $W(z, t)$ denote a meromorphic function whose real part is the Green's function $g(z, t)$ with pole at $t \in S$. The differential $\text{id } W(z, t)$ is positive along ∂S and has $N = 2g + m - 1$ zeros $\{t_\nu\}$ in S . We assume that the points t_ν are simple and they are not on $\{C_\nu\}_{\nu=1}^{2g+m-1}$; the other cases will require only a slight modification. For simplicity, we do not distinguish between points $z \in S \cup \partial S$ and local parameters z . For an arbitrary integer q and for any positive continuous function $\rho(z)$ on ∂S , we let $H_{p,\rho}^q(S)$ [$p \geq 1$] be the Banach space of analytic differentials $f(z)(dz)^q$ of order q on S with a finite norm

$$\left\{ \frac{1}{2\pi} \int_{\partial S} |f(z)(dz)^q|^p \rho(z) [\text{id } W(z, t)]^{1-p} \right\}^{1/p} < \infty,$$

where $f(z)$ means the Fatou boundary value of f at $z \in \partial S$. Let $K_{q,t,\rho}(z, \bar{u})(dz)^q$ be the reproducing kernel of $H_{2,\rho}^q(S)$ which is characterized by the reproducing property

$$f(u) = \frac{1}{2\pi} \int_{\partial S} f(z)(dz)^q \overline{K_{q,t,\rho}(z, \bar{u})(dz)^q} \rho(z) [\text{id } W(z, t)]^{1-2q}$$

for all $f(z)(dz)^q \in H_{2,\rho}^q(S)$ (see [5]). Let $L_{q,t,\rho}(z, u)(dz)^{1-q}$ denote the adjoint L -kernel of $K_{q,t,\rho}(z, \bar{u})(dz)^q$. Then, $L_{q,t,\rho}(z, u)(dz)^{1-q}$ is a meromorphic differential on S of order $1 - q$ with a simple pole at u having residue 1.

Received by the editors March 13, 1978.

AMS (MOS) subject classifications (1970). Primary 30A04, 30A31; Secondary 30A40, 30A52.

Key words and phrases. Bergman kernel, kernel of Szegő type, compact bordered Riemann surface, critical points of the Green's function, direct product of two spaces of Szegő type, Dirichlet integral of analytic function.

¹Dedicated to my father on his 75th birthday.

Moreover:

$$\overline{K_{q,t,\rho}(z, \bar{u})(dz)^q} \rho(z) [\text{id } W(z, t)]^{1-2q} = (1/i) L_{q,t,\rho}(z, u)(dz)^{1-q} \quad \text{along } \partial S. \quad (1.1)$$

We note that $K_{q,t,\rho}(z, \bar{u})$ and $L_{q,t,\rho}(z, u)$ are continuous along ∂S . If S is a bounded regular region in the plane, we can define these kernels for arbitrary real values of q (cf. [5, §§2 and 6]).

Next, let $K^E(z, \bar{u})$ and $L^E(z, u)$ denote the exact Bergman kernel and its adjoint L -kernel on S , respectively (cf. [8, p. 117]). $L^E(z, u)dz$ is analytic on $S \cup \partial S$ except for u , where it has a double pole

$$\left\{ \frac{1}{\pi} \frac{1}{(z-u)^2} + \text{regular terms} \right\} dz. \quad (1.2)$$

Furthermore, it satisfies the relation

$$- \overline{K^E(z, \bar{u})} dz = L^E(u, z) dz \quad \text{along } \partial S. \quad (1.3)$$

Let $Z_\nu(z) = \int_C L(\zeta, z) d\zeta$. Then $\{Z_\nu(z) dz\}_{\nu=1}^N$ is a basis for the analytic differentials on S which are real along ∂S . Here $L(\zeta, z)$ is the adjoint L -kernel of the usual Bergman kernel $K(\zeta, \bar{z})$ on S (cf. [8, §§4.3, 4.5 and 4.10]). Then from (1.1) and (1.3), we obtain

$$K_{q,t,\rho}(z, \bar{u}) K_{1-q,t,\rho^{-1}}(z, \bar{u}) = \pi K^E(z, \bar{u}) + \sum_{\nu=1}^N \sum_{\mu=1}^N C_{\nu,\mu} \overline{Z_\nu(u)} Z_\mu(z) \quad (1.4)$$

and

$$L_{q,t,\rho}(z, u) L_{1-q,t,\rho^{-1}}(z, u) = \pi L^E(u, z) - \sum_{\nu=1}^N \sum_{\mu=1}^N \overline{C_{\nu,\mu}} Z_\nu(u) Z_\mu(z), \quad (1.5)$$

where the constants $C_{\nu,\mu}$ are uniquely determined as in [5].

2. The main theorem. Let $\{\Phi_j(z)(dz)^q\}_{j=1}^\infty$ and $\{\Psi_j(z)(dz)^{1-q}\}_{j=1}^\infty$ be complete orthonormal systems for $H_{2,\rho}^q(S)$ and $H_{2,\rho}^{1-q}(S)$, respectively. Let $H = H_{2,\rho}^q(S) \otimes H_{2,\rho}^{1-q}(S)$ denote the direct product of $H_{2,\rho}^q(S)$ and $H_{2,\rho}^{1-q}(S)$. The space H is composed of differentials $f(z_1, z_2)(dz_1)^q(dz_2)^{1-q}$ on $S \times S$ such that

$$f(z_1, z_2) = \sum_{j=1}^\infty \sum_{k=1}^\infty A_{j,k} \Phi_j(z_1) \Psi_k(z_2), \quad \sum_{j=1}^\infty \sum_{k=1}^\infty |A_{j,k}|^2 < \infty. \quad (2.1)$$

The scalar product $(\cdot, \cdot)_H$ is introduced as follows:

$$(f, h)_H = \sum_{j=1}^\infty \sum_{k=1}^\infty A_{j,k} \overline{B_{j,k}},$$

where

$$h(z_1, z_2) = \sum_{j=1}^\infty \sum_{k=1}^\infty B_{j,k} \Phi_j(z_1) \Psi_k(z_2) \quad \text{and} \quad \sum_{j=1}^\infty \sum_{k=1}^\infty |B_{j,k}|^2 < \infty$$

(cf. [1, §8]).

THEOREM 2.1. Suppose that

$$f(z_1, z_2)(dz_1)^q(dz_2)^{1-q} = \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} A_{j,k} \Phi_j(z_1) \Psi_k(z_2) (dz_1)^q (dz_2)^{1-q} \in H.$$

Then $f(z, z)$ can be uniquely decomposed as follows:

$$f(z, z) = h'(z) + \sum_{\nu=1}^N d_{\nu} Z_{\nu}(z) \quad \text{for } z \in S. \quad (2.2)$$

It is understood that the d_{ν} are constants, $h(z)$ is analytic on S , and

$$\iint_S |h'(z)|^2 dx dy < \infty \quad (z = x + iy).$$

In addition,

$$\begin{aligned} & \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} |A_{j,k}|^2 \\ & > \min \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} \left\{ \frac{1}{2\pi} \int_{\partial S} \varphi_j(z_1) (dz_1)^q \overline{\varphi_k(z_1) (dz_1)^q} \right. \\ & \quad \cdot \rho(z_1) [\text{id } W(z_1, t)]^{1-2q} \cdot \frac{1}{2\pi} \int_{\partial S} \psi_j(z_2) (dz_2)^{1-q} \\ & \quad \cdot \overline{\psi_k(z_2) (dz_2)^{1-q}} (\rho(z_2))^{-1} [\text{id } W(z_2, t)]^{2q-1} \Big\} \\ & = \frac{1}{\pi} \iint_S |h'(z)|^2 dx dy + \sum_{\nu=1}^N \sum_{\mu=1}^N D_{\nu,\mu} d_{\nu} \overline{d_{\mu}}, \end{aligned} \quad (2.3)$$

where $\|D_{\nu,\mu}\|$ is the inverse of $\|C_{\nu,\mu}\|$. The minimum is taken here over all analytic functions $\sum_{j=1}^{\infty} \varphi_j(z_1) \psi_j(z_2)$ on $S \times S$ satisfying

$$f(z, z) = \sum_{j=1}^{\infty} \varphi_j(z) \psi_j(z) \quad \text{on } S, \quad (2.4)$$

$$\varphi_j(z)(dz)^q \in H_{2,\rho}^q(S) \quad \text{and} \quad \psi_j(z)(dz)^{1-q} \in H_{2,\rho}^{1-q}(S).$$

PROOF. The crucial ingredient in this proof is the observation that $\|C_{\nu,\mu}\|$ is positive definite (cf. equation (1.4) and [5, p. 549]). Refer to the proof of Theorem 2.1 in [6]. The positive definiteness of $\|C_{\nu,\mu}\|$ implies that

$$k(z, \bar{u}) = \sum_{\nu=1}^N \sum_{\mu=1}^N C_{\nu,\mu} \overline{Z_{\nu}(u)} Z_{\mu}(z)$$

is a reproducing kernel for the finite dimensional class $\mathfrak{F}_2$ which is generated by $\{Z_{\nu}(z)\}_{\nu=1}^N$ (see [1, pp. 346–347]). The scalar product is given by

$$(f, h)_2 = \sum_{\nu=1}^N \sum_{\mu=1}^N D_{\nu,\mu} \xi_{\nu} \overline{\eta_{\mu}}$$

for $f(z) = \sum_{\nu=1}^N \zeta_{\nu} Z_{\nu}(z)$ and $h(z) = \sum_{\nu=1}^N \eta_{\nu} Z_{\nu}(z)$. Note that $K_{q,t,\rho}(z, \bar{u}) K_{1-q,t,\rho^{-1}}(z, \bar{u}) dz$ is the reproducing kernel of the space $\mathcal{F}$ which is formed by restricting the functions in H to the diagonal set $D = \{(z, z) | z \in S\}$ (cf. [1, p. 361, Theorem II]). For $f \in \mathcal{F}$, the norm $\|f\|_{\mathcal{F}}$ is given by $\min \|h\|_H$ where $h(z_1, z_2)$ ranges over all elements of H whose restriction to D is $f(z)$. Of course, $\|h\|_H$ denotes the norm of h in H .

On the other hand, the space $\mathcal{F}$ must coincide with the class corresponding kernel function $K_{q,t,\rho}(z, \bar{u}) K_{1-q,t,\rho^{-1}}(z, \bar{u})$ when it is considered as the sum of the kernel functions $\pi K^E(z, \bar{u})$ and $k(z, \bar{u})$ (see [1, pp. 352–357]). We thus obtain the decomposition (2.2). The uniqueness follows from [8, pp. 104 and 108].

Finally, from the definition of the norm in H [1, pp. 357–361], we have the inequality (2.3).

Using [1] and the preceding remark about $\mathcal{F}$, we immediately obtain

COROLLARY 2.1. *Any analytic function $f(z)$ on S with a finite Dirichlet integral can be represented as a series*

$$f'(z) = \sum_{j=1}^{\infty} \varphi_j(z) \psi_j(z) \quad \text{on } S, \quad (2.5)$$

where $\varphi_j(z)(dz)^q \in H_{2,\rho}^q(S)$ and $\psi_j(z)(dz)^{1-q} \in H_{2,\rho^{-1}}^{1-q}(S)$.

Furthermore, the equation

$$\begin{aligned} & \frac{1}{\pi} \iint_S |f'(z)|^2 dx dy \\ &= \min \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} \left\{ \frac{1}{2\pi} \int_{\partial S} \varphi_j(z_1)(dz_1)^q \overline{\varphi_k(z_1)(dz_1)^q} \right. \\ & \quad \cdot \rho(z_1) [\text{id } W(z_1, t)]^{1-2q} \cdot \frac{1}{2\pi} \int_{\partial S} \psi_j(z_2)(dz_2)^{1-q} \\ & \quad \cdot \overline{\psi_k(z_2)(dz_2)^{1-q}} (\rho(z_2))^{-1} [\text{id } W(z_2, t)]^{2q-1} \left. \right\} \quad (2.6) \end{aligned}$$

is valid. The minimum is taken here over all analytic functions $\sum_{j=1}^{\infty} \varphi_j(z_1) \psi_j(z_2)$ satisfying (2.5).

Conversely, if the jk sum in (2.6) is finite, then the exact differential $f'(z)dz$ defined by the series (2.5) has a finite Dirichlet integral.

3. Some inequalities. As an application of the main theorem, we derive some inequalities. To start with, consider the case $f(z_1, z_2) = \varphi(z_1) \psi(z_2)$. This leads to

THEOREM 3.1. For any $\varphi(z)(dz)^q \in H_{2,p}^q(S)$ and $\psi(z)(dz)^{1-q} \in H_{2,p}^{1-q}(S)$, we have

$$\begin{aligned} & \frac{1}{2\pi} \int_{\partial S} |\varphi(z_1)(dz_1)^q|^2 \rho(z_1) [\text{id } W(z_1, t)]^{1-2q} \\ & \quad \cdot \frac{1}{2\pi} \int_{\partial S} |\psi(z_2)(dz_2)^{1-q}|^2 (\rho(z_2))^{-1} [\text{id } W(z_2, t)]^{2q-1} \\ & > \min \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} \left\{ \frac{1}{2\pi} \int_{\partial S} \varphi_j(z_1)(dz_1)^q \overline{\varphi_k(z_1)(dz_1)^q} \right. \\ & \quad \cdot \rho(z_1) [\text{id } W(z_1, t)]^{1-2q} \cdot \frac{1}{2\pi} \int_{\partial S} \psi_j(z_2)(dz_2)^{1-q} \\ & \quad \cdot \overline{\psi_k(z_2)(dz_2)^{1-q}} (\rho(z_2))^{-1} [\text{id } W(z_2, t)]^{2q-1} \Big\} \\ & = \frac{1}{\pi} \iint_S |h'(z)|^2 dx dy + \sum_{\nu=1}^N \sum_{\mu=1}^N D_{\nu,\mu} d_{\nu} \overline{d_{\mu}}, \end{aligned} \quad (3.1)$$

where $\varphi(z)\psi(z) = h'(z) + \sum_{\nu=1}^N d_{\nu} Z_{\nu}(z)$ on S , and where the minimum is taken over all analytic functions $\sum_{j=1}^{\infty} \varphi_j(z_1)\psi_j(z_2)$ on $S \times S$ such that

$$\varphi(z)\psi(z) = \sum_{j=1}^{\infty} \varphi_j(z)\psi_j(z). \quad (3.2)$$

Of course, $\varphi_j(z)(dz)^q \in H_{2,p}^q(S)$ and $\psi_j(z)(dz)^{1-q} \in H_{2,p}^{1-q}(S)$.

Equality holds in (3.1) if and only if $\varphi(z)\psi(z)$ is expressible in the form $CK_{q,t,p}(z, \bar{u})K_{1-q,t,p^{-1}}(z, \bar{u})$ for some point $u \in S$ and for some constant C .

The equality statement in Theorem 3.1 will be proved in §5.

We can now take $q = 0$, $\rho(z) \equiv 1$, $\varphi(z) \equiv 1$, $\psi(z) \equiv f'(z)$. This yields

COROLLARY 3.1. For any analytic function $f(z)$ on $S \cup \partial S$, we have

$$\iint_S |f'(z)|^2 dx dy < \frac{1}{2} \int_{\partial S} \frac{|f'(z)dz|^2}{\text{id } W(z, t)}. \quad (3.3)$$

Equality holds in (3.3) if and only if S is simply connected and $f'(z)$ is expressible in the form $C\pi K^E(z, \bar{t}) = CK_{1,t,1}(z, \bar{t})$ for some constant C .

Regarding the equality statement in Corollary 3.1, we note that $K_{0,t,1}(z, \bar{u}) \equiv 1$ if and only if $u = t$ [3]. Furthermore, we can compare (3.3) with the inequality

$$\begin{aligned} \left(\frac{1}{\pi} \iint_S |f'(z)|^2 dx dy \right)^2 &= \left(\frac{1}{2\pi i} \int_{\partial S} \overline{f(z)} f'(z) dz \right)^2 \\ &< \frac{1}{2\pi} \int_{\partial S} |f(z)|^2 \text{id } W(z, t) \frac{1}{2\pi} \int_{\partial S} \frac{|f'(z)dz|^2}{\text{id } W(z, t)}. \end{aligned}$$

Let $K_{1,t,1}^E(z, \bar{u})dz$ denote the reproducing kernel of the closed subspace of $H_{2,1}^1(S)$ composed of exact analytic differentials on S (cf. [2] and [4]). Since $L^E(z, u) = L^E(u, z)$ if and only if S is planar [8, pp. 114–120], Theorem 3.3 in [2] requires a modification when $g > 1$. But, this is not difficult. Using Corollary 3.1, we now obtain

COROLLARY 3.2. *For all t and $u \in S$, we have*

$$K_{1,t,1}^E(u, \bar{u}) < \pi K^E(u, \bar{u}). \quad (3.4)$$

Equality holds in (3.4) if and only if S is simply connected and $u = t$.

PROOF. From (3.3) and the extremal property of $K^E(z, \bar{u})$ [8, pp. 135–137], we have

$$\begin{aligned} \frac{1}{K^E(u, \bar{u})} &= \iint_S \left| \frac{K^E(z, \bar{u})}{K^E(u, \bar{u})} \right|^2 dx dy < \iint_S \left| \frac{K_{1,t,1}^E(z, \bar{u})}{K_{1,t,1}^E(u, \bar{u})} \right|^2 dx dy \\ &< \frac{1}{2} \int_{\partial S} \left| \frac{K_{1,t,1}^E(z, \bar{u})}{K_{1,t,1}^E(u, \bar{u})} \right|^2 \frac{|dz|^2}{\text{id } W(z, t)} = \frac{\pi}{K_{1,t,1}^E(u, \bar{u})}. \end{aligned} \quad (3.5)$$

We note that (3.3) is, in general, *not valid* for arbitrary analytic differentials $f(z) dz$. Indeed, suppose that inequality (3.3) were valid for $K_{1,t,1}(z, \bar{t}) dz$. Then, from the argument of Corollary 3.2, we would have

$$K_{1,t,1}(t, \bar{t}) < \pi K(t, \bar{t}), \quad (3.6)$$

which implies a contradiction for doubly connected regions S (cf. [6, §7]). The relationship between $K_{1,t,1}^E(z, \bar{u})$ and $\pi K^E(z, \bar{u})$ is, in general, quite complicated. (See [4, equation (2.6)].)

On the other hand, by setting $q = 0$ and $\rho(z) \equiv 1$ in Theorem 3.3, we obtain

COROLLARY 3.3. *For the critical points $\{t_\mu\}_{\mu=1}^N$ of the Green's function $g(z, t)$ of G , the matrix*

$$\left\| \sum_{\mu=1}^N \frac{Z_\nu(t_\mu) Z_\nu(t_\mu)}{W''(t_\mu, t)} - D_{\nu, \nu'} \right\|^{N \times N}$$

is positive definite. A slight modification is required here when the t_μ are not simple.

PROOF. We consider the decomposition $1 \cdot \sum_{\nu=1}^N d_\nu Z_\nu(z)$ for arbitrary constants d_ν . Then, we have

$$\frac{1}{2\pi} \int_{\partial S} \frac{|\sum_{\nu=1}^N d_\nu Z_\nu(z) dz|^2}{\text{id } W(z, t)} > \sum_{\nu=1}^N \sum_{\nu'=1}^N D_{\nu, \nu'} d_\nu \bar{d}_{\nu'}. \quad (3.7)$$

Using the residue theorem, we deduce the desired result.

4. Integral transform by $K_{q,t,\rho}(z, \bar{u})K_{1-q,t,\rho^{-1}}(z, \bar{v})$. As another application of the main theorem, we show that all the results of [7] are valid for the present H .

THEOREM 4.1. *Assume that $p > 1$. Then, for $\sigma \in L_p(\partial S)$,*

$$F_\sigma(z_1, z_2) = \int_{\partial S} \sigma(\xi) \overline{K_{q,t,\rho}(\xi, \bar{z}_1) K_{1-q,t,\rho^{-1}}(\xi, \bar{z}_2)} d\xi \quad (4.1)$$

belongs to H if and only if the projection $h_1(z)$ of $\sigma(z)$ onto $H_{p,1}^0(S)$ belongs to the Bergman space of S ; that is,

$$\iint_S |h'_1(z)|^2 dx dy < \infty. \quad (4.2)$$

PROOF. Refer to the proof of [7, Theorem 2.1]. The necessity is now apparent from Theorem 2.1. The crucial step in the sufficiency is to show that

$$\begin{aligned} & \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} \left| \int_{\partial S} h_1(z) \overline{\Phi_j(z) \Psi_k(z)} dz \right|^2 \\ &= \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} \left| -2i \iint_S h'_1(z) \overline{\Phi_j(z) \Psi_k(z)} dx dy \right|^2 \end{aligned} \quad (4.3)$$

converges. For any double sequence $\{A_{j,k}\}$ satisfying

$$\sum_{j=1}^{\infty} \sum_{k=1}^{\infty} |A_{j,k}|^2 < \infty \quad (4.4)$$

we consider the function $f(z_1, z_2)$ defined by (2.1). Then, using Theorem 2.1, we have

$$f(z, z) = \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} A_{j,k} \Phi_j(z) \Psi_k(z) = \tilde{f}'(z) + \sum_{\nu=1}^N \tilde{C}_\nu Z_\nu(z) \quad \text{on } S \quad (4.5)$$

and

$$\iint_S |f(z, z)|^2 dx dy < \infty. \quad (4.6)$$

Note that

$$f_M(z, z) = \sum_{j=1}^M \sum_{k=1}^M A_{j,k} \Phi_j(z) \Psi_k(z) \quad (4.7)$$

also can be uniquely decomposed as follows:

$$f_M(z, z) = \tilde{f}'_M(z) + \sum_{\nu=1}^N \tilde{C}_\nu^{(M)} Z_\nu(z) \quad \text{on } S. \quad (4.8)$$

From the main theorem, we see that the convergence of $f_M(z, z) dz$ to $f(z, z) dz$ in $\mathcal{F}$ implies both the convergence of $\tilde{f}'_M(z)$ to $\tilde{f}'(z)$ in the Dirichlet norm and $\lim_{M \rightarrow \infty} \tilde{C}_\nu^{(M)} = \tilde{C}_\nu$, for each ν . We thus obtain

$$\begin{aligned}
\iint_S h'_1(z) \overline{f(z, z)} \, dx \, dy &= \iint_S h'_1(z) \overline{\left(\lim_{M \rightarrow \infty} f_M(z, z) \right)} \, dx \, dy \\
&= \iint_S h'_1(z) \overline{\left(\lim_{M \rightarrow \infty} \tilde{f}'_M(z) + \lim_{M \rightarrow \infty} \sum_{\nu=1}^N \tilde{C}_\nu^{(M)} Z_\nu(z) \right)} \, dx \, dy \\
&= \lim_{M \rightarrow \infty} \iint_S h'_1(z) \overline{\left(\tilde{f}'_M(z) + \sum_{\nu=1}^N \tilde{C}_\nu^{(M)} Z_\nu(z) \right)} \, dx \, dy \\
&= \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} \left(\iint_S h'_1(z) \overline{\Phi_j(z) \Psi_k(z)} \, dx \, dy \right) \overline{A_{j,k}} \quad (4.9)
\end{aligned}$$

for any $\{A_{j,k}\}$ satisfying (4.4). Hence, from the Landau theorem, we obtain the desired result.

The proof of this theorem shows that all the results of [7] are, in general, valid for the present H .

Let $H_{D(0)}$ denote the subspace of H formed by those functions in H which vanish along the diagonal set D . Let $(H_{D(0)})^\perp$ denote the orthocomplement of $H_{D(0)}$ in H . As in [7], we obtain

THEOREM 4.2. *Any $f(z_1, z_2)(dz_1)^q(dz_2)^{1-q} \in (H_{D(0)})^\perp$ is expressible in the form*

$$f(z_1, z_2) = \frac{1}{2\pi} \int_{\partial S} \frac{h(\zeta) d\zeta \overline{K_{q,t,\rho}(\zeta, \bar{z}_1) K_{1-q,t,\rho^{-1}}(\zeta, \bar{z}_2)} d\bar{\zeta}}{\text{id } W(\zeta, t)} \quad (4.10)$$

for a uniquely determined $h(z) \, dz$ in $H_{2,1}^1(S)$.

Furthermore:

$$h(z) = -W'(z, t) \left\{ \int_t^z \left(\sum_{\nu=1}^N X_\nu(f) L_{q,t,\rho}(\zeta, t_\nu) L_{1-q,t,\rho^{-1}}(\zeta, t_\nu) + f(\zeta, \zeta) \right) d\zeta \right\}, \quad (4.11)$$

where the constants $X_\nu(f)$ are uniquely determined using the equations

$$\begin{aligned}
\sum_{\nu=1}^N X_\nu(f) \int_{C_\mu} L_{q,t,\rho}(\zeta, t_\nu) L_{1-q,t,\rho^{-1}}(\zeta, t_\nu) d\zeta \\
= - \int_{C_\mu} f(\zeta, \zeta) d\zeta, \quad \mu = 1, 2, \dots, N. \quad (4.12)
\end{aligned}$$

5. Proof of the equality statement in Theorem 3.1. As an application of Theorem 4.2, we now prove the equality statement in Theorem 3.1.

If equality holds in (3.1) for $\varphi(z_1)(dz_1)^q\psi(z_2)(dz_2)^{1-q} \in H$, then we have $\varphi(z_1)(dz_1)^q\psi(z_2)(dz_2)^{1-q} \in (H_{D(0)})^\perp$ (cf. [6, equation (3.2)]). Hence, by Theorem 4.2, we have

$$\varphi(z_1)\psi(z_2) = \frac{1}{2\pi} \int_{\partial S} \frac{h(\zeta) d\zeta \overline{K_{q,t,\rho}(\zeta, \bar{z}_1) K_{1-q,t,\rho^{-1}}(\zeta, \bar{z}_2)} d\bar{\zeta}}{\text{id } W(\zeta, t)} \quad (5.1)$$

with $h(\zeta) d\zeta \in H_{2,1}^1(S)$. Consider any $\varphi^{(0)}(z)(dz)^q$ such that: (i) $\varphi^{(0)}(z) \not\equiv 0$; (ii) $\varphi^{(0)}(z)$ is analytic on $S \cup \partial S$; (iii) $\varphi^{(0)}(z)(dz)^q$ is orthogonal to $\varphi(z)(dz)^q$ in $H_{2,\rho}^q(S)$. From (5.1), we obtain

$$\int_{\partial S} \frac{h(\zeta) d\zeta \overline{\varphi^{(0)}(\zeta) K_{1-q,t,\rho^{-1}}(\zeta, \bar{z}_2) d\zeta}}{\text{id } W(\zeta, t)} = 0 \quad \text{for all } z_2 \in S$$

and so

$$\int_{\partial S} \frac{h(\zeta) d\zeta \overline{\varphi^{(0)}(\zeta) f(\zeta) d\zeta}}{\text{id } W(\zeta, t)} = 0, \quad (5.2)$$

for all $f(\zeta)(d\zeta)^{1-q} \in H_{2,\rho}^{1-q}(S)$. Hence, from the theorem of Cauchy-Read, we obtain

$$\frac{\overline{h(\zeta) d\zeta \varphi^{(0)}(\zeta) (d\zeta)^q}}{\text{id } W(\zeta, t)} = g(\zeta) (d\zeta)^q \quad \text{a.e. along } \partial S, \quad (5.3)$$

with $g(\zeta)(d\zeta)^q \in H_{2,\rho}^q(S)$ (cf. [5, p. 549] and [6, equation (3.9)]). From (5.1), we deduce that

$$\varphi(z_1)\psi(z_2) = \frac{1}{2\pi} \int_{\partial S} \frac{g(\zeta)(d\zeta)^q K_{q,t,\rho}(\zeta, \bar{z}_1) K_{1-q,t,\rho^{-1}}(\zeta, \bar{z}_2) d\zeta}{\varphi^{(0)}(\zeta) (d\zeta)^q}. \quad (5.4)$$

Using (5.4) and the residue theorem, we obtain

$$\varphi(z_1)\psi(z_2) = \sum_{j=1}^a \sum_{k=0}^b X_{j,k} \frac{\partial^k (K_{q,t,\rho}(z_1, \bar{u}_j) K_{1-q,t,\rho^{-1}}(z_2, \bar{u}_j))}{\partial \bar{u}_j^k} \quad (5.5)$$

and so

$$\varphi(z_1) = \sum_{j=1}^a \sum_{k=0}^b Y_{j,k}^{(1)} \frac{\partial^k K_{q,t,\rho}(z_1, \bar{u}_j)}{\partial \bar{u}_j^k}, \quad (5.6)$$

$$\psi(z_2) = \sum_{j=1}^a \sum_{k=0}^b Y_{j,k}^{(2)} \frac{\partial^k K_{1-q,t,\rho^{-1}}(z_2, \bar{u}_j)}{\partial \bar{u}_j^k} \quad (5.7)$$

for some points $u_j \in S$ and some constants $\{X_{j,k}\}$, $\{Y_{j,k}^{(1)}\}$ and $\{Y_{j,k}^{(2)}\}$. From (1.1), we now obtain

$$\begin{aligned} & \sum_{j=1}^a \sum_{k=0}^b \overline{X_{j,k}} \frac{\partial^k (L_{q,t,\rho}(z_1, u_j) L_{1-q,t,\rho^{-1}}(z_2, u_j))}{\partial u_j^k} \\ &= \sum_{j'=1}^a \sum_{k'=0}^b \sum_{j''=1}^a \sum_{k''=0}^b \overline{Y_{j',k'}^{(1)}} \overline{Y_{j'',k''}^{(2)}} \frac{\partial^{k'} L_{q,t,\rho}(z_1, u_{j'})}{\partial u_{j'}^{k'}} \frac{\partial^{k''} L_{1-q,t,\rho^{-1}}(z_2, u_{j''})}{\partial u_{j''}^{k''}} \end{aligned} \quad (5.8)$$

for all z_1 and $z_2 \in S$. By setting $z_1 = z_2 = z$ and comparing the orders of the poles at each u_j , we see that $X_{j,k} = 0$ for all j and k such that $k \neq 0$, and so

$Y_{j,k}^{(1)} = Y_{j,k}^{(2)} = 0$ for all j and k such that $k \neq 0$. Thus

$$\begin{aligned} \sum_{j=1}^a \overline{X_{j,0}} L_{q,t,\rho}(z_1, u_j) L_{1-q,t,\rho^{-1}}(z_2, u_j) \\ = \left(\sum_{j'=1}^a \overline{Y_{j',0}^{(1)}} L_{q,t,\rho}(z_1, u_{j'}) \right) \left(\sum_{j''=1}^a \overline{Y_{j'',0}^{(2)}} L_{1-q,t,\rho^{-1}}(z_2, u_{j''}) \right) \quad (5.9) \end{aligned}$$

for all z_1 and $z_2 \in S$. Without loss of generality, some $X_{j_0,0}$ is *nonzero*. By considering (5.9) as the identity with respect to z_1 ,

$$\overline{X_{j_0,0}} L_{1-q,t,\rho^{-1}}(z_2, u_{j_0}) = \overline{Y_{j_0,0}^{(1)}} \left(\sum_{j''=1}^a \overline{Y_{j'',0}^{(2)}} L_{1-q,t,\rho^{-1}}(z_2, u_{j''}) \right) \quad (5.10)$$

for all $z_2 \in S$. Therefore $Y_{j'',0}^{(2)} = 0$ for all j'' except j_0 and so $Y_{j',0}^{(1)} = 0$ for all j' except j_0 . Hence $X_{j,0} = 0$ for all j except j_0 . This yields the desired result:

$$\varphi(z_1)\psi(z_2) = X_{j_0,0} K_{q,t,\rho}(z_1, \bar{u}_{j_0}) K_{1-q,t,\rho^{-1}}(z_2, \bar{u}_{j_0}). \quad (5.11)$$

Acknowledgement. The author gratefully acknowledges the help of the referee in improving the representation of this material.

REFERENCES

1. N. Aronszajn, *Theory of reproducing kernels*, Trans. Amer. Math. Soc. **68** (1950), 337–404.
2. S. Saitoh, *The kernel functions of Szegő type on Riemann surfaces*, Kōdai Math. Sem. Rep. **24** (1972), 410–421.
3. ———, *The Rudin kernel and the extremal functions in Hardy classes*, Kōdai Math. Sem. Rep. **25** (1973), 37–47.
4. ———, *On some completenesses of the Bergman kernel and the Rudin kernel*, Pacific J. Math. **56** (1975), 581–596.
5. ———, *The exact Bergman kernel and the kernels of Szegő type*, Pacific J. Math. **71** (1977), 545–557.
6. ———, *The Bergman norm and the Szegő norm*, Trans. Amer. Math. Soc. **249** (1979), 261–279.
7. ———, *Integral transform by the product of two Szegő kernel functions* (submitted).
8. M. Schiffer and D. C. Spencer, *Functionals of finite Riemann surfaces*, Princeton Univ. Press, Princeton, N. J., 1954.

DEPARTMENT OF MATHEMATICS, FACULTY OF ENGINEERING, GUNMA UNIVERSITY, 1-5-1, TENJIN-CHO, KIRYU 376, JAPAN