LIFTING CYCLES TO DEFORMATIONS OF TWO-DIMENSIONAL PSEUDOCONVEX MANIFOLDS

BY

HENRY B. LAUFER¹

ABSTRACT. Let M be a strictly pseudoconvex manifold with exceptional set A . Let $D > 0$ be a cycle on A . Let $\omega: \mathcal{M} \rightarrow Q$ be a deformation of M . Kodaira's theory for deforming submanifolds of $\mathcal{M}$ is extended to the subspace D . Let $\mathcal{T}$ be the sheaf of germs of infinitesimal deformations of D . Suppose that $H^1(D, \mathcal{T}) = 0$. If ω is the versal deformation, then D lifts to above a submanifold of Q . This lifting is a complete deformation of D with a smooth generic fiber.

If all of the fibers of $\mathcal{M}$ are isomorphic, then ω is the trivial deformation. If M has no exceptional curves of the first kind, then there exists ω such that only any given irreducible component of A disappears as part of the exceptional set.

I. Introduction. In [12], Kodaira develops a theory of deforming compact submanifolds within a given ambient manifold $\mathcal{M}$. In this paper we develop an analogous theory for some singular subvarieties of $\mathcal{M}$.

Let M be a given strictly pseudoconvex two-dimensional manifold. Let $\omega: \mathcal{M} \rightarrow Q$ be a 1-convex deformation of M . (See §II for the definition of 1-convex.) M in fact has a versal deformation [19]. We are interested in how the exceptional set A of M changes under ω . Essentially [18, §II, pp. 3–7] the new exceptional curves are liftings of cycles D on the original exceptional set A . So it is important to see how cycles on A lift to Q . Let $\Phi = \Theta_D(D)$. Obstructions to lifting D lie in $H^1(D, \Phi)$.

§II is devoted to first obstructions. As an application, we prove the analogue of [5] and [25, Theorem 2.1 and Proposition 2.3]. (Theorem 2.11): If all of the fibers of a deformation $\omega: \mathcal{M} \rightarrow Q$ of M are isomorphic, then ω is the trivial deformation.

§III is devoted to higher obstructions. We first do the formal theory. Convergence is then proved using Kodaira's methods, but there are some subtle points. Theorem 2.8 of [18] is proved here as Theorem 3.5. It says roughly the following: Suppose that ω is the versal deformation of M . Let $\mathcal{T}$ be the sheaf of germs of infinitesimal deformations of D . Suppose that $H^1(D, \mathcal{T}) = 0$. Then D lifts precisely to a submanifold S of Q of codimension $\dim H^1(D, \Phi)$. There is an induced deformation $\lambda: \mathcal{D} \rightarrow S$ of D as an abstract analytic space. λ is a complete deformation. The generic fiber of λ is a manifold. We also show (Theorem 3.6) that if M has no exceptional curves of the first kind, then M may be deformed so that only any given irreducible component A_k of A disappears as part of the exceptional set.

Received by the editors May 27, 1980.

1980 *Mathematics Subject Classification*. Primary 32G05, 32G10; Secondary 14J15.

Key words and phrases. Deformation, pseudoconvex, exceptional set, obstruction.

¹This research was partially supported by NSF Grant MCS 760496A01.

© 1981 American Mathematical Society
0002-9947/81/0000-0307/\$06.00

Some of the results of this paper were obtained independently by Wahl [27], [28] in the algebraic category.

The author's manuscript [17] now essentially appears as three papers: [18], [19] and this paper.

II. First obstructions to lifting cycles. Recall that a map $\omega: \mathfrak{N} \rightarrow Q$ is 1-convex if there is a real C^∞ function φ on $\mathfrak{N}$ and a real number c_* such that

- (i) φ is strictly plurisubharmonic on $\{m \in \mathfrak{N} | \varphi(m) > c_*\}$ and
- (ii) for all c , ω is a proper map on $\{m \in \mathfrak{N} | \varphi(m) < c\}$.

Let $\omega: \mathfrak{N} \rightarrow Q$ be a 1-convex map which gives a deformation of the two-dimensional manifold $M = M_0 = \omega^{-1}(0)$. Let $A = A_0$ be the exceptional set in M . Let A_q be the exceptional set in $M_q = \omega^{-1}(q)$, $q \in Q$. Let $W = \bigcup A_q$. Recall [23, p. 553] that W is a subvariety of $\mathfrak{N}$. Let V be an irreducible component of W such that $A \cap V \neq \emptyset$. Let $P = \omega(V)$. Then by [18, Theorem 2.1, p. 3], there is a cycle $B = B_0$ on A such that $V = \bigcup \text{supp}(B_q)$, $q \in P$, with B_q the unique cycle on A_q which is homologous to B in $\mathfrak{N}$. We shall say that B weakly lifts to P locally. More generally and more formally,

DEFINITION 2.1. Let $\omega: \mathfrak{N} \rightarrow Q$ be a 1-convex map which gives a deformation of the two-dimensional manifold $M = \omega^{-1}(0)$. Let $D = D_0 > 0$ be a cycle on A and let R be a reduced subvariety of Q , $0 \in R$. D weakly lifts to R locally if for each $r \in R_1$, some neighborhood of 0 in R , there is a (necessarily unique) cycle $D_r > 0$ on A_r , the exceptional set $M_r = \omega^{-1}(r)$, such that D and D_r are homologous in $\mathfrak{N}$.

D lifts to R locally if there is a (nonreduced) subspace $\mathfrak{D}$ of $\mathfrak{N}$ such that $\lambda = \omega|_{\mathfrak{D}} \rightarrow R$ is a flat deformation of D near 0.

Of course, if D lifts, then D also weakly lifts. Namely, connect 0 and r by a real curve $\gamma \subset R$. Then, suitably counting multiplicities, $\lambda^{-1}(\gamma)$ is a chain which shows that D is homologous to D_r .

Recall [18, Proposition 2.7, p. 6],

PROPOSITION 2.2. Let $\omega: \mathfrak{N} \rightarrow Q$ be a 1-convex map which gives a deformation of the two-dimensional manifold $M = \omega^{-1}(0)$. Let $D > 0$ be a cycle on A , the exceptional set. Let $R = \{r \in Q | \text{there exists a cycle } D_r > 0 \text{ on } A_r \text{ such that } D_r \text{ is homologous to } D \text{ in } M\}$. Then there is a sufficiently small neighborhood Q_1 of 0 in Q such that $R_1 = R \cap Q_1$ is a subvariety of Q_1 . In particular, there is a maximal subvariety R of Q to which D weakly lifts locally.

Let S be any subvariety of Q to which D weakly lifts locally. Then $\bigcup D_s$, $s \in S$, is a subvariety of $\omega^{-1}(S)$. Let $\pi: T \rightarrow S$ be a resolution of S . Let $\tau: \mathfrak{U} \rightarrow T$ be the pull-back of ω via π . Then near any $t_0 \in \tau^{-1}(0)$, $D \subset M_{t_0}$ lifts to T locally.

Proposition 2.2 implies that for R smooth, the two notions of lifting in Definition 2.1 coincide. The author does not know if weak lifting always implies lifting in the case that R is singular. However, in case M is of dimension 3 and A is of dimension 1, even for R smooth, A can weakly lift, but not lift; see Example 2.2, $k > 2$, especially the end of §II, in [20].

We shall now examine how $D > 0$ lifts by extending to the singular case some ideas of Kodaira. See [12], [1, Lemma 3.2, p. 339]. We shall use Kodaira's notation

in [12] as much as possible. Until after Proposition 2.4 below, M may be a manifold of arbitrary dimension and D any compact divisorial cycle. D need not lie on an exceptional set.

On M , D locally has a single defining equation which may be reducible. Let $\mathfrak{U} = \{U_i\}$ be a cover of M such that, on U_i , $\text{Id } D = (\tilde{w}_i)$. On $U_i \cap U_j$, there is a nonzero holomorphic function $\tilde{F}_{ij}$ such that

$$\tilde{w}_i(z) = \tilde{F}_{ij}(z)\tilde{w}_j(z), \quad z \in U_i \cap U_j. \quad (2.1)$$

On $U_i \cap U_j \cap U_k$, $\tilde{F}_{ij} \cdot \tilde{F}_{jk} \cdot \tilde{F}_{ki} \equiv 1$. Let D also denote the nonreduced subspace of M defined on each U_i by $(D, \mathcal{O}/(\tilde{w}_i))$. Then the $\{\tilde{F}_{ij}\}$ give transition functions as follows for a line bundle L over D . L is trivial over U_i with fiber coordinate y_i .

$$y_i = \tilde{F}_{ij}(z)y_j, \quad z \in U_i \cap U_j. \quad (2.2)$$

Let Φ be the sheaf of germs of sections of the line bundle L . Then Φ is canonically isomorphic to $\mathcal{O}_D(D)$ and also coincides with the normal sheaf to D should D be a submanifold.

Let $\omega: \mathfrak{M} \rightarrow Q$ be a deformation of $M = \omega^{-1}(0)$ with Q smooth of dimension d . Deformations of the manifold M are locally trivial. So taking Q to be a small neighborhood of 0 and shrinking M , we may cover $\mathfrak{M}$ by the $U_i \times Q$.

We take the U_i to be local coordinate patches on M with local coordinate z_i . Then the change of coordinates on $\mathfrak{M}$ between $U_j \times Q$ and $U_i \times Q$ may be given by holomorphic maps $g_{ij}(z_j, q)$, i.e. $(z_i, q) = (g_{ij}(z_j, q), q)$. Let $f_{ij}(z_j) = g_{ij}(z_j, 0)$. Then $z_i = f_{ij}(z_j)$ is the change of coordinates between U_i and U_j as coordinate patches on M .

We shall use the unsubscripted letter z to denote a point in M . If also $z \in U_i$, then z has coordinate z_i on U_i . Recall from (2.1) the functions $\tilde{w}_i(z)$ on U_i and $\tilde{F}_{ij}(z)$ on $U_i \cap U_j$. The product structure on $U_i \times Q$ allows an extension of $\tilde{w}_i$ from $U_i \times \{0\}$ to $U_i \times Q$. We denote this extension by $w_i(z_i)$. Note that $w_i(z_i)$ is strictly speaking a function both of z_i and of q . $w_i(z_i) = \tilde{w}_i(z)$. Now look at $\tilde{F}_{ij}(z)$, defined on $U_i \cap U_j$. We wish to extend $\tilde{F}_{ij}$ to $(U_i \times Q) \cap (U_j \times Q)$ as a subset of $U_j \times Q$. Initially, $\tilde{F}_{ij}$ only extends to $(U_i \cap U_j) \times Q$. However, this initial extension suffices for a shrunken $\{U_i\}$ and Q . So there is a holomorphic function $F_{ij}(z_j)$ on $(U_i \times Q) \cap (U_j \times Q) \subset U_j \times Q$ such that for $z \in U_i \cap U_j$, $F_{ij}(z_j) = \tilde{F}_{ij}(z)$. Again, $F_{ij}(z_j)$ is strictly speaking a function both of z_j and of q . Observe that we do not claim that, for $q \neq 0$ and $z_i = g_{ij}(z_j, q)$, $F_{ij}(z_j)F_{ji}(z_i) \equiv 1$. Let T be smooth and $\pi: T \rightarrow Q$ a holomorphic map, $\pi(0) = 0$. Let $t = (t_1, \dots, t_n)$ be local coordinates near 0 for T . In local coordinates, $q = \pi(t) = u(t_1, \dots, t_n)$, where u is d -tuple of holomorphic functions. The map u will be important in the proof of Theorem 3.5 below. Let $\tau: \mathfrak{U} \rightarrow T$ be the pull-back of ω via π . Then D lifts to T if and only if for some cover $\mathfrak{U} = \{U_i\}$ of M , on each $U_i \times T$ there exists a holomorphic function $\varphi_i(z_i, t)$ and on each $(U_i \times T) \cap (U_j \times T)$ there exists a holomorphic function $h_{ij}(z_j, t)$ such that

$$\begin{aligned} \varphi_i(z_i, 0) &\equiv 0, \quad h_{ij}(z_j, 0) \equiv 0, \quad \text{and with } z_i = g_{ij}(z_j, u(t)), \\ w_i(z_i) - \varphi_i(z_i, t) &= [F_{ij}(z_j) + h_{ij}(z_j, t)][w_j(z_j) - \varphi_j(z_j, t)]. \end{aligned} \quad (2.3)$$

Namely, if (2.3) is satisfied, let $\mathfrak{D}$ be defined by $w_i(z_i) - \varphi_i(z_i, t)$ on $U_i \times T$. Conversely, the existence of the lifting $\mathfrak{D}$ gives defining equation $w_i(z_i) - \varphi_i(z_i, t)$ on U_i .

We shall solve (2.3) via power series in t . Thus let

$$\varphi_i(z_i, t) = \varphi_{i|1}(z_i, t) + \cdots + \varphi_{i|p}(z_i, t) + \cdots$$

with $\varphi_{i|p}(z_i, t)$ a homogeneous function of degree p in t with coefficients which are holomorphic functions of z_i . Let $h_{ij}(z_j, t) = h_{ij|1}(z_j, t) + \cdots + h_{ij|p}(z_j, t) + \cdots$ with $h_{ij|p}(z_j, t)$ defined similarly. The coefficients of $h_{ij|p}$ are holomorphic on $U_i \cap U_j$. Letting $\equiv_m$ denote equality modulo t^m , to first order (2.3) is

$$w_i(z_i) - \varphi_{i|1}(z_i, t) \equiv_2 [F_{ij}(z_j) + h_{ij|1}(z_j, t)][w_j(z_j) - \varphi_{j|1}(z_j, t)]. \quad (2.4)$$

To see if (2.4) can be satisfied, we shall transform the left side to z_j coordinates, modulo t^2 . Let $\rho_0: {}_Q T_0 \rightarrow H^1(M, \Theta)$ be the Kodaira-Spencer map for ω . It will suffice to identify ${}_Q T_0$ with $\{t\}$. Then, there is an n -tuple $\lambda_{ij}(z)$ of vector fields on $U_i \cap U_j$ such that $\rho_0(t) = \text{cls}[t \cdot \lambda_{ij}]$ in $H^1(M, \Theta)$. Vector fields map functions to functions. $\lambda_{ij}(z)$ may also be thought of as a vector field depending on z_j or on z_i . $\tilde{w}_i(z)$ may also be thought of as a function of z_i . Modulo t^2 ,

$$z_i = g_{ij}(z_j, t) \equiv_2 f_{ij}(z_j) + (t \cdot \lambda_{ij} f_{ij})(z_j),$$

$$w_i(z_i) \equiv_2 \tilde{w}_i(f_{ij}(z_j)) + (t \cdot \lambda_{ij} \tilde{w}_i)(f_{ij}(z_j)).$$

Using (2.1), (2.4) becomes

$$\begin{aligned} (t \cdot \lambda_{ij} \tilde{w}_i)(z_i) - \varphi_{i|1}(z_i, t) &= -\tilde{F}_{ij}(z_j) \varphi_{j|1}(z_j, t) + h_{ij|1}(z_j, t) \tilde{w}_j(z_j); \\ z_i &= f_{ij}(z_j). \end{aligned} \quad (2.5)$$

The terms in (2.5) are all linear functions in t and equality means that the coefficients are equal. Recall Φ of (2.2). Let $\Gamma(U_i, \Phi)t$ denote the group of linear functions in t with coefficients in $\Gamma(U_i, \Phi)$. Recall that $(\tilde{w}_i) = (\tilde{w}_j)$ on $U_i \cap U_j$. Define $\psi_{ij}(z, t) \in \Gamma(U_i \cap U_j, \Phi)t$ by

$$y_i = \psi_{ij}(z, t) = -(t \cdot \lambda_{ij} \tilde{w}_i)(z) = (t \cdot \lambda_{ji} \tilde{w}_j)(z), \quad \text{mod } (\tilde{w}_j(z)).$$

$\{\psi_{ij}\}$ is in fact a cocycle. Indeed the cocycle condition is on $U_i \cap U_j \cap U_k$,

$$\psi_{ik}(z, t) = \psi_{ij}(z, t) + \tilde{F}_{ij}(z) \psi_{jk}(z, t).$$

The cocycle condition is easily verified modulo $\text{Id } D$, or see (3.1) and its following below. The class of $[\psi_{ij}]$ is independent of choice of the cocycle $[\lambda_{ij}]$ in $\text{cls}[\lambda_{ij}]$ in $H^1(M, \Theta)$.

DEFINITION 2.3. For $t \in {}_Q T_0$, $\delta(t)$, defined as the class of $[t \cdot \lambda_{ji} \tilde{w}_j]$ in $H^1(M, \Phi)$, is the first obstruction to lifting D in the direction t .

PROPOSITION 2.4. $D > 0$ lifts infinitesimally to order 1 exactly on the subspace S of ${}_Q T_0$ where $\delta(t) = 0$.

PROOF. D lifts to order 1 exactly if (2.5) is satisfied. $\varphi_{i|1}$ and $\tilde{F}_{ij} \varphi_{j|1}$ act as coboundaries, modulo $\tilde{w}_j$, in (2.5). So (2.5) can be satisfied if and only if $\delta(t) = 0$.

The reason for the notation δ in Definition 2.3 is as follows. Using Kodaira's notation [12, p. 149], there is in the manifold case, and also in our codimension one

case, an exact sequence over D

$$0 \rightarrow \Phi \rightarrow \Psi \rightarrow \mathcal{O}^n \rightarrow 0 \quad (2.6)$$

where Ψ is the normal sheaf to D in $\mathfrak{N}$. $\mathcal{O}^n$ corresponds to sections over D of the pull-back to D via ω of the tangent bundle Q at 0. Each $t \in {}_Q T_0$ gives an element of $\Gamma(D, \mathcal{O}^n) \approx \bigoplus \Gamma(D, \mathcal{O})$ which is a direct sum of constants. The coboundary map $\delta: \Gamma(D, \mathcal{O}^n) \rightarrow H^1(D, \Phi)$ of (2.6) and δ of Definition 2.3 agree on t .

PROPOSITION 2.5. *Let $D > 0$ be a divisor on an exceptional set A in the two-dimensional manifold M . Let $\Phi = \mathcal{O}_D(D)$ from (2.2). Then $\Gamma(M, \Phi) = 0$.*

PROOF. The line bundle L of which Φ is the sheaf of germs of sections restricts to a line bundle L_i of Chern class $A_i \cdot D$ over an irreducible component A_i of A with $A_i \subset \text{supp}(D)$. Let $\mathcal{L}_i$ be the sheaf of germs of sections of L_i . Since the intersection matrix $(A_i \cdot A_j)$ is negative definite, there is an A_1 such that $A_1 \cdot D < 0$. Let $\mathcal{G}_1$ be the ideal sheaf of A_1 in M .

$$0 \rightarrow \mathcal{G}_1 \cdot \mathcal{O}_D(D) \rightarrow \mathcal{O}_D(D) \rightarrow \mathcal{L}_1 \rightarrow 0 \quad (2.7)$$

is an exact sheaf sequence. $\Gamma(M, \mathcal{L}_1) = \Gamma(A_1, \mathcal{L}_1) = 0$ since L_1 has negative Chern class. If $D = A_1$, then we are done. Otherwise, let $D' = D - A_1$. $\mathcal{O}_D(D)$ is the sheaf of germs of meromorphic functions on M which have poles at most D modulo the sheaf of germs of holomorphic functions on M . Then, via the inclusion map on meromorphic functions, $\mathcal{O}_D(D') \approx \mathcal{G}_1 \cdot \mathcal{O}_D(D)$. Now use left exactness for sections of (2.7) and an induction argument to complete the proof.

We want to develop more information about the map δ of Definition 2.3. Φ is defined by the $\tilde{F}_{ij}$ of (2.1). The following map $\beta: \Theta \rightarrow \Phi$ is easily seen to be a well-defined map of coherent sheaves of $\mathcal{O}$ -modules on M . If λ is a vector field near a point $z_0 \in M$ and $\tilde{w}(z)$ is a local defining equation for D near z_0 , then $\beta(\lambda) = (\lambda\tilde{w})(z)$. β induces $\beta_*: H^1(M, \Theta) \rightarrow H^1(M, \Phi) = H^1(D, \Phi)$. Let $\rho_0: {}_Q T_0 \rightarrow H^1(M, \Phi)$ be the Kodaira-Spencer map. Then, from Definition 2.3, $\delta = \beta_* \circ \rho_0$.

Let A_i be an irreducible component of A . Let Φ_i be the Φ associated to A_i . As in the proof of Proposition 2.5, there is an inclusion map $\iota: \Phi_i \rightarrow \Phi$. The proof of the following lemma is straightforward and will be omitted.

LEMMA 2.6. *Let A_i be an irreducible component of the exceptional set A . Let n be the coefficient of A_i in the cycle $D > 0$. Then near any point which is in A_i but is not in any other A_j in the support of D , the two maps $\beta: \Theta \rightarrow \Phi$ and $\beta_i: \Theta \rightarrow \Phi_i$ satisfy $\beta = n \cdot (\iota \circ \beta_i)$.*

Let $\mathcal{S}$ be the kernel of the map β . $\mathcal{S}$ is then the subsheaf of Θ of vector fields which take D to D . See [26]. Note that $\mathcal{S}$, but not β , depends in fact only on $\text{supp}(D)$, the support of D . Namely, near each regular point of $\text{supp}(D)$, by Lemma 2.6, β has the same kernel as some β_i . The regular points are dense in $\text{supp}(D)$, so $\mathcal{S}$ is the intersection of the kernels of the β_i .

Let $\mathcal{N}$ be the image of the map β . In case $\text{supp}(D)$ has normal crossings, $\mathcal{N} = \bigoplus \mathcal{N}_i$, where $\mathcal{N}_i$ is the sheaf of germs of sections of the normal bundle of A_i in M , with the direct sum taken over $A_i \subset \text{supp}(D)$ [26, Proposition 2.2, p. 333]. In

any event, $0 \rightarrow \mathcal{S} \xrightarrow{\alpha} \Theta \xrightarrow{\gamma} \mathcal{U} \rightarrow 0$ is an exact sequence. Since $\mathcal{U} \subset \Phi$ and $\Gamma(M, \Phi) = 0$ by Proposition 2.5, $\Gamma(M, \mathcal{U}) = 0$. So by [24]

$$0 \rightarrow H^1(M, \mathcal{S}) \xrightarrow{\alpha_*} H^1(M, \Theta) \xrightarrow{\gamma_*} H^1(M, \mathcal{U}) \rightarrow 0 \quad (2.8)$$

is an exact sequence.

Let $\mathcal{T}$ be the sheaf of germs of infinitesimal deformations of D . See [8, p. 109]. $\mathcal{T}$ is defined locally in $z = (x, y)$ coordinates by $\Theta / (\tilde{w}, \partial \tilde{w} / \partial x, \partial \tilde{w} / \partial y)$, where $\tilde{w}(z)$ is a local defining equation for D . Since β maps $\partial / \partial x$ and $\partial / \partial y$ to $\partial \tilde{w} / \partial x$ and $\partial \tilde{w} / \partial y$ respectively, there is an exact sheaf sequence on D , $0 \rightarrow \mathcal{U} \xrightarrow{\beta} \Phi \rightarrow \mathcal{T} \rightarrow 0$. Hence (2.9) below is also exact, using Proposition 2.5.

$$0 \rightarrow \Gamma(D, \mathcal{T}) \rightarrow H^1(D, \mathcal{U}) \xrightarrow{\beta_*} H^1(D, \Phi) \xrightarrow{\sigma_*} H^1(D, \mathcal{T}) \rightarrow 0. \quad (2.9)$$

Recall [19, Definition 1].

DEFINITION 2.7. Let M be a strictly pseudoconvex manifold. A *special cover* $\mathcal{U} = \{U_i\}$, $0 \leq i \leq m$, is a finite cover of M such that each U_i is Stein and such that $\overline{U}_i \cap \overline{U}_j \cap \overline{U}_k = \emptyset$ for $1 \neq j \neq k$. ($\bar{}$ denotes closure in M .)

Obstructed cycles can now be seen to be most common, namely:

THEOREM 2.8. Let $D > 0$ be a cycle on the exceptional set A in the strongly pseudoconvex two-dimensional manifold M . Suppose that no irreducible component A_i in $\text{supp}(D)$ is an exceptional curve of the first kind. Then $\beta_*: H^1(M, \Theta) \rightarrow H^1(D, \Phi)$ is not the zero map.

PROOF. Look at the proof of Proposition 2.5. For each D' (and Φ'), $H^1(M, \Phi')$ is injected into $H^1(M, \Phi)$. Continue the argument until reaching the last D' , which is an irreducible component A_1 of $\text{supp}(D)$. Then $\iota_*: H^1(M, \Phi_1) \rightarrow H^1(M, \Phi)$ is an injection.

Since A_1 is reduced, the sheaf $\mathcal{T}_1$ of germs of infinitesimal deformations of A_1 is supported at only finitely many points. So $H^1(A_1, \mathcal{T}_1) = 0$. So by (2.8) and (2.9), $\beta_{1,*}: H^1(M, \Theta) \rightarrow H^1(A_1, \Phi_1)$ is surjective. By Serre duality and the Riemann-Roch theorem,

$$\dim H^1(A, \Phi_1) = g_1 + \delta_1 - A_1 \cdot A_1 - 1,$$

where δ_1 is the "number" of nodes and cusps in the singular points of A_1 . Since A_1 is, by the hypothesis, not an exceptional curve of the first kind, $H^1(A_1, \Phi_1) \neq 0$.

We can find a nonzero element in the image of β_* as follows. Restriction is an isomorphism on $H^1(M, \Theta)$ [14, Lemma 3.1, p. 599]. So by [19, Theorem 2], we may assume that M has a special cover $\mathcal{U} = \{U_i\}$. Also, the construction of [19, Theorem 2] allows us to assume that each singular point of A is an element of only one of the U_i . On $N(\mathcal{U})$, take a nonzero element of $H^1(A_1, \Phi_1)$ as the image of a 1-cocycle c from Θ . Change the cocycle c to be 0 on the $U_i \cap U_j$ which do not meet A_1 . $\beta_1(c)$ is unchanged and still nonzero. Since ι_* is an inclusion, by Lemma 2.6, $\beta(c)$ is also nonzero.

Let ${}_D\Theta$ be the sheaf of germs of vector fields on D . If $\mathfrak{n}$ denotes the ideal sheaf of D , ${}_D\Theta$ is the tangent sheaf ${}_n\Theta$ of Grauert [6, p. 357]. ${}_D\Theta$ is denoted by ${}_D\theta$ in [15] and by just Θ in [16]. Let $\mathcal{G}$, rather than $\mathfrak{n}$, denote the ideal sheaf of D in M . Then also ${}_D\Theta = \mathcal{S} / \mathcal{G}\mathcal{S}$.

LEMMA 2.9. *Let M be a strictly pseudoconvex two-dimensional manifold. Then the map $\tau: H^1(M, \mathcal{S}) \rightarrow H^1(M, {}_D\Theta)$ is surjective for all cycles D on the exceptional set. For all sufficiently large cycles D , τ is an isomorphism.*

PROOF. $H^1(M, \mathcal{S}) \xrightarrow{\sigma} H^1(M, \mathcal{S}) \rightarrow H^1(M, {}_D\Theta) \rightarrow 0$ is exact since $H^2(M, \mathcal{S}) = 0$. By [6, Satz 1, p. 355], σ is the zero map for all sufficiently large D .

PROPOSITION 2.10. *Let $\omega: \mathcal{M} \rightarrow Q$ be a deformation of the strictly pseudoconvex two-manifold $M = \omega^{-1}(0)$ such that $\rho_0: {}_Q T_0 \rightarrow H^1(M, \Theta)$ is surjective. Let D be a cycle on the exceptional set of M . Let S be the subspace of Proposition 2.4. Then the infinitesimal lifting of D to S is an infinitesimally complete deformation of D .*

PROOF. By (2.8) and (2.9), $H^1(M, \Theta)$ maps onto $\Gamma(D, \mathcal{T})$. By Lemma 2.9, $H^1(M, \mathcal{S})$, a subspace of $H^1(M, \Theta)$, maps onto $H^1(D, {}_D\Theta)$. The lemma now follows from the description [8] of the infinitesimal deformations of D .

In the case of normal crossings, $H^1(M, \Theta)$ can often be conveniently computed by letting D be the reduced exceptional set. Since $\mathcal{U} = \bigoplus \mathcal{U}_i$, $H^1(M, \mathcal{U})$ in (2.8) is easily computed by Serre duality and the Riemann-Roch theorem. $H^1(M, \mathcal{S}) \approx H^1(M, {}_D\Theta) = H^1(D, {}_D\Theta)$ for large D by Lemma 2.8. $H^1(D, {}_D\Theta)$ can often be computed as in [16].

THEOREM 2.11. *Let M be a strictly pseudoconvex manifold of dimension two. Let A be the exceptional set of M . Let $\pi: \mathcal{M} \rightarrow P$ be a deformation of $M = \pi^{-1}(0)$ such that P is reduced, π is 1-convex and for each $p \in P$, some neighborhood of A_p in $M_p = \pi^{-1}(p)$ is biholomorphic to some neighborhood of A in M . Then given any compact set $K \subset M$, π is the trivial deformation near K in $\mathcal{M}$.*

PROOF. Let $\omega: \mathcal{U} \rightarrow Q$ be the versal deformation of M [19, Theorem 8]. Let $f: P_1 \rightarrow Q_1$, with P_1 and Q_1 neighborhoods of 0 in P and Q , respectively, induce π from ω near K . We want that $f(P_1) = 0$. Since some neighborhood of A_p is biholomorphic to some neighborhood of A , $\rho_q: {}_Q T_q \rightarrow H^1(M_q, \Theta)$ is an isomorphism for any $q \in f(P_1)$. Thus it suffices to prove Theorem 2.11 for smooth P . For on the regular points of P_1 , df will then be 0. Then f is a constant map on P_1 . Hence $f(P_1) = 0$ as desired.

We now assume that P is smooth. By [18, Proposition 2.3, p. 4] every irreducible component $C_{i,p}$ of A_p , for small p , is homologous to an integral cycle $\sum a_{ij} A_j$, $a_{ij} > 0$, on A ; we shall write $C_{i,p} \sim \sum a_{ij} A_j$. By hypothesis, the $C_{i,p}$ have an intersection matrix isomorphic to $(A_i \cdot A_j)$. It then follows that $a_{ij} = 0$ for all j except for one j_0 and $a_{ij_0} = 1$. For if not, choosing subscripts so that the isomorphism has $C_{i,p} \approx A_i$, form $D_i = \sum a_{ij} C_{j,p} \sim \sum a_{ij} (\sum a_{jk} A_k)$. $(D_i \cdot D_j) \approx (A_i \cdot A_j)$. Iterating this process would give cycles E_i on A , some of whose coefficients are arbitrarily large, with the same intersection matrix as $(A_i \cdot A_j)$. But then $(\sum E_i) \cdot (\sum E_j)$ is both arbitrarily large and equal to $(\sum A_i) \cdot (\sum A_j)$, a contradiction. Then each A_i lifts weakly to P locally. By Proposition 2.2, each A_i lifts to P locally.

Multiplying the equations for the deformations of the A_i gives a deformation λ of A . Let $\mu(s)$ be the Milnor number of a singular point $s \in A$. Under deformation, each $\mu(s)$ is upper-semicontinuous. Since $A \approx A_p$, each $\mu(s)$ is constant for small p

and the deformation of A by λ is equisingular [21]. So we may simultaneously blow up at the singular points of A_p and still satisfy the hypothesis of this theorem. In particular, we may assume without loss of generality that A has nonsingular irreducible components and normal crossings. Then, also, multiplying the equations for the deformations of the A_i gives a locally trivial deformation of any cycle D on A . Choose D to have equal coefficients for all A_i and sufficiently large so that τ of Lemma 2.9 is an isomorphism.

Let $f: P_1 \rightarrow Q_1$ be as in the first paragraph of this proof. Then by [15, Theorem 3.6, p. 76] $df(p) = 0$ for all $p \in P_1$. Hence $f \equiv 0$, as required to complete the proof of the theorem.

Recall that any isolated singularity $p \in V$ has a versal deformation [7], [22], [11], [4]. If there are several isolated singularities, we may form the Cartesian product of the parameter spaces to get a versal deformation.

COROLLARY 2.12 [2], [1]. *Let $\omega: \mathfrak{M} \rightarrow Q$ be a 1-convex map which is the versal deformation of the two-dimensional strictly pseudoconvex manifold $M = \omega^{-1}(0)$. Suppose that M has no exceptional curves of the first kind. Let $\pi: M \rightarrow V$ be the blow-down of M . Let T be the simultaneous-blow-down subspace of Q . Let $\pi_a: \mathfrak{X} \rightarrow T$ be the blow-down of ω over T . Let $\tau: \mathfrak{V} \rightarrow U$ be the versal deformation of the singularities of V . Let $f: T_1 \rightarrow U_1$ be a holomorphic map which induces π_a from τ near the singularities of V ; T_1 and U_1 are neighborhoods of 0 in T and U respectively. Then there are arbitrarily small T_1 and U_1 for which f is finite and proper.*

PROOF. By Theorem 2.11, for small T_1 , $f^{-1}(0) = 0$. The corollary now follows as below. We first reduce to the case where T and U are manifolds of the same dimension. f is induced near $0 \in T$ by an ambient map $g: \Delta \rightarrow U$ with Δ a polydisc. $g^{-1}(0) \cap T = 0$. Let D be an ambient neighborhood of 0 in U . $g = (g_1, \dots, g_n): \Delta \rightarrow D$. There are additional holomorphic functions $g_{n+1}, \dots, g_m$ on Δ such that $g_i(T) = 0$, $n+1 \leq i \leq m$, and $\{g_1 = \dots = g_m = 0\} = 0$. Note that $\{g_1, \dots, g_m\}$ induces the map f . Let r be the dimension of Δ . Then there are r linear combinations of the g_i , $1 \leq i \leq m$, whose common zeroes W have codimension r . Then there is a smaller Δ' with $W \cap \Delta' = 0$. Using these r linear combinations, we may assume that T and U are equidimensional manifolds.

Let K be a small sphere about 0 in T . Let $\nu > 1$ be the multiplicity of the map $f: K \rightarrow U - \{0\}$. Let U_1 be a connected neighborhood of 0 in U with $U_1 \cap f(K) = \emptyset$. Let B be the ball in T having K as its boundary. Let $T_1 = f^{-1}(U_1) \cap B$. Then $f: T_1 \rightarrow U_1$ is a ν -fold branched covering map; for $t \in T_1$, let the multiplicity be the degree of f on a small sphere about t . $f: T_1 \rightarrow U_1$ is necessarily proper. This completes the proof of Corollary 2.12.

III. Higher order obstructions. Let us now return to solving (2.3). In order for D to lift to T , by Proposition 2.4, the tangential map for u must map to the kernel of $\delta: {}_Q T_0 \rightarrow H^1(D, \Phi)$. Suppose that (2.3) is satisfied modulo t^{m+1} , i.e. let $u^m(t) = u_1(t) + \dots + u_p(t) + \dots + u_m(t)$ where $u_p(t)$ is a homogeneous polynomial of order p in t .

$$\begin{aligned}\varphi_i^m(z_i, t) &= \varphi_{i|1}(z_i, t) + \cdots + \varphi_{i|p}(z_i, t) + \cdots + \varphi_{i|m}(z_i, t), \\ h_{ij}^m(z_j, t) &= h_{ij|1}(z_j, t) + \cdots + h_{ij|p}(z_j, t) + \cdots + h_{ij|m}(z_j, t).\end{aligned}$$

Let $\equiv_{m+1}$ denote equality modulo t^{m+1} . Then we are supposing that for $z_i = g_{ij}(z_j, u^m(t))$, $z_j \in U_i \cap U_j$,

$$w_i(z_i) - \varphi_i^m(z_i, t) \equiv_{m+1} [F_{ij}(z_j) + h_{ij}^m(z_j, t)][w_j(z_j) - \varphi_j^m(z_j, t)]. \quad (3.1)_m$$

We want to find $u_{m+1}(t)$, $\varphi_{i|m+1}(z_i, t)$ and $h_{ij|m+1}(z_j, t)$ such that $(3.1)_{m+1}$ is satisfied. Suppose that $u_{m+1}(t)$ is already determined. Let $\{ \}^{m+1}$ denote truncation of a power series so as to include terms in t up to and including terms of degree $m+1$. For $z_j \in U_i \cap U_j$, let $z_i = g_{ij}(z_j, u^{m+1}(t))$ and

$$\psi_{ij}(z_j, t) = \{ \varphi_i^m(z_i, t) - w_i(z_i) + [F_{ij}(z_j) + h_{ij}^m(z_j, t)][w_j(z_j) - \varphi_j^m(z_j, t)] \}^{m+1}. \quad (3.2)$$

Since $(3.1)_m$ is satisfied, $\psi_{ij}(z_j, t)$ is in fact homogeneous of degree $m+1$.

Recall Φ of (2.2). Let $\Gamma(U_i \cap U_j, \Phi)t^{m+1}$ denote homogeneous polynomials of degree $m+1$ in t with coefficients in $\Gamma(U_i \cap U_j, \Phi)$. Letting $y_i = \psi_{ij}(z_j, t)$, modulo $\text{Id } D$, gives an element of $\Gamma(U_i \cap U_j, \Phi)t^{m+1}$. We claim that $\{\psi_{ij}\}$, modulo $\text{Id } D$, is in fact a cocycle. Let $z_k \in U_i \cap U_j \cap U_k$ and $z_j = f_{jk}(z_k)$. Then the cocycle condition may be written as

$$\psi_{ik}(z_k, t) \equiv_{m+2} \psi_{ij}(z_j, t) + \tilde{F}_{ij}(z_j)\psi_{jk}(z_k, t) \pmod{(\tilde{w}_k(z_k))}. \quad (3.3)$$

Since the ψ_{ij} are homogeneous of degree $m+1$, we may in fact verify (3.3) with the change of coordinates $z_j \equiv_{m+2} g_{jk}(z_k, u^{m+1}(t))$. Then

$$\begin{aligned}z_i &\equiv_{m+2} g_{ij}(z_j, u^{m+1}(t)) \equiv_{m+2} g_{ij}(g_{jk}(z_k, u^{m+1}(t)), u^{m+1}(t)) \\ &\equiv_{m+2} g_{ik}(z_k, u^{m+1}(t)).\end{aligned}$$

We may also replace $\tilde{F}_{ij}$ and $\tilde{w}_k$ of (3.3) by F_{ij} and w_k respectively. Then, substituting from (3.2) and cancelling identical terms, we see that (3.3) is equivalent to

$$\begin{aligned}-F_{ik}(z_k)\varphi_k^m(z_k, t) - h_{ik}^m(z_k, t)\varphi_k^m(z_k, t) \\ \equiv_{m+2} h_{ij}^m(z_j, t)w_j(z_j) - h_{ij}^m(z_j, t)\varphi_j^m(z_j, t) \\ - F_{ij}(z_j)[F_{jk}(z_k) + h_{jk}^m(z_k, t)]\varphi_k^m(z_k, t) \pmod{(w_k(z_k))}.\end{aligned} \quad (3.4)$$

From $(3.1)_m$,

$$F_{ik}(z_k) + h_{ik}^m(z_k, t) \equiv_{m+1} [F_{ij}(z_j) + h_{ij}^m(z_j, t)][F_{jk}(z_k) + h_{jk}^m(z_k, t)]. \quad (3.5)$$

Since φ_k^m has terms in t only of order at least one, (3.5) shows that

$$\begin{aligned}F_{ik}(z_k)\varphi_k^m(z_k, t) + h_{ik}^m(z_k, t)\varphi_k^m(z_k, t) \\ \equiv_{m+2} F_{ij}(z_j)[F_{jk}(z_k) + h_{jk}^m(z_k, t)]\varphi_k^m(z_k, t) \\ + h_{ij}^m(z_j, t)[F_{jk}(z_k) + h_{jk}^m(z_k, t)]\varphi_k^m(z_k, t).\end{aligned}$$

Since h_{ij}^m has terms in t only of order at least one, $(3.1)_m$ shows that

$$\begin{aligned} h_{ij}^m(z_j, t)w_j(z_j) - h_{ij}^m(z_j, t)\varphi_j^m(z_j, t) \\ \equiv_{m+2} h_{ij}^m(z_j, t)[F_{jk}(z_k) + h_{jk}^m(z_k, t)][w_k(z_k) - \varphi_k^m(z_k, t)]. \end{aligned}$$

Hence (3.4) and (3.3) hold.

Our numbering of obstructions is one higher than Kodaira's [12].

Let $H^1(D, \Phi)t^{m+1}$ denote the group of polynomials of degree $m+1$ in t with coefficients in $H^1(D, \Phi)$.

DEFINITION 3.1. Given $u^{m+1}(t)$ and $(3.1)_m$, $\text{cls}[\psi_{ij}] \in H^1(D, \Phi)t^{m+1}$ is the $(m+1)$ st obstruction to lifting D .

$m=0$ in Definition 3.1 agrees with our previous definition in Definition 2.3.

PROPOSITION 3.2. Given $(3.1)_m$ and $u_{m+1}(t)$, there exist $\varphi_{i|m+1}(z_i, t)$ and $h_{ij|m+1}(z_j, t)$ satisfying $(3.1)_{m+1}$ if and only if the $(m+1)$ st obstruction vanishes.

PROOF. The proof is the same as the proof of Proposition 2.4.

PROPOSITION 3.3. Given D and $u^{m+1}(t)$, $\text{cls}[\psi_{ij}] \in H^1(D, \Phi)t^{m+1}$ is independent of the choice of the $w_i(z_i)$ and of the $\varphi_i^m(z_i, t)$ satisfying $(3.1)_m$.

PROOF. Different $w_i(z_i)$ may be regarded as additional $w_i(z_i)$ on a refinement of the original cover. This will not change any cohomology classes. So we always use the given $w_i(z_i)$.

Use primes ' to denote a different choice. We will first show, by induction on p , that for all U_i , there exist holomorphic θ_i^m on U_i ,

$$\theta_i^m(z_i, t) = \theta_{i|1}(z_i, t) + \cdots + \theta_{i|p}(z_i, t) + \cdots + \theta_{i|m}(z_i, t)$$

where $\theta_{i|p}(z_i, t)$ is homogeneous of degree p in t , such that

$$w_i(z_i) - \varphi_i^m(z_i, t) \equiv_{m+1} (w_i(z_i) - \varphi_i^m(z_i, t))(1 + \theta_i^m(z_i, t)). \quad (3.6)_m$$

$(3.6)_0$ holds trivially. So let us assume $(3.6)_p$ and prove $(3.6)_{p+1}$, $p+1 \leq m$. $z_i = g_{ij}(z_j, u^{m+1}(t))$. $(3.7)_{p+1}$ below says that the $(p+1)$ st obstruction vanishes.

$$\begin{aligned} \psi_{ij|p+1}(z_j) &= \{ \varphi_i^p(z_i, t) - w_i(z_i) + [F_{ij}(z_j) + h_{ij}^p(z_j, t)][w_j(z_j) - \varphi_j^p(z_j, t)] \}^{p+1} \\ &= \{ -\varphi_{i|p+1}(z_i, t) + F_{ij}(z_j)\varphi_{j|p+1}(z_j, t) - h_{ij|p+1}(z_j, t)w_j(z_j) \}^{p+1}, \end{aligned} \quad (3.7)_{p+1}$$

$1/(1 + \theta_j^m(z_j, t)) = \Sigma(-1)^r[\theta_j^m(z_j, t)]^r$, $r \geq 0$. Since $\psi_{ij|p+1}$ is in fact homogeneous of degree $p+1$ in t ,

$$\begin{aligned} \psi_{ij|p+1}(z_j) &= \{ [\varphi_i^p(z_i, t) - w_i(z_i)][1 + \theta_i^p(z_i, t)] \\ &\quad + [F_{ij}(z_j) + h_{ij}^p(z_j, t)][1 + \theta_i^p(z_i, t)][w_j(z_j) - \varphi_j^p(z_j, t)] \}^{p+1} \\ &= \{ [\varphi_i^p(z_i, t) - w_i(z_i)][1 + \theta_i^p(z_i, t)] \\ &\quad + [F_{ij}(z_j) + h_{ij}^p(z_j, t)][1 + \theta_i^p(z_i, t)]/[1 + \theta_j^p(z_j, t)] \\ &\quad \cdot [w_j(z_j) - \varphi_j^p(z_j, t)][1 + \theta_j^p(z_j, t)] \}^{p+1}. \end{aligned}$$

From (3.6)_p,

$$\begin{aligned}
 F_{ij}(z_j) + h'_{ij}(z_j, t) &= \{ [F_{ij}(z_j) + h_{ij}^p(z_j, t)] \cdot [1 + \theta_i^p(z_i, t)] / [1 + \theta_j^p(z_j, t)] \}^p, \\
 \psi_{ij|p+1}(z_j) - \psi'_{ij|p+1}(z_j) \\
 &\equiv \{ \varphi_{i|1}(z_i, t)\theta_{i|p}(z_i, t) + \cdots + \varphi_{i|p}(z_i, t)\theta_{i|1}(z_i, t) \\
 &\quad - F_{ij}(z_j)[\varphi_{j|1}(z_j, t)\theta_{j|p}(z_j, t) + \cdots + \varphi_{j|p}(z_j, t)\theta_{j|1}(z_j, t)] \}^{p+1} \\
 &\quad \text{mod } (w_j(z_j)).
 \end{aligned} \tag{3.8}_{p+1}$$

Since $\Gamma(D, \Phi) = 0$, $\psi'_{ij|p+1}(z_j)$ is a coboundary in a unique way, modulo $(w_j(z_j))$. From (3.7)_{p+1} and (3.8)_{p+1},

$$\begin{aligned}
 \psi'_{ij|p+1}(z_j, t) &\equiv \{ -\varphi'_{i|p+1}(z_i, t) + F_{ij}(z_j)\varphi'_{j|p+1}(z_j) \}^{p+1} \\
 &\equiv \{ -\varphi_{i|p+1}(z_i, t) - \varphi_{i|1}(z_i, t)\theta_{i|p}(z_i, t) - \cdots - \varphi_{i|p}(z_i, t)\theta_{i|1}(z_i, t) \\
 &\quad + F_{ij}(z_j)[\varphi_{j|p+1}(z_j, t) + \varphi_{j|1}(z_j, t)\theta_{j|p}(z_j, t) \\
 &\quad + \cdots + \varphi_{j|p}(z_j, t)\theta_{j|1}(z_j, t)] \}^{p+1} \\
 &\quad \text{mod } (w_j(z_j)).
 \end{aligned}$$

Then for each i , there is a $\theta_{i|p+1}(z_i, t)$, homogeneous in t of degree $p + 1$, such that

$$\begin{aligned}
 \varphi'_{i|p+1}(z_i, t) &= \varphi_{i|p+1}(z_i, t) + \varphi_{i|1}(z_i, t)\theta_{i|p}(z_i, t) \\
 &\quad + \cdots + \varphi_{i|p}(z_i, t)\theta_{i|1}(z_i, t) + w_i(z_i)\theta_{i|p+1}(z_i, t).
 \end{aligned}$$

This is precisely what is needed to derive (3.6)_{p+1} from (3.6)_p. So (3.6)_m is true. Also (3.8)_{m+1} is true. Thus, $\text{cls}[\psi_{ij}] = \text{cls}[\psi'_{ij}]$, which completes the proof of the proposition.

Since Proposition 2.2 allows the possibility of nonsmooth parameter spaces for weak liftings, we would like to say something directly in this case. For R a subvariety of Q , let $C(R, 0) = C_3(R, 0)$ be the Whitney tangent cone [29]. Recall δ of Definition 2.3.

PROPOSITION 3.4. *Let $\omega: \mathfrak{U} \rightarrow Q$ be a 1-convex map which gives a deformation of the two-dimensional manifold $M = \omega^{-1}(0)$. Let R be a subvariety of Q to which a cycle $D > 0$ on A weakly lifts locally. Let $\nu \in C(R, 0)$. Then $\delta(\nu) = 0$.*

PROOF. By [29, Lemma 11.2, p. 517] there is an irreducible one-dimensional subvariety S of R near 0 such that $\nu \in C(S, 0)$. Let $\pi: T \rightarrow S$ be a resolution of S . Let t be a local parameter for T near $0 = \pi^{-1}(0)$. Choose coordinates $q = (q_1, \dots, q_n)$ for Q so that ν is in the direction of the q_1 -axis. Then, since S is irreducible and of dimension 1, there is an integer $p > 1$ so that $\pi(t) \equiv_{p+1} (ct^p, 0, \dots, 0)$ with $c \in \mathbb{C} - \{0\}$. By multiplying t by $c^{1/p}$, we may assume that $c = 1$. Let $\tau: \mathfrak{U} \rightarrow T$ be the pull-back of ω via π . Then on $\mathfrak{U}$, modulo t^{p+1} , the change of coordinates is given by $z_1 = g_{ij}(z_j, t^p)$. In (2.3), $\varphi_{i|m}$ and $h_{ij|m}$ may be

chosen to be 0 for $1 \leq m \leq p-1$. Let $\{\lambda_{ij}\}$ be the cycle in $H^1(N(\mathcal{U}), \Theta)$ corresponding to ν . (2.5) is replaced by

$$(t^p \cdot \lambda_{ij} \tilde{w}_i)(z_i) - \varphi_{i|p}(z_i, t) = -\tilde{F}_{ij}(z_j) \varphi_{j|p}(z_j, t) + h_{ij|p}(z_j, t) \tilde{w}_j(z_j); \quad z_i = f_i(z_j).$$

D lifts to T locally. By Proposition 3.3, the p th obstruction to lifting D is well defined and vanishes. So the above replacement for (2.5) can be solved. Hence $\delta(\nu) = 0$.

THEOREM 3.5. *Let D be a cycle on the exceptional set of a strictly pseudoconvex manifold M . Then*

(i) $\beta_*: H^1(M, \Theta) \rightarrow H^1(D, \Phi)$ is onto if and only if $H^1(D, \mathfrak{T}) = 0$.

Let $\omega: \mathfrak{N} \rightarrow Q$ be a deformation of M with Q smooth. Suppose that $\delta: {}_Q T_0 \rightarrow H^1(D, \Phi)$ is onto. Then

(ii) There is a submanifold S of Q with ${}_S T_0 = \ker \delta$ and a subspace $\mathfrak{D}$ of $\mathfrak{N}$ such that $\lambda = \omega|_{\mathfrak{D}}: \mathfrak{D} \rightarrow S$ is a lifting of D locally.

(iii) S is the maximal subspace of Q to which D locally lifts weakly.

Suppose that additionally $\rho_0: {}_Q T_0 \rightarrow H^1(M, \Theta)$ is onto. Then

(iv) $\lambda: \mathfrak{D} \rightarrow S$ is a complete deformation of D as an abstract (nonembedded) analytic space.

(v) Off a proper subvariety of S , $D_s = \lambda^{-1}(s)$ consists of finitely many disjoint (reduced, nonsingular) Riemann surfaces.

PROOF. (i) follows immediately from (2.8) and (2.9).

We shall first construct S of (ii) formally and then prove convergence. S will be given as the biholomorphic image of a manifold T under a map $u: T \rightarrow Q$, $u(0) = 0$. Let n be the dimension of $\ker \delta$. Let $t = (t_1, \dots, t_n)$ be coordinates near 0 in T . Let $u_i(t)$ be any linear map whose image is $\ker \delta$. We may choose coordinates $q = (q_1, \dots, q_n, q_{n+1}, \dots, q_{n+d})$, $d = \dim H^1(D, \Phi)$ so that $u_i(t_1, \dots, t_n) = (t_1, \dots, t_n, 0, \dots, 0)$. Identify Q with its tangent space at 0. Then $\delta: \{(0, \dots, 0, q_{n+1}, \dots, q_{n+d})\} \rightarrow H^1(D, \Phi)$ is an isomorphism. Proposition 2.4 says that (3.1)₁ can be satisfied. We now continue by induction on m , assuming that $u^m(t)$, $\varphi_i^m(z_i, t)$ and $h_{ij}^m(z_j, t)$ have been found for all U_i and $U_i \cap U_j$ and that (3.1)_m holds.

Let $\text{cls}[\psi_0] \in H^1(D, \Phi)t^{m+1}$ be the $(m+1)$ st obstruction to lifting D to T for $u_{m+1}(t) = 0$. For each coefficient c in $\text{cls}[\psi_0]$, there is a unique $q = (0, \dots, 0, q_{n+1}, \dots, q_{n+d}) \in Q$ such that $\delta(q) = c$. Let $u_{m+1}(t)$ be redefined so that the corresponding coefficient is $-q$. Since $\varphi_i^m(z_i, t)$ has terms in t only of order at least one, (3.2) shows that for the new $u_{m+1}(t)$, $\text{cls}[\psi_{ij}] = 0$. Hence (3.1)_{m+1} can be satisfied.

We shall now prove convergence. We will be modifying Kodaira's proof for nonsingular D [12, Theorem 1, pp. 150–158]. We first reduce to the case where the A_i , the irreducible components of the exceptional set A of D , are nonsingular and have normal crossings. So, if $x \in A$ is any point where an A_i is singular or there are not normal crossings, take any section of $\mathfrak{N}$ above Q and through x . Perform a monoidal transformation $\pi: \mathfrak{N}' \rightarrow \mathfrak{N}$ along the image of the section. This induces

a quadratic transformation on each M_q and in particular on $M = M_0$ at x . $\omega' = \omega \circ \pi: \mathfrak{M}' \rightarrow Q$ is a deformation of $M' = (\omega')^{-1}(0)$. Let D' be the total transform on M' of D on M . Then by [18, Proposition 4.7, p. 26] $\delta': {}_Q T_0 \rightarrow H^1(D', \Phi')$ is onto. Suppose that we have proved (ii) for D' . Look at $\mathfrak{D}'$ near $\pi^{-1}(x) = A'_0$. ω' is a trivial deformation near A'_0 . There is a small neighborhood U' of A'_0 in M' such that $H^1(U', \Theta) = 0$. Let V' be a Cartesian product neighborhood of U' in $(\omega')^{-1}(S)$. We may take $H^1(V', \Theta) = 0$. Since $\mathfrak{D}'$ is a topologically trivial divisor in V' , it is also analytically trivial in V' with a single defining equation w' . Let $\pi: V' \rightarrow V$ be the blow down of V' under π . Then $w' = \pi^*(w)$ for a holomorphic function on V . Also, D is the divisor of w on M near x . Thus, we can find equations which define $\mathfrak{D}'$ on $(\omega')^{-1}(S)$ which blow down to define $\mathfrak{D}$ on $\omega^{-1}(S)$. So it suffices to prove (ii) for D' . After finitely many quadratic transformations, we may assume nonsingular A_i and normal crossings.

We first construct a particular special cover $\mathfrak{U} = \{U_i\}$, $0 \leq i \leq s$, of a neighborhood of A . Each U_i will be privileged for Φ and $U_i \cap U_j$ will also behave well. Choose a finite set $B = \{b_i\}$, $1 \leq i \leq s$, of points in A such that B includes all singular points of A and at least one point in each A_i . Choose a small neighborhood U_i of each b_i such that U_i is a polydisc in local coordinates $z_i = (x_i, y_i)$ and either $A \cap U_i = \{x_i y_i = 0\}$ or $A \cap U_i = \{y_i = 0\}$. Let V_i be a smaller polydisc with $b_i \in V_i$ and $V_i \subset \subset U_i$, $1 \leq i \leq s$. We choose U_0 as a neighborhood in M of $R = A - \bigcup \bar{V}_i$, $1 \leq i \leq s$, $\bar{R} \cap B = \emptyset$, which is isomorphic to the Cartesian product of R and a disc as follows. Perform quadratic transformations at sufficiently many b_i , increasing the number of b_i if necessary, so that the proper transforms A'_j of the A_j satisfy $A'_j \cap A'_k = \emptyset$, $j \neq k$, and have very large self-intersections, i.e. so that in [6, Corollary, p. 363], all obstructions to extending isomorphisms near the A'_j vanish. Then each A'_j has a neighborhood isomorphic to a neighborhood of the zero section of its normal bundle N_j . We may regard $A_j - B$ as an open Riemann surface and so Stein [10, Theorem IX.C.10, p. 270]. Moreover, since $H^1(A_j - B, \Theta) = 0$ and $H^2(A_j - B; \mathbf{Z}) = 0$, N_j is the trivial bundle over $A_j - B$. So we may choose U_0 as a Cartesian product of R and a disc. Let x_0 denote a typical point in R and y_0 a coordinate for the disc. $z_0 \in U_0$ is given by $z_0 = (x_0, y_0)$. So that we may add small quantities to x_0 , immerse R in $\mathbf{C}$ [9]. We then have a coordinate cover for R in which each change of coordinates is the identity.

Following Kodaira, we let

$$A(t) = \frac{a}{16b} \sum \frac{1}{m^2} b^m (t_1 + \cdots + t_n)^m, \quad 1 \leq m < \infty,$$

where a and b are positive real constants to be chosen later. If $\xi(z, u) = \sum \xi_{lm \dots n}(z) u_1^l u_2^m \dots u_n^n$ is a power series in $u_1, u_2, \dots, u_n$ whose coefficients carry a norm $\|\xi_{lm \dots n}(z)\|$ and if $a(u) = \sum a_{lm \dots n} u_1^l u_2^m \dots u_n^n$ has $a_{lm \dots n} > 0$, we shall write $\xi(z, u) \ll a(u)$ to mean $\|\xi_{lm \dots n}(z)\| < a_{lm \dots n}$ for all $l, m, \dots, n$. Note that

$$A(t)^\nu \ll (a/b)^{\nu-1} A(t) \quad \text{for } \nu \geq 2. \quad (3.9)$$

We wish to solve (3.1)_m for all m , as before, and show that for all m, i, ij , we may choose $u^m(t)$, $\varphi_i^m(z_i, t)$ and $h_{ij}^m(z_j, t)$ such that

$$u^m(t) \ll A(t), \quad \varphi_i^m(z_i, t) \ll A(t), \quad h_{ij}^m(z_j, t) \ll A(t). \quad (3.10)_m$$

In (3.10)_m, the norm for $u_m(t)$ is the maximum over the absolute value of the components. For $\varphi_{i|m}(z_i, t)$ and $h_{ij|m}(z_j, t)$, it is the supremum over $z_i \in U_i$ and $z_j \in U_i \cap U_j$ respectively. Working initially on a larger cover than $\mathfrak{U}$ and then shrinking, we may assume that $\varphi_{i|1}(z_i, t)$ and $h_{ij|1}(z_j, t)$ have bounded coefficients. Then for sufficiently large a , (3.10)₁ holds. We now assume by induction that (3.10)_m holds.

We use the following norm $|\cdot|$ on sections of Φ . Different local coordinates will give equivalent norms. Let V'_k be a local polydisc coordinate patch with coordinates $z_k = (x_k, y_k)$ and suppose $\text{Id } D = (y_k^d) \cdot \Phi$ is the sheaf of germs of sections of the trivial line bundle on V'_k . Let $V_k \subset \subset V'_k$ be a smaller polydisc. Any $\varphi \in \Gamma(V_k, \Phi)$ corresponds to a $\varphi_1 \in \Gamma(V_k, \Theta / (y_k^d))$. Let $\varphi_2 \in \Gamma(V_k, \Theta)$ be an ambient representative for φ_1 . Let ∂_ν denote the ν th partial derivative with respect to y_k . Let

$$|\varphi|_{V_k} = \max \sup |\varphi_2^{(\nu)}(z_k)|, \quad z_k = (x_k, 0) \in V_k, \quad 0 \leq \nu \leq d-1.$$

$|\varphi|_{V_k}$ is independent of the choice of φ_2 . The other case for $\text{Id } D$ is $\text{Id } D = (x_k^e y_k^d)$. Let $V_1 = V_k - \{x = 0\}$ and $V_2 = V_k - \{y = 0\}$. Then let $|\varphi|_{V_k} = \max(|\varphi|_{V_1}, |\varphi|_{V_2})$. For an arbitrary set U , cover $\bar{U}$ by finitely many coordinate patches V_k . Then $|\varphi|_U = \max |\varphi|_{V_k \cap U}$. Different covers give equivalent norms.

We first let $u_{m+1}(t) = 0$ and estimate $|\psi_0|$ for ψ_0 given by (3.2) as a cocycle in $C^1(N(\mathfrak{U}), \Phi)$. Let $\psi_{ij0}(z_j)$ denote ψ_0 on $U_i \cap U_j$. From the Cartesian product structure, Cauchy's integral formula for derivatives, and the maximum principle, there is a constant c_1 such that if $\Psi_{ij}(z_j) \in \Gamma(U_i \cap U_j, \Theta) t^{m+1}$ restricts to $\psi_{ij0}(z_j)$ on D ,

$$|\psi_{ij0}(z_j)| \leq c_1 \|\Psi_{ij}(z_j)\|, \quad z_j \in U_i \cap U_j. \quad (3.11)$$

This c_1 and future c_k will be independent of m . Let $\{ \}_{m+1}$ denote the terms of order $m+1$ in t . Then we may take

$$\begin{aligned} \Psi_{ij}(z_j) &= \{ \varphi_i^m(z_i, t) - w_i(z_i) - h_{ij}^m(z_j, t) \varphi_j^m(z_j, t) \}_{m+1}, \\ z_i &= g_{ij}(z_j, u^m(t)). \end{aligned} \quad (3.12)$$

By (3.10)_m and (3.9),

$$h_{ij}^m(z_j, t) \varphi_j^m(z_j, t) \ll ab^{-1} A(t), \quad z_j \in U_i \cap U_j. \quad (3.13)$$

By starting with a larger cover and then shrinking, we may assume that ω is a trivial deformation on each U_i . Recall our notation $z_i = g_{ij}(z_j, q)$, $q \in Q$, for the transition maps; $f_{ij}(z_j) = g_{ij}(z_j, 0)$. Recall that $r = n + d$ is the dimension of Q . By shrinking the cover, we may also assume that there are constants c_2 and c_3 such that, with $\eta = (\eta_1, \eta_2)$

$$g_{ij}(z_j, q) - f_{ij}(z_j) \ll \sum c_2^\nu (q_1 + \cdots + q_\nu)^\nu, \quad 1 \leq \nu < \infty, z_j \in U_i \cap U_j, \quad (3.14)$$

$$w_i(z_i + \eta) - w_i(z_i) \ll \sum c_3^\tau (\eta_1 + \eta_2)^\tau, \quad 1 \leq \tau < \infty, z_i \in U_i. \quad (3.15)$$

Since $q = u^m(t) \ll A(t)$,

$$g_{ij}(z_j, u^m(t)) - f_{ij}(z_j) \ll \frac{c_2 r A(t)}{1 - c_2 r(a/b)}.$$

By choosing b to be sufficiently large with respect to a so that $1 - c_2 r a/b > \frac{1}{2}$ we may assume that

$$g_{ij}(z_j, u^m(t)) - f_{ij}(z_j) \ll 2c_2 r A(t). \quad (3.16)$$

Let $\eta = g_{ij}(z_j, u^m(t)) - f_{ij}(z_j)$. From (3.15), $w_i(g_{ij}(z_j, u^m(t))) - w_i(f_{ij}(z_j)) \ll \Sigma c_3^\tau(\eta_1 + \eta_2)$, $1 \leq \tau < \infty$, $z_j \in U_i \cap U_j$. Since q contains no terms in t of order $m+1$, in estimating $\{w_i(z_i)\}_{m+1}$ from (3.14) and (3.15), for $\tau = 1$ we may start the summation in ν from $\nu = 2$. Then, also using (3.16)

$$\{w_i(g_{ij}(z_j, u^m(t)))\}_{m+1} \ll 4c_2^2 c_3 r a b^{-1} A(t) + \frac{16c_2^2 c_3^2 r^2 (a/b) A(t)}{1 - 4c_2 c_3 r(a/b)}.$$

With $1 - 4c_2 c_3 r(a/b) > \frac{1}{2}$, there is a constant c_4 with

$$\{w_i(g_{ij}(z_j, u^m(t)))\}_{n+1} \ll c_4 a b^{-1} A(t). \quad (3.17)$$

Finally, we must estimate $\varphi_i^m(z_i, t)$ from (3.12). Let $U'_i \subset U_i$ and $\delta > 0$ be chosen so that, for $z_i \in U'_i$, the polydisc Δ centered at z_i with polyradius δ satisfies $\Delta \subset U_i$, and so that $\{U'_i\}$ is a cover for A . Let $\eta = (\eta_1, \eta_2)$. Then for $z_i \in U'_i$, $\varphi_i^m(z_i + \eta, t) \ll A(t)(1 - \eta_1/\delta)^{-1}(1 - \eta_2/\delta)^{-1}$. Let $\eta = g_{ij}(z_j, u^m(t)) - f_{ij}(z_j)$. Then calculating as in [12, p. 156], there is a constant c_5 such that, for sufficiently large b with respect to a ,

$$\{\varphi_i^m(g_{ij}(z_j, u^m(t)), t)\}_{m+1} \ll c_5 a b^{-1} A(t), \quad z_j \in U'_i \cap U_j. \quad (3.18)$$

From (3.11), (3.12), (3.13), (3.17) and (3.18), $\psi_{i0}(z_j) \ll c_1(1 + c_4 + c_5)ab^{-1}A(t)$, $z_j \in U'_i \cap U_j$. From the cocycle condition on ψ_{i0} , there is a constant c_6 so that, as in [12, pp. 156–157],

$$\psi_{i0}(z_j) \ll c_1(1 + c_4 + c_5)c_6 ab^{-1}A(t), \quad z_j \in U_i \cap U_j.$$

We may compute $H^1(D, \Phi)$ using bounded cochains from $N(\mathbb{U})$; see [13, §6, pp. 499–500]. Give $H^1(D, \Phi)$ the quotient norm. Let $c_7 = c_1(1 + c_4 + c_5)c_6$. Then for $\text{cls}[\psi_0] \in H^1(D, \Phi)t^{m+1}$, $\text{cls}[\psi_0] \ll c_7 ab^{-1}A(t)$. There is a constant c_8 such that

$$u_{m+1}(t) \ll c_7 c_8 ab^{-1}A(t). \quad (3.19)$$

Now let ψ_{ij} be computed with $u^{m+1}(t) = u^m(t) + u_{m+1}(t)$. Then there is a constant c_9 such that

$$\psi_{ij}(z_j) \ll (c_7 + c_7 c_8 c_9) ab^{-1}A(t). \quad (3.20)$$

$\text{cls}(\psi_{ij}) = 0$ in $H^1(D, \Phi)$. Then, as in [12, Lemma, p. 157], there is a constant c_{10} so that we may choose $\varphi'_{i|m+1}(z_i, t) \in \Gamma(U_i, \Phi)t^{m+1}$ such that

$$\psi_{ij}(z_j) = -F_{ij}(z_j)\varphi'_{j|m+1}(z_j, t) + \varphi'_{i|m+1}(z_i, t), \quad z_i = f_{ij}(z_j),$$

and $\varphi'_{i|m+1} \ll c_{10}(c_7 + c_7 c_8 c_9)ab^{-1}A(t)$. From the Cartesian product choices for our cover $\mathbb{U}$, there is a constant c_{11} such that each $\varphi'_{i|m+1}(z_i, t)$ has an ambient representative $\varphi_{i|m+1}(z_i, t) \in \Gamma(U_i, \Theta)t^{m+1}$ with

$$\varphi_{i|m+1}(z_i, t) \ll c_{11}c_{10}(c_7 + c_7 c_8 c_9)ab^{-1}A(t). \quad (3.21)$$

$(3.1)_{m+1}$ is now satisfied modulo $(w_j(z_j))$. Let $z_i = g_{ij}(z_j, u^{m+1}(t))$ and

$$h_{ij|m+1}(z_j, t) = \frac{1}{w_j(z_j)} \{ w_i(z_i) - \varphi_i^{m+1}(z_i, t) - [F_{ij}(z_j) + h_{ij}^m(z_j, t)] [w_j(z_j) - \varphi_j^{m+1}(z_j, t)] \}^{m+1}. \quad (3.22)$$

Since $(3.1)_{m+1}$ is satisfied, $h_{ij|m+1}(z_j, t)$ is homogeneous of degree $m+1$ as desired. Since $\varphi_{i|m+1}(z_i, t)$ is homogeneous of degree $m+1$ in t ,

$$\{ \varphi_{i|m+1}(g_{ij}(z_j, u^{m+1}(t)), t) \}^{m+1} = \varphi_{i|m+1}(f_{ij}(z_j), t) \quad \text{for } z_i = f_{ij}(z_j) \in U_i \cap U_j.$$

Then by (3.20) and (3.21), there is a constant c_{12} such that the numerator of (3.22) is dominated by $c_{12}ab^{-1}A(t)$. By the construction of $\mathfrak{U}$ and the maximum principle, functions on $U_i \cap U_j$ may be estimated by their values away from A . Then there is a constant c_{13} such that

$$h_{ij|m+1}(z_j, t) \ll c_{13}c_{12}ab^{-1}A(t). \quad (3.23)$$

We now complete the induction step and the proof of (ii) by choosing ab^{-1} sufficiently small so that the coefficients of $A(t)$ in (3.19), (3.21) and (3.23) are all less than 1.

By [23, Satz 1, p. 547], we may assume that ω is 1-convex.

Let R be the maximum subspace of Q , given by Proposition 2.2, to which D weakly lifts locally. We need that $R \subset S$. If not, there is a one-dimensional disc T and a holomorphic map $u: T \rightarrow Q$, $u(0) = 0$, $u(T) \not\subset S$, and such that D lifts locally to T on the deformation of M induced from ω by u . Choose local coordinates on Q so that S is a linear subspace of Q . $u(t) = u_1(t) + \cdots + u_p(t) + \cdots$. Since $u(T) \not\subset S$, there is a least $m \geq 0$ such that the image of $u_{m+1}(t)$ is not in S . Since D lifts to S , the $(m+1)$ st obstruction $[\psi_0]$, well defined by Proposition 3.4, to lifting D to T with $u = u^m(t)$ vanishes. Since D lifts to T , the $(m+1)$ st obstruction $[\psi]$ to lifting D to T with $u = u^{m+1}(t)$ also vanishes. But, as in the proof of (ii), $[\psi - \psi_0] = \delta(u_{m+1}(t))$. Since ${}_S T_0 = \ker \delta$, the image of $u_{m+1}(t)$ is contained in S . This contradicts our minimal choice of m . Hence $R \subset S$ as needed to prove (iii).

By Proposition 2.10, S is an infinitesimally complete deformation of D . Let S_1 be a submanifold of S such that ${}_S T_0$ is isomorphic to the infinitesimal deformations of D . Then the map from S_1 to the parameter space for the versal deformation of D [8], [3] is an isomorphism since S_1 is smooth. Hence S_1 and also S is a complete deformation of D .

By the openness of completeness [8, Hauptsatz, p. 140], λ is a complete deformation of D_s for any $s \in S$ near 0. Let W be the reduction of $\mathfrak{D}$. Let $\lambda_1: W \rightarrow S$ be the induced deformation from λ . Then, as in the proofs of [18, Theorem 2.1, p. 3 and Theorem 2.2, p. 4], off a proper subvariety S_1 of S , W_s is reduced and λ_1 gives an equisingular deformation of the singular points of W_s . We will first show that in fact for $s \notin S_1$, W_s is nonsingular. Suppose that x is a singular point of W_s , $s \notin S_1$. Choose local coordinates z with $x = 0$. Let $f(z)$ generate $\text{Id}(D_s)$ near 0. Let

$$f(z) = f_1^{q_1}(z) \cdots f_k^{q_k}(z) \quad (3.24)$$

be a factorization of f into prime factors; f_i and f_j are not associates for $i \neq j$. Since (3.24) may be obtained by decomposing D_s into irreducible components, since λ_1 is an equisingular deformation, and since $\mathfrak{D}$ differs from W only in that the local defining equations for $\mathfrak{D}$ may have multiple factors, $\mathfrak{D}$ has a local defining equation

$$g(z, s) = g_1^{q_1}(z, s) \cdots g_k^{q_k}(z, s) \quad (3.25)$$

with $g_i(z, 0) = f_i(z)$.

Consider the following nonzero infinitesimal deformation λ_2 of D_s , with a parameter space $\varepsilon/\varepsilon^2$, ε one-dimensional. In z -coordinates, λ_2 is given by

$$h(z, \varepsilon) = f_1^{q_1}(z) \cdots f_k^{q_k}(z) + \varepsilon f_1^{q_1-1}(z) \cdots f_k^{q_k-1}(z). \quad (3.26)$$

Since λ is a complete deformation of D_s , (3.26) may be induced from (3.25), modulo ε^2 and up to multiplication by a unit $u(z, \varepsilon)$. $u(z, \varepsilon)$ may be assumed to be of the form $u(z, \varepsilon) = 1 + \varepsilon u_1(z, \varepsilon)$, with $u_1(z, \varepsilon)$ holomorphic. Looking at the order of the zero of the coefficient of ε at $z = 0$, one sees that (3.26) may be induced from (3.25) only if $k = 1$ and moreover $g_1(z, s(\varepsilon)) = f_1(z) + \varepsilon u_2(z)$, modulo ε^2 , with $u_2(0) \neq 0$.

We can see that, for small $\varepsilon \neq 0$ and small z , $\{z | g_1(z, s(\varepsilon)) = 0\}$ is nonsingular as follows. Multiplying by $1/u_2(z)$, we may assume that $g_1(z, s(\varepsilon))$ is given by $F(z) + \varepsilon + \varepsilon^2 G(z, \varepsilon)$ with $\{F(z) = 0\}$ having a singularity only at the origin. For $i = 1, 2$, let the subscript i denote the partial derivative with respect to z_1 and z_2 respectively. We want that $(0, 0)$ is the only solution for small z and ε to the equations (3.27) and (3.28) below.

$$F(z) + \varepsilon + \varepsilon^2 G(z, \varepsilon) = 0, \quad (3.27)$$

$$F_i(z) + \varepsilon^2 G_i(z, \varepsilon) = 0, \quad i = 1, 2. \quad (3.28)$$

From (3.27), $\varepsilon = -F(z)[1 + H(z)]$ with $H(z)$ holomorphic. Substituting into (3.28) gives

$$F_i(z) + F^2(z)H^{(i)}(z) = 0, \quad i = 1, 2, \quad (3.29)$$

for holomorphic $H^{(i)}(z)$. Let V be the set of z satisfying (3.29). We need that $V = \{0\}$. If not, V is one-dimensional. Let $\tau: T \rightarrow V$ be a resolution of a branch of V at 0, $z = z(t)$. From (3.29), $V \cap \{F(z) = 0\} = \emptyset$. So F is not identically 0 on T . An application of the chain rule shows that the order of the zero of $F(z(t))$ at $t = 0$ is strictly greater than the order of the zero of $F_i(z(t))$ at $t = 0$ for either $i = 1$ or $i = 2$. Then (3.29) cannot hold. Hence $V = \{0\}$ as desired. Hence the assumption that x is a singular point of W_s contradicts the fact that λ_1 is an equisingular deformation off S_1 .

So to complete the proof of Theorem 3.5, we need that $\mathfrak{D}_s$ is reduced. $\delta = \delta_0: \varrho T_0 \rightarrow H^1(D, \Phi)$ is given as onto. Then as in the proof of [19, Theorem 3], $\delta_s: \varrho T_s \rightarrow H^1(D_s, \Phi_s)$ is onto for all small $s \in S$. Then by part (i) of this theorem, $H^1(D, \mathfrak{T}_s) = 0$. Let $\mathcal{I}_s$ be the ideal sheaf of a component C of W_s . If $q_1 > 1$, then $\mathfrak{T}_s/\mathcal{I}_s\mathfrak{T}_s$ is the sheaf of germs of sections of a line bundle over C of Chern class $q_1 C \cdot C$. Then $H^1(C, \mathfrak{T}_s/\mathcal{I}_s\mathfrak{T}_s) \neq 0$. This contradicts $H^1(D_s, \mathfrak{T}_s) = 0$ and completes the proof of Theorem 3.5.

We shall now give an example of a nonzero second obstruction to lifting a cycle. The example was suggested by work of Wahl [27]. Consider a rational singularity whose minimal resolution has the dual graph

Let E_1 be the central vertex and E_2, E_3, E_4, E_5 the outer vertices. All E_i are nonsingular rational curves. Let $E_1 \cdot E_1 = -3$; $E_i \cdot E_i = -2$, $i = 2, 3, 4, 5$. The fundamental cycle is given by $Z = 2E_1 + E_2 + E_3 + E_4 + E_5$. We may choose coordinates (x, y) near $E_1 \cap E_2$, (x_1, y_1) near $E_2 - E_1$ and (x_2, y_2) near $E_1 - E_2$ with coordinate transitions $x_1 = xy^2, y_1 = 1/x, x_2 = 1/x, y_2 = x^3y$. Consider the following infinitesimally nontrivial deformation, with one-dimensional parameter ϵ :

$$\begin{aligned} x_1 &= xy^2 + 2\epsilon y, & y_1 &= 1/x, \\ x_1 &= 1/x, & y_2 &= x^3y + \epsilon x^2. \end{aligned}$$

The deformation is trivial on the other changes of coordinates. The cycle Z may be lifted to first order by the defining equations

x_1 in (x_1, y_1, ϵ) coordinates,

$xy^2 + 2\epsilon y$ in (x, y, ϵ) coordinates,

y_2^2 in (x_2, y_2, ϵ) coordinates away from E_3, E_4 and E_5 .

(3.1)₁ between (x, y, ϵ) and (x_2, y_2, ϵ) coordinates is given by $y_2^2 = x^6y^2 + 2\epsilon x^5y + \epsilon^2x^4 \equiv x^5(xy^2 + 2\epsilon y)$. Then $\text{cls}[x^4] \in H^1(Z, \Phi)$ is the second obstruction. Let $\mathcal{G}$ be the ideal sheaf of E_1 . Let $\mathcal{T}$ be the sheaf of germs of infinitesimal deformations of Z . Then $\mathcal{T}/\mathcal{G}\mathcal{T}$ is isomorphic to the sheaf of germs of sections of a line bundle of Chern class $E_1 \cdot Z = -2$ over E_1 . Then via (2.9), the image of $\text{cls}[x^4]$ in $H^1(E_1, \mathcal{T}/\mathcal{G}\mathcal{T})$ is nonzero.

Observe that Z is a candidate from [18, Proposition 2.3, p. 4] for weak lifting. Also, in the construction of Theorem 3.5, we are really concerned with higher obstructions in $H^1(Z, \Phi)/H^1(Z, \mathcal{U})$. $\text{cls}[x^4]$ is nonzero in $H^1(Z, \Phi)/H^1(Z, \mathcal{U})$.

THEOREM 3.6. *Let M be a strictly pseudoconvex two-dimensional manifold. Let $A = \bigcup A_i$, $1 \leq i \leq n$, be the decomposition of the exceptional set into irreducible components. Suppose that no A_i is an exceptional curve of the first kind. Given any A_k , let $A' = \bigcup A_i$, $i \neq k$. Then there exists a deformation $\omega: \mathcal{U} \rightarrow T$ of M with T smooth and one-dimensional such that for $t \neq 0$, M_t has an exceptional set A_t with exactly $n - 1$ irreducible components, $A_{t,i}$, $1 \leq i \leq n$ and $i \neq k$, with $A_{t,i}$ homologous in $\mathcal{U}$ to A_i , $i \neq k$. Moreover, ω induces a trivial deformation of the (reduced) analytic space A' .*

PROOF. Let $\tau: \mathcal{U} \rightarrow Q$ be the versal deformation of $M = N_0 = \tau^{-1}(0)$ [19, Theorem 8]. Proposition 2.2 says that there is a maximal subspace S of Q to which A' weakly lifts. Also by Proposition 2.2, S is also maximal for all q sufficiently near to 0. Since A' is a reduced space, $\mathcal{T}$ has support only at the isolated singular points of A' . Then $H^1(D, \mathcal{T}) = 0$. Let $\lambda: \mathcal{D} \rightarrow S$ be the complete deformation of A' from Theorem 3.5(iv).

λ may be induced from the versal deformation of A' [8]. So there is a maximal submanifold T_1 of S where λ is the trivial deformation of A' near 0. T_1 is also maximal at all points $t \in T_1$.

By [18, Theorem 2.2, p. 4], off of a proper subvariety V of T_1 , τ induces an equisingular deformation of the full exceptional set. For $t \in T_1 - V$, we shall show that the existence of an additional component B_t of the exceptional set A_t , with B_t not homologous to an A_i , $i \neq k$, leads to a contradiction. B_t is not an exceptional curve of the first kind [1, Lemma 2.1, p. 334]. As in the proof of Theorem 2.8, we can find a special cover $\mathcal{U}$ of $N_t = \tau^{-1}(t)$ and a cocycle c with $[c] \in H^1(N_t, \Theta_t)$ such that the lifting of B_t is obstructed in the direction of c . Moreover, c will vanish near A' . As in the proof of [19, Theorem 1], we may construct a deformation π of N_t in the direction of c which induces the trivial deformation on A' . Since τ is a complete deformation at t [19, Theorem 8] and since T_1 is the maximal subspace of Q near t over which there is a trivial deformation of A'_t , π may be induced from τ restricted to T_1 . But then B_t cannot exist on the exceptional set over $T_1 - V$.

To find the desired T of the theorem, just take a 1-dimensional submanifold of T_1 such that $T \cap V = \{0\}$.

REFERENCES

1. M. Artin, *Algebraic construction of Brieskorn's resolutions*, J. Algebra **29** (1974), 330–348.
2. M. Artin and M. Schlessinger, *Algebraic construction of Brieskorn's resolutions*, preprint.
3. A. Douady, *Le problème des modules locaux pour les espaces C -analytiques compacts*, Ann. Sci. École Norm. Sup. (4) **7** (1974), 569–602 (1975).
4. R. Elkik, *Algébrisation du module formel d'une singularité isolée*, Astérisque **16** (1974), 133–144.
5. W. Fischer and H. Grauert, *Lokal-triviale Familien kompakter komplexer Mannigfaltigkeiten*, Nachr. Akad. Wiss. Göttingen Math.-Phys. Kl. II **1965**, 89–94.
6. H. Grauert, *Über Modifikationen und exzeptionelle analytische Mengen*, Math. Ann. **146** (1962), 331–368.
7. ———, *Über die Deformation isolierter Singularitäten analytischer Mengen*, Invent. Math. **15** (1972), 171–199.
8. ———, *Der Satz von Kuranishi für kompakte komplexe Räume*, Invent. Math. **25** (1974), 107–142.
9. R. Gunning and R. Narasimhan, *Immersion of open Riemann surfaces*, Math. Ann. **174** (1967), 103–108.
10. R. Gunning and H. Rossi, *Analytic functions of several complex variables*, Prentice-Hall, Englewood Cliffs, N. J., 1965.
11. A. Kas and M. Schlessinger, *On the versal deformation of a complex space with an isolated singularity*, Math. Ann. **196** (1972), 23–29.
12. K. Kodaira, *A theorem of completeness of characteristic systems for analytic families of compact submanifolds of complex manifolds*, Ann. of Math. **75** (1962), 146–162.
13. K. Kodaira and D. Spencer, *A theorem of completeness of characteristic systems of complete continuous systems*, Amer. J. Math. **81** (1959), 577–500.
14. H. Laufer, *On rational singularities*, Amer. J. Math. **94** (1972), 597–608.
15. ———, *Deformations of resolutions of two-dimensional singularities*, Rice Univ. Studies **59** (1973), 53–96.

16. ———, *Taut two-dimensional singularities*, Math. Ann. **205** (1973), 131–164.
17. ———, *Ambient deformations for one-dimensional exceptional sets*, 1978, preprint.
18. ———, *Ambient deformations for exceptional sets in two-manifolds*, Invent. Math. **55** (1979), 1–36.
19. ———, *Versal deformations for two-dimensional pseudoconvex manifolds*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **7** (1980), 511–521.
20. ———, *On $\mathbb{CP}^1$ as an exceptional set*, Recent Developments in Several Complex Variables (J. Fornæss, Ed.), Ann. of Math. Studies No. 100, Princeton Univ. Press, Princeton, N. J., 1981, pp. 261–275.
21. D. Lê and C. Ramanujan, *The invariance of Milnor's number implies the invariance of the topological type*, Amer. J. Math. **98** (1976), 67–78.
22. G. Pourcin, *Déformation de singularités isolées*, Quelques Problèmes de Modules (Sem. Géométrie Analytique, Ecole Norm. Sup., Paris, 1971–1972), Astérisque, no. 16, Soc. Math. France, Paris, 1974, pp. 161–173.
23. O. Riemenschneider, *Familien komplexer Räume mit streng pseudokonvexer spezieller Faser*, Comment. Math. Helv. **51** (1976), 547–565.
24. Y.-T. Siu, *Analytic sheaf cohomology groups of dimension n of n -dimensional complex spaces*, Trans. Amer. Math. Soc. **143** (1969), 77–94.
25. B. Teissier, *Deformations à type topologique constant. I*, Astérisque **16** (1974), 215–227.
26. J. Wahl, *Equisingular deformations of normal surface singularities. I*, Ann. of Math. **104** (1976), 325–356.
27. ———, *The discriminant locus of a rational singularity*, informal research announcement, 1976.
28. ———, *Simultaneous resolution and discriminantal loci*, Duke Math. J. **46** (1979), 341–374.
29. H. Whitney, *Tangents to an analytic variety*, Ann. of Math. **81** (1965), 496–549.

DEPARTMENT OF MATHEMATICS, STATE UNIVERSITY OF NEW YORK, STONY BROOK, NEW YORK 11794