

EPIMORPHICALLY CLOSED PERMUTATIVE VARIETIES

BY

N. M. KHAN

ABSTRACT. We show that for semigroups all permutation identities are preserved under epis and that all subvarieties of the permutative variety defined by any permutation identity

$$x_1 x_2 \cdots x_n = x_{i_1} x_{i_2} \cdots x_{i_n},$$

with $n \geq 3$ and such that $i_n \neq n$ or $i_1 \neq 1$, are closed under epis. Finally we find some sufficient conditions that an identity be preserved under epis in conjunction with any nontrivial permutation identity.

1. Introduction and summary. We establish that for semigroups all permutation identities are preserved under epis. A stronger result for commutativity has long been known, namely that the semigroup dominion of a commutative semigroup is also commutative; we show by a counterexample due to P. M. Higgins that this stronger result is false for each (nontrivial) permutation identity other than commutativity. Next we show that all subvarieties of the permutative variety defined by any permutation identity

$$(1) \quad x_1 x_2 \cdots x_n = x_{i_1} x_{i_2} \cdots x_{i_n},$$

with $n \geq 3$ and such that $i_n \neq [i_1 \neq 1]$, are closed under epis, thus generalizing Theorem 4.1 of the author [10] which states that all commutative varieties are closed under epis. Finally we find some sufficient conditions that an identity be preserved under epis in conjunction with any nontrivial permutation identity.

2. Preliminaries. Let U, S be semigroups with U a subsemigroup of S . We say that U *dominates* an element d of S if for every semigroup T and for all homomorphisms $\beta, \gamma: S \rightarrow T$, $u\beta = u\gamma$ for all $u \in U$ implies $d\beta = d\gamma$. The set of all elements of S dominated by U is called the *dominion* of U in S , and we denote it by $\text{Dom}_S(U)$. It can be easily verified that $\text{Dom}_S(U)$ is a subsemigroup of S containing U . Following Howie and Isbell [8], we call a semigroup U *saturated* if $\text{Dom}_S(U) \neq S$ for every properly contained semigroup S .

A morphism $\alpha: A \rightarrow B$ in the category \mathcal{C} of semigroups is called an *epimorphism* (*epi* for short) if for all $C \in \mathcal{C}$ and for all morphisms $\beta, \gamma: B \rightarrow C$, $\alpha\beta = \alpha\gamma$ implies $\beta = \gamma$. It can be easily verified that a morphism $\alpha: S \rightarrow T$ is epi if and only if the inclusion map $i: S\alpha \rightarrow T$ is epi, and the inclusion map $i: U \rightarrow S$ from any

Received by the editors June 30, 1983 and, in revised form, November 15, 1983.

1980 *Mathematics Subject Classification*. Primary 20M07.

Key words and phrases. Semigroup, variety, epimorphism, dominion.

©1985 American Mathematical Society

0002-9947/85 \$1.00 + \$.25 per page

subsemigroup U of S is epi if and only if $\text{Dom}_S(U) = S$. In such a case S will be called an *epimorphic extension* of U .

A most useful characterization of semigroup dominions is provided by Isbell's Zigzag Theorem.

RESULT 1 [9, Theorem 2.3 or 7, Theorem VII.2.13]. Let U be a subsemigroup of any semigroup S , and let d be any element of S . Then $d \in \text{Dom}_S(U)$ if and only if either $d \in U$ or there are elements $a_0, a_1, a_2, \dots, a_{2m} \in U, t_1, t_2, \dots, t_m, y_1, y_2, \dots, y_m \in S$ such that

$$(2) \quad \begin{aligned} d &= a_0 t_1, & a_0 &= y_1 a_1, \\ y_i a_{2i} &= y_{i+1} a_{2i+1}, & a_{2i-1} t_i &= a_{2i} t_{i+1} \quad (i = 1, 2, \dots, m-1), \\ a_{2m-1} t_m &= a_{2m}, & y_m a_{2m} &= d. \end{aligned}$$

These equations are called a *zigzag of length m over U with value d and with spine $a_0, a_1, a_2, \dots, a_{2m}$* .

An identity of the form of equation (1) for some permutation i of the set $\{1, 2, \dots, n\}$ is called a *permutation identity*. The permutation identity (1) is said to be *nontrivial* if the permutation i is different from the identity permutation.

RESULT 2 [11, Result 3]. Let U and S be any semigroups with U a subsemigroup of S . For any $d \in \text{Dom}_S(U) \setminus U$, if (2) is a zigzag of shortest possible length m over U with value d , then $t_j, y_j \in S \setminus U$ for $j = 1, 2, \dots, m$.

RESULT 3 [11, Proposition 3.1]. Let S be any semigroup satisfying the identity (1) with $n \geq 3$.

(i) For each $j \in \{2, 3, \dots, n\}$ such that $x_{j-1}x_j$ is not a subword of $x_{i_1}x_{i_2} \cdots x_{i_n}$, S also satisfies the permutation identity

$$x_1 x_2 \cdots x_{j-1} x y x_j \cdots x_n = x_1 x_2 \cdots x_{j-1} y x x_j \cdots x_n.$$

(ii) If $x_1 \neq x_{i_1}$, then S also satisfies the permutation identity

$$x y x_1 x_2 \cdots x_n = y x x_1 x_2 \cdots x_n.$$

In the following results, let U and S be any semigroups with U a subsemigroup of S and such that $\text{Dom}_S(U) = S$.

RESULT 4 [11, Result 4]. If $d \in S \setminus U$, then for any positive integer k , there exist $a_1, a_2, \dots, a_k \in U$ and $d_k \in S \setminus U$ such that $d = a_1 a_2 \cdots a_k d_k$.

RESULT 5 [11, Corollary 4.4]. Let U satisfy a permutation identity (1) with $i_n \neq n$. Then, for each positive integer k ,

$$s x_1 x_2 \cdots x_k = s x_{j_1} x_{j_2} \cdots x_{j_k}$$

for all $x_1, x_2, \dots, x_k \in S, s \in S \setminus U$, and for any permutation j of the set $\{1, 2, \dots, k\}$.

RESULT 6 [11, Corollary 4.2]. If U satisfies a nontrivial permutation identity, then for each positive integer k ,

$$s x_1 x_2 \cdots x_k t = s x_{j_1} x_{j_2} \cdots x_{j_k} t$$

for all $s, t \in S \setminus U, x_1, x_2, \dots, x_k \in S$, and for any permutation j of the set $\{1, 2, \dots, k\}$.

The notations and conventions of Clifford and Preston [3] and Howie [7] will be used throughout without explicit mention.

The general question of which varieties are closed under epis has been studied in semigroup theory, ring theory and elsewhere [2]. For example, in [4] Gardner has shown that certain identities weaker than commutativity are not preserved under epis of rings although the variety of commutative rings is closed under epis [1]. P. M. Higgins [5] has shown that identities for which both sides contain a repeated variable are not preserved under epis of semigroups. In showing that all varieties of commutative semigroups are closed under epis [10], the author has generalized the classic result of Isbell [9, Corollary 2.5] that commutativity is preserved under epis. However, finding a complete determination of all identities which are preserved under epis of semigroups still remains an open problem.

3. Epimorphisms of semigroups and permutation identities. An identity $u = v$ is said to be *preserved under epis* if for all semigroups U and S with U a subsemigroup of S and such that $\text{Dom}_S(U) = S$, U satisfying $u = v$ implies S satisfies $u = v$.

THEOREM 3.1. *All permutation identities are preserved under epis.*

PROOF. Let equation (1) be any permutation identity with $n \geq 3$. Without loss we can assume that (1) is nontrivial. Take any semigroup U satisfying (1), and any semigroup S containing U properly and such that $\text{Dom}_S(U) = S$. We shall show that S also satisfies (1).

For $k = 1, 2, \dots, n$, consider the word $x_{i_1}x_{i_2} \cdots x_{i_k}$ of length k . We shall prove the theorem by induction on the length of these words, assuming that the remaining elements $x_{i_{k+1}}, \dots, x_{i_n} \in U$.

First for $k = 1$, that is, when $x_{i_1} \in S$, and $x_{i_2}, \dots, x_{i_n} \in U$, we wish to show that equation (1) holds. When $x_{i_1} \in U$, (1) holds so we assume that $x_{i_1} \in S \setminus U$. By Result 1, we may let (2) be a zigzag of shortest possible length m over U with value x_{i_1} .

First we introduce some notation:

$$(3) \quad \begin{aligned} w_1(x_{i_1}, x_{i_2}, \dots, x_{i_n}) &= x_{i_1}x_{i_2} \cdots x_{i_n} = u_1(x_1, x_2, \dots, x_n), \\ w_2(x_{i_1}, x_{i_2}, \dots, x_{i_n}) &= x_1x_2 \cdots x_n = u_2(x_1, x_2, \dots, x_n). \end{aligned}$$

Case (i). $i_1 = 1$. Now

$$\begin{aligned} x_{i_1}x_{i_2} \cdots x_{i_n} &= y_m a_{2m} x_{i_2} \cdots x_{i_n} = y_m w_1(a_{2m}, x_{i_2}, \dots, x_{i_n}) \\ &= y_m w_2(a_{2m}, x_{i_2}, \dots, x_{i_n}) \quad (\text{since } U \text{ satisfies (1)}) \\ &= y_m a_{2m} x_2 \cdots x_n = x_1 x_2 \cdots x_n, \end{aligned}$$

as required.

Case (ii). $1 < i_1 < n$. Now, putting $j = i_1$, we have

$$\begin{aligned} (4) \quad x_{i_1}x_{i_2} \cdots x_{i_n} &= y_m a_{2m} x_{i_2} \cdots x_{i_n} \quad (\text{from equations (2)}) \\ &= y_m w_1(a_{2m}, x_{i_2}, \dots, x_{i_n}) \\ &= y_m w_2(a_{2m}, x_{i_2}, \dots, x_{i_n}) \quad (\text{since } U \text{ satisfies (1)}) \\ &= y_m x_1 x_2 \cdots x_{j-1} a_{2m} x_{j+1} \cdots x_n \\ &= y_m x_1 x_2 \cdots x_{j-1} a_{2m-1} t_m z \quad (\text{from equations (2)}), \\ &\quad \text{where } z = x_{j+1}, \dots, x_n \end{aligned}$$

(Continues)

$$\begin{aligned}
&= y_m x_1 x_2 \cdots x_{j-1} a_{2m-1} b_{j+1}^{(m)} \cdots b_n^{(m)} t'_m z \\
&\quad \text{(by Result 4, for some } b_{j+1}^{(m)}, \dots, b_n^{(m)} \in U, \\
&\quad \text{and } t'_m \in S \setminus U, \text{ since } t_m \in S \setminus U) \\
&= y_m u_2(x_1, x_2, \dots, x_{j-1}, a_{2m-1}, b_{j+1}^{(m)}, \dots, b_n^{(m)}) t'_m z \\
&= y_m u_1(x_1, x_2, \dots, x_{j-1}, a_{2m-1}, b_{j+1}^{(m)}, \dots, b_n^{(m)}) t'_m z \\
&\quad \text{(since } U \text{ satisfies (1)).}
\end{aligned}$$

Now $u_1(z_1, z_2, \dots, z_n)$ begins with $z_{i_1} = z_j$, so the product (4) in S contains $y_m a_{2m-1}$ which equals $y_{m-1} a_{2m-2}$ (from equations (2)). Thus the product (4) above equals

$$\begin{aligned}
(5) \quad & y_{m-1} u_1(x_1, x_2, \dots, x_{j-1}, a_{2m-2}, b_{j+1}^{(m)}, \dots, b_n^{(m)}) t'_m z \\
&= y_{m-1} u_2(x_1, x_2, \dots, x_{j-1}, a_{2m-2}, b_{j+1}^{(m)}, \dots, b_n^{(m)}) t'_m z \quad \text{(since } U \text{ satisfies (1))} \\
&= y_{m-1} x_1 x_2 \cdots x_{j-1} a_{2m-2} b_{j+1}^{(m)} \cdots b_n^{(m)} t'_m z \\
&= y_{m-1} x_1 x_2 \cdots x_{j-1} a_{2m-2} t'_m z \quad \text{(since } t_m = b_{j+1}^{(m)} \cdots b_n^{(m)} t'_m z) \\
&= y_{m-1} x_1 x_2 \cdots x_{j-1} a_{2m-3} t'_{m-1} z \quad \text{(from equations (2))} \\
&\vdots \\
&= y_1 x_1 x_2 \cdots x_{j-1} a_1 t'_1 z \\
&= y_1 x_1 x_2 \cdots x_{j-1} a_1 b_{j+1}^{(1)} \cdots b_n^{(1)} t'_1 z \\
&\quad \text{(by Result 4, for some} \\
&\quad b_{j+1}^{(1)}, \dots, b_n^{(1)} \in U, \text{ and } t'_1 \in S \setminus U, \text{ since } t_1 \in S \setminus U) \\
&= y_1 u_2(x_1, x_2, \dots, x_{j-1}, a_1, b_{j+1}^{(1)}, \dots, b_n^{(1)}) t'_1 z \\
&= y_1 u_1(x_1, x_2, \dots, x_{j-1}, a_1, b_{j+1}^{(1)}, \dots, b_n^{(1)}) t'_1 z \quad \text{(since } U \text{ satisfies (1)).}
\end{aligned}$$

Again as before, product (5) in S contains $y_1 \hat{a}_1$ which equals a_0 (from equations (2)). Thus the product (5) above equals

$$\begin{aligned}
& u_1(x_1, x_2, \dots, x_{j-1}, a_0, b_{j+1}^{(1)}, \dots, b_n^{(1)}) t'_1 z \\
&= u_2(x_1, x_2, \dots, x_{j-1}, a_0, b_{j+1}^{(1)}, \dots, b_n^{(1)}) t'_1 z \quad \text{(since } U \text{ satisfies (1))} \\
&= x_1 x_2 \cdots x_{j-1} a_0 b_{j+1}^{(1)} \cdots b_n^{(1)} t'_1 z \\
&= x_1 x_2 \cdots x_{j-1} a_0 t'_1 z \quad \text{(since } t_1 = b_{j+1}^{(1)} \cdots b_n^{(1)} t'_1) \\
&= x_1 x_2 \cdots x_n \quad \text{(since } a_0 t_1 = x_{i_1} = x_j, \text{ and } z = x_{j+1} \cdots x_n),
\end{aligned}$$

which proves the result for $k = 1$ in Case (ii).

Case (iii). $i_1 = n$. Now

$$\begin{aligned}
 x_{i_1} x_{i_2} \cdots x_{i_n} &= y_m a_{2m} x_{i_2} \cdots x_{i_n} \\
 &= y_m x_1 x_2 \cdots x_{n-1} a_{2m} \quad (\text{since } U \text{ satisfies (1)}) \\
 &= y_m x_1 x_2 \cdots x_{n-1} a_{2m-1} t_m \quad (\text{from equations (2)}) \\
 &= y_m a_{2m-1} x_{i_2} \cdots x_{i_n} t_m \quad (\text{since } U \text{ satisfies (1)}) \\
 &= y_{m-1} a_{2m-2} x_{i_2} \cdots x_{i_n} t_m \quad (\text{from equations (2)}) \\
 &= y_{m-1} x_1 x_2 \cdots x_{n-1} a_{2m-2} t_m \quad (\text{since } U \text{ satisfies (1)}) \\
 &= y_{m-1} x_1 x_2 \cdots x_{n-1} a_{2m-3} t_{m-1} \quad (\text{from equations (2)}) \\
 &\vdots \\
 &= y_1 x_1 x_2 \cdots x_{n-1} a_1 t_1 \\
 &= y_1 a_1 x_{i_2} \cdots x_{i_n} t_1 \quad (\text{since } U \text{ satisfies (1)}) \\
 &= a_0 x_{i_2} \cdots x_{i_n} t_1 \quad (\text{from equations (2)}) \\
 &= x_1 x_2 \cdots x_{n-1} a_0 t_1 \\
 &= x_1 x_2 \cdots x_n \quad (\text{from equations (2), since } i_1 = n),
 \end{aligned}$$

as required.

REMARK 1. A proof for Case (iii) could also be obtained from the proof for Case (ii) above by making the following conventions:

- (a) the word $x_{j+1} \cdots x_n = 1$;
- (b) $b_{j+1}^{(k)} = \cdots = b_n^{(k)} = 1$ and $t'_k = t_k$ for $k = 1, 2, \dots, m$;
- (c) the vector

$$(x_1, x_2, \dots, x_{j-1}, a_{2k-1}, b_{j+1}^{(k)}, \dots, b_n^{(k)}) = (x_1, x_2, \dots, x_{n-1}, a_{2k-1})$$

for $k = 1, 2, \dots, m$;

- (d) the vector

$$(x_1, x_2, \dots, x_{j-1}, a_{2k-2}, b_{j+1}^{(k)}, \dots, b_n^{(k)}) = (x_1, x_2, \dots, x_{n-1}, a_{2k-2})$$

for $k = 1, 2, \dots, m$.

So assume now that (1) is true for all $x_{i_1}, x_{i_2}, \dots, x_{i_{q-1}} \in S$ and all $x_{i_q}, x_{i_{q+1}}, \dots, x_{i_n} \in U$. We prove from this assumption that (1) is true for all $x_{i_1}, x_{i_2}, \dots, x_{i_q} \in S$ and for all $x_{i_{q+1}}, x_{i_{q+2}}, \dots, x_{i_n} \in U$. We need not consider the case where $x_{i_q} \in U$, so we assume that $x_{i_q} \in S \setminus U$. As $x_{i_q} \in S \setminus U$ and $\text{Dom}_S(U) = S$, by Result 1, we may let (2) be a zigzag of shortest possible length m over U with value x_{i_q} .

Put $j = i_q$ and $l = i_{q-1}$.

Case (i). $l = j - 1$. Now

$$\begin{aligned}
 x_{i_1} x_{i_2} \cdots x_{i_n} &= x_{i_1} x_{i_2} \cdots x_{i_{q-1}} (y_m a_{2m}) x_{i_{q+1}} \cdots x_{i_n} \\
 &= x_{i_1} x_{i_2} \cdots (x_{i_{q-1}} y_m) a_{2m} x_{i_{q+1}} \cdots x_{i_n} \\
 &= x_1 x_2 \cdots x_{j-2} (x_{i_{q-1}} y_m) a_{2m} x_{j+1} \cdots x_n
 \end{aligned}$$

(by the inductive hypothesis)

(Continues)

$$\begin{aligned}
&= x_1 x_2 \cdots x_{j-2} x_{i_{q-1}} (y_m a_{2m}) x_{j+1} \cdots x_n \\
&= x_1 x_2 \cdots x_{j-2} x_{j-1} x_j x_{j+1} \cdots x_n \\
&\quad \left(\text{since } x_{i_q} = x_j \text{ and } x_{i_{q-1}} = x_{j-1} \right),
\end{aligned}$$

as required.

Case (ii). $l < j - 1$ and $j < n$. Now

$$\begin{aligned}
(6) \quad & x_{i_1} \cdots x_{i_{q-1}} x_{i_q} \cdots x_{i_n} \\
&= x_{i_1} \cdots x_{i_{q-1}} y_m a_{2m} \cdots x_{i_n} \quad (\text{from equations (2)}) \\
&= w_1(x_{i_1}, x_{i_2}, \dots, x_{i_{q-1}} y_m, a_{2m}, \dots, x_{i_n}) \\
&= w_2(x_{i_1}, x_{i_2}, \dots, x_{i_{q-1}} y_m, a_{2m}, \dots, x_{i_n}) \quad (\text{by the inductive hypothesis}) \\
&= w_2(x_{i_1}, \dots, x_{i_{q-1}} y_m, a_{2m-1} t_m, \dots, x_{i_n}) \quad (\text{from equations (2)}) \\
&= x_1 x_2 \cdots x_{l-1} (x_{i_{q-1}} y_m) x_{l+1} \cdots x_{j-1} (a_{2m-1} t_m) z \quad (\text{where } z = x_{j+1} \cdots x_n) \\
&= x_1 x_2 \cdots x_{l-1} (x_{i_{q-1}} y_m) x_{l+1} \cdots x_{j-1} a_{2m-1} b_{j+1}^{(m)} \cdots b_n^{(m)} t'_m z \\
&\quad (\text{by Result 4 for some } b_{j+1}^{(m)}, \dots, b_n^{(m)} \in U, \\
&\quad \text{and } t'_m \in S \setminus U, \text{ since } t_m \in S \setminus U) \\
&= u_2(x_1, x_2, \dots, x_{l-1}, x_{i_{q-1}} y_m, x_{l+1}, \dots, x_{j-1}, a_{2m-1}, b_{j+1}^{(m)}, \dots, b_n^{(m)}) t'_m z \\
&= u_1(x_1, \dots, x_{l-1}, x_{i_{q-1}} y_m, x_{l+1}, \dots, x_{j-1}, a_{2m-1}, b_{j+1}^{(m)}, \dots, b_n^{(m)}) t'_m z \\
&\quad (\text{by the inductive hypothesis}).
\end{aligned}$$

Since $u_1(z_1, z_2, \dots, z_n)$ contains as a subword $z_{i_{q-1}} z_{i_q}$, the product (6) in S contains $(x_{i_{q-1}} y_m) a_{2m-1}$ which equals $(x_{i_{q-1}} y_{m-1}) a_{2m-2}$ (from equations (2)). Thus the product (6) above equals

$$\begin{aligned}
(7) \quad & u_1(x_1, \dots, x_{l-1}, x_{i_{q-1}} y_{m-1}, x_{l+1}, \dots, x_{j-1}, a_{2m-2}, b_{j+1}^{(m)}, \dots, b_n^{(m)}) t'_m z \\
&= u_2(x_1, \dots, x_{l-1}, x_{i_{q-1}} y_{m-1}, x_{l+1}, \dots, x_{j-1}, a_{2m-2}, b_{j+1}^{(m)}, \dots, b_n^{(m)}) t'_m z \\
&\quad (\text{by the inductive hypothesis}) \\
&= x_1 x_2 \cdots x_{l-1} x_{i_{q-1}} y_{m-1} x_{l+1} \cdots x_{j-1} a_{2m-2} b_{j+1}^{(m)} \cdots b_n^{(m)} t'_m z \\
&= x_1 x_2 \cdots x_{l-1} x_{i_{q-1}} y_{m-1} x_{l+1} \cdots x_{j-1} a_{2m-2} t_m z \quad (\text{since } t_m = b_{j+1}^{(m)} \cdots b_n^{(m)} t'_m) \\
&= x_1 x_2 \cdots x_{l-1} x_{i_{q-1}} y_{m-1} x_{l+1} \cdots x_{j-1} a_{2m-3} t_{m-1} z \quad (\text{from equations (2)}) \\
&\quad \vdots \\
&= x_1 x_2 \cdots x_{l-1} x_{i_{q-1}} y_1 x_{l+1} \cdots x_{j-1} a_1 t_1 z \\
&= x_1 x_2 \cdots x_{l-1} x_{i_{q-1}} y_1 x_{l+1} \cdots x_{j-1} a_1 b_{j+1}^{(1)} \cdots b_n^{(1)} t'_1 z \\
&\quad (\text{by Result 4 for some } b_{j+1}^{(1)}, \dots, b_n^{(1)} \in U, \quad t'_1 \in S \setminus U, \text{ since } t_1 \in S \setminus U)
\end{aligned}$$

(Continues)

$$= u_2(x_1, \dots, x_{l-1}, x_{i_{q-1}}y_1, x_{l+1}, \dots, x_{j-1}, a_1, b_{j+1}^{(1)}, \dots, b_n^{(1)})t_1'z$$

$$= u_1(x_1, \dots, x_{l-1}, x_{i_{q-1}}y_1, x_{l+1}, \dots, x_{j-1}, a_1, b_{j+1}^{(1)}, \dots, b_n^{(1)})t_1'z$$

(by the inductive hypothesis).

Now as $u_1(z_1, z_2, \dots, z_n)$ contains $z_{i_{q-1}}z_{i_q}$ as a subword, the product (7) in S contains $(x_{i_{q-1}}y_1)a_1$ which equals $x_{i_{q-1}}a_0$ (from equations (2)). Thus the product (7) above equals

$$u_1(x_1, x_2, \dots, x_{l-1}, x_{i_{q-1}}, x_{l+1}, \dots, x_{j-1}, a_0, b_{j+1}^{(1)}, \dots, b_n^{(1)})t_1'z$$

$$= u_2(x_1, \dots, x_{l-1}, x_{i_{q-1}}, x_{l+1}, \dots, x_{j-1}, a_0, b_{j+1}^{(1)}, \dots, b_n^{(1)})t_1'z$$

(by the inductive hypothesis)

$$= x_1 \cdots x_{l-1}x_{i_{q-1}}x_{l+1} \cdots x_{j-1}a_0b_{j+1}^{(1)} \cdots b_n^{(1)}t_1'z$$

$$= x_1 \cdots x_{l-1}x_{i_{q-1}}x_{l+1} \cdots x_{j-1}a_0t_1x_{j+1} \cdots x_n \quad (\text{since } z = x_{j+1} \cdots x_n)$$

$$= x_1x_2 \cdots x_n \quad (\text{since } x_{i_{q-1}} = x_l, \text{ and } a_0t_1 = x_{i_q} = x_j),$$

as required.

Case (iii). $l < j - 1, j = n$. Now

$$x_{i_1}x_{i_2} \cdots x_{i_{q-1}}x_{i_q} \cdots x_{i_n}$$

$$= x_{i_1}x_{i_2} \cdots x_{i_{q-1}}y_ma_{2m} \cdots x_{i_n} \quad (\text{from equations (2)})$$

$$= x_{i_1}x_{i_2} \cdots (x_{i_{q-1}}y_m)a_{2m} \cdots x_{i_n}$$

$$= x_1x_2 \cdots x_{l-1}(x_{i_{q-1}}y_m)x_{l+1} \cdots x_{n-1}a_{2m} \quad (\text{by the inductive hypothesis})$$

$$= x_1x_2 \cdots x_{l-1}(x_{i_{q-1}}y_m)x_{l+1} \cdots x_{n-1}a_{2m-1}t_m \quad (\text{from equations (2)})$$

$$= x_{i_1}x_{i_2} \cdots (x_{i_{q-1}}y_m)a_{2m-1} \cdots x_{i_n}t_m \quad (\text{by the inductive hypothesis})$$

$$= x_{i_1}x_{i_2} \cdots x_{i_{q-1}}(y_ma_{2m-1}) \cdots x_{i_n}t_m$$

$$= x_{i_1}x_{i_2} \cdots x_{i_{q-1}}(y_{m-1}a_{2m-2}) \cdots x_{i_n}t_m \quad (\text{from equations (2)})$$

$$= x_{i_1}x_{i_2} \cdots (x_{i_{q-1}}y_{m-1})a_{2m-2} \cdots x_{i_n}t_m$$

$$= x_1x_2 \cdots x_{l-1}(x_{i_{q-1}}y_{m-1})x_{l+1} \cdots x_{n-1}a_{2m-2}t_m \quad (\text{by the inductive hypothesis})$$

$$= x_1x_2 \cdots x_{l-1}(x_{i_{q-1}}y_{m-1})x_{l+1} \cdots x_{n-1}a_{2m-3}t_{m-1} \quad (\text{from equations (2)})$$

\vdots

$$= x_1x_2 \cdots x_{l-1}(x_{i_{q-1}}y_1)x_{l+1} \cdots x_{n-1}a_1t_1$$

$$= x_{i_1}x_{i_2} \cdots (x_{i_{q-1}}y_1)a_1 \cdots x_{i_n}t_1 \quad (\text{by the inductive hypothesis})$$

$$= x_{i_1}x_{i_2} \cdots x_{i_{q-1}}(y_1a_1) \cdots x_{i_n}t_1$$

$$= x_{i_1}x_{i_2} \cdots x_{i_{q-1}}a_0 \cdots x_{i_n}t_1$$

$$= x_1x_2 \cdots x_{l-1}x_{i_{q-1}} \cdots x_{n-1}a_0t_1 \quad (\text{by the inductive hypothesis})$$

$$= x_1x_2 \cdots x_n \quad (\text{from equations (2), since } i_q = n \text{ and } i_{q-1} = l),$$

as required.

REMARK 2. A proof for Case (iii) could also be obtained from the proof for Case (ii) by making the following conventions:

- (a) the word $x_{j+1} \cdots x_n = 1$,
- (b) $b_{j+1}^{(k)} = \cdots = b_n^{(k)} = 1$ and $t'_k = t_k$ for $k = 1, 2, \dots, m$;
- (c) the vector

$$\begin{aligned} & \left(x_1, x_2, \dots, x_{i_{q-1}} y_k, x_{l+1}, \dots, x_{j-1}, a_{2k-1}, b_{j+1}^{(k)}, \dots, b_n^{(k)} \right) \\ &= \left(x_1, x_2, \dots, x_{i_{q-1}} y_k, \dots, x_{n-1}, a_{2k-1} \right) \quad \text{for } k = 1, 2, \dots, m; \end{aligned}$$

and

$$\begin{aligned} & \left(x_1, x_2, \dots, x_{i_{q-1}} y_{k-1}, x_{l+1}, \dots, x_{j-1}, a_{2k-2}, b_{j+1}^{(k)}, \dots, b_n^{(k)} \right) \\ &= \left(x_1, x_2, \dots, x_{i_{q-1}} y_{k-1}, \dots, x_{n-1}, a_{2k-2} \right) \\ & \quad \text{for } k = 1, 2, \dots, m \text{ and where } y_0 = 1. \end{aligned}$$

Case (iv). $j + 1 < l < n$. We have

(8)

$$\begin{aligned} x_{i_1} x_{i_2} \cdots x_{i_n} &= x_{i_1} x_{i_2} \cdots x_{i_{q-1}} x_{i_q} \cdots x_{i_n} \\ &= x_{i_1} x_{i_2} \cdots x_{i_{q-1}} y_m a_{2m} x_{i_{q+1}} \cdots x_{i_n} \quad (\text{from equations (2)}) \\ &= w_1(x_{i_1}, x_{i_2}, \dots, x_{i_{q-1}} y_m, a_{2m}, x_{i_{q+1}}, \dots, x_{i_n}) \\ &= w_2(x_{i_1}, \dots, x_{i_{q-1}} y_m, a_{2m}, x_{i_{q+1}}, \dots, x_{i_n}) \quad (\text{by the inductive hypothesis}) \\ &= x_1 x_2 \cdots x_{j-1} a_{2m} x_{j+1} \cdots x_{l-1} x_{i_{q-1}} y_m x_{l+1} \cdots x_n \\ &= x_1 x_2 \cdots x_{j-1} a_{2m-1} t_m x_{j+1} \cdots x_{l-1} x_{i_{q-1}} y_m x_{l+1} \cdots x_n \\ & \quad (\text{from equations (2)}) \\ &= x_1 x_2 \cdots x_{j-1} a_{2m-1} b_{j+1}^{(m)} \cdots b_{j+(l-j-1)}^{(m)} t'_m x_{j+1} \cdots x_{l-1} x_{i_{q-1}} y_m x_{l+1} \cdots x_n \\ & \quad (\text{by Result 4 for some } b_{j+1}^{(m)}, \dots, b_{j+(l-j-1)}^{(m)} \in U, \\ & \quad \text{and } t_m \in S \setminus U \text{ since } t_m \in S \setminus U) \\ &= u_2(x_1, x_2, \dots, x_{j-1}, a_{2m-1}, b_{j+1}^{(m)}, \dots, b_{j+(l-j-1)}^{(m)}, t'_m x_{j+1} \cdots x_{i_{q-1}} y_m, \\ & \quad x_{l+1}, \dots, x_n) \\ &= u_1(x_1, x_2, \dots, x_{j-1}, a_{2m-1}, b_{j+1}^{(m)}, \dots, b_{j+(l-j-1)}^{(m)}, t'_m x_{j+1} \cdots x_{i_{q-1}} y_m, \\ & \quad x_{l+1}, \dots, x_n) \\ & \quad (\text{by the inductive hypothesis}). \end{aligned}$$

Now since the word $u_1(z_1, z_2, \dots, z_n)$ contains $z_{i_{q-1}} z_{i_q}$ as a subword, the product (8) in S contains $(x_{i_{q-1}} y_m) a_{2m-1}$ which equals $(x_{i_{q-1}} y_{m-1}) a_{2m-2}$ (from equations (2)). Thus the product (8) above equals

$$\begin{aligned} (9) \quad & u_1(x_1, x_2, \dots, x_{j-1}, a_{2m-2}, b_{j+1}^{(m)}, \dots, b_{j+(l-j-1)}^{(m)}, t'_m x_{j+1} \\ & \quad \cdots x_{l-1} x_{i_{q-1}} y_{m-1}, x_{l+1}, \dots, x_n) \end{aligned}$$

(Continues)

$$\begin{aligned}
 &= u_2(x_1, x_2, \dots, x_{j-1}, a_{2m-2}, b_{j+1}^{(m)}, \dots, b_{j+(l-j-1)}^{(m)}, t'_m x_{j+1} \\
 &\quad \dots x_{l-1} x_{i_{q-1}} y_{m-1}, x_{l+1}, \dots, x_n) \\
 &\quad \text{(by the inductive hypothesis)} \\
 &= x_1 x_2 \dots x_{j-1} a_{2m-2} b_{j+1}^{(m)} \dots b_{j+(l-j-1)}^{(m)} t'_m x_{j+1} \dots x_{l-1} x_{i_{q-1}} y_{m-1} x_{l+1} \dots x_n \\
 &= x_1 x_2 \dots x_{j-1} a_{2m-2} t'_m x_{j+1} \dots x_{l-1} x_{i_{q-1}} y_{m-1} x_{l+1} \dots x_n \\
 &\quad \text{(since } t_m = b_{j+1}^{(m)} \dots b_{j+(l-j-1)}^{(m)} t'_m) \\
 &= x_1 x_2 \dots x_{j-1} a_{2m-3} t'_{m-1} x_{j+1} \dots x_{l-1} x_{i_{q-1}} y_{m-1} x_{l+1} \dots x_n \\
 &\quad \text{(from equations (2))} \\
 &\vdots \\
 &= x_1 x_2 \dots x_{j-1} a_1 t_1 x_{j+1} \dots x_{l-1} x_{i_{q-1}} y_1 x_{l+1} \dots x_n \\
 &= x_1 x_2 \dots x_{j-1} a_1 b_{j+1}^{(1)} \dots b_{j+(l-j-1)}^{(1)} t'_1 x_{j+1} \dots x_{l-1} x_{i_{q-1}} y_1 x_{l+1} \dots x_n \\
 &\quad \text{(by Result 4 for some } b_{j+1}^{(1)}, \dots, b_{j+(l-j-1)}^{(1)} \in U, \\
 &\quad \text{and } t'_1 \in S \setminus U, \text{ since } t_1 \in S \setminus U)
 \end{aligned}$$

$$\begin{aligned}
 &= u_2(x_1, \dots, x_{j-1}, a_1, b_{j+1}^{(1)}, \dots, b_{j+(l-j-1)}^{(1)}, t'_1 x_{j+1} \dots x_{i_{q-1}} y_1, x_{l+1}, \dots, x_n) \\
 &= u_1(x_1, \dots, x_{j-1}, a_1, b_{j+1}^{(1)}, \dots, b_{j+(l-j-1)}^{(1)}, t'_1 x_{j+1} \dots x_{i_{q-1}} y_1, x_{l+1}, \dots, x_n).
 \end{aligned}$$

As $u_1(z_1, z_2, \dots, z_n)$ contains $z_{i_{q-1}} z_{i_q}$ as a subword, the product (9) in S contains $(x_{i_{q-1}} y_1) a_1$ which equals $x_{i_{q-1}} a_0$ (from equations (2)). Thus the product (9) above equals

$$\begin{aligned}
 &u_1(x_1, \dots, x_{j-1}, a_0, b_{j+1}^{(1)}, \dots, b_{j+(l-j-1)}^{(1)}, t'_1 x_{j+1} \dots x_{i_{q-1}}, x_{l+1}, \dots, x_n) \\
 &= u_2(x_1, \dots, x_{j-1}, a_0, b_{j+1}^{(1)}, \dots, b_{j+(l-j-1)}^{(1)}, t'_1 x_{j+1} \dots x_{i_{q-1}}, x_{l+1}, \dots, x_n) \\
 &\quad \text{(by the inductive hypothesis)} \\
 &= x_1 x_2 \dots x_{j-1} a_0 b_{j+1}^{(1)} \dots b_{j+(l-j-1)}^{(1)} t'_1 x_{j+1} \dots x_{l-1} x_{i_{q-1}} x_{l+1} \dots x_n \\
 &= x_1 x_2 \dots x_{j-1} a_0 t_1 x_{j+1} \dots x_{l-1} x_{i_{q-1}} x_{l+1} \dots x_n \\
 &\quad \text{(since } t_1 = b_{j+1}^{(1)} \dots b_{j+(l-j-1)}^{(1)} t'_1) \\
 &= x_1 x_2 \dots x_n \quad \text{(since } x_{i_{q-1}} = x_l, \text{ and } x_{i_q} = a_0 t_1 \doteq x_j),
 \end{aligned}$$

as required.

Case (v). $j + 1 = l$. Now

$$x_{i_1} x_{i_2} \dots x_{i_n}$$

$$= x_{i_1} x_{i_2} \dots x_{i_{q-1}} y_m a_{2m} \dots x_{i_n} \quad \text{(from equations (2))}$$

$$= x_{i_1} x_{i_2} \dots (x_{i_{q-1}} y_m) a_{2m} \dots x_{i_n}$$

$$= x_1 x_2 \dots x_{j-1} a_{2m} (x_{i_{q-1}} y_m) x_{l+1} \dots x_n$$

(by the inductive hypothesis; if $l = n$, the product $x_{l+1} \dots x_n = 1$)

(Continues)

$$\begin{aligned}
&= x_1 x_2 \cdots x_{j-1} a_{2m-1} t_m (x_{i_{q-1}} y_m) x_{l+1} \cdots x_n \quad (\text{from equations (2)}) \\
&= x_1 x_2 \cdots x_{j-1} a_{2m-1} (t_m x_{i_{q-1}} y_m) x_{l+1} \cdots x_n \\
&= x_{i_1} x_{i_2} \cdots x_{i_{q-2}} (t_m x_{i_{q-1}} y_m) a_{2m-1} x_{i_{q+1}} \cdots x_{i_n} \quad (\text{by the inductive hypothesis}) \\
&= x_{i_1} x_{i_2} \cdots x_{i_{q-2}} (t_m x_{i_{q-1}} y_{m-1}) a_{2m-2} x_{i_{q+1}} \cdots x_{i_n} \quad (\text{from equations (2)}) \\
&= x_1 x_2 \cdots x_{j-1} a_{2m-2} (t_m x_{i_{q-1}} y_{m-1}) x_{l+1} \cdots x_n \quad (\text{by the inductive hypothesis}) \\
&= x_1 x_2 \cdots x_{j-1} a_{2m-3} (t_{m-1} x_{i_{q-1}} y_{m-1}) x_{l+1} \cdots x_n \quad (\text{from equations (2)}) \\
&\vdots \\
&= x_1 x_2 \cdots x_{j-1} a_1 (t_1 x_{i_{q-1}} y_1) x_{l+1} \cdots x_n \\
&= x_{i_1} x_{i_2} \cdots x_{i_{q-2}} (t_1 x_{i_{q-1}} y_1) a_1 x_{i_{q+1}} \cdots x_{i_n} \quad (\text{by the inductive hypothesis}) \\
&= x_{i_1} x_{i_2} \cdots x_{i_{q-2}} (t_1 x_{i_{q-1}}) a_0 x_{i_{q+1}} \cdots x_{i_n} \quad (\text{from equations (2)}) \\
&= x_1 x_2 \cdots x_{j-1} a_0 (t_1 x_{i_{q-1}}) x_{l+1} \cdots x_n \quad (\text{by the inductive hypothesis}) \\
&= x_1 x_2 \cdots x_n \quad (\text{from equations (2) and } i_{q-1} = l = j + 1),
\end{aligned}$$

as required.

Finally, a proof in the remaining Case (vi), namely when $j + 1 < l$ and $l = n$, can be obtained from the proof for Case (iv) above by making the following conventions:

- (a) the word $x_{l+1} \cdots x_n = 1$;
- (b) the vector

$$\begin{aligned}
&\left(x_1, x_2, \dots, x_{j-1}, a_{2k-1}, b_{j+1}^{(k)}, \dots, b_{j+(l-j-1)}^{(k)}, t'_k x_{j+1} \cdots x_{l-1} x_{i_{q-1}} y_k, x_{l+1}, \dots, x_n \right) \\
&= \left(x_1, x_2, \dots, x_{j-1}, a_{2k-1}, b_{j+1}^{(k)}, \dots, b_{n-1}^{(k)}, t'_k x_{j+1} \cdots x_{n-1} x_{i_{q-1}} y_k \right)
\end{aligned}$$

for $k = 1, 2, \dots, m$;

- (c) the vector

$$\begin{aligned}
&\left(x_1, x_2, \dots, x_{j-1}, a_{2k-2}, b_{j+1}^{(k)}, \dots, b_{j+(l-j-1)}^{(k)}, t'_k x_{j+1} \cdots x_{l-1} x_{i_{q-1}} y_{k-1}, x_{l+1}, \dots, x_n \right) \\
&= \left(x_1, x_2, \dots, x_{j-1}, a_{2k-2}, b_{j+1}^{(k)}, \dots, b_{n-1}^{(k)}, t'_k x_{j+1} \cdots x_{n-1} x_{i_{q-1}} y_{k-1} \right)
\end{aligned}$$

for $k = 1, 2, \dots, m$ and where $y_0 = 1$.

This completes the proof of Theorem 3.1.

The following corollary gives a sufficient condition for $\text{Dom}_S(U)$ to satisfy any permutation identity that U satisfies and, thus, generalizes [9, Corollary 2.5] from commutativity to any permutation identity.

COROLLARY 3.2 (TO THE PROOF OF THEOREM 3.1). *Let U and S be any semigroups with U a subsemigroup of S . Let U satisfy a permutation identity (1). If for all $s \in S \setminus U$, $s = as'$ for some $a \in U$ and $s' \in S$, then $\text{Dom}_S(U)$ also satisfies the permutation identity (1) satisfied by U .*

REMARK 3. Theorem 3.1 generalizes [9, Corollary 2.5], which stated that commutativity is preserved under epis of semigroups.

EXAMPLE (P. M. HIGGINS, VERBAL COMMUNICATION). This shows that the nontrivial permutation identities other than commutativity are not carried over to dominions.

Let F_X be the free semigroup on a countable infinite set $X = \{x_1, x_2, \dots\}$. Let $T = \langle Y \rangle$, the subsemigroup of F_X generated by the set Y , where

$$Y = \bigcup_{n=0}^{\infty} \{x_{3n+1}x_{3n+2}, x_{3n+2}, x_{3n+2}x_{3n+3}\}.$$

Put $S = F_{X/\rho}$ and $\bar{T} = T^{\rho^h}$, where ρ is the congruence generated by the relation ρ_0 which consists of the pairs $(u_1u_2 \cdots u_n, u_{i_1}u_{i_2} \cdots u_{i_n})$ with $u_j \in T$ for $j = 1, 2, \dots, n$, and where i is a fixed nontrivial permutation of the set $\{1, 2, \dots, n\}$ with $n \geq 3$. It is easy to see that for each $n = 0, 1, 2, \dots$, $(x_{3n+1}x_{3n+2}x_{3n+3})\rho \in \text{Dom}_S(\bar{T})$. Now we show that $\text{Dom}_S(\bar{T})$ does not satisfy the permutation identity corresponding to the permutation i .

To see this consider the product $(x_1x_2x_3)(x_4x_5x_6) \cdots (x_{3n+1}x_{3n+2}x_{3n+3})$ in F_X . Since no n members of T occur consecutively in this word, no elementary ρ_0 transition is possible from this base and hence $\text{Dom}_S(\bar{T})$ does not satisfy the permutation identity corresponding to the permutation i .

4. Epimorphically closed permutative varieties. In Theorems 4.1 and 4.4 the bracketed statements are dual to the other statements.

THEOREM 4.1. *Let equation (1) be any permutation identity with $n \geq 3$ and such that $i_n \neq n$ [$i_1 \neq 1$]. Then all identities, in conjunction with (1), are preserved under epis.*

PROOF. Take any identity

$$(10) \quad u(x_1, x_2, \dots, x_p) = v(x_1, x_2, \dots, x_p)$$

and any semigroups U and S such that U is a subsemigroup of S , U satisfies (1) and (10), and $\text{Dom}_S(U) = S$.

By Theorem 3.1, S satisfies (1). Now we show that S satisfies (10). Since S satisfies (1), by the dual of Result 3, S also satisfies the permutation identity

$$(11) \quad x_1x_2, \dots, x_nxy = x_1x_2 \cdots x_nyx.$$

LEMMA 4.2. *Take any word w in variables x_1, x_2, \dots, x_k say, any $a_1, a_2, \dots, a_k \in U$, and any $t_1, t_2, \dots, t_k \in S^1$ such that if $t_i \in S$, then $a_i = y_i b_i$ for some $y_i \in S \setminus U$, $b_i \in S$ ($i = 1, 2, \dots, k$). Then*

$$w(a_1t_1, a_2t_2, \dots, a_kt_k) = w(a_1, a_2, \dots, a_k)w(t_1, t_2, \dots, t_k).$$

PROOF. Let x_q be the first variable appearing in w for which $t_q \in S$ (whence $a_q = y_q b_q$ for some $y_q \in S \setminus U$, $b_q \in S$). Then

$$\begin{aligned} w(a_1t_1, a_2t_2, \dots, a_kt_k) &= w(a_1t_1, a_2t_2, \dots, y_q b_q t_q, \dots, a_kt_k) \\ &= w(a_1, a_2, \dots, y_q b_q, \dots, a_k)w(t_1, t_2, \dots, t_k) \quad (\text{by Result 5}) \\ &= w(a_1, a_2, \dots, a_k)w(t_1, t_2, \dots, t_k), \end{aligned}$$

as required.

We return to the proof of Theorem 4.1. Take any $d_1, d_2, \dots, d_p \in S$. If some $d_i \in U$, there is a zigzag in S^1 over u with value d_i , namely

$$d_i = d_i 1 = 1 d_i 1 = 1 d_i.$$

Now d_1, d_2, \dots, d_p all have zigzags over U in S^1 of some common length [10, Lemma 4.2], say

$$(12) \quad \begin{aligned} d_i &= a_0^{(i)} t_1^{(i)}, & a_0^{(i)} &= y_1^{(i)} a_1^{(i)}, \\ y_k^{(i)} a_{2k}^{(i)} &= y_{k+1}^{(i)} a_{2k+1}^{(i)}, & a_{2k-1}^{(i)} t_k^{(i)} &= a_{2k}^{(i)} t_{k+1}^{(i)} \\ & & (i &= 1, 2, \dots, p, k = 1, 2, \dots, m-1), \\ a_{2m-1}^{(i)} t_m^{(i)} &= a_{2m}^{(i)}, & y_m^{(i)} a_{2m}^{(i)} &= d_i, \end{aligned}$$

where $a_j^{(i)} \in U$ ($i = 1, 2, \dots, p, j = 0, 1, 2, \dots, 2m$) and $t_q^{(i)}, y_q^{(i)} \in S^1$ ($i = 1, 2, \dots, p, q = 1, 2, \dots, m$), and further, for each $d_i \in S \setminus U$ we can assume that $t_q^{(i)}, y_q^{(i)} \in S \setminus U$ (from the proof of [10, Lemma 4.2]).

In the following, we shall make free use of Lemma 4.2 without explicit mention. We put $\tilde{x} = (x_1, x_2, \dots, x_p)$. In this notation, the identity (10) is simply $u(\tilde{x}) = v(\tilde{x})$.

Put

$$(13) \quad \begin{aligned} \tilde{d} &= (d_1, d_2, \dots, d_p), \\ \tilde{a}_k &= (a_k^{(1)}, a_k^{(2)}, \dots, a_k^{(p)}) \quad (k = 0, 1, 2, \dots, 2m), \\ \tilde{t}_q &= (t_q^{(1)}, t_q^{(2)}, \dots, t_q^{(p)}) \quad (q = 1, 2, \dots, m), \\ \tilde{y}_q &= (y_q^{(1)}, y_q^{(2)}, \dots, y_q^{(p)}) \quad (q = 1, 2, \dots, m). \end{aligned}$$

We wish to show that $u(\tilde{d}) = v(\tilde{d})$.

By [10, Lemma 4.3], $\tilde{d} \in S^p$ is in the dominion of U^p in $(S^1)^p$, where T^γ , for any semigroup T and any integer $\gamma \geq 2$, denotes the cartesian product of γ -copies of the semigroup T ; \tilde{d} has the following zigzag of length m :

$$(14) \quad \begin{aligned} \tilde{d} &= \tilde{a}_0 \tilde{t}_1, & \tilde{a}_0 &= \tilde{y}_1 \tilde{a}_1, \\ \tilde{y}_k \tilde{a}_{2k} &= \tilde{y}_{k+1} \tilde{a}_{2k+1}, & \tilde{a}_{2k-1} \tilde{t}_k &= \tilde{a}_{2k} \tilde{t}_{k+1} \quad (k = 1, 2, \dots, m-1), \\ \tilde{a}_{2m-1} \tilde{t}_m &= \tilde{a}_{2m}, & \tilde{y}_m \tilde{a}_{2m} &= \tilde{d}, \end{aligned}$$

where $\tilde{a}_t \in U^p$ ($t \in 0, 1, 2, \dots, 2m$), and $\tilde{y}_q, \tilde{t}_q \in (S^1)^p$ ($q = 1, 2, \dots, m$).

LEMMA 4.3. *Let the word v in (10) begin with x_j , say. If $d_j \in S \setminus U$, then $u(\tilde{d}) = v(\tilde{d})$.*

PROOF.

$$\begin{aligned} u(\tilde{d}) &= u(\tilde{a}_0 \tilde{t}_1) \quad (\text{from equations (14)}) \\ &= u(\tilde{a}_0) u(\tilde{t}_1) \quad (\text{by Lemma 4.2, since each } a_0^{(i)} = y_1^{(i)} a_1^{(i)}) \\ &= v(\tilde{y}_1 \tilde{a}_1) u(\tilde{t}_1) \quad (\text{since } U \text{ satisfies (10)}) \\ &= v(\tilde{y}_1) v(\tilde{a}_1) u(\tilde{t}_1) \quad (\text{by Result 5, since } y_1^{(j)} \in S \setminus U) \\ &= v(\tilde{y}_1) u(\tilde{a}_1) u(\tilde{t}_1) \quad (\text{since } U \text{ satisfies (10)}) \end{aligned}$$

(Continues)

$$\begin{aligned}
 &= v(\tilde{y}_1)u(\tilde{a}_1\tilde{t}_1) \quad (\text{by Result 5, since } y_1^{(j)} \in S \setminus U) \\
 &= v(\tilde{y}_1)u(\tilde{a}_2\tilde{t}_2) \quad (\text{from equation (14)}) \\
 &\vdots \\
 &= v(\tilde{y}_{m-1})u(\tilde{a}_{2m-2}\tilde{t}_m) \\
 &= v(\tilde{y}_{m-1})u(\tilde{a}_{2m-2})u(\tilde{t}_m) \quad (\text{by Result 5, since } y_{m-1}^{(j)} \in S \setminus U) \\
 &= v(\tilde{y}_{m-1})v(\tilde{a}_{2m-2})u(\tilde{t}_m) \quad (\text{since } U \text{ satisfies (10)}) \\
 &= v(\tilde{y}_{m-1}\tilde{a}_{2m-2})u(\tilde{t}_m) \quad (\text{by Result 5, since } y_{m-1}^{(j)} \in S \setminus U) \\
 &= v(\tilde{y}_m\tilde{a}_{2m-1})u(\tilde{t}_m) \quad (\text{from equations (14)}) \\
 &= v(\tilde{y}_m)v(\tilde{a}_{2m-1})u(\tilde{t}_m) \quad (\text{by Result 5, since } y_m^{(j)} \in S \setminus U) \\
 &= v(\tilde{y}_m)u(\tilde{a}_{2m-1}\tilde{t}_m) \quad (\text{by Result 5, since } y_m^{(j)} \in S \setminus U, \\
 &\hspace{15em} \text{and since } U \text{ satisfies (10)}) \\
 &= v(\tilde{y}_m)u(\tilde{a}_{2m}) \quad (\text{from equations (14)}) \\
 &= v(\tilde{y}_m\tilde{a}_{2m}) \quad (\text{by Result 5, since } y_m^{(j)} \in S \setminus U \text{ and } U \text{ satisfies (10)}) \\
 &= v(\tilde{d}) \quad (\text{from equations (14)}).
 \end{aligned}$$

This completes the proof of Lemma 4.3.

We return again to the proof of Theorem 4.1. We regard the variables x_1, x_2, \dots, x_n as being “replaced by” d_1, d_2, \dots, d_n respectively, and it will be convenient for us to use the phrase “replaced by” in our proof. If all the variables in u and v are replaced from U , then $u(\tilde{d}) = v(\tilde{d})$ as required; hence we assume that in v , say, not every variable is replaced from U .

By Lemma 4.3, if the first variable of v is replaced by an element of $S \setminus U$, then we have the required result again. Hence we consider now the case where further the first variable of v is replaced by an element of U . Then

$$(15) \quad v(\tilde{x}) = v_1(\tilde{x})v_2(\tilde{x})$$

for some words v_1 and v_2 in the variables x_1, x_2, \dots, x_p , where v_1 is of the maximum length such that all the variables of v_1 are replaced by elements of U (the word v_1 is nonempty and not all the variables x_1, x_2, \dots, x_p appear in v_1). Let the first variable of $v_2(\tilde{x})$ be x_i , say (that is, x_i is the first variable appearing in $v(\tilde{x})$ which is replaced by an element of $S \setminus U$).

For any i , if $d_i \in S \setminus U$ then $y_j^{(i)} \in S \setminus U$ for $j = 1, 2, \dots, m$. Therefore, by Results 1 and 2, for $d_i \in S \setminus U$, we can write

$$(16) \quad y_j^{(i)} = b_j^{(i)}\bar{y}_j^{(i)} \quad \text{and} \quad b_j^{(i)} = z_j^{(i)}c_j^{(i)} \quad \text{for } j = 1, 2, \dots, m,$$

for some $b_j^{(i)}, c_j^{(i)} \in U, \bar{y}_j^{(i)}, z_j^{(i)} \in S \setminus U$. For each $d_i \in U$, we put

$$(17) \quad b_j^{(i)} = c_j^{(i)} = \bar{y}_j^{(i)} = z_j^{(i)} = 1.$$

In addition to the notations (13), we shall also use the following:

$$(18) \quad \begin{aligned} \tilde{b}_q &= (b_q^{(1)}, b_q^{(2)}, \dots, b_q^{(p)}) & (q = 1, 2, \dots, m), \\ \bar{y}_q &= (\bar{y}_q^{(1)}, \bar{y}_q^{(2)}, \dots, \bar{y}_q^{(p)}) & (q = 1, 2, \dots, m), \\ \tilde{c}_q &= (c_q^{(1)}, c_q^{(2)}, \dots, c_q^{(p)}) & (q = 1, 2, \dots, m), \\ \tilde{z}_q &= (z_q^{(1)}, z_q^{(2)}, \dots, z_q^{(p)}) & (q = 1, 2, \dots, m). \end{aligned}$$

Now from equations (17) and (18) we have

$$(19) \quad \tilde{y}_q = \tilde{b}_q \bar{y}_q = \tilde{z}_q \tilde{c}_q \bar{y}_q.$$

Now

$$\begin{aligned} u(\tilde{d}) &= u(\tilde{a}_0 \tilde{t}_1) \quad (\text{from equations (14)}) \\ &= u(\tilde{a}_0) u(\tilde{t}_1) \quad (\text{by Lemma 4.2 since } a_0^{(i)} = y_1^{(i)} a_1^{(i)} \text{ for } i = 1, 2, \dots, p) \\ &= v(\tilde{a}_0) u(\tilde{t}_1) \quad (\text{since } U \text{ satisfies (10)}) \\ &= v(\tilde{y}_1 \tilde{a}_1) u(\tilde{t}_1) \quad (\text{from equations (14)}) \\ &\vdots \\ &= v(\tilde{y}_i \tilde{a}_{2i-1}) u(\tilde{t}_i) \quad (\text{this equality is essentially an inductive assumption;}) \\ &\quad \text{we now obtain equality with } v(\tilde{y}_{i+1} \tilde{a}_{2i+1}) u(\tilde{t}_{i+1})) \\ &= v_1(\tilde{y}_i \tilde{a}_{2i-1}) v_2(\tilde{y}_1 \tilde{a}_{2i-1}) u(\tilde{t}_1) \quad (\text{from equation (15)}) \\ &= v_1(\tilde{a}_{2i-1}) v_2(\tilde{y}_i \tilde{a}_{2i-1}) u(\tilde{t}_1) \\ &\quad (\text{since all variables of } v_1 \text{ are replaced from } U) \\ &= v_1(\tilde{a}_{2i-1}) v_2(\tilde{y}_i) v_2(\tilde{a}_{2i-1}) u(\tilde{t}_1) \quad (\text{by Result 5, since } v_2(\tilde{y}_i \tilde{a}_{2i-1}) \\ &\quad \text{begins with } y_i^{(l)} a_{2i-1}^{(l)} \text{ and } y_i^{(l)} \in S \setminus U) \\ &= v_1(\tilde{a}_{2i-1}) v_2(\tilde{b}_i y_i) v_2(\tilde{a}_{2i-1}) u(\tilde{t}_1) \quad (\text{from equation (19)}) \\ &= v_1(\tilde{a}_{2i-1}) v_2(\tilde{b}_i) v_2(\bar{y}_i) v_2(\tilde{a}_{2i-1}) u(\tilde{t}_i) \quad (\text{by Result 5, since } b_i^{(l)} = z_i^{(l)} c_i^{(l)} \\ &\quad \text{and } z_i^{(l)} \in S \setminus U \text{ from equation (16)}) \\ &= v_1(\tilde{a}_{2i-1}) v_2(\tilde{b}_i \tilde{a}_{2i-1}) u(\tilde{t}_i) v_2(\bar{y}_i) \quad (\text{by Result 5, since } b_i^{(l)} = z_i^{(l)} c_i^{(l)} \\ &\quad \text{and } z_i^{(l)} \in S \setminus U \text{ from equation (16)}) \\ &= v_1(\tilde{b}_i \tilde{a}_{2i-1}) v_2(\tilde{b}_i \tilde{a}_{2i-1}) u(\tilde{t}_i) v_2(\bar{y}_i) \\ &\quad (\text{since all variables of } v_1 \text{ are replaced from } U) \\ &= v(\tilde{b}_i \tilde{a}_{2i-1}) u(\tilde{t}_i) v_2(\bar{y}_i) \quad (\text{from equation (15)}) \\ &= u(\tilde{b}_i \tilde{a}_{2i-1}) u(\tilde{t}_i) v_2(\bar{y}_i) \quad (\text{since } U \text{ satisfies (10)}) \\ &= u(\tilde{b}_i \tilde{a}_{2i-1} \tilde{t}_i) v_2(\bar{y}_i) \quad (\text{by Lemma 4.2, since if any } t_i^{(j)} \in S \setminus U \text{ for any } j, \\ &\quad \text{then } b_i^{(j)} = z_i^{(j)} c_i^{(j)} \text{ with } z_i^{(j)} \in S \setminus U \text{ from equation (16)}) \\ &= u(\tilde{b}_i \tilde{a}_{2i} \tilde{t}_{i+1}) v_2(\bar{y}_i) \quad (\text{from equations (14)}) \end{aligned}$$

(Continues)

$$\begin{aligned}
 &= u(\tilde{b}_i \tilde{a}_{2i}) u(\tilde{t}_{i+1}) v_2(\tilde{y}_i) \quad (\text{by Lemma 4.2, since if any } t_{i+1}^{(j)} \in S \setminus U \text{ for any } j, \\
 &\quad \text{then } b_i^{(j)} = z_i^{(j)} c_i^{(j)} \text{ with } z_i^{(j)} \in S \setminus U \text{ from equation (16)}) \\
 &= v(\tilde{b}_i \tilde{a}_{2i}) u(\tilde{t}_{i+1}) v_2(\tilde{y}_i) \quad (\text{since } U \text{ satisfies (10)}) \\
 &= v_1(\tilde{b}_i \tilde{a}_{2i}) v_2(\tilde{b}_i \tilde{a}_{2i}) u(\tilde{t}_{i+1}) v_2(\tilde{y}_i) \quad (\text{from equation (15)}) \\
 &= v_1(\tilde{a}_{2i}) v_2(\tilde{b}_i \tilde{a}_{2i}) u(\tilde{t}_{i+1}) v_2(\tilde{y}_i) \\
 &\quad (\text{since all variables of } v_1 \text{ are replaced from } U) \\
 &= v_1(\tilde{a}_{2i}) v_2(\tilde{b}_i) v_2(\tilde{y}_i) v_2(\tilde{a}_{2i}) u(\tilde{t}_{i+1}) \quad (\text{by Result 5, since } b_i^{(l)} = z_i^{(l)} c_i^{(l)} \\
 &\quad \text{and } z_i^{(l)} \in S \setminus U \text{ from equation (16)}) \\
 &= v_1(\tilde{a}_{2i}) v_2(\tilde{b}_i \tilde{y}_i \tilde{a}_{2i}) u(\tilde{t}_{i+1}) \quad (\text{by Result 5, since } b_i^{(l)} = z_i^{(l)} c_i^{(l)} \\
 &\quad \text{and } z_i^{(l)} \in S \setminus U \text{ from equation (16)}) \\
 &= v_1(\tilde{a}_{2i}) v_2(\tilde{y}_i \tilde{a}_{2i}) u(\tilde{t}_{i+1}) \quad (\text{from equation (19)}) \\
 &= v_1(\tilde{y}_i \tilde{a}_{2i}) v_2(\tilde{y}_i \tilde{a}_{2i}) u(\tilde{t}_{i+1}) \quad (\text{since all variables of } v_1 \text{ are replaced from } U) \\
 &= v(\tilde{y}_i \tilde{a}_{2i}) u(\tilde{t}_{i+1}) \\
 &= v(\tilde{y}_{i+1} \tilde{a}_{2i+1}) u(\tilde{t}_{i+1}) \quad (\text{if } i \leq m-1) \\
 &\quad \vdots \\
 &= v(\tilde{y}_m \tilde{a}_{2m}) = v(\tilde{d}),
 \end{aligned}$$

as required. This completes the proof of Theorem 4.1.

A restatement of Theorem 4.1 in terms of permutative varieties gives us a generalization of the author's result [10, Theorem 4.1] which states that all commutative varieties are closed under epis.

THEOREM 4.4. *Let \mathcal{V} be the permutative variety defined by a permutation identity (1) such that $i_n \neq n$ [$i_1 \neq 1$]. Then all subvarieties of the variety \mathcal{V} are closed under epis.*

Call an identity $u = v$ *epimorphically stable* or *stable under epis* if all identities in conjunction with it are preserved under epis, by which we mean that if U is any semigroup satisfying $u = v$ and S is any epimorphic extension of U , then S satisfies all the identities satisfied by U .

In his paper [6], P. M. Higgins has provided an example showing that some permutation identities are not epimorphically stable, namely those permutation identities which are consequences of the normality identity $xyzw = xzyw$. Theorem 4.4 gives a sufficient condition for permutation identities to be epimorphically stable. So as a joint result, in the following theorem, we determine all the permutation identities which are epimorphically stable.

THEOREM 4.5. *A permutation identity (1) is epimorphically stable if and only if $i_n \neq n$ or $i_1 \neq 1$.*

PROPOSITION 4.6. *Let U and S be any semigroups with U a subsemigroup of S and such that $\text{Dom}_S(U) = S$. Take any $d \in S \setminus U$. Let (2) be a zigzag of length m over U with value d with $y_1 \in S \setminus U$ (for example if the zigzag is of shortest possible length). If U satisfies any nontrivial permutation identity, then $d^k = a_0^k t_1^k$ for any positive integer k .*

PROOF. We have

$$\begin{aligned}
 d^k &= (a_0 t_1)^k = a_0 t_1 (a_0 t_1)^{k-2} a_0 t_1 \quad (\text{if } k-2 = 0, (a_0 t_1)^{k-2} = 1) \\
 &= y_1 a_1 t_1 (a_0 t_1)^{k-2} a_0 t_1 \quad (\text{from equations (2)}) \\
 &= y_1 a_1 a_0^{k-1} t_1^k \quad (\text{by Result 5, since } y_1, t_1 \in S \setminus U) \\
 &= a_0^k t_1^k,
 \end{aligned}$$

as required.

In general, for any nontrivial permutation identity I , say, we have not yet been able to determine completely which identities are preserved under epis in conjunction with I . However, we have the following

THEOREM 4.7. *Let equation (1) be any nontrivial permutation identity. Then a nontrivial semigroup identity I (one which is not satisfied by the class of all semigroups) is preserved under epis in conjunction with (1) if I has one of the following forms:*

- (i) *at least one side of I has no repeated variable;*
- (ii) $x^p = y^q, p, q > 0$;
- (iii) $x_1^p x_2^p \cdots x_l^p = x_1^q x_2^q \cdots x_l^q, p, q > 0, l \geq 1$;
- (iv) $x^p y^q = y^r x^s, p, q, r, s > 0$;
- (v) $x^p = 0, p > 0$;
- (vi) $x^p y^q = 0, p, q > 0$.

REMARK 4. We regard $u = 0$ (for some nonempty word u) as a semigroup identity: We define it to mean the conjunction of the two identities $uy = u = yu$ (in each case y is a variable not occurring in the word u).

PROOF. Take any semigroups U and S with U epimorphically embedded in S , and such that U (and hence, S , by Theorem 3.1) satisfies the identity (1). We show that each of the identities (i) to (vi) satisfied by U is also satisfied by S .

(i) That S satisfies (i), if U does, follows from [11, Theorem 3.5].

(ii) Assume U satisfies (ii). Then for all $u, v \in U$ we have $u^p = v^q = v^p = u^q$.

Take any $x, y \in S$. We assume first that $x \in S \setminus U$. By Result 1, we may let (2) be a zigzag of shortest possible length m over U with value x . Then

$$\begin{aligned}
 x^p &= a_0^p t_1^p \quad (\text{by Proposition 4.6 and equations (2)}) \\
 &= (y_1 a_1^2)^p t_1^p \quad (\text{since } y_1 a_1^2 = y_1 a_1 a_1 = a_0 a_1 \in U) \\
 &= y_1^p a_1^p (a_1 t_1)^p \quad (\text{by Result 6, since } y_1, t_1 \in S \setminus U) \\
 &= y_1^p a_1^p (a_2 t_2)^p \quad (\text{from equations (2)}) \\
 &= y_1^p a_1^p a_2^p t_2^p \quad (\text{by Result 6, since } y_1, t_2 \in S \setminus U) \\
 &= y_1^p a_1^p a_3^p t_2^p \quad (\text{since } a_2^p = a_3^p) \\
 &\vdots \\
 &= y_1^p a_1^p a_{2m-1}^p t_m^p \\
 &= (y_1 a_1 a_{2m-1} t_m)^p \quad (\text{by Result 6, since } y_1, t_m \in S \setminus U) \\
 &= (y_1 a_1 a_{2m})^p \quad (\text{from equations (2)}) \\
 &= (a_0 a_{2m})^p = u^p \quad \text{for all } u \in U.
 \end{aligned}$$

Hence $x^p = u^p$ for all $x \in S$, $u \in U$ and likewise $y^q = u^q$ for all $y \in S$ and $u \in U$. Therefore $x^p = u^p = u^q = y^q$, as required.

(iii) Assume U satisfies (iii). For $k = 1, 2, \dots, l$, consider the word $x_1^p x_2^p \cdots x_k^p$ of length kp . We shall prove that S satisfies (iii) by induction on the length of these words, assuming that the remaining elements $x_{k+1}, \dots, x_l \in U$.

First for $k = 0$, the equation (iii) is satisfied vacuously. So assume next that (iii) is true for all $x_1, x_2, \dots, x_{k-1} \in S$ and all $x_k, x_{k+1}, \dots, x_l \in U$. We prove from this assumption that (iii) is true for all $x_1, x_2, \dots, x_k \in S$ and for all $x_{k+1}, x_{k+2}, \dots, x_l \in U$. We need not consider the case where $x_k \in U$, so we assume that $x_k \in S \setminus U$. As $x_k \in S \setminus U$ and $\text{Dom}_S(U) = S$, by Result 1, we may let (2) be a zigzag of shortest possible length m over U with value x_k . Assume first that $1 < k < l$. Then

$$\begin{aligned}
 x_1^p x_2^p \cdots x_l^p &= x_1^p x_2^p \cdots a_0^p t_1^p x_{k+1}^p \cdots x_l^p \quad (\text{by Proposition 4.6 and equations (2)}) \\
 &= x_1^p x_2^p \cdots a_0^p b_{k+1}^{(1)p} \cdots b_l^{(1)p} t_1'^p z \\
 &\quad (\text{by Result 4 and Proposition 4.6}) \\
 &\quad \text{for some } b_{k+1}^{(1)}, \dots, b_l^{(1)} \in U, \text{ and } t_1' \in S \setminus U, \\
 &\quad \text{since } t_1 \in S \setminus U, \text{ and where } z = x_{k+1}^p \cdots x_l^p) \\
 &= x_1^q x_2^q \cdots a_0^q b_{k+1}^{(1)q} \cdots b_l^{(1)q} t_1'^p z \quad (\text{by the inductive hypothesis}) \\
 &= w y_1^q a_1^q b_{k+1}^{(1)q} \cdots b_l^{(1)q} t_1'^p z \quad (\text{by Result 6 and equations (2), since } y_1, t_1' \\
 &\quad \in S \setminus U, \text{ and where } w = x_1^q \cdots x_{k-1}^q) \\
 &= w y_1^q c_1^{(1)q} c_2^{(1)q} \cdots c_{k-1}^{(1)q} a_1^q b_{k+1}^{(1)q} \cdots b_l^{(1)q} t_1'^p z \\
 &\quad (\text{by Result 4 for some } c_1^{(1)}, \dots, c_{k-1}^{(1)} \in U, \\
 &\quad \text{and } y_1' \in S \setminus U, \text{ since } y_1 \in S \setminus U) \\
 &= w y_1^q c_1^{(1)p} \cdots c_{k-1}^{(1)p} a_1^p b_{k+1}^{(1)p} \cdots b_l^{(1)p} t_1'^p z \quad (\text{since } U \text{ satisfies (iii)}) \\
 &= w y_1^q c_1^{(1)p} \cdots c_{k-1}^{(1)p} a_1^p t_1'^p z \quad (\text{since } t_1^p = b_{k+1}^{(1)p} \cdots b_l^{(1)p} t_1'^p) \\
 &= w y_1^q c_1^{(1)p} \cdots c_{k-1}^{(1)p} a_2^p t_2'^p z \quad (\text{by Result 6 and equations (2),} \\
 &\quad \text{since } y_1', t_1, t_2 \in S \setminus U) \\
 &\vdots \\
 &= w y_{m-1}^q c_1^{(m-1)p} \cdots c_{k-1}^{(m-1)p} a_{2m-2}^p t_m'^p z \quad (\text{for some } c_1^{(m-1)}, \dots, c_{k-1}^{(m-1)} \in U \\
 &\quad \text{and } y_{m-1}' \in S \setminus U) \\
 &= w y_{m-1}^q c_1^{(m-1)p} \cdots c_{k-1}^{(m-1)p} a_{2m-2}^p b_{k+1}^{(m)p} \cdots b_l^{(m)p} t_m'^p z \\
 &\quad (\text{by Result 4 and Proposition 4.6 for some } b_{k+1}^{(m)}, \dots, b_l^{(m)} \in U, \\
 &\quad \text{and } t_m' \in S \setminus U, \text{ since } t_m \in S \setminus U) \\
 &= w y_{m-1}^q c_1^{(m-1)q} \cdots c_{k-1}^{(m-1)q} a_{2m-2}^q b_{k+1}^{(m)q} \cdots b_l^{(m)q} t_m'^p z \\
 &\quad (\text{since } U \text{ satisfies (iii)}) \\
 &= w y_{m-1}^q a_{2m-2}^q b_{k+1}^{(m)q} \cdots b_l^{(m)q} t_m'^p z \quad (\text{since } y_{m-1}^q = y_{m-1}'^q c_1^{(m-1)q} \cdots c_{k-1}^{(m-1)q}) \\
 &\quad (\text{Continues})
 \end{aligned}$$

$$\begin{aligned}
&= wy_m^q a_{2m-1}^q b_{k+1}^{(m)q} \cdots b_l^{(m)q} t_m'^p z \quad (\text{by Result 6 and equations (2)}, \\
&\hspace{25em} \text{since } y_{m-1}, y_m, t_m' \in S \setminus U) \\
&= wy_m'^q c_1^{(m)q} \cdots c_{k-1}^{(m)q} a_{2m-1}^q b_{k+1}^{(m)q} \cdots b_l^{(m)q} t_m'^p z \\
&\hspace{15em} (\text{by Result 4 and Proposition 4.6 for some } c_1^{(m)}, \dots, c_{k-1}^{(m)} \in U, \\
&\hspace{25em} \text{and } y_m' \in S \setminus U, \text{ since } y_m \in S \setminus U) \\
&= wy_m'^q c_1^{(m)p} \cdots c_{k-1}^{(m)p} a_{2m-1}^p b_{k+1}^{(m)p} \cdots b_l^{(m)p} t_m'^p z \quad (\text{since } U \text{ satisfies (iii)}) \\
&= wy_m'^q c_1^{(m)p} \cdots c_{k-1}^{(m)p} a_{2m-1}^p t_m^p z \quad (\text{since } b_{k+1}^{(m)p} \cdots b_l^{(m)p} t_m'^p = t_m^p) \\
&= wy_m'^q c_1^{(m)p} \cdots c_{k-1}^{(m)p} a_{2m}^p x_{k+1}^p \cdots x_l^p \\
&\hspace{15em} (\text{by Result 6 and equations (2)}, \\
&\hspace{25em} \text{since } y_m', t_m \in S \setminus U, \text{ and since } z = x_{k+1}^p \cdots x_l^p) \\
&= wy_m'^q c_1^{(m)q} \cdots c_{k-1}^{(m)q} a_{2m}^q x_{k+1}^q \cdots x_l^q \quad (\text{since } U \text{ satisfies (iii)}) \\
&= wy_m^q a_{2m}^q x_{k+1}^q \cdots x_l^q \quad (\text{since } y_m'^q c_1^{(m)q} \cdots c_{k-1}^{(m)q} = y_m^q) \\
&= x_1^q \cdots x_{k-1}^q x_k^q x_{k+1}^q \cdots x_l^q \quad (\text{by Proposition 4.6 and equations (2)}),
\end{aligned}$$

as required.

Finally, a proof in the remaining cases, namely when $k = 1$ or $k = l$, can be obtained from the proof above by making the following conventions:

First when $k = 1$,

- (i) the word $w = 1$,
- (ii) the word $c_1^{(i)p} \cdots c_{k-1}^{(i)p} = c_1^{(i)q} \cdots c_{k-1}^{(i)q} = 1$ and $y_i' = y_i$ for $i = 1, 2, \dots, m$.

Dually when $k = l$,

- (i) the word $z = 1$,
- (ii) the word $b_{k+1}^{(i)p} \cdots b_l^{(i)p} = b_{k+1}^{(i)q} \cdots b_l^{(i)q} = 1$ and $t_m' = t_m$ for $i = 1, 2, \dots, m$.

(iv) Assume U satisfies (iv) and take any $x, y \in S$. First we consider the case where $x \in S \setminus U$ and $y \in U$ (the case where $x \in U$ and $y \in S \setminus U$ is symmetric to this case).

Since $x \in S \setminus U$, we may let (2), by Result 1, be a zigzag for x of shortest possible length m over U . Now

$$\begin{aligned}
x^p y^q &= y_m^p a_{2m}^p y^q \quad (\text{by Proposition 4.6 and equations (2)}) \\
&= y_m^p y^r a_{2m}^s \quad (\text{since } U \text{ satisfies (iv)}) \\
&= y_m^p y^r (a_{2m-1} t_m)^s \quad (\text{from equations (2)}) \\
&= y_m^p y^r a_{2m-1}^s t_m^s \quad (\text{by Result 6, since } y_m, t_m \in S \setminus U) \\
&= y_m^p a_{2m-1}^p y^q t_m^s \quad (\text{since } U \text{ satisfies (iv)}) \\
&= (y_m a_{2m-1})^p y^q t_m^s \quad (\text{by Result 6, since } y_m, t_m \in S \setminus U) \\
&= (y_{m-1} a_{2m-2})^p y^q t_m^s \quad (\text{from equations (2)}) \\
&= y_{m-1}^p a_{2m-2}^p y^q t_m^s \quad (\text{by Result 6, since } y_{m-1}, t_m \in S \setminus U) \\
&= y_{m-1}^p y^r a_{2m-2}^s t_m^s \quad (\text{since } U \text{ satisfies (iv)})
\end{aligned}$$

(Continues)

$$\begin{aligned}
 &= y_{m-1}^p y^r (a_{2m-2} t_m)^s \quad (\text{by Result 6, since } y_{m-1}, t_m \in S \setminus U) \\
 &= y_{m-1}^p y^r (a_{2m-3} t_{m-1})^s \quad (\text{from equations (2)}) \\
 &\vdots \\
 &= y_1^p y^r (a_1 t_1)^s \\
 &= y_1^p y^r a_1^s t_1^s \quad (\text{by Result 6, since } y_1, t_1 \in S \setminus U) \\
 &= y_1^p a_1^p y^q t_1^s \quad (\text{since } U \text{ satisfies (iv)}) \\
 &= (y_1 a_1)^p y^q t_1^s \quad (\text{by Result 6, since } y_1, t_1 \in S \setminus U) \\
 &= a_0^p y^q t_1^s \quad (\text{from equations (2)}) \\
 &= y^r a_0^s t_1^s \quad (\text{since } U \text{ satisfies (iv)}) \\
 &= y^r x^s \quad (\text{by Proposition 4.6 and equations (2)}),
 \end{aligned}$$

as required.

So we assume next that $x, y \in S \setminus U$. By Result 1, we may let (2) be a zigzag for x of shortest possible length m over U . Then

$$\begin{aligned}
 x^p y^q &= y_m^p a_{2m}^p y^q \quad (\text{by Proposition 4.6 and equations (2)}) \\
 &= y_m^p y^r a_{2m}^s \quad (\text{by the first part of the proof}) \\
 &= y_m^p y^r (a_{2m-1} t_m)^s \quad (\text{from equations (2)}) \\
 &= y_m^p y^r a_{2m-1}^s t_m^s \quad (\text{by Result 6, since } y_m, t_m \in S \setminus U) \\
 &= y_m^p a_{2m-1}^p y^q t_m^s \quad (\text{by the first part of the proof}) \\
 &= (y_m a_{2m-1})^p y^q t_m^s \quad (\text{by Result 6, since } y_m, t_m \in S \setminus U) \\
 &= (y_{m-1} a_{2m-2})^p y^q t_m^s \quad (\text{from equations (2)}) \\
 &= y_{m-1}^p a_{2m-2}^p y^q t_m^s \quad (\text{by Result 6, since } y_{m-1}, t_m \in S \setminus U) \\
 &= y_{m-1}^p y^r a_{2m-2}^s t_m^s \quad (\text{by the first part of the proof}) \\
 &= y_{m-1}^p y^r (a_{2m-2} t_m)^s \quad (\text{by Result 6, since } y_{m-1}, t_m \in S \setminus U) \\
 &= y_{m-1}^p y^r (a_{2m-3} t_{m-1})^s \quad (\text{from equations (2)}) \\
 &\vdots \\
 &= y_1^p y^r (a_1 t_1)^s \\
 &= y_1^p y^r a_1^s t_1^s \quad (\text{by Result 6, since } y_1, t_1 \in S \setminus U) \\
 &= y_1^p a_1^p y^q t_1^s \quad (\text{by the first part of the proof}) \\
 &= (y_1 a_1)^p y^q t_1^s \quad (\text{by Result 6, since } y_1, t_1 \in S \setminus U) \\
 &= a_0^p y^q t_1^s \quad (\text{from equations (2)}) \\
 &= y^r a_0^s t_1^s \quad (\text{by the first part of the proof}) \\
 &= y^r x^s \quad (\text{by Proposition 4.6 and equations (2)}),
 \end{aligned}$$

as required. This completes the proof of part (iv).

(v) Assume U satisfies $x^p = 0$ and take any $x, y \in S$; we show that $x^p y = y x^p = x^p$.

Case (a). $x \in S \setminus U, y \in U$. Let (2), by Result 1, be a zigzag for x over U of shortest possible length m . Then

$$\begin{aligned} x^p y &= y_m^p a_{2m}^p y \quad (\text{by Proposition 4.6 and equations (2)}) \\ &= y_m^p a_{2m}^p \quad (\text{since } U \text{ satisfies (v)}) \\ &= x^p. \end{aligned}$$

Similarly $y x^p = x^p$, as required.

Case (b). $x \in U, y \in S \setminus U$. Since $y \in S \setminus U$, we may let (2), by Result 1, be a zigzag of length m over U with value y . Then

$$\begin{aligned} x^p y &= x^p a_0 t_1 = x^p a_1 t_1 \quad (\text{since } U \text{ satisfies (v)}) \\ &= x^p a_2 t_2 \quad (\text{from equations (2)}) \\ &= x^p a_3 t_2 \quad (\text{since } U \text{ satisfies (v)}) \\ &\vdots \\ &= x^p a_{2m-1} t_m \\ &= x^p a_{2m} = x^p \quad (\text{since } U \text{ satisfies (v)}). \end{aligned}$$

Similarly $y x^p = x^p$, as required.

Case (c). $x, y \in S \setminus U$. By Result 1, let (2) be a zigzag for x of shortest possible length m over U . Then

$$\begin{aligned} x^p y &= y_m^p a_{2m}^p y \quad (\text{by Proposition 4.6 and equations (2)}) \\ &= y_m^p a_{2m}^p \quad (\text{from case (b) above}) \\ &= x^p. \end{aligned}$$

Similarly $y x^p = x^p$, are required.

(vi) Assume U satisfies (vi) and take any $x, y, z \in S$; we prove that $x^p y^q z = z x^p y^q = x^p y^q$.

Case (a). $x, y \in U, z \in S \setminus U$. Let (2), by Result 1, be a zigzag of shortest possible length m over U with value z . Then

$$\begin{aligned} x^p y^q z &= x^p y^q a_0 t_1 \quad (\text{from equations (2)}) \\ &= x^p y^q a_1 t_1 \quad (\text{since } U \text{ satisfies (vi)}) \\ &= x^p y^q a_2 t_2 \quad (\text{from equations (2)}) \\ &\vdots \\ &= x^p y^q a_{2m-2} t_m \\ &= x^p y^q a_{2m-1} t_m \quad (\text{since } U \text{ satisfies (vi)}) \\ &= x^p y^q a_{2m} \quad (\text{from equations (2)}) \\ &= x^p y^q \quad (\text{since } U \text{ satisfies (vi)}). \end{aligned}$$

By a similar argument we can show easily that $z x^p y^q = x^p y^q$. Therefore $x^p y^q z = z x^p y^q = x^p y^q$, as required.

Case (b). $y, z \in U, x \in S \setminus U$. As $x \in S \setminus U$, by Result 1, we may let (2) be a zigzag of shortest possible length m over U with value x . Then

$$\begin{aligned} x^p y^q z &= y_m^p a_{2m}^p y^q z \quad (\text{by Proposition 4.6 and equations (2)}) \\ &= y_m^p a_{2m}^p y^q \quad (\text{since } U \text{ satisfies (vi)}) \\ &= x^p y^q \quad (\text{by Proposition 4.6 and equations (2)}). \end{aligned}$$

Also

$$\begin{aligned} zx^p y^q &= zy_m^p a_{2m}^p y^q \quad (\text{by Proposition 4.6 and equations (2)}) \\ &= y_m^p a_{2m}^p y^q \quad (\text{from Case (a)}) \\ &= x^p y^q \quad (\text{by Proposition 4.6 and equations (2)}). \end{aligned}$$

Therefore $x^p y^q z = zx^p y^q = x^p y^q$, as required.

Case (c). $x, z \in U, y \in S \setminus U$. This case is dual to Case (b).

Case (d). $z \in U, x, y \in S \setminus U$. Let (2), by Result 1, be a zigzag of shortest possible length m over U with value x . Now

$$\begin{aligned} x^p y^q z &= y_m^p a_{2m}^p y^q z \quad (\text{by Proposition 4.6 and equations (2)}) \\ &= y_m^p a_{2m}^p y^q \quad (\text{from Case (c)}) \\ &= x^p y^q \quad (\text{by Proposition 4.6 and equations (2)}). \end{aligned}$$

Since $y \in S \setminus U$, by Result 1, we may let $y = b_0 z_1 = s_1 b_1 z_1$ be the first two lines of a zigzag for y with $b_0, b_1 \in U$, and $s_1, z_1 \in S \setminus U$. Now

$$\begin{aligned} x^p y^q &= zy_m^p a_{2m}^p b_0^q z_1^q \quad (\text{by Proposition 4.6 and equations (2)}) \\ &= y_m^p a_{2m}^p b_0^q z_1^q \quad (\text{from Case (a)}) \\ &= x^p y^q \quad (\text{by Proposition 4.6 and equations (2)}). \end{aligned}$$

Therefore $x^p y^q z = zx^p y^q = x^p y^q$, as required.

Case (e). $y \in U, x, z \in S \setminus U$ or $x \in U, y, z \in S \setminus U$ or $x, y, z \in S \setminus U$. As $z \in S \setminus U$, by Result 1, we may let (2) be a zigzag of length m over U with value z . Now

$$\begin{aligned} x^p y^q z &= x^p y^q a_0 t_1 \quad (\text{from equations (2)}) \\ &= x^p y^q a_1 t_1 \quad (\text{from Cases (b), (c) and (d)}) \\ &= x^p y^q a_2 t_2 \quad (\text{from equations (2)}) \\ &\vdots \\ &= x^p y^q a_{2m-2} t_m \\ &= x^p y^q a_{2m-1} t_m \quad (\text{from Cases (b), (c) and (d)}) \\ &= x^p y^q a_{2m} \quad (\text{from equations (2)}) \\ &= x^p y^q \quad (\text{from Case (d)}). \end{aligned}$$

The dual argument shows that $zx^p y^q = x^p y^q$. Therefore

$$x^p y^q z = zx^p y^q = x^p y^q,$$

as required, thus completing the proof of Theorem 4.7.

ACKNOWLEDGEMENT. I thank Tom Hall, my supervisor, for his help and encouragement throughout the writing of this paper.

REFERENCES

1. A. Bulazewska and J. Krempka, *On epimorphisms in the category of all associative rings*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. **23** (1975).
2. W. Burgess, *The meaning of mono and epi in some familiar categories*, Canad. Math. Bull. **8** (1965).
3. A. H. Clifford and G. B. Preston, *The algebraic theory of semigroups*, Vol. 1, Math. Surveys, No. 7, Amer. Math. Soc., Providence, R. I., 1961; Vol. II, 1967.
4. B. J. Gardner, *A note on ring epimorphisms and polynomial identities*, Comment. Math. Univ. Carolin. **20** (1979).
5. P. M. Higgins, *Saturated and epimorphically closed varieties of semigroups*, J. Austral. Math. Soc. Ser. A **36** (1984), 153–175.
6. ———, *Epimorphisms, permutation identities and finite semigroups*, Semigroup Forum (to appear).
7. J. M. Howie, *An introduction to semigroup theory*, London Math. Soc. Monographs, vol. 7, Academic Press, 1976.
8. J. M. Howie and J. R. Isbell, *Epimorphisms and dominions*. II, J. Algebra **6** (1967), 7–21.
9. J. R. Isbell, *Epimorphisms and dominions*, Proc. Conf. on Categorical Algebra (La Jolla, 1965), Lange & Springer, Berlin, 1966, pp. 232–246.
10. N. M. Khan, *Epimorphisms, dominions and varieties of semigroups*, Semigroup Forum **25** (1982), 331–337.
11. ———, *On saturated varieties and consequences of permutation identities*. J. Austral. Math. Soc. (to appear).

DEPARTMENT OF MATHEMATICS, MONASH UNIVERSITY, CLAYTON, VICTORIA, AUSTRALIA 3168

Current address: Department of Mathematics, Aligarh Muslim University, Aligarh, India 202001